Evolutionary insights into the genomic organization of major ribosomal DNA in ant chromosomes

Gisele Amaro Teixeira, Hilton Jeferson Alves Cardoso de Aguiar, Frédéric Petitclerc, Jérôme Orivel, Denilce Meneses Lopes, Luisa Antonia Campos Barros

► To cite this version:

Gisele Amaro Teixeira, Hilton Jeferson Alves Cardoso de Aguiar, Frédéric Petitclerc, Jérôme Orivel, Denilce Meneses Lopes, et al.. Evolutionary insights into the genomic organization of major ribosomal DNA in ant chromosomes. *Insect Molecular Biology*, 2021. hal-03369943

HAL Id: hal-03369943

<https://hal.science/hal-03369943>

Submitted on 7 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

**Evolutionary insights into the genomic organization of major ribosomal
DNA in ant chromosomes**

Gisele Amaro Teixeira^{1, 2}, Hilton Jeferson Alves Cardoso de Aguiar^{3*}, Frédéric
Petitclerc⁴, Jérôme Orivel⁴, Denilce Meneses Lopes², Luísa Antônia Campos Barros³

1 Programa de Pós-graduação em Biologia Celular e Estrutural, Universidade Federal
de Viçosa, Viçosa, MG 36570-000, Brazil.

2 Laboratório de Citogenética de Insetos, Departamento de Biologia Geral,
Universidade Federal de Viçosa, Viçosa, MG 36570-900, Brazil.

3-Universidade Federal do Amapá, Campus Binacional – Oiapoque, n°3051, Bairro
Universidade, 68980-000, Oiapoque, Amapá, Brazil.

4 CNRS, UMR EcoFoG, AgroParisTech, CIRAD, INRA, Université de Guyane,
Université des Antilles, Campus Agronomique, BP 316, 97379 Kourou Cedex, France.

Short running title: Evolution of rDNA genes in ant chromosomes

* Corresponding author

E-mail: hilton.aguiar@hotmail.com

Abstract

The major rDNA genes are composed of tandem repeats and are part of the nucleolus organizing regions (NORs). They are highly conserved and therefore useful in understanding the evolutionary patterns of chromosomal locations. The evolutionary dynamics of the karyotype may affect the organization of rDNA genes within chromosomes. In this study, we physically mapped 18S rDNA genes in 13 Neotropical ant species from four subfamilies using fluorescence *in situ* hybridization. Furthermore, a survey of published rDNA cytogenetic data for 50 additional species was performed, which allowed us to detect the evolutionary patterns of these genes in ant chromosomes. Species from the Neotropical, Palearctic, and Australian regions, comprising a total of 63 species from 19 genera within six subfamilies, were analyzed. Most of the species (48 out of 63) had rDNA genes restricted to a single chromosome pair in their intrachromosomal regions. The position of rDNA genes within the chromosomes appears to hinder their dispersal throughout the genome, as translocations and ectopic recombination are uncommon in intrachromosomal regions because they can generate meiotic abnormalities. Therefore, rDNA genes restricted to a single chromosome pair seem to be a plesiomorphic feature in ants, while multiple rDNA sites, observed in distinct subfamilies, may have independent origins in different genera.

Keywords: Evolution - ribosomal DNA - gene dispersion - Formicidae - chromosome rearrangements - physical mapping

1 **Introduction**

2 Eukaryotic genomes have repetitive tandem sequences such as in the major
3 ribosomal RNA genes (45S = 18S + 5.8S + 28S), herein denominated rDNA, which
4 contain highly conserved genic sequences and are therefore useful as molecular genetic
5 markers, allowing comparisons across distant taxa. However, intergenic spacers vary in
6 both sequence and length (Long and Dawid, 1980; Sumner, 2003; Symonová, 2019). The
7 45S ribosomal genes are part of the nucleolus organizing regions (NORs) and are located
8 in portions of the DNA that, after their condensation, usually appear as secondary
9 constrictions on metaphase chromosomes (Sumner, 2003).

10 The silver nitrate impregnation technique (Ag-NOR banding) shows only
11 transcriptionally active NORs by staining nuclear acid proteins involved in transcription
12 (Howell and Black, 1980; reviewed by Sumner, 2003). Species with multiple rDNA
13 clusters do not usually exhibit silver-staining in all clusters (reviewed by Sumner, 2003;
14 Vicari *et al.*, 2008). However, it must be assumed that if a species has only a single NOR
15 (or single 45S rDNA gene site), then it will be transcriptionally active, as evidenced for
16 different organisms (Dobigny *et al.*, 2002; Barros *et al.*, 2015; Falcione *et al.*, 2018;
17 Cholak *et al.*, 2020; Malimpensa *et al.*, 2020).

18 Intercellular and interindividual variations are frequently detected by the Ag-NOR
19 banding method (Zurita *et al.*, 1997; Cross *et al.*, 2003; Walker *et al.*, 2014; Schmid *et al.*,
20 2017). However, the use of this technique in ant species has not produced reliable
21 results owing to low repeatability, difficulty in obtaining good-quality markings, and the
22 appearance of unspecified marks on heterochromatic regions of many chromosomes,
23 which makes using this procedure unsatisfactory and inconclusive (Imai *et al.*, 1992;
24 Hirai *et al.*, 1994; Lorite *et al.*, 1997; Barros *et al.*, 2009, 2015; Aguiar *et al.*, 2017).

1 Since the 1980s, molecular cytogenetic tools have been used to study karyotypes.
2 For example, fluorescence *in situ* hybridization (FISH) has proven to be an effective and
3 precise tool for physically mapping specific DNA sequences within chromosomes
4 (reviewed by Levsky and Singer, 2003; Liehr, 2017). The rDNA genes can be located in
5 single or multiple chromosome pairs (Sochorová *et al.*, 2018). In several organisms,
6 studies of these genes have pointed to chromosomal differences within species complexes
7 (Mantovani *et al.*, 2005; Barbosa *et al.*, 2017; Dutra *et al.*, 2020) and between species
8 with similar karyotypes (Panzera *et al.*, 2012; Golub *et al.*, 2015; Gokhman *et al.*, 2016).
9 As a consequence, inferences can be made based on chromosomal rearrangements that
10 shape the chromosomal evolution of a species (Roy *et al.*, 2005; Nguyen *et al.*, 2010;
11 Britton-Davidian *et al.*, 2011; Cabral-de-Mello *et al.*, 2011; Dutrillaux and Dutrillaux,
12 2012; Roa and Guerra, 2012; Menezes *et al.*, 2019; Degrandi *et al.*, 2020).

13 In ants, the physical mapping of rDNA genes using the FISH technique was first
14 described in Australian ants by Hirai *et al.* (1994, 1996). Since then, the number of studies
15 mapping these genes has increased (Mariano *et al.*, 2008; Santos *et al.*, 2016; Micolino
16 *et al.*, 2019a; Teixeira *et al.*, 2020) and other repetitive sequences, such as telomeres
17 (Meyne *et al.*, 1995; Pereira *et al.*, 2018; Micolino *et al.*, 2020; Castro *et al.*, 2020),
18 satellite DNA (Lorite *et al.*, 2004; Huang *et al.*, 2016), 5S ribosomal genes (Aguar *et al.*,
19 2017), and microsatellites (Barros *et al.*, 2018; Micolino *et al.*, 2019b), have been mapped
20 in the chromosomes using the FISH technique. To date, molecular cytogenetic studies on
21 rDNA genes in ants have improved understanding of chromosomal evolution and
22 phylogeny, and provided taxonomic resolutions for different ant groups (Hirai *et al.*,
23 1994, 1996; Santos *et al.*, 2010, 2016).

24 In this study, we physically mapped 18S rDNA clusters using the FISH technique
25 and verified if they were GC-rich in 13 Neotropical ant species from four subfamilies

(Ectatomminae, Formicinae, Myrmicinae, and Ponerinae). In addition, we reviewed previous molecular cytogenetic data related to rDNA gene clusters (45S, 18S, or 28S) in ants. Using these data, we investigated whether the number and location of the ribosomal gene clusters followed a specific pattern or were randomly distributed in order to understand the genomic organization and evolutionary dynamics of these genes in ants.

Results

1 - Chromosome mapping of 18S rDNA clusters in 13 Neotropical ant species

All of the studied ant species presented only a single 18S rDNA site that was co-localized with GC-rich regions (CMA₃⁺) (Table 2), while AT-rich regions (DAPI⁺) were not detected in any species. Most of the species presented GC-rich rDNA genes in the intrachromosomal regions (pericentromeric or interstitial), including *Pseudoponera gilberti* (Kempf, 1960) (Fig. 1A, S1A), *Anochetus targionii* Emery, 1894 (Fig. 1B, S1B), *Odontomachus haematodus* (Linnaeus, 1758) (Fig. 1C, D, S1C), *Odontomachus bauri* Emery, 1892 (Fig. 1E, S1D), *Pheidole germaini* Emery, 1896 (Fig. 2A, S2A), *Crematogaster longispina* Emery, 1890 (Fig. 2B, S2B), *Solenopsis geminata* (Fabricius, 1804) (Fig. 2C, D, S2C), *Myrmicocrypta* sp. (Fig. 2E, S2D), and *Acromyrmex echinator* (Forel, 1899) (Fig. 2F). Additional GC-rich bands only occurred in *P. gilberti* in the pericentromeric region of the 3rd and 4th metacentric chromosome pairs (Fig. S1B). Differences in rDNA cluster sizes between homologous chromosomes were observed in heterozygous *O. bauri* individuals. One of the homologous chromosomes showed clusters approximately twice the size of those in the other chromosome (Fig. 1F, S1E). A male

individual bearing a chromosome with minor GC-rich bands was analyzed (Fig. S1F).
Homozygous individuals with duplicated clusters were not observed.

Figure 1. Fluorescence *in situ* hybridization with 18S rDNA probe (red blocks) of Ponerinae ants: (A) *Pseudoponera gilberti* ($2n=12$), (B) *Anochetus targionii* ($2n=30$), (C, D) *Odontomachus haematodus* ($2n=44$, Brazil and French Guiana, respectively), (E, F) *Odontomachus bauri* ($2n=44$). In the last species, the 18S rDNA site was located on the long arm of the polymorphic chromosomal pair. The boxes show a polymorphic subtelocentric pair in (E) in the homozygous state, and (F) in the heterozygous state. Bars = 5 μm .

Figure 2. Fluorescence *in situ* hybridization with 18S rDNA probe (red blocks) of Myrmicinae ants: (A) *Pheidole germaini* (2n=22), (B) *Crematogaster longispina* (2n=24), (C) *Solenopsis geminata* (2n=32) and (D) *Solenopsis geminata* (n=16), (E) *Myrmicocrypta* sp. (2n=30), (F) *Acromyrmex echinator* (2n=38), (G) *Strumigenys diabolus* (2n=40). Bars = 5 μm.

The remaining species showed GC-rich 18S rDNA clusters across the entire chromosome arm, occupying either the long arm, as in *Gnamptogenys tortuolosa* (Smith, 1858) (Fig. 3A, S3A), or residing in the short arm, as in *Strumigenys diabolus* Bolton, 2000 (Fig. 2G, S2E), *Camponotus atriceps* (Smith, 1858) (Fig. 3B, S3B), and *Gigantiops destructor* (Fabricius, 1804) (Fig. 3C, S3C, D). Heteromorphism of 18S rDNA clusters was detected in all of the analyzed *C. atriceps* and *G. tortuolosa* individuals. In the latter species, the heteromorphism of the NOR resulted in differences in total size between

homologous chromosomes, which changed their morphology such that one was submetacentric while the other was subtelocentric. In *G. destructor*, additional GC-rich bands were located in the interstitial region of the long arm of the largest subtelocentric chromosome pair (Fig. S3C, D).

Figure 3. Fluorescence *in situ* hybridization with 18S rDNA probe (red blocks) of an Ectatomminae ant (A) *Gnamptogenys tortuolosa* (2n=44), and Formicinae ants (B) *Camponotus atriceps* (2n=40) and (C) *Gigantiops destructor* (n=39). In *G. tortuolosa*, the box shows a remarkable size heteromorphism of 18S rDNA clusters between homologs. Bars: 5 μ m.

2 - Chromosome mapping review of rDNA clusters in ants

Cytogenetic data available in the literature related to the rDNA genes of 50 ant species from 12 genera and six subfamilies were reviewed (Table 2; Fig. 4). Most data were concentrated on Neotropical ants, with information on 33 species, while the Palearctic

1 and Australian regions had data on only one and 16 species, respectively. A single rDNA
2 site localized in the intrachromosomal region was observed in most species (Fig. 4A, B).
3 However, *Camponotus renggeri* Emery, 1894, *Dinoponera gigantea* (Perty, 1833), and
4 most of the studied *Myrmecia* species presented multiple rDNA sites over the entire short
5 chromosome arm. The subfamily Myrmicinae possessed most of the rDNA data, and the
6 Myrmeciinae subfamily showed a different pattern in relation to other subfamilies with
7 multiple rDNA sites observed in the majority of species (Fig. 4C). *Dolichoderus*
8 *voraginosus* Mackay, 1993 did not show any co-localization of 18S rDNA clusters and
9 GC-rich bands. Figure 5 summarizes the available data on the number and position of
10 NORs in ant species and the phylogenetic relationships among these species, and is based
11 on the published molecular phylogenies.

13 **Discussion**

15 **1 - Patterns of rDNA clusters in the karyotypes of specific ant groups**

17 Specific rDNA patterns can be observed in the karyotypes of some ant groups
18 where several species have been studied. In species of the genus *Dolichoderus*, the
19 chromosome numbers range from $2n=10$ to 58, and a single rDNA site in the interstitial
20 region has been observed in the majority of the investigated species. However, there are
21 two exceptions: *D. voraginosus* and *Dolichoderus attelaboides* (Fabricius, 1775), where
22 the rDNA clusters are located in the terminal region of the long arm and in the short arm,
23 respectively (Santos *et al.*, 2016).

24 Despite the chromosomal variation observed in *Gnamptogenys* spp. ($2n=24$ to 44),
25 they all have a single rDNA site in the intrachromosomal region (Teixeira *et al.*, 2020).

Even *Gnamptogenys moelleri* (Forel, 1912), a species with differing chromosome numbers between two populations ($2n=34$ and 44), has a single intrachromosomal rDNA site. *Gnamptogenys tortuolosa* is an exception, with rDNA clusters occurring over the entire long arm (this study). In the genus *Anochetus*, the chromosome number ranges from $2n=30$ to 46 , and a single pericentromeric rDNA site has been observed (Santos *et al.*, 2010).

All analyzed fungus-farming ants (Attina) had a single rDNA site located in the intrachromosomal region (Table 2). In the genus *Mycetophylax*, although chromosome number differs among species ($2n=26$ to 36), a single rDNA site is located in the pericentromeric or terminal region in all the species of the genus (Micolino *et al.*, 2019a). The leaf-cutting ants are considered the most derived among the Attina species (Schultz and Brady, 2008). *Atta* spp. ($2n=22$) have a single rDNA site in the interstitial region and *Acromyrmex* spp. ($2n=38$) have a single rDNA site in the terminal region (Barros *et al.*, 2015, 2016; Teixeira *et al.*, 2017). *Acromyrmex striatus* (Roger, 1863), a sister group of the leaf-cutting ants, has $2n=22$ and pericentromeric rDNA clusters that are not located on the same chromosome pair relative to that in *Atta* spp. (Cristiano *et al.*, 2013; Teixeira *et al.*, 2017). In *A. echinator* ($2n=38$), rDNA clusters are located in the interstitial region of the same pair as observed in other *Acromyrmex* spp. (this study).

The chromosome number of the Australian bulldog ants is highly variable, ranging from $2n=2$ to 76 (Imai *et al.*, 1994). In addition, *Myrmecia* spp. present remarkable patterns of multiple 28S rDNA clusters that are highly dispersed throughout their genomes. The number of rDNA sites increases with the chromosome number of the species. This suggests several ribosomal gene amplification events have occurred in the different species of *Myrmecia* and that they have accumulated in karyotypes throughout the evolution of the genus. Only four out of 16 species from this monophyletic genus

1 have the entire arm or intrachromosomal rDNA clusters restricted to a single pair of
2 chromosomes (Hirai *et al.*, 1994, 1996; Hirai, 2020). This pattern is observed in species
3 with small chromosome numbers, which suggests that a single NOR is plesiomorphic
4 among *Myrmecia* (Hirai, 2020).

5 Carpenter ants (*Camponotus*) from the subgenus *Myrmothrix* have $2n=40$
6 chromosomes, with its studied species having a single rDNA site in the terminal region.
7 *Camponotus renggeri* is the only exception, having an additional rDNA cluster at the
8 terminal region of a medium-sized subtelocentric pair (Aguar *et al.*, 2017).

9 In the giant ants of the genus *Dinoponera*, two contrasting patterns have been
10 observed: *D. gigantea* ($2n=82$) has multiple rDNA sites located on its short chromosome
11 arms (Aguar *et al.*, 2011), whereas *Dinoponera lucida* Emery, 1901 has a higher
12 chromosome number ($2n=120$), but only a single rDNA site restricted to the
13 intrachromosomal region of its largest chromosome pair (Mariano *et al.*, 2008).

14 Homeology patterns among chromosomal pairs bearing ribosomal genes can be
15 detected in a few ant genera, such as *Gnamptogenys* (*striatula* group), *Camponotus*
16 (*Myrmothrix*), *Atta*, and *Acromyrmex* (Table 2). However, it is speculative to infer such
17 homeology patterns for the entire Formicidae family. Ants constitute an ultra-diverse
18 monophyletic group with more than 13,800 described species (Bolton, 2020). They show
19 a wide range of karyotype variation both in number ($2n=2$ to $2n=120$) and chromosomal
20 morphology (reviewed by Lorite and Palomeque, 2010; Mariano *et al.*, 2019).

21
22 **2 - Insights concerning the organizational patterns of ribosomal gene clusters in the**
23 **ant genome**
24

1 The mapping of the ribosomal gene clusters of 63 species distributed in 19 genera
2 and six subfamilies, together with information on their phylogenetic relationships,
3 demonstrated that a single pair of chromosomes bearing the GC-rich rDNA clusters is the
4 most frequent trait among the studied species, regardless of the chromosome number
5 (Table 1, Fig. 4A). Genomes carrying rDNA clusters in more than a single chromosome
6 pair have been observed in non-related taxa, such as *D. gigantea* (Aguiar *et al.*, 2011), *C.*
7 *renggeri* (Aguiar *et al.*, 2017), and *Myrmecia* spp. (Hirai *et al.*, 1994, 1996). We
8 hypothesize that having a single rDNA site should be considered a plesiomorphic trait
9 because multiple rDNA sites were observed in different non-related lineages that do not
10 share exclusive common ancestry and appear *de novo* throughout the Formicidae family.

Figure 4. Summary of available molecular cytogenetic data (this study and in the literature) concerning rDNA genes in ant species: (A) number of species with single and multiple rDNA sites in a karyotype, (B) location of rDNA genes in chromosomes of ant species, considering the following chromosomal regions: c, centromeric region; p, pericentromeric region; i, interstitial region; t, terminal region; wsa, whole short arm; wla,

whole long arm, and (C) number of species with single and multiple rDNA sites by subfamilies of Formicidae.

In eukaryotes, it is common to observe variations in the number of rDNA clusters and the location of these genes in the chromosomes within genera (Sánchez-Gea *et al.*, 2000; Gross *et al.*, 2010; Cabral-de-Mello *et al.*, 2011; Gokhman *et al.*, 2014; Mazzoleni *et al.*, 2018), among populations (Panzera *et al.*, 2014; Ferreti *et al.*, 2019; Menezes *et al.*, 2019), and sexes of the same species (Nakayama *et al.*, 2001; Šťáhlavský *et al.*, 2018). The possession of terminal rDNA clusters seems to be a common trait among mammals, fish, and mollusks, but less so in arthropods (Sochorová *et al.*, 2018). Within the ultra-diverse insect group, the location of rDNA clusters may follow distinct patterns in its two largest orders; terminal rDNA sites are more abundant in Coleoptera, whereas pericentromeric rDNA clusters are more frequent in Orthoptera (Sochorová *et al.*, 2018). In Formicidae, terminal rDNA sites are a less common feature and all species with multiple rDNA clusters show these genes in the entire short chromosome arms including terminal/subterminal regions, such as *C. renggeri*, *D. gigantea*, and *Myrmecia* spp. (Aguilar *et al.*, 2011, 2017; Hirai *et al.*, 1994, 1996).

Different hypotheses have been proposed to explain the cytogenetic pattern (conservative or variable) of these rDNA clusters in the genome of several organisms. Such hypotheses are linked to the specific locations of these rDNA clusters on the chromosomes. Rearrangements, such as translocations, unequal exchange, and ectopic recombination mechanisms (i.e., between non-homologous chromosomes), which can lead to gene dispersion or increases in number in the genome, are more likely in the terminal/subterminal regions of chromosomes and are uncommon in the intrachromosomal regions (Martins and Wasko, 2004; Mantovani *et al.*, 2005; Nguyen *et al.*, 2010; Roy and Guerra, 2012; Hirai, 2020).

Based on rDNA mapping data in fish karyotypes, Martins and Wasko (2004) proposed that translocations are more likely to occur in telomeric regions because of their proximity within the interphase nucleus, which originates from the ordering of chromosomes according to Rabl's model. Effects due to the location of ribosomal genes in relation to their dispersion in the karyotype were also observed in Coleoptera (Dutrillaux *et al.*, 2016) and primates (Gerbault-Seureau *et al.*, 2017). These authors argued that translocations in the interstitial position could result in abnormal meiosis and, therefore, unbalanced gametes. In contrast, translocations in terminal positions may increase the number of rDNA genes in the genome. This would lead to fewer meiotic abnormalities and highlights the selection for interstitial rDNA site stability (Dutrillaux *et al.*, 2016; Gerbault-Seureau *et al.*, 2017).

Ectopic recombination is another mechanism suggested to explain the rDNA patterning in moths and butterflies (Nguyen *et al.*, 2010) and plants (Roa and Guerra, 2012). It is also included in the recent model proposed by Hirai (2020). In this model, two mechanisms are important: the "site effect" and the "molecular effect." The former allows terminal region associations due to the proximity of these regions in a meiotic bouquet. The "site effect" is a precondition for the "molecular effect," which refers to systems of affinity/non-affinity due to the similarity between rDNA sequences with other repetitive sequences. Thus, rDNA clusters in the terminal regions tend to associate with other repetitive sequences of non-homologous chromosomes more easily, facilitating the occurrence of ectopic recombination and dispersion of these genes in the genome (Hirai, 2020).

There are reports of species with multiple rDNA clusters associated with the centromeres of acrocentric chromosomes (Cazaux *et al.*, 2011). In the recent model proposed by Hirai (2020), the centromeric region of acrocentric chromosomes

(chromosomes with a short and heterochromatic arm) that have rDNA genes associated with the centromere may behave as subterminal regions. Therefore, such an arrangement would also facilitate eventual associations of rDNA genes with other repetitive sequences and the occurrence of ectopic recombination, which leads to their dispersal in the genome (for details, see Hirai, 2020).

The ant rDNA chromosome evolution seems to be in accordance with the above-mentioned hypothesis about dispersal and NOR location because the single rDNA clusters of most studied species are interstitial or pericentromeric (Fig. 4B). In ant species, terminal rDNA clusters are prone to rearrangements that lead to their dispersal. *Camponotus renggeri* (Aguilar *et al.*, 2017), *D. gigantea* (Aguilar *et al.*, 2011), and *Myrmecia* spp. present multiple NORs in the entire short chromosome arms including terminal/subterminal regions, which facilitates the association of these genes with the heterochromatic sequences of other non-homologous acrocentric chromosomes during meiosis and the subsequent occurrence of ectopic recombination (Hirai, 2020). This pattern may be applicable to different ant groups. In addition, inversions have been shown to change the position of rDNA genes in *A. echinator* (this study), *Dolichoderus* spp. (Santos *et al.*, 2016), and *Myrmecia* spp. (Hirai *et al.*, 1996).

1

2 **Figure 5.** Summary of available data (this study and literature) concerning the number
3 and position of rDNA genes in ant species, along with their degree of relatedness and
4 diploid chromosome numbers. Ideograms of rDNA-bearing chromosomes show number

and location (terminal, interstitial, pericentromeric or centromeric) of rDNA clusters (red) in the haploid complement. Phylogenetic relationships are based on 1- Moreau and Bell (2013), 2 - Schmidt (2013), 3 - Larabee *et al.* (2016), 4 - Santos *et al.* (2016), 5 - Hasegawa and Crozier (2006), 6 - Ward *et al.* (2015), 7 - Solomon *et al.* (2019), 8 - Micolino *et al.* (2019a), 9 - Bacci *et al.* (2009), 10 - Queiroz (2015). Colours of phylogenetic branches indicate the following subfamilies: red, Ponerinae; purple, Dolichoderinae; pink, Myrmeciinae; blue, Formicinae; green, Myrmicinae; gray, Ectatomminae.

A single rDNA site located in the terminal region or entire chromosomal arm was observed in some ant species. The repetitive sequences in the subterminal/terminal chromosome regions probably do not form affinity systems with ribosomal genes (the so-called molecular effect; for details, see Hirai, 2020) in these species. Therefore, rDNA clusters are restricted to a single chromosomal pair. Future studies focusing on the characterization of repetitive sequences that make up the heterochromatin of these species will help clarify this hypothesis.

Size heteromorphisms are frequent in karyotypes where the rDNA clusters have terminal positions in the chromosomes, as reported in this study (*G. tortuolosa* and *C. atriceps*) as well as in other ants (Aguiar *et al.*, 2017) and insects in general (Cabral-de-Mello *et al.*, 2011; Maryńska-Nadachowska *et al.*, 2016; Andrade-Souza *et al.*, 2018). Subtle variations in the size of the rDNA clusters between homologous chromosomes can be observed as a result of late condensation during cell division (Sumner, 2003). However, large variations, such as those mentioned above, at the terminal region on the chromosome are usually related to duplications/deletions as a result of unequal exchange (Schubert and Lysack, 2011). It is believed that exchanges are less common in intrachromosomal regions (Hirai, 2020).

1 **Table 1.** Summary of the available molecular cytogenetic data concerning major ribosomal genes (45S, 28S or 18S) detected by FISH in ants

Ant species	2n/(n)	Location of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
Neotropical ants					
Subfamily Myrmicinae					
<i>Acromyrmex aspersus</i>	38	Largest subtelocentric pair		Yes	Teixeira et al. (2017)
<i>Acromyrmex coronatus</i>	38	Largest subtelocentric pair		Yes	Barros et al. (2016)
<i>Acromyrmex disciger</i>	38	Largest subtelocentric pair		Yes	Barros et al. (2016)
<i>Acromyrmex echinator</i>	38	Largest subtelocentric pair		Yes	Barros et al. (2016)/ Present study
<i>Acromyrmex niger</i>	38	Largest subtelocentric pair		Yes ^a	Barros et al. (2016)
<i>Acromyrmex striatus</i>	22	2nd metacentric pair		Yes ^a	Cristiano et al. (2013)/ Teixeira et al. (2017)
<i>Acromyrmex subterraneus molestans</i>	38	Largest subtelocentric pair		Yes	Teixeira et al. (2017)
<i>Atta bisphaerica</i>	22	4th metacentric pair		Yes	Barros et al. (2014)/ Teixeira et al. (2017)
<i>Atta laevigata</i>	22	4th metacentric pair		Yes	Barros et al. (2014)/ Teixeira et al. (2017)
<i>Atta robusta</i>	22	4th metacentric pair		Yes	Barros et al. (2015)
<i>Atta sexdens rubropilosa</i>	22	4th metacentric pair		Yes	Barros et al. (2014)/ Teixeira et al. (2017)

1 **Table 1. Cont.**

Ant species	2n/(n)	Localization of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
<i>Crematogaster longispina</i>	24	Largest metacentric pair		Yes	Present study
<i>Mycetophylax conformis</i>	30	11th metacentric pair		Yes	Cardoso et al. (2014)/ Micolino et al. (2019a)
<i>Mycetophylax morschi</i>	26	2nd submetacentric pair		–	Micolino et al. (2019a)
<i>Mycetophylax morschi</i>	28	7th metacentric pair		–	Micolino et al. (2019a)
<i>Mycetophylax morschi</i>	30	Acrocentric pair		–	Micolino et al. (2019a)
<i>Mycetophylax simplex</i>	36	Smallest metacentric pair		Yes	Cardoso et al. (2014)/ Micolino et al. (2019a)
<i>Mycocepurus goeldii</i>	8	2nd metacentric pair		Yes ^a	Barros et al. (2010)/ Barros et al. (2012)
<i>Myrmicocrypta</i> sp.	30	9th metacentric pair		Yes	Present study
<i>Pheidole germaini</i>	22	Subtelocentric pair		Yes	Present study
<i>Solenopsis geminata</i>	32	Smallest submetacentric pair		Yes	Present study
<i>Strumigenys diabolus</i>	40	4th submetacentric pair		Yes	Present study
<i>Trachymyrmex holmgreni</i>	20	4th metacentric pair		Yes	Barros et al. (2018)
Subfamily Formicinae					
<i>Camponotus atriceps</i> ^c	40	2nd submetacentric pair		Yes	Present study

2
3
4

1 **Table 1. Cont.**

Ant species	2n/(n)	Localization of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
<i>Camponotus cingulatus</i> ^c	40	2nd submetacentric pair		Yes	Aguiar et al. (2017)
<i>Camponotus renggeri</i>	40	2nd submetacentric pair and medium-sized subtelocentric pair		Yes	Aguiar et al. (2017)
<i>Camponotus rufipes</i> ^c	40	2nd submetacentric pair		Yes	Aguiar et al. (2017)
<i>Gigantiops destructor</i>	(39)	8th metacentric pair		Yes ^a	Present study
Subfamily Ponerinae					
<i>Anochetus altisquamis</i> ^c	30	3rd submetacentric pair		–	Santos et al. (2010)
<i>Anochetus horridus</i> ^c	46	4th telocentric pair		–	Santos et al. (2010)
<i>Anochetus targionii</i>	30	7th metacentric pair		Yes	Present study
<i>Dinoponera gigantea</i>	82	Multiple pairs	-	–	Aguiar et al. (2011)
<i>Dinoponera lucida</i>	(59)	Largest pair (A ^{Mt})		Yes	Mariano et al. (2008)
<i>Odontomachus bauri</i> ^d	44/(22)	2nd subtelocentric pair		Yes	Present study
<i>Odontomachus haematodus</i>	44	3rd subtelocentric pair		Yes	Present study
<i>Pseudoponera gilberti</i>	22	Largest metacentric pair		Yes ^a	Present study

2

1 **Table 1.** Cont.

Ant species	2n/(n)	Localization of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
Subfamily Ectatomminae					
<i>Gnamptogenys moelleri</i>	34, 44	4th metacentric pair		Yes	Teixeira et al. (2020)
<i>Gnamptongeys regularis</i> ^d	26	Metacentric and submetacentric homologous		Yes	Teixeira et al. (2020)
<i>Gnamptogenys striatula</i>	32	5th metacentric pair		Yes	Teixeira et al. (2020)
<i>Gnamptogenys striatula</i>	34	4th metacentric pair		Yes	Teixeira et al. (2020)
<i>Gnamptogenys triangularis</i>	24	Largest metacentric pair		Yes	Teixeira et al. (2020)
<i>Gnamptogenys tortuolosa</i> ^c	44	Subtelocentric and submetacentric homologous		Yes	Present study
Subfamily Dolichoderinae					
<i>Dolichoderus attelaboides</i>	58	Largest submetacentric pair		Yes	Santos et al. (2016)
<i>Dolichoderus bidens</i>	18	Largest metacentric pair		Yes ^a	Santos et al. (2016)
<i>Dolichoderus decollatus</i>	38	2nd metacentric pair		Yes	Santos et al. (2016)
<i>Dolichoderus diversus</i>	22	Largest metacentric pair		Yes ^a	Santos et al. (2016)

2

1 **Table 1.** Cont.

Ant species	2n/(n)	Localization of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
<i>Dolichoderus imitator</i>	38	Largest metacentric pair		Yes	Santos et al. (2016)
<i>Dolichoderus lutosus</i>	10	2nd metacentric pair		Yes ^a	Santos et al. (2016)
<i>Dolichoderus voraginosus</i>	20	Largest metacentric pair		No ^a	Santos et al. (2016)
Australian ants					
Subfamily Myrmeciinae					
<i>Myrmecia banksi</i> ^c	10	Smallest acrocentric pair		—	Hirai et al. (1994), (1996)
<i>Myrmecia chasei</i>	47	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia croslandi</i> ^d	2, 3, 4 ^b	Acrocentric pair	,	—	Hirai et al. (1994), (1996)
<i>Myrmecia forficata</i>	52 ^b , 54	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia fulvipes</i>	48, 50	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia gulosa</i>	38 ^b	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia haskinsorum</i>	18	Multiple pairs	-	—	Hirai et al. (1994), (1996)

2
3
4

1 **Table 1.** Cont.

2

Ant species	2n/(n)	Localization of rDNA genes in karyotype	Ideogram	Co-localization CMA ₃ /rDNA	Reference CMA ₃ /rDNA
<i>Myrmecia imaii</i> ^c	8	Largest acrocentric pair		—	Hirai et al. (1994), (1996)
<i>Myrmecia mandibularis</i>	56 ^b	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia michaelseni</i>	27	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia occidentalis</i>	64 ^b	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia pavidia</i>	44	Multiple pairs	-	—	Hirai et al. 1996
<i>Myrmecia pilosula</i>	22 ^b	Acrocentric pair		—	Hirai et al. (1996)
<i>Myrmecia pilosula</i>	19, 23 ^b , 27, 32	Multiple pairs	-	—	Hirai et al. (1994), (1996)
<i>Myrmecia simillima</i>	70 ^b	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia arnoldi</i>	53, 55, 57, 59, 60 ^b , 64, 66	Multiple pairs	-	—	Hirai et al. (1996)
<i>Myrmecia vindex</i>	74, 76	Multiple pairs	-	—	Hirai et al. (1996)
Palearctic ant					
Subfamily Dolichoderinae					
<i>Tapinoma nigerrimum</i>	18	Largest submetacentric pair		Yes	Lorite et al. (1997)

3 ^a: Additional markers CMA₃⁺. ^b: Data showed by Hirai et al. (1996). ^c: Heteromorphism of rDNA clusters between the homologous.

4 ^d: Polymorphism of rDNA clusters between the homologous. — unavailable data.

1 **Table 2.** The relationship between Neotropical ant species studied using 18S rDNA FISH and their chromosome number
2

Ant species	Subfamily	Locality	2n	3
<i>Acromyrmex echinator</i> (Forel, 1899)	Myrmicinae	Barro Colorado – Panamá	38	4
<i>Anochetus targionii</i> Emery 1894	Ponerinae	Campus Agronomique, Kourou – FG	30	5
<i>Camponotus atriceps</i> (Smith, 1858)	Formicinae	Viçosa – Minas Gerais – BR	40	6
<i>Crematogaster longispina</i> Emery, 1890	Myrmicinae	La Montagne des Singes – FG	24	7
<i>Gigantiops destructor</i> (Fabricius, 1804)	Formicinae	Sinnamary – FG	78	8
<i>Gnamptogenys tortuolosa</i> (Smith, 1858)	Ectatominae	Sinnamary – FG	44	9
<i>Myrmicocrypta</i> sp.	Myrmicinae	Sinnamary – FG	30	10
<i>Odontomachus bauri</i> Emery, 1892	Ponerinae	Açailândia – Maranhão – BR	44	11
<i>Odontomachus haematodus</i> (Linnaeus, 1758)	Ponerinae	Campus Agronomique, Kourou – FG	44	12
		Ubá – Minas Gerais – BR	44	13
<i>Pheidole germaini</i> Emery, 1896	Myrmicinae	Viçosa – Minas Gerais – BR	22	14
<i>Pseudoponera gilberti</i> (Kempf, 1960)	Ponerinae	Sinnamary – FG	12	15
<i>Solenopsis geminata</i> (Fabricius, 1804)	Myrmicinae	Sinnamary – FG	32	16
<i>Strumigenys diabolus</i> Bolton 2000	Myrmicinae	Sinnamary – FG	40	17
				18
				19
				20

21 FG: French Guiana, BR: Brazil.

Size variations in the rDNA clusters can be observed when these genes are located in the interstitial/pericentromeric region of the chromosomes, as seen in *Gnamptogenys regularis* Mayr, 1870 (Teixeira *et al.*, 2020) and in *O. bauri* (this study). A different path seems to be involved in the evolution of these karyotypes compared to the rearrangements involved in terminal rDNA heteromorphisms. In these cases, the mechanism may be associated with the formation of extrachromosomal circular DNA (eccDNA), which is likely to form tandem repetitive sequences similar to the rDNA genes (Cohen and Segal, 2009). These eccDNAs may be lost, leading to deletions in the original rDNA sequences, or they may be replicated via a rolling circle mechanism and reintegrated into the original chromosome, producing duplications of these repetitive sequences (Cohen and Segal, 2009).

In ants as well as in other eukaryotes, rDNA clusters are located in GC-rich regions (Symonová, 2019) and, therefore, usually coincide with CMA₃⁺ bands, possibly as a result of GC-biased gene conversion (gBGC) over the course of evolutionary time (Escobar *et al.*, 2011). This co-localization of GC-rich regions and rDNA was observed for all the ants studied to date (Table 2), with the exception of *D. voraginosus* (Santos *et al.*, 2016). However, GC-rich chromatin is not always an indication of ribosomal genes, as seen here in *G. destructor*, *P. gilberti*, and in some *Dolichoderus* spp. and fungus-farming ants (Table 2).

In insects, a compilation of previous data concerning rDNA genes has been conducted, for example, in moths and butterflies (Lepidoptera; Nguyen *et al.*, 2010), beetles (Coleoptera; Dutrillaux and Dutrillaux, 2012), and kissing bugs (Heteroptera; Panzera *et al.*, 2012). However, this is the first survey of Hymenoptera species. We have compiled available information and new data on 13 Neotropical ant species. In different organisms, including ants, the number and location of chromosomes bearing rDNA

clusters within the genome follow general patterns that govern the modes of evolution for these genes (Martins and Wasko, 2004; Nguyen *et al.*, 2010; Dutrillaux *et al.*, 2016; Gerbault-Seureau *et al.*, 2017; Hirai, 2020; this study). We can conclude that having only a single pair of chromosomes bearing rDNA clusters is more common in the ant genome because of the pericentromeric/interstitial location of these genes on the chromosomes. Intrachromosomal regions are sites with low frequencies of rearrangements, such as non-Robertsonian translocations and ectopic recombination, and are therefore less prone to meiotic abnormalities. It should be assumed that the chromosomal location of rDNA clusters influences the dispersion of these genes within the karyotype.

Future studies will allow the mapping of rDNA genes in more ant taxa, including the other remaining subfamilies. Other repetitive sequences, such as 5S rDNA and histone genes, may also be mapped in ant species as a tool to investigate further patterns that reflect the relationship between chromosomal location and dispersion in the genome. Finally, a solid understanding of the evolutionary patterns of ribosomal gene dispersal in ant chromosomes may provide a comparative model for other insects.

Experimental Procedures

1 - Obtaining samples for analysis

Field surveys to collect ant colonies were performed in French Guiana, Brazil, and Panama (Table 1) from the following locations: *La Montagne des Singes*, Kourou (5.07225, -52.69407), Campus Agronomique, Kourou (5.17312, -52.65480), and Sinnamary (5.28482, -52.91403), all in French Guiana; Viçosa (-20.757041, -42.873516)

1 and Ubá (-21.128880, -42.937646), both in Minas Gerais, Brazil, and Açailândia (-
2 4.84200, -47.29667) in Maranhão, Brazil; and Barro Colorado Island (9.150000, -
3 79.83333) in Panama. Sampling permit in Brazil was provided by the Instituto Chico
4 Mendes de Conservação da Biodiversidade (ICMBio) (SISBIO accession numbers 32459
5 and 62598). Adult specimens were identified by Dr. Jacques H. C. Delabie and deposited
6 in the ant collection at the Laboratório de Mirmecologia do Centro de Pesquisas do Cacau
7 (CPDC/Brazil), in Bahia, Brazil, under records #5802, #5803, and #5804.

8 In this study, previous molecular cytogenetic data related to rDNA genes (45S,
9 18S, or 28S) mapped by FISH from 50 ant species were used. For comparative analysis,
10 the following traits were considered for each species: chromosome number, number of
11 rDNA-bearing chromosomes, location of rDNA clusters in the karyotype, and co-
12 localization of CMA₃ fluorochrome and rDNA clusters. Ag-NOR data were disregarded
13 due to the highly variable patterns and unreliable results (for details, see Introduction).

14 15 **2 – Chromosome preparation**

16
17 Mitotic metaphase chromosomes were obtained from the cerebral ganglia of
18 larvae after meconium elimination, using colchicine hypotonic solution (0,005%) and
19 fixatives according to the methods described by Imai *et al.* (1988).

20 21 **3 - Staining with fluorochromes chromomycin A₃ (CMA₃) and 4'-diamidino-2-** 22 **phenylindole (DAPI)**

23
24 Metaphase chromosomes of all the species, except *Ac. echinator*, were stained
25 with the fluorochromes CMA₃ and DAPI for the detection of GC and AT-rich regions,

respectively, based on the technique proposed by Schweizer (1980). The *Ac. echinator* samples studied here correspond to the same colonies studied by Barros *et al.* (2016) and the CMA₃/DAPI staining in this species was performed by these authors.

4 - Fluorescent *in situ* hybridization with rDNA 18S probe

The 18S rDNA probes were obtained by amplification via polymerase chain reaction (PCR) using *Melipona quiquefasciata* Lepeletier, 1836, rDNA primers 18SF1 (5'-GTC ATA GCT TTG TCT CAA AGA-3') and 18SR1.1 (5'-CGC AAA TGA AAC TTT AAT CT-3') (Pereira, 2006) in the genomic DNA from the ant *Camponotus rufipes* (Fabricius, 1775). Gene amplification was performed following Pereira (2006). The probes were labeled by an indirect method using digoxigenin-11-dUTP (Roche Applied Science, Mannheim, Germany), and the FISH signals were detected with anti-digoxigenin-rhodamine (Roche Applied Science), following the manufacturer's protocol.

The rDNA 18S genes were mapped by FISH, following the protocol of Pinkel *et al.* (1986). The slides were treated with RNase A (100 µg/mL) and kept in a moist chamber at 37°C for 1h. After that, they were washed in 2×SSC for 5 min, incubated in 5 µg/ml pepsin in 0.01 N HCl for 10 min, washed in 1×PBS for 5 min and dehydrated in 50%, 70% and 100% alcohol series for 2 min each. After this pretreatment, metaphase chromosomes were denatured in 70% formamide/2×SSC at 75°C for 3 min, and 20 µL of hybridization mix including 200 ng of labeled probe, 2×SSC, 50% formamide, and 10% dextran sulfate was denatured for 10 min at 85°C and added on preparations. The slides were kept in a moist chamber at 37°C overnight. Then, the slides were washed in 2×SSC for 5 min; the detection solution including anti-digoxigenin-rhodamine was added on slides that were kept in a moist chamber at 37°C for 1h. The slides were washed in

4×SSC/Tween, and dehydrated in an alcohol series. Finally, counterstaining with DAPI (DAPI Fluoroshield, Sigma Aldrich) was performed.

5 - Chromosomal analysis

Chromosomes were arranged in order of decreasing size and based on the ratio of the chromosomes arm lengths ($r = \text{long arm/short arm}$), according to the classification proposed by Levan *et al.* (1964). The chromosomes were classified as m = metacentric ($r = 1-1.7$), sm = submetacentric ($r = 1.7-3$), st = subtelocentric ($r = 3-7$), and a = acrocentric ($r > 7$); they were organized using Adobe Photoshop® 21.1.1 and measured using Image Pro Plus®. Ideograms of the NOR-bearing chromosome/chromosomes (i.e., graphical representation of the chromosomes concerning the rDNA clusters) of the ant species were then designed with the Easy Idio software (Diniz and Xavier, 2006).

For the fluorochrome staining and FISH 18S rDNA technique, 30 metaphases from at least three individuals of each species were analyzed. In the case of *O. bauri*, which presented a chromosomal polymorphism involving rDNA clusters, seven individuals were analyzed (six females and one male). The metaphases were analyzed and photographed using a fluorescence microscope, Olympus BX60, attached to an image system, QColor Olympus®, with the filters WB (450–480 nm), WU (330-385 nm), and WG (510-550 nm) for the fluorochromes CMA₃, DAPI, and rhodamine, respectively.

6 - Phylogenetic relationships

The phylogenetic relationship among ant species was determined by associating with previously published molecular phylogenies. The resultant cladogram topology at

the subfamily level was determined following Moreau and Bell (2013). The Poneroid clade topology was determined according to Schmidt (2013) and Larabee *et al.* (2016); the clade topology for the subfamily Dolichoderinae was determined according to Santos *et al.* (2016) and that for the subfamily Myrmicinae was determined according to Bacci *et al.* (2009), Ward *et al.* (2015), Queiroz (2015), Solomon *et al.* (2019), and Micolino *et al.* (2019a). The topology of the species groups within the subfamily Myrmeciinae was determined according to Hasegawa and Crozier (2006).

Acknowledgments

We would like to thank Dr. Cléa S. F. Mariano for kindly providing *Acromyrmex echinator* samples, Dr. Jacques H. C. Delabie for species identification, and Dr. Luiz Fernando Gomes for his valuable assistance in the laboratory with *Odontomachus bauri* chromosome preparations. We also acknowledge Laboratório de Biologia Molecular de Insetos of the Universidade Federal de Viçosa (UFV) for technical support. GAT thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the scholarships granted. Financial support for this study was provided by "Investissement d'Avenir" grants managed by the French Agence Nationale de la Recherche (DRIIHM ref. ANR-11-LABX-0010 and CEBA, ref. ANR-10-LABX-25-01), by the PO-FEDER 2014-2020, Région Guyane (BiNG, ref. GY0007194), and by the Programa de Auxílio ao Pesquisador – PAPESQ/UNIFAP/2019.

Figure S1. Staining with chromomycin A₃ showing GC-rich regions (arrows) of Ponerinae ants: (A) *Pseudoponera gilberti* (2n=12), (B) *Anochetus targionii* (2n=30), (C) *Odontomachus haematodus* (2n=44, French Guiana), (D, E) *Odontomachus bauri* showing the homozygous and heterozygous states, respectively (2n=44), and (F) male of *Odontomachus bauri* with the smaller size regions rich in GC base pairs. Bars = 5 μm.

Figure S2. Staining with chromomycin A₃ showing GC-rich regions (arrows) of Myrmicinae ants: (A) *Pheidole germani* (2n=22), (B) *Crematogaster longispina* (2n=24), (C) *Solenopsis geminata* (n=16), (D) *Myrmicocrypta* sp. (2n=30), (E) *Strumigenys diabolus* (2n=40). Bars = 5 μm.

Figure S3 Staining with chromomycin A₃ showing GC-rich regions (arrows) of an Ectatomminae ant (A) *Gnamptogenys tortuolosa* (2n=44), and Formicinae ants (B) *Camponotus atriceps* (2n=40) and (C, D) *Gigantiops destructor* (2n=78, n=39). Bars = 5 μ m.

References

- Aguiar, H.J.A.C., Barros, L.A.C., Alves, D.R., Mariano, C.S.F., Delabie, J.H.C. and Pompolo, S.G. (2017) Cytogenetic studies on populations of *Camponotus rufipes* (Fabricius, 1775) and *Camponotus renggeri* Emery, 1894 (Formicidae: Formicinae). *PLoS ONE*, **12**(5), e0177702.
- Aguiar, H.J.A.C., Barros, L.A.C., Mariano, C.S.F., Delabie, J.H.C. and Pompolo, S.G. (2011) 45S rDNA localization for the giant Ant *Dinoponera gigantea* with Evolutionary inferences for the *Dinoponera* Genus (Formicidae: Ponerinae). *Sociobiology*, **57**(3), 607–620.
- Andrade-Souza, V., Duarte, O.M.P., Martins, C.C.C., Santos, I.S., Costa, M.G.C. and Costa, M.A. (2018) Comparative molecular cytogenetics of *Melipona* Illiger species (Hymenoptera: Apidae). *Sociobiology*, **65**(4), 696–705.
- Bacci, Jr. M., Solomon, S.E., Mueller, U.G., Martins, V.G., Carvalho, A.O.R., Vieira, L.G.E. and Silva-Pinhati, A.C.O. (2009). Phylogeny of leafcutter ants in the genus *Atta* Fabricius (Formicidae: Attini) based on mitochondrial and nuclear DNA sequences. *Molecular Phylogenetics and Evolution*, **51**(3), 427–437.
- Barbosa, P., Leal, E.V., Silva, M., Almeida, M.C., Moreira-Filho, O. and Artoni, R.F. (2017). Variability and evolutionary implications of repetitive DNA dynamics in genome of *Astyanax scabripinnis* (Teleostei, Characidae). *Comparative Cytogenetics*, **11**(1), 143–162.
- Barros, L.A.C., Aguiar, H.J.A.C., Andrade-Souza, V., Mariano, C.S.F., Delabie, J.H.C. and Pompolo, S.G. (2012). Occurrence of pre-nucleolar bodies and 45S rDNA location

1 on the chromosomes of the ant *Mycocepurus goeldii* (Forel) (Formicidae, Myrmicinae,
2 Attini). *Hereditas*, **149**, 50–54.

3 Barros, L.A.C., Aguiar, H.J.A.C., Mariano, C.S., Andrade-Souza, V., Costa, M.A.,
4 Delabie, J.H.C. and Pompolo, S.G. (2016). Cytogenetic data on six leafcutter ants of the
5 genus *Acromyrmex* Mayr, 1865 (Hymenoptera, Formicidae, Myrmicinae): insights into
6 chromosome evolution and taxonomic implications. *Comparative Cytogenetics*, **10**(2),
7 229–243.

8 Barros, L.A.C., Aguiar, H.J.A.C., Mariano, C.S.F., Delabie, J.H.C. and Pompolo, S.G.
9 (2010). Cytogenetic characterization of the lower-Attine *Mycocepurus goeldii*
10 (Formicidae: Myrmicinae: Attini). *Sociobiology*, **56**, 57–66.

11 Barros, L.A.C., Aguiar, H.J.A.C., Teixeira, G.A., Mariano, C.S.F., Teixeira, M.C.,
12 Delabie, J.H.C. and Pompolo, S.G. (2015). Cytogenetic data on the threatened leafcutter
13 ant *Atta robusta* Borgmeier, 1939 (Formicidae: Myrmicinae: Attini). *Comptes Rendus*
14 *Biologies*, **338**(10), 660–665.

15 Barros, L.A.C., Mariano, C.S.F., Pompolo, S.G., and Delabie, J.H.C. (2009). Hsc-FA
16 and NOR bandings on chromosomes of the giant ant *Dinoponera lucida* Emery, 1901
17 (Hymenoptera: Formicidae). *Comparative Cytogenetics*, **3**, 97-102.

18 Barros, L.A.C., Teixeira, G.A., Aguiar, H.J.A.C., Lopes, D.M. and Pompolo, S.G.
19 (2018). Cytogenetic studies in *Trachymyrmex holmgreni* Wheeler, 1925 (Formicidae:
20 Myrmicinae) by conventional and molecular methods. *Sociobiology*, **65**(2), 185–190.

21 Barros, L.A.C., Teixeira, G.A., Aguiar, H.J.A.C., Mariano, C.S.F, Delabie, J.H.C. and
22 Pompolo, S.G. (2014). Banding patterns of three leafcutter ant species of the genus *Atta*
23 (Formicidae: Myrmicinae) and Chromosomal Inferences. *Florida Entomologist*, **97**(4),
24 1694–1701.

1 Bolton, B. 2020. An online catalog of the ants of the world. Available from
2 <https://antcat.org>. (accessed 15th May, 2020)

3 Britton-Davidian, J., Cazaux, B. and Catalan, J. (2012). Chromosomal dynamics of
4 nucleolar organizer regions (NORs) in the house mouse: micro-evolutionary insights.
5 *Heredity*, **108**(1), 68-74.

6 Cabral-de-Mello, D.C., Oliveira, S.G., Moura, R.C. and Martins, C. (2011).
7 Chromosomal organization of the 18S and 5S rRNAs and histone H3 genes in
8 Scarabaeinae coleopterans: insights into the evolutionary dynamics of multigene
9 families and heterochromatin. *BMC Genetics*, **12**, 88.

10 Cardoso, D.C., Pompolo, S.G., Cristiano, M.P. and Tavares, M.G. (2014). The role of
11 fusion in ant chromosome evolution: insights from cytogenetic analysis using a
12 molecular phylogenetic approach in the genus *Mycetophylax*. *PLoS ONE*, **9**(4), e95408.

13 Castro, C.P.M., Cardoso, D.C., Micolino, R., and Cristiano, M.P. (2020). Comparative
14 FISH-mapping of TTAGG telomeric sequences to the chromosomes of leafcutter ants
15 (Formicidae, Myrmicinae): is the insect canonical sequence conserved? *Comparative*
16 *Cytogenetics*, **14**(3), 369–385.

17 Cazaux, B., Catalan, J., Veyrunes, F., Douzery, E.J. and Britton-Davidian, J. (2011).
18 Are ribosomal DNA clusters rearrangement hotspots? A case study in the genus *Mus*
19 (Rodentia, Muridae). *BMC Evolutionary Biology*, **11**(1), 124.

20 Cholak, L.R., Haddad, C.F. and Parise-Maltempi, P.P. (2020). Cytogenetic analysis of
21 the genus *Thoropa* Cope, 1865 (Anura-Cycloramphidae) with evolutionary inferences
22 based on repetitive sequences. *Genetics and Molecular Biology*, **43**(3).

1 Cohen, S. and Segal, D. (2009). Extrachromosomal Circular DNA in Eukaryotes:
2 possible involvement in the plasticity of tandem repeats. *Cytogenetic and Genome*
3 *Research*, **124**, 327–338.

4 Cristiano, M.P., Cardoso, D.C. and Fernandes-Salomão, T.M. (2013). Cytogenetic and
5 molecular analyses reveal a divergence between *Acromyrmex striatus* (Roger, 1863) and
6 other congeneric species: taxonomic implications. *PLoS ONE*, **8**(3), e59784.

7 Cross, I., Vega, L., and Rebordinos, L. (2003). Nucleolar organizing regions in
8 *Crassostrea angulata*: chromosomal location and polymorphism. *Genetica*, **119**(1), 65-
9 74.

10 Degrandi, T.M., Gunski, R.J., Garnerio, A.D.V., Oliveira, E.H.C.D., Kretschmer, R.,
11 Souza, M.S.D., Barcellos S.A. and Hass, I. (2020). The distribution of 45S rDNA sites
12 in bird chromosomes suggests multiple evolutionary histories. *Genetics and Molecular*
13 *Biology*, **43**(2).

14 Diniz, D. and Xavier, P.M. (2006). Easy Idio.

15 Dobigny, G., Ozouf-Costaz, C., Bonillo, C. and Volobouev, V. (2002). “Ag-NORs” are
16 not always true NORs: new evidence in mammals. *Cytogenetic and Genome Research*,
17 **98**(1), 75-77.

18 Dutra, R.T., Bitencourt, J., Netto, M.R.C.B., Paim, F.G., Sarmiento-Soares. L.M. and
19 Afonso, P.R.A.M. (2020). Chromosomal markers are useful to species identification in
20 Rivulidae (Cyprinodontiformes, Aplocheiloidei), including the resolution of taxonomic
21 uncertainties in a vulnerable species complex. *Zebrafish*, **17**(1), 48–55.

22 Dutrillaux, A.M. and Dutrillaux, B. (2012). Chromosome analysis of 82 species of
23 Scarabaeoidea (Coleoptera), with special focus on NOR localization. *Cytogenetic and*
24 *Genome Research*, **136**(3), 208–219.

- 1 Dutrillaux, A.M., Carton, B., Cacheux, L and Dutrillaux, B. (2016). Interstitial NORs,
2 fragile sites, and chromosome evolution: a not so simple relationship-the example of
3 *Melolontha melolontha* and genus *Protaetia* (Coleoptera: Scarabaeidae). *Cytogenetic*
4 *and Genome Research*, **149**(4), 304–311.
- 5 Escobar, J.S., Glémin, S. and Galtier, N. (2011). GC-biased gene conversion impacts
6 ribosomal DNA evolution in vertebrates, angiosperms, and other eukaryotes. *Molecular*
7 *Biology and Evolution*, **28**(9), 2561–2575.
- 8 Falcione, C., Hernando, A. and Bressa, M.J. (2018). Comparative cytogenetic analysis
9 in *Erythrolamprus* snakes (Serpentes: Dipsadidae) from Argentina. *Anais da Academia*
10 *Brasileira de Ciências*, **90**(2), 1417-1429.
- 11 Ferreti, A.B.S.M., Ruiz-Ruano, F.J., Milani, D., Loreto, V., Martí, D.A., Ramos, E.,
12 Martins, C. and Cabral-de-Mello, D.C. (2019). How dynamic could be the 45S rDNA
13 cistron? An intriguing variability in a grasshopper species revealed by integration of
14 chromosomal and genomic data. *Chromosoma*, **128**, 165–175.
- 15 Gerbault-Seureau, M., Cacheux, L. and Dutrillaux, B. (2017). The relationship between
16 the (In-) stability of NORs and their chromosomal location: the example of
17 Cercopithecidae and a short review of other primates. *Cytogenetic and Genome*
18 *Research*, **153**(3), 138–146.
- 19 Gokhman, V.E., Anokhin, B.A. and Kuznetsova, V.G. (2014). Distribution of 18S
20 rDNA sites and absence of the canonical TTAGG insect telomeric repeat in parasitoid
21 Hymenoptera. *Genetica*, **142**, 317–322.
- 22 Gokhman, V.E., Bolsheva, N.L., Govind, S. and Muravenko, O.V. (2016). A
23 comparative cytogenetic study of *Drosophila* parasitoids (Hymenoptera, Figitidae)

1 using DNA-binding fluorochromes and FISH with 45S rDNA probe. *Genetica*, **144**(3),
2 335–339.

3 Golub, N.V., Golub, V.B. and Kuznetsova, V.G. (2015). Variability of 18rDNA loci in
4 four lace bug species (Hemiptera, Tingidae) with the same chromosome number.
5 *Comparative Cytogenetics*, **9**(4), 513–522.

6 Gross, M.C., Schneider, C.H., Valente, G.T., Martins, C. and Feldberg E. (2010).
7 Variability of 18S rDNA locus among *Symphysodon* fishes: chromosomal
8 rearrangements. *Journal of Fish Biologies*, **76**(5), 1117–1127.

9 Hasegawa, E, and Crozier, R.H. (2006). Phylogenetic relationships among species
10 groups of the ant genus *Myrmecia*. *Molecular Phylogenetics and Evolution*, **38**, 575–
11 582.

12 Hirai, H. (2020). Chromosome dynamics regulating genomic dispersion and alteration
13 of Nucleolus Organizer Regions (NORs). *Cells*, **9**(4), 971–984.

14 Hirai, H., Yamamoto, M.T., Ogura, K., Satta, Y., Yamada, M., Taylor, R.W. and Imai,
15 H.T. (1994). Multiplication of 28S rDNA and NOR activity in chromosome evolution
16 among ants of the *Myrmecia pilosula* species complex. *Chromosoma*, **103**(3), 171–178.

17 Hirai, H., Yamamoto, M.T., Taylor, R.W. and Imai, H.T. (1996). Genomic dispersion of
18 28S rDNA during karyotypic evolution in the ant genus *Myrmecia* (Formicidae).
19 *Chromosoma*, **105**, 190–196.

20 Howell, W.T., and Black, D.A. (1980). Controlled silver-staining of nucleolus organizer
21 regions with a protective colloidal developer: a 1-step method. *Experientia*, **36**(8),
22 1014-1015.

1 Huang, Y.C., Lee, C.C., Kao, C.Y., Chang, N.C., Lin, C.C., Shoemaker, D. and Wang,
2 J. (2016). Evolution of long centromeres in fire ants. *BMC Evolutionary Biology*, **16**,
3 189.

4 Imai, H., Taylor, R.W., Crosland, M.W. and Crozier, R.H. (1988). Modes of
5 spontaneous chromosomal mutation and karyotype evolution in ants with reference to
6 the minimum interaction hypothesis. *Japanese Journal of Genetics*, **63**, 159–185.

7 Imai, H.T., Hirai, H., Satta, Y., Shiroishi, T., Yamada, M., and Taylor, R.W. (1992).
8 Phase specific Ag-staining of nucleolar organizer regions (NORs) and kinetochores in
9 the Australian ant *Myrmecia croslandi*. *The Japanese Journal of Genetics*, **67**(6), 437-
10 447.

11 Imai, H.T., Taylor, R.W., and Crozier, R.H. (1994). Experimental bases for the
12 minimum interaction theory. I. Chromosome evolution in ants of the *Myrmecia pilosula*
13 species complex (Hymenoptera: Formicidae: Myrmeciinae). *The Japanese Journal of*
14 *Genetics*, **69**(2), 137-182.

15 Larabee, F.J., Fisher, B.K., Schmidt, C.A., Matos-Maraví, P., Janda, M. and Suarez,
16 A.V. (2016). Molecular phylogenetics and diversification of trap-jaw ants in the genera
17 *Anochetus* and *Odontomachus* (Hymenoptera: Formicidae). *Molecular Phylogenetics*
18 *and Evolution*, **103**, 143–154.

19 Levsky, J.F. and Singer, R.H. (2003). Fluorescence *in situ* hybridization: past, present
20 and future. *Journal of Cell Science*, **116**, 2833-2838.

21 Liehr, T. (2017). “Classical cytogenetics” is not equal to “banding cytogenetics”.
22 *Molecular Cytogenetics*, **10**, 1–3.

23 Long, E.O. and Dawid, I.B. (1980). Repeated genes in eukaryotes. *Annual Review of*
24 *Biochemistry*, **49**(1), 727–764.

- 1 Lorite, P., and Palomeque, T. (2010). Karyotype evolution in ants (Hymenoptera:
2 Formicidae), with a review of the known ant chromosome numbers. *Myrmecological*
3 *News*, **13**(1), 89-102.
- 4 Lorite, P., Aránega, A.E., Luque, F. and Palomeque, T. (1997). Analysis of the
5 nucleolar organizing regions in the ant *Tapinoma nigerrimum* (Hymenoptera,
6 Formicidae). *Heredity*, **78**, 578–582.
- 7 Lorite, P., Carrillo, J.A., Tinaut, A. and Palomeque, T. (2004). Evolutionary dynamics
8 of satellite DNA in species of the genus *Formica* (Hymenoptera, Formicidae). *Gene*,
9 **332**, 159–168.
- 10 Malimpensa, G.D.C., Traldi, J.B., Martinez, J.D.F., Deon, G., Azambuja, M., Nogaroto,
11 V., Vicari M.R. and Moreira-Filho, O. (2020). Chromosomal Diversification in Two
12 Species of *Pimelodus* (Siluriformes: Pimelodidae): Comparative Cytogenetic Mapping
13 of Multigene Families. *Zebrafish*, **17**(4), 278-286.
- 14 Mantovani, M., Abel. L.D. and Moreira-Filho, O. (2005). Conserved 5S and variable
15 45S rDNA chromosomal localization revealed by FISH in *Astyanax scabripinnis*
16 (Pisces, Characidae). *Genetica*, **123**, 211–216.
- 17 Mariano, C.S.F., Barros, L.A.C., Velasco, Y.M., Guimarães, I.N., Pompolo, S.G. and
18 Delabie, J.H.C. (2019). Citogenética de hormigas de la región neotropical, In Hormigas
19 de Colombia (Fernández, F., Guerrero, R. and Delsinne, T., eds.). pp. 131–157.
20 Universidad Nacional de Colombia, Bogotá.
- 21 Mariano, C.S.F., Pompolo, S.G., Barros, L.A.C., Mariano-Neto, E., Campiolo, S. and
22 Delabie, J.C.H. (2008). A biogeographical study of the threatened ant *Dinoponera*
23 *lucida* Emery (Hymenoptera: Formicidae: Ponerinae) using a cytogenetic approach.
24 *Insect Conservation and Diversity*, **1**(3), 161–168.

1 Martins, C. and Wasko, A.P. (2004). Organization and evolution of 5S ribosomal DNA
2 in the fish genome. In Focus on genome research (Williams, C.R., ed.). pp. 335–363.
3 Nova Science Publishers Inc, New York.

4 Maryańska-Nadachowska, A., Anokhin, B.A., Gnezdilov, V.M. and Kuznetsova, V.G.
5 2016. Karyotype stability in the family Issidae (Hemiptera, Auchenorrhyncha) revealed
6 by chromosome techniques and FISH with telomeric (TTAGG)_n and 18S rDNA probes.
7 *Comparative Cytogenetics*, **10(3)**, 347–369.

8 Mazzoleni, S., Rovatsos, M., Schillaci, O. and Dumas, F. (2018). Evolutionary insight
9 on localization of 18S, 28S rDNA genes on homologous chromosomes in Primates
10 genomes. *Comparative Cytogenetics*, **12(1)**, 27–40.

11 Menezes, R.S.T., Gazoni, T. and Costa, M.A. (2019). Cytogenetics of warrior wasps
12 (Vespidae: *Synoeca*) reveals intense evolutionary dynamics of ribosomal DNA clusters
13 and an unprecedented number of microchromosomes in Hymenoptera. *Biological*
14 *Journal of the Linnean Society*, **126(4)**, 925–935.

15 Meyne, J., Hirai, H. and Imai, H.T. (1995). FISH analysis of the telomere sequences of
16 bulldog ants (*Mymecia*: Formicidae). *Chromosoma*, **104**, 14–18.

17 Micolino, R., Cristiano, M.P., Travenzoli, N.M., Lopes, D.M. and Cardoso, D.C.
18 (2019a). Chromosomal dynamics in space and time: evolutionary history of
19 *Mycetophylax* ants across past climatic changes in the Brazilian Atlantic coast.
20 *Scientific Reports*, **9(1)**, 1–13.

21 Micolino, R., Cristiano, M. P. and Cardoso, D. C. (2019b). Population-based
22 cytogenetic banding analysis and phylogenetic relationships of the neotropical fungus-
23 farming ant *Trachymyrmex holmgreni* Wheeler, 1925. *Cytogenetic and Genome*
24 *Research*, **159(3)**, 151-162.

- 1 Micolino, R., Cristiano, M.P., and Cardoso, D.C. (2020). Karyotype and putative
2 chromosomal inversion suggested by integration of cytogenetic and molecular data of
3 the fungus-farming ant *Mycetomoellerius iheringi* Emery, 1888. *Comparative*
4 *Cytogenetics*, **14**(2), 197–210.
- 5 Moreau, C.S. and Bell, C.D. (2013). Testing the museum versus cradle tropical
6 biological diversity hypothesis: phylogeny, diversification, and ancestral biogeographic
7 range evolution of the ants. *Evolution*, **67-8**, 2240–2257.
- 8 Nguyen, P., Sahara, K., Yoshido, A. and Marec, F. (2010). Evolutionary dynamics of
9 rDNA clusters on chromosomes of moths and butterflies (Lepidoptera). *Genetica*, **138**,
10 343–354.
- 11 Panzera, F., Ferreiro, M.J., Pita, S., Calleros, L., Pérez, R., Basmadjian, Y., Guevara,
12 Y., Brenière, S.F. and Panzera, Y. (2014). Evolutionary and dispersal history of
13 *Triatoma infestans*, main vector of Chagas disease, by chromosomal markers. *Infection*,
14 *Genetics and Evolution*, **27**, 105–113.
- 15 Panzera, Y.I., Pita, S., Ferreiro, M.J., Ferrandis, I., Lages, C., Pérez, R., Silva, A.E.,
16 Guerra, M. and Panzera, F. (2012). High dynamics of rDNA cluster location in kissing
17 bug holocentric chromosomes (Triatominae, Heteroptera). *Cytogenetics and Genome*
18 *Research*, **38**(1), 56–67.
- 19 Pereira, J.O.P. (2006). Diversidade genética da abelha sem ferrão *Melipona*
20 *quinquefasciata* baseada no sequenciamento das regiões ITS1 parcial e 18S do DNA
21 ribossômico nuclear. Unpublished D. Phil. Thesis, Universidade Federal do Ceará,
22 Fortaleza, Brazil.
- 23 Pereira, T.T.P., Reis, A.C.C., Cardoso, D.C. and Cristiano, M.P. (2018). Molecular
24 phylogenetic reconstruction and localization of the (TTAGG)_n telomeric repeats in the

1 chromosomes of *Acromyrmex striatus* (Roger, 1863) suggests a lower ancestral
2 karyotype for leafcutter ants (Hymenoptera). *Comparative Cytogenetics*, **12**(1), 13–26.

3 Pinkel, D., Straume, T. and Gray, J.W. (1986). Cytogenetic analysis using quantitative,
4 high-sensitivity, fluorescence hybridization. *Proceedings of the National Academy of*
5 *Sciences of the United States of America*, **83**, 2934–2938.

6 Queiroz, E.C. (2015). Estudos moleculares em *Acromyrmex* Mayr, 1865. Unpublished
7 D. Phil. Thesis, Universidade Federal do Paraná, Curitiba, Brazil.

8 Roa, F. and Guerra, M. (2012). Distribution of 45S rDNA sites in chromosomes of
9 plants: structural and evolutionary implications. *BMC Evolutionary Biology*, **12**(1), 1-
10 13.

11 Roy, V, Monti-Dedieu, L., Chaminade, N., Siljak-Yakovlev, S., Aulard, S., Lemeunier,
12 F., Montchamp-Moreau, C. (2005). Evolution of the chromosomal location of rDNA
13 genes in two *Drosophila* species subgroups: *ananassae* and *melanogaster*. *Heredity*,
14 **94**(4), 388–395.

15 Sánchez-Gea, J.F., Serrano, J. and Galián, J. (2000). Variability in rDNA loci in Iberian
16 species of the genus *Zabrus* (Coleoptera: Carabidae) detected by fluorescence *in situ*
17 hybridization. *Genome* **43**, 22–28.

18 Santos, I.S., Mariano, C.S.F., Andrade, V., Costa, M.A., Delabie, J.H.C. and Silva, J.G.
19 (2010). A Cytogenetic Approach to the Study of Neotropical *Odontomachus* and
20 *Anochetus* Ants (Hymenoptera: Formicidae). *Annals of the Entomological Society of*
21 *America*, **103**(3), 424–429.

22 Santos, I.S., Mariano, C.S.F., Delabie, J.H.C., Costa, M.A., Carvalho, A.F. and Silva,
23 JG. (2016). "Much more than a neck": karyotype differentiation between *Dolichoderus*
24 *attelaboides* (Fabricius, 1775) and *Dolichoderus decollatus* F. Smith, 1858

1 (Hymenoptera: Formicidae) and karyotypic diversity of five other Neotropical species
2 of *Dolichoderus* Lund, 1831. *Myrmecological News*, **23**, 61–69.

3 Schmid, M., Steinlein, C., Feichtinger, W. and Nanda, I. (2017). Chromosome banding
4 in Amphibia. XXXV. Highly mobile nucleolus organizing regions in *Craugastor*
5 *fitzingeri* (Anura, Craugastoridae). *Cytogenetic and Genome Research*, **152**(4), 180-193.

6 Schmidt, C. (2013). Molecular phylogenetics of ponerine ants (Hymenoptera:
7 Formicidae: Ponerinae). *Zootaxa*, **3647**(2), 201–250.

8 Schubert, I. and Lysack, M.A. (2011). Interpretation of karyotype evolution should
9 consider chromosome structural constraints. *Trends in Genetics*, **27**(6), 207–216.

10 Schultz, T.R. and Brady S.G. (2008). Major evolutionary transitions in ant agriculture.
11 *Proceedings of the National Academy of Sciences of the United States of America*,
12 **105**(14), 5435–5440.

13 Schweizer, D. (1980). Simultaneous fluorescent staining of R bands and specific
14 heterocromatic regions (DA/DAPI-bands) in human chromosomes. *Cytogenetics and*
15 *Cell Genetics*, **27**, 190–193.

16 Sochorová, J., Garcia, S., Gálvez, F., Symonová, R. and Kovařík, A. (2018).
17 Evolutionary trends in animal ribosomal DNA loci: introduction to a new online
18 database. *Chromosoma*, **127**, 141–150.

19 Solomon, S.E., Rabeling, C., Sosa-Calvo, J., Lopes, C.T., Rodrigues, A., Vasconcelos,
20 H.L., Bacci, Jr.M., Mueller, U.G. and Schultz, T.R. (2019). The molecular
21 phylogenetics of *Trachymyrmex* Forel ants and their fungal cultivars provide insights
22 into the origin and coevolutionary history of ‘higher attine’ ant agriculture. *Systematic*
23 *Entomology*, **44**(4), 939–956.

- 1 Šťáhlavský, F., Opatova, V., Just, P., Lotz, L.N. and Haddad, C.R. (2018). Molecular
2 technique reveals high variability of 18S rDNA distribution in harvestmen (Opiliones,
3 Phalangidae) from South Africa. *Comparative Cytogenetics*, **12**(1), 41–59.
- 4 Sumner, A.T. (2003). Chromosomes: Organization and Function. Blackwell Publishing,
5 North Berwick, United Kingdom.
- 6 Symonová, R. (2019). Integrative rDNAomics - Importance of the oldest repetitive
7 fraction of the eukaryote genome. *Genes*, **10**(5), 345–360.
- 8 Teixeira, G.A., Barros, L.A.C., Aguiar, H.J.A.C. and Pompolo, S.G. (2017).
9 Comparative physical mapping of 18S rDNA in the karyotypes of six leafcutter ant
10 species of the genera *Atta* and *Acromyrmex* (Formicidae: Myrmicinae). *Genetica*, **145**,
11 351–357.
- 12 Teixeira, G.A., Barros, L.A.C., Lopes, D.M. and Aguiar, H.J.A.C. (2020). Cytogenetic
13 variability in four species of *Gnamptogenys* Roger, 1863 (Formicidae: Ectatomminae)
14 showing chromosomal polymorphisms, species complex, and cryptic species.
15 *Protoplasma*, **257**, 549–560.
- 16 Vicari, M.R., Artoni, R.F., Moreira-Filho, O. and Bertollo, L.A.C. (2008).
17 Diversification of a ZZ/ZW sex chromosome system in *Characidium* fish (Crenuchidae,
18 Characiformes). *Genetica*, **134**(3), 311.
- 19 Walker, L.I., Soto, M.A. and Spotorno, Á.E. (2014). Similarities and differences among
20 the chromosomes of the wild guinea pig *Cavia tschudii* and the domestic guinea pig
21 *Cavia porcellus* (Rodentia, Caviidae). *Comparative Cytogenetics*, **8**(2), 153.
- 22 Ward, P.S., Brady, S.G., Fisher, B.L. and Schultz, T.R. (2015). The evolution of
23 myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade
24 (Hymenoptera: Formicidae). *Systematic Entomology*, **40**, 61–81

- 1 Zurita, F., Sánchez, A., Burgos, M., Jiménez, R. and de la Guardia, R.D. (1997).
- 2 Interchromosomal, intercellular and interindividual variability of NORs studied with
- 3 silver staining and *in situ* hybridization. *Heredity*, **78**(3), 229-234.