Impact of Pleistocene Eustatic Fluctuations on Evolutionary Dynamics in Southeast Asian Biodiversity Hotspots

Arni Sholihah^{1,2,*}, Erwan Delrieu-Trottin^{2,3}, Fabien L. Condamine², Daisy Wowor⁴, Lukas Rüber^{5,6}, Laurent Pouyaud², Jean-François Agnès², Nicolas Hubert²

¹*Institut Teknologi Bandung, School of Life Sciences and Technology, Jalan Ganesha 10, Bandung 40132, Indonesia.*

²*UMR 5554 ISEM (IRD, UM, CNRS, EPHE), Université de Montpellier, Place Eugène Bataillon, 34095 Montpellier cedex 05, France*

³*Museum für Naturkunde, Leibniz-Institut für Evolutions und Biodiversitätsforschung an der Humboldt-Universität zu Berlin*

⁴*Division of Zoology, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Jalan Raya Jakarta Bogor Km 46, Cibinong 16911, Indonesia.*

⁵*Naturhistorisches Museum Bern, Bernastrasse 15, 3005 Bern, Switzerland*

⁶*Aquatic Ecology and Evolution, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland*

**Correspondence to be sent to UMR 5554 ISEM (IRD, UM, CNRS, EPHE), Université de Montpellier, Bâtiment 22, Place Eugène Bataillon, 34095 Montpellier cedex 05, France; E-mail: arni.sholihah@gmail.com*

ABSTRACT

Pleistocene Climatic Fluctuations (PCF) are frequently highlighted as important evolutionary engines that triggered cycles of biome expansion and contraction. While there is ample evidence of the impact of PCF on biodiversity of continental biomes, the consequences

in insular systems depend on the geology of the islands and the ecology of the taxa inhabiting them. The idiosyncratic aspects of insular systems are exemplified by the islands of the Sunda Shelf in Southeast Asia (Sundaland), where PCF-induced eustatic fluctuations had complex interactions with the geology of the region, resulting in high species diversity and endemism. Emergent land in Southeast Asia varied drastically with sea level fluctuations during the Pleistocene. Climate-induced fluctuations in sea level caused temporary connections between insular and continental biodiversity hotspots in Southeast Asia. These exposed lands likely had freshwater drainage systems that extended between modern islands: the Paleoriver Hypothesis. Built upon the assumption that aquatic organisms are among the most suitable models to trace ancient river boundaries and fluctuations of landmass coverage, the present study aims to examine the evolutionary consequences of PCF on the dispersal of freshwater biodiversity in Southeast Asia. Time-calibrated phylogenies of DNA-delimited species were inferred for six species-rich freshwater fish genera in Southeast Asia (*Clarias*, *Channa*, *Glyptothorax*, *Hemirhamphodon*, *Dermogenys*, *Nomorhamphus*). The results highlight rampant cryptic diversity and the temporal localization of most speciation events during the Pleistocene, with 88% of speciation events occurring during this period. Diversification analyses indicate that sea level-dependent diversification models poorly account for species proliferation patterns for all clades excepting *Channa*. Ancestral area estimations point to Borneo as the most likely origin for most lineages, with two waves of dispersal to Sumatra and Java during the last 5 Myr. Speciation events are more frequently associated with boundaries of the paleoriver watersheds, with 60%, than islands boundaries, with 40%. In total, one-third of speciation events are inferred to have occurred within paleorivers on a single island, suggesting that habitat heterogeneity and factors other than allopatry between islands substantially affected diversification of Sundaland fishes. Our results suggest that species proliferation in Sundaland is not wholly reliant on Pleistocene sea-level fluctuations isolating

populations on different islands. [Dispersal; diversification; eustatic fluctuations; freshwater fishes; insular systems; Milankovitch cycles; paleoenvironments; vicariance]

INTRODUCTION

Pleistocene Climatic Fluctuations (PCF) are widely considered to be drivers of species diversification and distribution patterns (Dodson et al. 1995; Haffer 1997; Lohman et al. 2011; de Bruyn et al. 2014) that contributed to the development of highly heterogeneous patterns of species richness and endemism (Voris 2000; Nores 2004; Hubert and Renno 2006; Pellissier et al. 2014). Temporally located between 2.58-0.012 million years ago (Ma) (Walker et al. 2018), PCFs result from dynamic interactions between tectonic movements and oscillations of Earth's orbit, leading to fluctuations of average temperature and global climate. These Milankovitch cycles have impacted biomes through paleoenvironmental dynamics such as eustatic fluctuations. This caused physical fragmentation of terrestrial and freshwater biotas and shaped contemporary distribution patterns (Dodson et al. 1995; Gathorne-Hardy et al. 2002; Currie et al. 2004; Bird et al. 2005; Cannon et al. 2009; Beck et al. 2017). Climate differences are frequently invoked to account for heterogeneous speciation and extinction patterns in temperate and tropical biomes. For instance, the Late Pleistocene Hypothesis pinpoints glacial cycles and associated sea-level fluctuations as the main drivers of increased speciation in the Northern Hemisphere (Barraclough and Nee 2001; Wiens and Donoghue 2004; Mittelbach et al. 2007; Beck et al. 2017). In the tropics, repeated marine incursions and lowland drying resulted in fragmentation of highlands (Haffer 1997; Nores 1999; Hubert and Renno 2006) and shrinking of terrestrial biotas into refugia, leading to extinction of coastal organisms (Condamine et al. 2015b) or the divergence of isolated populations across refugia (Bates et al. 1998; Nores 1999; Hubert and Renno 2006; Condamine et al. 2015). Pleistocene Climatic Fluctuations are frequently considered to be a “species pump” that offered increased

opportunities of speciation through cycles of alternating species' ranges contraction and populations' fragmentation during glacial maxima with dispersal and species' ranges expansion during inter-glacial times (Esselstyn and Brown 2009; April et al. 2012; Brown et al. 2013; Papadopoulou and Knowles 2015a, 2015b; Li and Li 2018). As such, PCFs created opportunities for speciation in otherwise physically stable landscapes in the short-term. Over the long-term, PCFs interacted with geology, resulting in intricate outcomes. Some of these constitute model systems to explore the consequences of PCFs and associated sea-level fluctuations on spatial biodiversity patterns (Weigelt et al. 2016). The Pleistocene Aggregate Island Complex (PAIC) model, for instance, describes the species pump hypothesis on islands, most notably in the Philippines, Caribbean and Mediterranean archipelagos (Esselstyn and Brown 2009; Brown et al. 2013; Papadopoulou and Knowles 2015a, 2015b). Understanding the impact of PCFs on diversity patterns is particularly challenging for the implementation of science-based conservation in the most diverse and threatened biotas globally *i.e.* biodiversity hotspots (Myers et al. 2000; Hoffmann et al. 2010).

The biodiversity hotspots in Southeast Asia occur in an area where Pleistocene eustatic fluctuations interact with geology (Voris 2000; Woodruff 2010; Lohman et al. 2011), including the landmasses of Sundaland (Java, Sumatra, Borneo and peninsular Malaysia). Sundaland emerged during the early Cenozoic (ca. 66 Ma) as a promontory at the southern end of Eurasia (Lohman et al. 2011). The isolation of Sundaland, however, is much more recent (Fig. 1). Complex tectonic movements and subduction activities during the Miocene triggered the formation of Borneo between 20 and 10 Ma (Figs. 1a & 1b) and the subsequent emergence of Sumatra and Java between 10 and 5 Ma (Figs. 1b & 1c). From a geological perspective, Sundaland's configuration has been stable for the past 5 Ma (Lohman et al. 2011; Hall 2013). However, insular Sundaland was not completely separated from mainland Southeast Asia until the Pliocene (5.33-2.58 Ma) (Hall 2009; Lohman et al. 2011). Upon

entering the early Quaternary (2.58 Ma), sea level dropped from 40 to 60 m below present levels over a ca. 41 kyr period (Lisiecki and Raymo 2005; Miller et al. 2005; Cannon et al. 2009; Lohman et al. 2011; de Bruyn et al. 2014) forming land bridges among the islands of Sundaland and the mainland (Fig. 1d, -60 m). Around 800 kyr, Sundaland experienced longer glacial cycles of 100 kyr and lower sea level during eustasy (Lisiecki and Raymo 2005; Miller et al. 2005) which enlarged land bridges and increased fragmentation during interglacial times. During the Last Glacial Maximum (LGM) ca. 17 kyr, sea levels dropped to 120 m below presents (Voris 2000; Sathiamurthy and Voris 2006) (Fig. 1d), resulting in the widest landmass extension to date. At the end of the LGM, rising sea levels shrank terrestrial habitats, and Sundaland biotas are currently considered to be refugial (Cannon et al. 2009; Woodruff 2010; Lohman et al. 2011). Interactions between the geological history of subduction and PCFs are expected to have significantly affected the diversification of terrestrial and freshwater biodiversity on Sundaland (Voris 2000; Sathiamurthy and Voris 2006; Lohman et al. 2011; de Bruyn et al. 2013).

Eustatic fluctuations in Sundaland not only altered land exposure over time but also impacted the configuration of Sundaland's watersheds. During glacial maxima, sea levels were especially low, allowing riverine watersheds to expand to newly exposed land that was previously sea floor (Voris 2000; Sathiamurthy and Voris 2006; Lohman et al. 2011). These temporary paleoriver systems connected Sundaland's rivers and created temporary linkages between Sundaland and continental Southeast Asia. Particularly well-documented for the Pleistocene LGM, the lower sea level created four large paleoriver systems that connected insular rivers across Sundaland's landmasses (Kottelat et al. 1993; Voris 2000; Woodruff 2010) including the paleorivers of East Sunda (1 in Fig. 1d), North Sunda (2 in Fig. 1d), Malacca Straits (3 in Fig. 1d) and Siam (4 in Fig. 1d). For freshwater-dependent organisms, the existence of these paleoriver systems has been long considered to be one of the main

drivers of contemporary distribution patterns of freshwater organismal in the region (Kottelat et al. 1993; Voris 2000; Lohman et al. 2011). Formally known as the Paleoriver Hypothesis, the impact of paleorivers on freshwater species distribution has been investigated in *Hemibagrus nemurus* (Dodson et al. 1995), viviparous halfbeaks (de Bruyn et al. 2013), snakeheads (Tan et al. 2012), killifishes (Beck et al. 2017) and *Macrobrachium* shrimps (De Bruyn et al. 2004). These studies detected some congruence between lineages distribution and boundaries of paleoriver watersheds. However, the determinants of these correlations are still debatable due to the limited number of taxa examined (Dodson et al. 1995; Lohman et al. 2011; Tan et al. 2012; de Bruyn et al. 2013; Beck et al. 2017). Furthermore, the ages of the taxa in these studies were generally older than the Pleistocene, casting doubt on the role of PCF in freshwater diversification (Dodson et al. 1995; Beck et al. 2017).

Here, we explore the evolutionary dynamics of species proliferation and spatial patterns of speciation among Southeast Asian freshwater fishes to address the following questions: (1) Did paleorivers fragment populations of widely distributed taxa? (2) Did paleorivers enable dispersal between islands during glacial maxima? (3) Did PCFs impact the pace of diversification?

From this set of initial questions, we derived several predictions made by the Paleoriver Hypothesis regarding the evolution of species range boundaries during speciation (*e.g.*, congruence between species range distribution and boundaries of paleoriver watersheds) and the timing of speciation (*e.g.* Pleistocene species pump) to address the evolutionary response of Southeast Asian freshwater biotas to eustatic fluctuations. To test predictions made by the Paleoriver Hypothesis, we compiled DNA sequences from widely distributed and well-studied freshwater fish lineages with a variety of life history traits (Kottelat et al. 1993). Our molecular phylogenetic dataset of 1,511 individuals was used to reconstruct the phylogenetic relationships and estimate the divergence times of 110 morphological species

belonging to four of the most species-rich freshwater fish lineages in Southeast Asia: *Channa* (30 morphological species), Zenarchopteridae (17 morphological species), *Glyptothorax* (50 morphological species), and *Clarias* (13 morphological species). Southeast Asian freshwater fishes are poorly known taxonomically and cryptic species are common (de Bruyn et al. 2013; Hubert et al. 2015; Lim et al. 2016; Beck et al. 2017; Nurul Farhana et al. 2018). Thus, all macroevolutionary inferences recognize Molecular Operational Taxonomic Units (MOTUs) delineated through DNA-based species delimitation methods (Blaxter et al. 2005; Ruane et al. 2014). Distribution ranges were characterized for these MOTUs and further used to explore spatial and temporal patterns of diversification through ancestral state reconstructions and birth-death modeling approaches.

MATERIALS AND METHODS

Hypothesis Testing, Taxa Selection and Sampling

The Paleoriver Hypothesis (Kottelat et al. 1993; Voris 2000; Woodruff 2010; de Bruyn et al. 2013) makes several predictions about the geography and timing of speciation. First, the paleorivers might be expected to promote allopatric speciation between populations of species that were once widespread or expanding. If so, range boundaries between sister species that diverged during the Pleistocene might be expected to coincide with boundaries of the paleoriver watersheds. Second, paleorivers might have enabled dispersal between different islands, such as South Borneo and Java in the East-Sunda system during periods of low sea level (Fig. 1d). This hypothesis predicts that species assemblages will be most similar within paleoriver drainage basins, even when those basins span two or more present-day islands. Assemblages between paleoriver drainages on the same present-day island are likely to be markedly different in comparison. Finally, the interactions between geological history and PCFs have intensified during the last 5 Ma with the elevation of Sundaland predicting

increased diversification. Estimating speciation events in different paleorivers would allow testing of these hypotheses. However, different species interact with their environment in different ways, which will further influence their dispersal and colonization abilities. Thus, dispersal traits might be expected to influence the evolutionary responses of biotas to environmental changes (Barracough and Nee 2001; Currie et al. 2004; McPeck 2008; Hubert et al. 2015; Burbrink et al. 2016; Liu et al. 2019). To explore the three predictions made by the Paleoriver Hypothesis, we selected taxa conforming to the following requirements: (1) widespread distribution of lineages across Southeast Asia including Sundaland, (2) comprehensive sampling of described species within the taxon, (3) availability of nuclear and mitochondrial sequences, and (4) varying life history traits such as body size and dispersal ability.

Four lineages fulfilled these requirements: (1) the snakehead genus *Channa* (Anabantiformes, Channidae) widely distributed in Asia (Conte-Grand et al. 2017), (2) the walking catfish genus *Clarias* (Siluriformes, Clariidae) widely distributed in Asia (Pouyaud et al. 2009), (3) the catfish genus *Glyptothorax* (Siluriformes, Sisoridae) widely distributed in Asia (Kottelat et al. 1993; Jiang et al. 2011) and, (4) the halfbeak genera of Southeast Asia *Dermogenys*, *Hemirhamphodon* and *Nomorhamphus* (Beloniformes, Zenarchopteridae) (de Bruyn et al. 2013; Lim et al. 2016; Nurul Farhana et al. 2018). Both *Clarias* and *Channa* can disperse easily. *Clarias* species, for instance, have a unique accessory air-breathing organ in the upper part of the branchial cavity that allows survival in oxygen-poor water and on land (Munshi 1961; Maina and Maloiy 1986). Meanwhile, *Channa* is a genus of air-breathing, predatory freshwater fishes with an extended swim bladder, paired supra-branchial chambers, and an interior labyrinth organ that enables them to live out of the water for several days. Several species can move on land and burrow into mud during droughts to keep their bodies wet (Kottelat et al. 1993; Berra 2001; Adamson et al. 2010; Serrao et al. 2014; Rüber et al.

201 2019). These capabilities make *Clarias* and *Channa* resilient to environmental variation, and
202 several species are even considered invasive outside their native range (Serrao et al. 2014;
203 Conte-Grand et al. 2017). The other genera have lower dispersal abilities. *Glyptothorax*, the
204 most species-rich genus in the catfish family Sisoridae, is smaller than *Clarias* and *Channa*,
205 and is not equipped with an air-breathing organ (Berra 2001). Distribution of *Glyptothorax*
206 species are mostly fragmented and confined within isolated mountain rapid streams (Ng and
207 Kottelat 2016; Hutama et al. 2017). The healfbeak genera are viviparous with limited larval
208 dispersal abilities (de Bruyn et al. 2013; Nurul Farhana et al. 2018). Although these genera are
209 geographically widespread, their genetic lineages have more restricted distributions (Downing
210 Meisner 2001; Tan and Lim 2013; Lim et al. 2016; Nurul Farhana et al. 2018). Our dataset
211 comprised 2,211 sequences dataset from 1,511 individuals belonging to 110 nominal species
212 and representing four lineages. Outgroups were then selected for each group, ranging from
213 closely related to distantly related taxa following the phylogenetic classification of bony
214 fishes by Betancur-R et al. (2017). Outgroup selection was further refined according to
215 previously published molecular studies for Clariidae (Pouyaud et al. 2009), Sisoridae (Jiang et
216 al. 2011) and Channidae (Conte-Grand et al. 2017).

218 ***Genetic Species Delimitation***

219 Genetic studies have identified substantial cryptic diversity in Southeast Asian
220 freshwater fishes (Nguyen et al. 2008; Pouyaud et al. 2009; Hubert et al. 2015, 2019;
221 Dahruddin et al. 2017; Hutama et al. 2017; Nurul Farhana et al. 2018; Sholihah et al. 2020).
222 Cryptic diversity could lead to biases in phylogenetic reconstruction and diversification
223 analyses, particularly due to the overestimation of divergence age estimates in the absence of
224 the closest relatives (Esselstyn et al. 2009; Patel et al. 2011; Ruane et al. 2014). To avoid
225 these shortcomings, genetic species delimitation methods were used to define MOTUs.

Several methods of species delineation have been developed, but each has pitfalls. Agreement among different methods suggest robust delimitation (Kekkonen and Hebert 2014; Kekkonen et al. 2015). Thus, we used four methods of species delimitation including a distance-based method with Automatic Barcode Gap Discovery (ABGD, Puillandre et al. 2012), a network-based method with Refined Single Linkage (RESL, Ratnasingham and Hebert 2013), and two tree-based methods including Poisson Tree Processes (PTP, Zhang et al. 2013) and the Generalized Mixed Yule Coalescent (GMYC, Pons et al. 2006). A final delimitation scheme was then established based on a 50% consensus (Hubert et al. 2019). All sequences used were aligned using MUSCLE (Edgar 2004) implemented in MEGA7 (Kumar et al. 2016) and manually edited.

These four delimitation methods use different input data. We used sequence alignments to carry out ABGD and RESL analyses. The ABGD analysis was performed through the online platform using the K2P substitution model (<https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html>). The RESL algorithm was implemented through BOLD systems version 4 (<http://www.boldsystems.org>, Ratnasingham and Hebert 2007), using Barcode Index Numbers (BIN, Ratnasingham and Hebert 2013). We used maximum likelihood (ML) phylogenetic trees for the PTP and GMYC analyses. For PTP, the analyses were carried out on the web service (<https://mptp.h-its.org/#/tree>). Only the single threshold version was used because preliminary analyses using the multiple threshold version of PTP were too conservative compared to other methods (*i.e.* too few species). Maximum Likelihood phylogenies were reconstructed using RAxML (GUI 1.5) with the GTR+I+ Γ model, and 5000 non-parametric bootstrap (BP) replicates were computed using RAxML-HPC Blackbox (Miller et al. 2010) with RAxML 8 (Stamatakis 2014). For GMYC, the multiple thresholds function (mGMYC) was implemented with the R-package SPLITS 1.0-19 (Fujisawa and Barraclough 2013). For GMYC, ultrametric trees were reconstructed

using the *chronopl* function in the R-package *ape* 5.3 (Paradis et al. 2004; Paradis 2012; Paradis and Schliep 2019). Time calibration was constrained with the widely accepted fish substitution rate of 1.2% of genetic divergence per million years (Myr) for mitochondrial protein-coding genes (Knowlton et al. 1992) and applied it to the maximum K2P distances, computed using mitochondrial protein coding-genes only, between descendant clades of selected nodes (Table 1). Reference nodes were selected to cover several levels of depth in the phylogenies and to cover a substantial portion of the total species diversity in the entire mitochondrial data set. A total of 13 nominal species were analyzed for *Clarias*, 50 nominal species for *Channa*, 30 nominal species for Zenarchopteridae and 17 nominal species for *Glyptothorax*.

Phylogenetic Reconstruction and Dating

Bayesian phylogenetic reconstructions were implemented with StarBEAST2 package (Ogilvie et al. 2017) of the BEAST2 suite (Heled and Drummond 2010; Bouckaert et al. 2014). This approach implements a mixed-model including a coalescent component within species and a diversification component between species that allows accounting for variations of substitution rates within and between species (Ho and Larson 2006). StarBEAST2 requires the designation of species, which were determined using the consensus of our species delimitation analyses. Best substitution models for each marker were estimated using jModelTest2 0.1.10 (Guindon and Gascuel 2003; Darriba et al. 2015). Preliminary analyses conducted on Zenarchopteridae indicated that estimating substitution model parameters jointly with node ages and topologies led to non-converging MCMC with widely fluctuating ages of the Most Recent Common Ancestor (MRCA). Hence, ML parameter estimates were used for MCMC searches in StarBEAST2 with no further estimation of the substitution model parameters. This analytical choice constrained the number of substitution models available in

StarBEAST2 and if the selected model was not supported in StarBEAST2, the next most likely model from jModelTest2 available in StarBEAST2 was used. An uncorrelated lognormal clock model was applied alongside estimation of clock rates for all analyses. The reference nodes calibrated for the species delimitation analyses were also used in the Bayesian analyses (Table 1) and further used with a calibrated Yule model. Age calibration was added under normal distribution priors, using a sigma of 1 in order to use wide distributions (Table 1). Yule birthrate value for the analysis was generated using the R-package *ape* by applying a Yule function on each ultrametric tree generated using the *chronopl* function for the GMYC delimitation. To yield enough statistical power (ESS values reaching >200), multiple MCMC runs with 10 million pre-burnin generations each were generated in parallel with tree states stored every 5,000 generations. Considering the number of individuals sampled and molecular heterogeneity within each taxon, the length of the MCMC was different for each group, ranging from 100 million for *Glyptothorax* to 250 million for *Channa*. The MCMC were run at least twice to check for the convergence of the chains and reach ESS values > 200 for MRCA age and parameters of the diversification and clock models. The statistics were visualized using Tracer 1.7.1 (Rambaut et al. 2018) to check the outputs and to determine the best converging combination of burnin for each StarBEAST2 run. Independent runs were then combined to a single concatenated dataset using LogCombiner 1.10.4 (Bouckaert et al. 2014) to compute the final Bayesian statistics and sampled trees, with or without resampling (depending on the chain lengths). The maximum clade credibility tree, age estimates and corresponding 95% highest posterior density (HPD) were summarized using TreeAnnotator 1.10.4 (Bouckaert et al. 2014).

Molecular dating constitutes a crucial step in macroevolutionary inferences (Condamine et al. 2015a) and divergence time estimates can widely vary according to calibration methods and clock models (Ho and Larson 2006; Duchêne et al. 2014). We

estimated the impact of clock rates on MOTUs age estimates and diversification trends. For each clade, we rescaled the StarBEAST2 maximum credibility tree according to an assortment of molecular clocks ranging from 0.4% to 1.6% per Ma, with an increment of 0.2%, to cover a wide range of known mitochondrial substitution rates (Orti Meyer, A. 1997; Hardman and Lundberg 2006; Read et al. 2006; Kadarusman et al. 2012). StarBEAST2 trees were rescaled using the Alter/Transform Branch Lengths function in Mesquite 3.61 (Maddison and Maddison 2019) resulting in a total of seven trees computed for 0.4, 0.6, 0.8, 1.0, 1.2, 1.4 and 1.6% per Myr clocks for each group.

Diversification Rates Estimations

We first plotted lineages through time (LTT) for each taxon using the R-package *ape* (Paradis and Schliep 2019) with confidence intervals computed with 1,000 random trees sampled from the StarBEAST2 MCMC file (Heled and Drummond 2010; Bouckaert et al. 2014). Then, to test the hypothesis that speciation was linked to past environmental changes (Condamine et al. 2013, 2019), we designed a ML framework including five types of diversification models: constant-rate, time-dependent, temperature-dependent, sea-level-dependent, and diversity-dependent models. These models rely on the ML framework originally developed by Morlon et al. (2011) and implemented in the R-package *RPANDA* 1.3 (Morlon et al. 2016). We accounted for missing species in the phylogeny by using a global sampling fraction that is the ratio of sampled species diversity over the total described species diversity for each clade. We designed and fit a total of 17 diversification models (Table S1). We first fit two constant-rate models, considered as references for model comparison: one with the speciation rate constant through time with no extinction (BCST) and another with speciation and extinction rates constant through time (BCSTDCST). Second, we fit four time-dependent models: a model with only speciation rate varying through time and no extinction

(BtimeVar), a model with speciation rate varying through time and constant extinction
 (BtimeVarDCST), a model with constant speciation and extinction rate varying through time
 (BCSTDtimeVar), and a model with speciation rate and extinction rate both varying through
 time (BtimeVarDtimeVar). We then fit eight models with speciation and extinction rates
 varying according to an external environmental variable of which four models had the
 temperature curve, and four had the sea level curve as follows: two models with speciation
 varying in function of the environmental variable (BtemperatureVar and Bsea-levelVar), two
 models with speciation varying in function of the variable with constant extinction rate
 (BtemperatureVarDCST and Bsea-levelVarDCST), four models with extinction rate varying
 as a function of the variable and constant speciation rate (BCSTDtemperatureVar and
 BCSTDsea-levelVar), and two models with both speciation and extinction rates varying as a
 function of the variable (BtemperatureVarDtemperatureVar and Bsea-levelVarDsea-
 levelVar). All diversification analyses were performed independently for each the four fish
 clades.

We chose an exponential dependence with time, temperature or sea level, and
 diversification rates because this exponential function is robust and can take a broad range of
 shapes depending on the strength and direction of the dependence to the fitted variable.
 Speciation rate (λ) and extinction rate (μ) are parameterized by the set of following
 parameters: When speciation and extinction rates are exponential functions of the sea level
 through time, we used the equations $\lambda(S_{(t)}) = \lambda_0 \times e^{\alpha S_{(t)}}$ and $\mu(S_{(t)}) = \mu_0 \times e^{\beta S_{(t)}}$, where λ_0
 and μ_0 are respectively the speciation rate and the extinction rate expected when sea level is at
 0 meters. The variables α (and β) are coefficients that measure the strength and sign of the
 relationship with sea level (*e.g.* $\alpha > 0$ and $\beta > 0$ indicate that speciation and extinction rates
 increase with sea level high stands). Similar equations can be written for an exponential
 relationship between temperature through time and speciation and extinction rates: $\lambda(T_{(t)}) =$

$\lambda_0 + \alpha T(t)$ or $\lambda(T(t)) = \lambda_0 \times e^{\alpha T(t)}$, where $T(t)$ is the temperature at time t and λ_0 is the speciation rate at the temperature of 0° C.

In addition, three diversity-dependent diversification models were fitted using a maximum likelihood function with the R-package *DDD* 3.7 (Etienne et al. 2012). In the diversity-dependent models, speciation rates or extinction rates vary as a function of the number of lineages in the clade (Etienne et al. 2012). We took this function to be linear as explained in Etienne et al. (2012). The diversity-dependent models are parameterized by λ_0 and μ_0 , the speciation and extinction rates in the absence of a competing lineage, and K that is the ‘carrying capacity’, and represents asymptotic clade size. All speciation and extinction rates were constrained to be positive. The models were compared with the corrected Akaike Information Criterion (AICc) and Akaike weight (AIC ω). The model with the lowest AICc and highest AIC ω was considered to be the best fitting model for the phylogeny. This model selection analysis was repeated for each of the seven dated trees based on different clock rate hypotheses (0.4% to 1.6% per Ma) for the groups departing from a constant diversification model and related to sea level or temperature.

Ancestral States Estimations

To infer the influence of Sundaland paleoriver and island systems on the diversification of Southeast Asian freshwater fishes, we estimated the ancestral areas of origin for each clade using the species trees obtained from StarBEAST2. Ancestral area estimations were performed with the R-package *BioGeoBEARS* 1.1 (Matzke 2013) using two sets of geographic delimitations based on: (1) paleorivers, and (2) islands. Then, geographic patterns of speciation were recorded as follows: (1) no dispersal, sister species co-occur within the same paleoriver and the same island; (2) dispersal between islands within a paleoriver forming sister species on different islands within the same paleoriver; (3) dispersal between

376 paleorivers within the same island forming sister species on different paleorivers within the
 377 same islands; and (4) dispersal between islands and between paleorivers forming sister
 378 species on different paleorivers and different islands. Ancestral area estimations involving
 379 paleorivers were based on the following geographic areas (Fig. 1): (1) the North Sunda river
 380 system, (2) the East Sunda river system, (3) the Malacca Straits river system, (4) the Siam
 381 river system, (5) the Bangka-Belitung, (6) the China-South Asia, (7) the Mekong and
 382 Irrawaddy river system, (8) the Northern Borneo and (9) the Philippines. Ancestral area
 383 estimations involving islands were based on the following areas: (1) Borneo, (2) Java-Bali,
 384 (3) Sumatra, (4) Bangka-Belitung, (5) China-South Asia, (6) Mainland Southeast Asia, (7)
 385 Philippines, and (8) Sulawesi. The occurrence of each MOTU in the areas were compiled in a
 386 presence/absence matrix for each taxon and served as input data together with the species
 387 tree. The range distribution of each MOTU from both approaches is provided in Table S2.
 388 Ancestral area estimations were then carried out using 6 models: Dispersal-Extinction
 389 Cladogenesis (DEC) and DEC+J (Matzke 2014); ML version of Dispersal-Vicariance
 390 analysis (DIVALIKE) and DIVALIKE+J; Bayesian biogeographical inference model
 391 (BAYAREALIKE) and BAYAREALIKE+J (Van Dam and Matzke 2016). The inclusion of
 392 the parameter J has been recently criticized from a conceptual and statistical perspective (Ree
 393 and Sanmartín 2018). The formalization of founder-effect speciation as “jump dispersal” has
 394 been developed for insular systems to account for divergence of a new lineage in a new
 395 geographic area established by colonization without an intermediate widespread ancestor
 396 (Clark et al. 2008; Ree and Sanmartín 2018). Considering the biogeographic scenario of
 397 Sundaland and the insularity of the system, jumping dispersal cannot be discarded *a priori*
 398 from a conceptual perspective and several studies have previously highlighted the importance
 399 of founder-effect speciation in insular systems (de Bruyn et al. 2013; Condamine et al. 2015b;
 400 Beck et al. 2017). In fact, the Paleoriver Hypothesis predicts that sea level low stands enabled

jump dispersal among islands through temporary land bridges. Models of ancestral area estimation including the J parameters were thus considered here and the best-fit model was estimated using the AICc.

RESULTS

Phylogenetic Reconstruction and Species Delimitation

A total of 2,211 sequences from 1,511 individuals were mined from *GenBank* and BOLD, representing 110 nominal species (Table S2). The alignment for the genus *Clarias* consisted of 5,322 base pairs (bp), including 3,093 bp from the mitochondrial genome (16S, COI and Cytb) and 2,229 bp from the nuclear genome (RAG1, RAG2), for 146 individuals belonging to 13 morphological species. The alignment for the genus *Channa* comprised of 2,856 bp, including 2,076 bp from the mitochondrial genome (16S, COI, Cytb) and 780 bp from the nuclear genome (RAG1), for 891 individuals belonging to 30 morphological species. The alignment for the genera *Dermogenys*, *Hemirhamphodon* and *Nomorhamphus* consisted of 4,109 bp, including 984 bp from the mitochondrial genome (COI, Control Region) and 3125 bp from the nuclear genomes (DP5, DP14, DP21, DP35, DP37, HP5, HP54, HPR56), for 266 individuals belonging to 17 morphological species. The alignment for the genus *Glyptothorax* data consisted of 2,776 bp, including 1,869 bp for the mitochondrial genome (COI, Cytb) and 907 bp for the nuclear genome (RAG2), for 208 individuals belonging to 50 morphological species. All alignments are available in TreeBASE (TB2:S26912).

Substitution models for each marker are provided in Table S3 (Online supplementary material). The four ML phylogenetic reconstructions generally provided well-supported clades with most internal nodes supported by BP > 80 except within the genus *Channa* (Fig. 2, Fig. S1). For the genus *Clarias*, Asian species constitute a monophyletic group (Fig. 2A, clades II + III) separated from the African species (Fig. 2A, clade I), with *Clarias gariepinus*

(Burchell 1822) as sister to all remaining species. All species are monophyletic except *C. nieuhofii* Valenciennes 1840, which is recovered as polyphyletic with three distinct lineages (Fig. S1). For the genus *Glyptothorax*, the inferred tree is imbalanced with continental species constituting a stem group at the root of the tree (Fig. 2B, Fig. S1) and Sundaland species constituting a monophyletic group (Fig. 2B, clade I). All species are monophyletic, except *G. platypogonides* (Bleeker 1855) and *G. fuscus* Fowler 1934, which are polyphyletic (Fig. S1). The monophyly of the Zenarchopterid genera *Hemirhamphodon* (Fig. 2C, clade I) *Nomorhamphus* (Fig. 2C, clade II), and *Dermogenys* (Fig. 2C, clade III) is recovered with a sister-relationship between *Dermogenys* and *Nomorhamphus* (Fig. 2C, clades II + III). All species are monophyletic, except *Nomorhamphus megarrhamphus* (Brembach 1982), which is paraphyletic (Fig. S1). For the genus *Channa*, internal relationships are poorly supported, but four main clades are observed (Fig. 2D, clades I to IV). All *Channa* species are monophyletic, except *Channa gachua* (Hamilton 1822), which is polyphyletic and includes at least three lineages distinctly associated to other species, and *C. lucius* (Cuvier 1831) and *C. marulius* (Hamilton 1822), which are paraphyletic (Fig. S1).

As expected, different species delimitation analyses yielded variable number of MOTUs (Table S2): (1) 47 *Clarias* using mGMYC, 32 using PTP, 7 using RESL and 24 using ABGD, resulting in a consensus of 29 MOTUs, (2) 107 *Glyptothorax* using mGMYC, 58 using PTP, 58 using RESL and 72 using ABGD, resulting in a consensus of 61 MOTUs, (3) 48 Zenarchopteridae using mGMYC, 27 using PTP, 35 using RESL and 49 using ABGD, resulting in a consensus of 43 MOTUs, and (4) 521 *Channa* using mGMYC, 120 OTUs using PTP, 57 using RESL and 133 using ABGD resulting in a consensus of 120 MOTUs. The consensus delimitation scheme resulted in between 1.22 (*Glyptothorax*) and 4.00 (*Channa*) more MOTUs than morphological species (Table S2).

The final Bayesian analyses as implemented in StarBEAST2 were performed using an MCMC of length required to reach convergence ($ESS > 200$). For *Clarias*, two parallel runs were run with a burnin of 65.01 million generations and 15 million generations, respectively, and both runs were concatenated into a 220-million-generation dataset of 44,000 sampled trees. For *Glyptothorax*, three parallel runs were launched with burnins of 5, 10 and 5 million generations, respectively, and then concatenated into a 280-million-generation dataset with 56,000 sampled trees. For *Channa*, four parallel runs were used with burnins of 182.5, 27.52, 47 and 43 million generations, respectively, and concatenated into a 700-million-generation dataset with 35,000 sampled trees. For Zenarchopteridae, four parallel runs were used with burnin of 70, 50, 38 and 60 million generations, respectively, and concatenated into a 182-million-generation dataset with 18,200 sampled trees. The StarBEAST2 chronograms were largely congruent with the topologies estimated with the ML approach (Fig. S2). The ages of the MRCA vary between clades with a MRCA dated at 7.09 Ma (95% HPD = 4.49-10.31 Ma,) for *Clarias*, 7.1 Ma (95% HPD = 4.51-9.54 Ma) for *Glyptothorax*, 12.63 Ma (95% HPD = 8.24-18.68 Ma) for Zenarchopteridae, and 14.35 Ma (95% HPD = 10.54-18.41 Ma) for *Channa* (Fig S2). For *Clarias*, the MRCA of the Asian species is dated around 6.37 Ma (95% HPD = 3.99-9.19 Ma) and the Zenarchopteridae genera are dated around 9.88 Ma (95% HPD = 6.02-14.70 Ma) for *Hemirhamphodon*, 5.88 Ma (95% HPD = 4.02-8.09 Ma) for *Dermogenys*, and 3.90 Ma (95% HPD = 2.14-5.94 Ma) for *Nomorhamphus*.

Diversification Rates

Most of the MOTUs delineated in the consensus scheme originated in the Pleistocene with divergence time estimates younger than 2.5 Myr (Fig. 3). More than half (53%) of the morphological species and 88% MOTUs are younger than 2.5 Myr (Fig. 3A). The LTT plots indicate that 84% of extant fish diversity occurred during the Pleistocene. On Sundaland,

Pleistocene speciation peaks during sea level low stands with 88% of speciation events inferred during the last 2.5 million years and 76% during the last 1.5 million years. This peak of Pleistocene speciation is observed for a wide range of clock rates, including some of the slowest clock rate hypothesis with 72% and 57% of speciation events during the last 2.5 and 1.5 Myr, respectively, for the 0.6% per million years (Fig. S3). Faster rates yielded a higher percentage of young MOTUs. The 1.6% per Myr rate yielded 93% and 83% of MOTUs younger than 2.5 and 1.5 Myr, respectively. The only exception was observed for the 0.4% per million years with 60% and 38% of the MOTUs younger than 2.5 and 1.5 Myr, respectively.

Likelihood scores and AIC_w for 17 diversification models indicate clade-specific patterns of diversification (Table S4). For *Clarias* and *Glyptothorax*, the constant-rate speciation model is the most likely according to AIC_w, respectively 0.267 and 0.212, with high speciation rates (λ) of 0.4556 and 0.5444, respectively. For Zenarchopteridae, three models are equally likely, including: (1) the time-dependent speciation model (AIC_w = 0.145, λ_0 = 0.432) with a speciation increasing through time (α = -0.0971); (2) the constant-rate speciation model (AIC_w = 0.119, λ_0 = 0.3316); and (3) the temperature-dependent speciation model (AIC_w = 0.116, λ_0 = 0.5132) with speciation increasing as temperatures cooled (α = -0.1444). While diversification patterns of the previous three groups show no dependency on sea-level eustasy, the most likely model for *Channa* diversification through time is the sea-level-dependent speciation model (AIC_w = 0.415, λ_0 = 0.2279) with speciation increasing as sea level dropped (α = -0.0182). These clade-dependent diversification patterns are reflected by the heterogeneous trends of speciation rates through time for the four lineages (Fig. 3B), including constant speciation rates through time for *Glyptothorax* and *Clarias*, a sea-level dependent speciation rate for *Channa*, and a time-dependent speciation rate for Zenarchopteridae. The dependency of *Channa* diversification on sea level eustasy was

recovered for all clock rates except for the two slowest clocks (Table S5). The most likely model for the 0.4% per Myr rate includes a positive relationship between speciation and sea level, speciation increased as sea level increased ($\alpha = 0.0119$). The most likely model for the 0.6% per Myr clock includes a positive relationship between extinction rates and sea levels eustasy, extinction increased when sea level increased ($\beta = 0.0351$).

Geography of Diversification

Ancestral area estimations with models incorporating the J parameter were more likely for all groups on islands or paleorivers than models lacking that parameter (Table S6). The DEC+J model is the most likely and was further used in subsequent analyses. Most colonization was initiated before the Pleistocene and in Borneo (the North Sunda and East Sunda paleoriver systems, Fig. 4), while several subsequent colonizations happened through Sumatra, and Java is typically the last island colonized. This pattern was observed within *Clarias*, with a first vicariance event inferred between Borneo and the Southeast Asian mainland at 6.37 Ma (95% HPD = 3.99-9.19 Ma) and associated with the first split in the Asian *Clarias* (Fig. 4A, clade II vs clade III), the colonization of Sumatra happening subsequently twice from Borneo (Fig. 4A, clade III). The same pattern is observed for *Glyptothorax* with a vicariance inferred between Borneo and mainland around 3.64 Ma (95% HPD = 2.34-5.17 Ma; Fig. 4B, clade I). Different patterns are observed for Zenarchopteridae (Fig. 4C) and *Channa* (Fig. 4D), indicating that the islands of Sundaland were colonized multiple times.

Most speciation events are inferred to happen within islands, as exemplified by the diversification of three of the four lineages (*Clarias*, *Glyptothorax* and Zenarchopteridae) in Borneo and Sumatra (Figs. 4A, 4B and 4C) or Java for *Channa* (Fig. 4D). The same trend is observed for paleorivers; most speciation events are inferred to occur within them (Fig. 4).

Four major geographic patterns of speciation were identified: (1) sister species occupy the same paleoriver on the same island, *e.g.* *Glyptothorax platypogon* 1-4 from Java (East Sunda River System), (2) sister species occupy the same paleoriver on different islands, *e.g.* *Glyptothorax robustus* (Boeseman 1966) 1-4 from Java and Sumatra (East Sunda River System), (3) sister species occupy different paleorivers within the same island, as in *G. amnestus* (North Sunda) and *G. fuscus* 4 (Malacca Strait), and (4) sister species occupy different islands and different paleorivers *e.g.* *Clarias pseudoleiacanthus* Sudarto, Teugels and Pouyaud 2003 (North Sunda in Borneo) and *C. leiakanthus* Bleeker 1851 (Malacca Strait in Sumatra).

In total, 66.3 % of the inferred speciation events involve dispersal either between islands or between paleorivers (Table 2). In Sundaland, 59.2% of speciation events occurred within island (40.8% between islands) while 39.8% of speciation events occurred within paleorivers (60.2% between paleorivers). These trends are observed for all groups except Zenarchopteridae, in which speciation events within islands represent 81.7% and speciation within paleorivers 63.6% (Table 2). The proportion of speciation events within or between islands is stable through time for speciation resulting from dispersal between paleorivers (Fig. 5A). However, this pattern was not observed for Zenarchopteridae (Fig. 5B), where speciation was primarily between islands. In *Glyptothorax*, most speciation events between paleorivers involve dispersal between islands (Fig. 5E). Within paleorivers, speciation events involving dispersal between islands are scarce for all groups (Figs. 5 B, C, D, E).

DISCUSSION

Diversification of Southeast Asian Aquatic Biotas

The "Late Pleistocene Hypothesis" (Barracclough and Nee 2001; Wiens and Donoghue 2004; Mittelbach et al. 2007; Beck et al. 2017) and the "Pleistocene Species Pump

Hypothesis" (Esselstyn and Brown 2009; Brown et al. 2013; Papadopoulou and Knowles 2015a, 2015b; Li and Li 2018) suggest that sea-level fluctuations accelerated diversification of insular biotas. However, we found poor support for these hypotheses and signature of sea-level fluctuations on diversification was apparent in only one of the four fish groups studied here. The genus *Channa* is the only group in which diversification rate was associated with sea-level fluctuations: its speciation rate increased as sea level dropped. This translates into increasing speciation rates through time since sea levels generally decreased through the Pleistocene (Miller et al. 2005). Hence, the genus *Channa* had the highest speciation rate of all studied taxa (Fig. 3B). The superior dispersal ability and environmental adaptability of *Channa* compared to the other taxa may enable them to colonize many different types of environments. With the closure of dispersal routes between islands during sea level highstands, gene flow was reduced, thereby increasing the probability of divergence between populations, eventually leading to speciation. Such speciation events likely occurred repeatedly over short time intervals (glaciation periods). Hence, the genus *Channa* is the only taxon we studied that supports the "Pleistocene Species Pump" hypothesis.

On the contrary, the diversification of the other clades does not support this hypothesis. Both silurid groups (*Clarias* and *Glyptothorax*) diversified at a constant rate despite different life history strategies. *Glyptothorax* had a higher speciation rate than *Clarias*, which might be related to the tendency of *Glyptothorax* to occupy hilly streams that are more isolated (Hutama et al. 2017). The dispersal abilities of *Clarias* likely limited allopatric speciation. Such habitat specificity for *Glyptothorax* lineages could have (1) inhibited inter-island dispersal during low seastand since the freshwater corridors between islands were unsuitable, as well as (2) facilitated isolation and *in situ* diversification due to restricted gene flow (Hubert et al. 2015; Hutama et al. 2017), hence generating high, constant speciation. In

the same way, none of the diversification trends for Zenarchopteridae were directly related to sea-level fluctuations.

Diversification trends show no tendency to decline following a diversity-dependent diversification (DDD) model (Seehausen 2007; Rabosky and Lovette 2008; Seehausen et al. 2008). The DDD model is based on the assumption that biotas are bounded by the carrying capacities of the environment, either in terms of habitat diversity (Kisel et al., 2011; Phillimore & Price, 2008; Schluter, 2000) or the total number of individuals that can be sustained (Alonso et al., 2006; Hubbell, 2001). This model predicts that diversification rates decelerate through time as a consequence of (1) declining speciation rates due to the increasing occupation of available niches by proliferating species (Schluter 2000; Gavrillets and Vose 2006; Hubert et al. 2015), or (2) increasing extinction rate through time over constant speciation rate as a consequence of the increasing importance of stochastic demographic dynamics within proliferating species, which are presumed to have reduced population sizes compared to their parental species (Alonso et al. 2006; Rabosky and Lovette 2008; Hubert et al. 2015). Here, however, we found no trend for declining diversification rates in any taxa. Two of the four clades have rates that increase through time (Fig. 3B). The high diversity of freshwater habitats in Sundaland likely provides a diversity of niches to sustain elevated and/or constant diversification. Water chemistry is widely diverse in tropical systems (Guyot 1993) and Sundaland is no exception, with water types ranging from clear and alkaline waters in karsts (Clements et al. 2006) and turbid sediment-rich waters of the floodplains to the humic acid-rich waters of the peat swamps (Ng et al. 1994; Giam et al. 2012). The large number of fish MOTUs distributed among multiple paleorivers and sister MOTUs diversifying within the same paleoriver seems consistent with the existence of such high diversity of habitat/niche diversity within paleorivers. However, documented examples of ecological speciation within freshwater rivers are scarce (Sullivan et al. 2002; Nolte et al.

2005; Alter et al. 2017). Most cases have been discovered in ancient lakes or other lentic water bodies (Nagel and Schluter 1998; Verheyen et al. 2003; Barluenga and Meyer 2004; Herder et al. 2006; Landry et al. 2007). Zenarchopteridae and *Clarias* species are distributed across a wide range of aquatic habitats, with some *Clarias* and *Hemirhamphodon* species occupying acidic waters of peat swamps while other *Clarias* and *Dermogenys* species occupy clear waters and floodplains (Pouyaud et al. 2009; Lim et al. 2016; Nurul Farhana et al. 2018). However, inspection of the phylogenies reconstructed here suggests that transitions between different types of aquatic habitats are scarce; peat swamp and floodplain-associated species of *Clarias* grouped in distinct clades. This trend suggests that adaptive divergence across freshwater habitats is limited in the groups studied here. By contrast, landscape fragmentation by geology has been widely document in the area (Nguyen et al. 2008; Lim et al. 2016; Dahrudin et al. 2017; Hutama et al. 2017; Nurul Farhana et al. 2018; Hubert et al. 2019).

Habitat diversity within paleorivers might be expected to have interacted with the availability of dispersal routes among paleorivers, which was affected by sea-level fluctuations (Brown et al. 2013). Similar trends have also been reported for the Pleistocene Aggregate Island Complex (PAIC) model in the Philippines archipelago. Brown et al (Brown et al. 2013) suggested an alternative "nested PAIC model" considering the interplay of the external sea level fluctuations causing cycles of island connection-isolation and changing local habitat diversity within an island. Due to both its paleoriver and archipelagic terrain, Sundaland is likely comparable to the Philippines with extensive and heterogeneous freshwater habitats (Brown et al., 2013; Esselstyn & Brown, 2009; Papadopoulou & Knowles, 2015b, 2015a). Frequent regional intermediate disturbance associated with PCF likely impacted biotic interactions among Southeast Asian freshwater fish species and probably triggered a reshuffling of their ecological communities, which may explain the higher species richness than expected under a DDD model.

Diversification Mostly Occurred Within Islands and Paleorivers

Our ancestral area estimations demonstrate that diversification frequently resulted from long-distance dispersal followed by *in situ* diversification (founder-effect speciation) as suggested by the high likelihood of the models with the J parameter (Matzke 2014). This pattern has been previously proposed through detection of trans-island dispersal of freshwater fishes (Dodson et al. 1995; Pouyaud et al. 2009; Adamson et al. 2010; Tan et al. 2012; Tan and Lim 2013; de Bruyn et al. 2013; Hubert et al. 2015, 2019; Lim et al. 2016; Beck et al. 2017; Dahrudin et al. 2017; Hutama et al. 2017; Nurul Farhana et al. 2018). The establishment of taxa from mainland Asia during the Miocene and Pliocene supports the pre-Pleistocene colonization hypothesis (Dodson et al. 1995; de Bruyn et al. 2013; Hendriks et al. 2019). These results agree that regional shallow seas only started to dry out during the Pliocene, offering the possibility for dispersal of freshwater ancestors from mainland Asia. During these periods, Borneo was certainly more connected to mainland Southeast Asia than Sumatra and Java (Figs. 1B & 1C), and Borneo probably played a role in the initial diversity build-up of Sundaland biotas (de Bruyn et al. 2013, 2014). From our inferences, Sumatra and Java were first colonized from Borneo around 2.96 Ma via the North Sunda river system and around 1.18 Ma via the East Sunda river system, respectively. Thus, our study reinforces the conclusion of De Bruyn et al. (2014) that Borneo was the origin of most insular Sundaland freshwater fish diversity.

Although the fish lineages we studied were present in Sundaland before the Pleistocene, most of the MOTUs delimited here result from Pleistocene diversification events that are expected to be affected by PCF (Esselstyn et al. 2009). Looking at the geography of speciation for each group, we estimated that speciation of *Clarias* and *Channa* involved more dispersal events compared to *Glyptothorax* and Zenarchopteridae, probably due to higher

649 dispersal ability (Kottelat et al. 1993; Berra 2001; Downing Meisner 2001; Pouyaud et al.
650 2009; Adamson et al. 2010; Jiang et al. 2011; Tan and Lim 2013; Serrao et al. 2014; Lim et
651 al. 2016; Ng and Kottelat 2016; Conte-Grand et al. 2017; Hutama et al. 2017; Nurul Farhana
652 et al. 2018). The dispersal patterns and absence of decline in the diversification rates suggest
653 that land bridges and the subsequent Pleistocene eustatic fluctuations drove further *in situ*
654 diversification with the interplay of the insular and paleoriver watersheds boundaries, as well
655 as habitat heterogeneity, which likely conditioned dispersal. Episodes of sea level rise during
656 interglacial periods might be expected to have fragmented Sundaland. Rising seas created
657 refugial insular areas in Borneo, Sumatra and Java. These saltwater barriers may have driven
658 further within-island diversification, explaining the numerous *in situ* diversification events
659 after long-distance dispersals in each of the four lineages. Yet, nearly half of Sundaland
660 Pleistocene speciation events involve dispersal between different islands. One possible
661 explanation is to consider the restored interconnectivity of both Sundaland land bridges and
662 paleoriver systems during Pleistocene glacial periods. Considering cooler and drier climate
663 during Pleistocene glaciations, it has been proposed that savanna and seasonal forest corridors
664 expanded through the interior of Sundaland, notably in East Sunda (Heaney 1992; Bird et al.
665 2005). Such inter-island bridges might be not easily penetrable by evergreen forest-dependent
666 taxa (Heaney 1992; Gorog et al. 2004; Bird et al. 2005; Pouyaud et al. 2009; Wurster et al.
667 2019). For example, despite its exemplary ability to disperse, *Clarias* was more likely to
668 speciate within an island (55.6%) than between islands (44.4%), which is supported by the
669 general phylogenetic division of the genus between black-water lineages and white-clear
670 water lineages. Black water lineages of *Clarias* are unlikely to switch habitats, inhibiting
671 them to easily penetrate freshwater habitats within the savanna/seasonal forest corridors
672 during glaciation periods (Pouyaud et al. 2009). Zenarchopteridae was the lineage least likely
673 to disperse between islands (18.3 %), most probably due to the poor dispersal ability of these

mostly ovoviviparous fishes (de Bruyn et al. 2013). Inter-island dispersal seems to be correlated not only with dispersal ability, but also with habitat specificity (Heaney 2007).

The substantial proportion of founder-effect speciation between islands coupled with dispersal between paleoriver systems and of speciation between paleorivers argues in favor of the importance of habitat heterogeneity/diversity in the diversification of Sundaland freshwater fishes. If paleoriver systems act as dispersal channels, within or between islands, one can also argue that the borders of both paleorivers and insular systems might not be as clearly delineated as previously supposed, due to biological aspects including the difference in vegetation cover and/or physical aspects such as island geomorphology (Pouyaud et al. 2009; Brown et al. 2013; Hutama et al. 2017). The Bangka-Belitung islands are located at the boundary between North Sunda and East Sunda river systems. We found no clear signal of its affiliation with either of the two paleorivers. Similarly, borders between the two river systems also could not be recovered strictly by biogeographic estimates in Lampung (eastern Sumatra) and the southwestern part of Borneo, probably because the topography of these lowlands facilitates gene flow between them. Notable examples for the indistinct boundaries between the North Sunda and East Sunda paleorivers are exemplified by: (1) the existence of sister lineages of *G. platypogonides* 1 and *G. stibaros* (Borneo), (2) dispersal during cladogenesis between the *G. pictus* and *G. major* groups in Borneo, and (3) the trans-paleoriver distributions of *Clarias leiacanthus* (Sumatra) and *C. meladerma* 1 (Sumatra and Borneo). The physical island geomorphology could also contribute to re-arrangements of paleoriver watersheds limits through headwater capture events among paleorivers through time. The Bornean part of the North Sunda and East Sunda river systems share the same headwater area in the center of the island.

Habitat fragmentation by mountainous terrain has been identified as a major geomorphological driver of *in situ* diversification in Java resulting in abundant cryptic

diversity (Dahrudin et al. 2017; Hutama et al. 2017; Hubert et al. 2019). The existence of a single paleoriver system may have driven ancestral lineages on the island to diversify further within isolated pockets of riparian environment. This suggests that extant freshwater fish diversity in Java resulted from high rampant *in situ* diversification rather than immigration from elsewhere (Nguyen et al. 2008; Pouyaud et al. 2009; Hubert et al. 2015, 2019; Kusuma et al. 2016; Dahrudin et al. 2017; Hutama et al. 2017). *In situ* diversification within islands are found in all lineages studied here (e.g. *Glyptothorax platypogon*, *G. robustus*, *Dermogenys pusilla*, and *Channa gachua*; Fig. 4, Fig. S2, Table S2).

Robustness of the Inferences, Limits and Perspectives

For all four taxa under study, we found that tree topologies are congruent with previously published phylogenetic hypotheses (Pouyaud et al. 2009; Jiang et al. 2011; de Bruyn et al. 2013; Conte-Grand et al. 2017). Both ML and Bayesian analyses retrieved similar and robust topologies, and each genus was found to be monophyletic, including the three genera of Zenarchopteridae, with *Nomorhamphus* being the sister group of *Dermogenys* and *Nomorhamphus* + *Dermogenys* as the sister group of *Hemirhamphodon*, supporting Meisner (2001) and Lovejoy et al. (2004). Minor differences concern the monophyly of *Dermogenys*, well supported now, contrary to previous studies (de Bruyn et al. 2013), and likely resulting from the incorporation of multiple outgroups here. For *Clarias*, our analysis supports the monophyly of Asian *Clarias* as previously suggested (Pouyaud et al. 2009). By contrast, ML phylogenetic reconstruction (Fig. 2A, Fig. S1) and the Bayesian species tree (Fig. 4A, Fig. S2) support the reciprocal monophyly of black water species (Clade II) and non black-water species (Clade III). Only *Clarias microstomus* Ng 2001 and *C. planiceps* Ng 1999 departed from this general trend. For *Glyptothorax*, differences with the reference phylogeny (Jiang et al. 2011) are only due to a recent revision of Sundaland *Glyptothorax* species that led to the

revision and description of new taxa (Ng and Kottelat 2016), that have been incorporated here as well as the addition of supplementary sequences (Hutama et al. 2017). Our findings are consistent with the occurrence of two distinct clades in Sundaland (Fig. 2B, Fig. 4B, Fig. S1-S2) as suggested earlier (Jiang et al. 2011). Finally, the species tree of *Channa* recovered here is consistent with the eight species groups previously recognized (Rüber et al. 2019). Our results differ from previous studies in placing the *C. punctata* group and *C. gachua* group in early-diverging position instead of the *C. micropeltes* group and *C. lucius* group (Conte-Grand et al. 2017; Rüber et al. 2019).

At the MOTUs level, several nominal species are not monophyletic, which is probably due to practical taxonomic limitations in which different lineages with no apparent morphological differences might have been lumped together into the same nominal species as in the polyphyletic *Clarias nieuhofii* (as previously observed by Pouyaud et al. 2009), *Glyptothorax platypogonides* and *Channa gachua*, as well as the paraphyletic *Nomorhamphus megarrhamphus* or several *Glyptothorax* species (Jiang et al. 2011; Ng and Kottelat 2016). These results are consistent with recent findings about the existence of high levels of cryptic diversity among Sundaland freshwater fishes (Nguyen et al. 2008; Pouyaud et al. 2009; Hubert et al. 2015, 2019; Conte-Grand et al. 2017; Dahrudin et al. 2017; Hutama et al. 2017; Sholihah et al. 2020) which further points to the necessity of using genetic delimitation methods for subsequent macroevolutionary analyses of complex biotas (Esselstyn et al. 2009; Patel et al. 2011; Ruane et al. 2014; Hubert et al. 2015, 2019; Hutama et al. 2017). The remarkable number of MOTUs recovered from nominal species (252 MOTUs vs. 110 morphological species) calls for more detailed taxonomic works on these taxa (Pouyaud et al. 2009; Hutama et al. 2017).

The divergence times we estimated using a molecular clock approach agree with (*Hemirhamphodon*, *Dermogenys*, *Nomorhamphus*) or are younger than (*Clarias* and *Channa*)

previous estimates based on geological and fossil calibrations. We also provided the first
 time-calibrated phylogeny for *Glyptothorax*. We found similar divergence times for
Hemirhamphodon (9.88 Ma, 95% HPD = 6.02-14.70 Ma, vs. 11.2 Ma, 95% HPD = 6.7-16
 Ma; de Bruyn et al. 2013) and for the *Dermogenys* + *Nomorhamphus* clade (6.28 Ma, 95%
 HPD = 4.28-8.66 Ma, vs. 13.3 Ma, 95% HPD = 8-18 Ma; de Bruyn et al. 2013). Younger
 divergence estimates are found in *Clarias* and *Channa*: we estimated the Asian *Clarias* clade
 to be 6.37 Ma (95% HPD = 3.99-9.19 Ma) while Pouyaud et al. (2009) dated the divergence
 at 33.4 Ma. In the same way, we found a younger divergence time for *Channa* (14.35 Ma,
 95% HPD = 10.54-18.41 Ma) than a previous estimate (28 Ma, 95% HPD = 24-32 Ma; Rüber
 et al. 2020). A potential cause of these discrepancies is likely our use of a mixed
 coalescent/diversification model here for the Bayesian species trees reconstruction that
 accounts for the heterogeneity of absolute substitution rates within and between species (Ho
 and Larson 2006). Within species, clock rates can be considerably higher than between
 species because observed substitution rates are close to mutation rates, genetic drift being
 later responsible for the loss of haplotypes leading to stationary substitution rates that are
 most commonly used in phylogenetic reconstructions and molecular age estimates (Ho and
 Larson 2006). Along the same lines, the use of alternative clock rates moderately impacted
 our main results: (1) most MOTUs originated during the Pleistocene regardless of the clock
 rate used; (2) diversification model selection was consistent across a wide range of clock
 rates. Furthermore, this study is based on genetic recognition of MOTUs instead of
 morphological species; the high proportion of cryptic diversity found here was not properly
 accounted for by the previous phylogenetic reconstructions (Pouyaud et al. 2009; de Bruyn et
 al. 2013). It can be expected that such a difference in taxon sampling has impacted absolute
 age estimates. The general concordance in the distribution through time of MOTUs age

among taxa with different life history traits and origins argue in favor of the robustness of the age estimates established here.

Finally, several issues can be highlighted that warrant further studies. First, although the samples generally cover a broad range of localities around Sundaland, we identify sampling gaps in large areas, including Sumatra for all taxa and in Peninsular Malaysia for *Clarias*. Second, lack of COI sequences deposited in BOLD limited the application of RESL algorithm and designation of Barcode Index Numbers (BINs) for older datasets (*Clarias* and *Channa*). While there was no significant challenge from missing BINs during species delimitation, more complete BIN would increase confidence in the species delimitation results. The subfamily Rasborinae (Brittan 1972; Sholihah et al. 2020) could be an additional biological model to assess the evolutionary history of Sundaland freshwater fishes. It has multiple species-rich genera of small size (Liao et al. 2011; Tan and Armbruster 2018). Lastly, the constant diversification rates through time inferred for two genera might result from inadequate sampling. It has also been suggested that the DDD model might be biased toward spurious detection of early explosive speciation dynamics resulting from the underestimation of the actual number of evolutionary lineages created by taxonomic and sampling bias (Barracough and Nee 2001; Rabosky and Lovette 2008). Alternatively, constant diversification rates have been a recurrent findings in tropical biomes (Esselstyn et al. 2009; Patel et al. 2011). In this case, we have emphasized our effort to avoid clade-specific taxonomic hypotheses (Esselstyn et al. 2009) by utilizing species delimitation methods for all clades. We have detected numerous cryptic lineages in all taxa which might explain the constant diversification rate found here, as proposed by Patel et al. (2011) to explain the constant diversification rates of Neotropical *Pteroglossus*.

CONCLUSIONS

Our results indicate that the diversification and biogeography of freshwater fishes on Sundaland are not solely dependent on Pleistocene sea-level fluctuations and associated paleoriver systems, but are also affected by: (1) pre-Pleistocene geological history of Sundaland, both at the origin of the development of paleoriver systems and responsible for ancient speciation events among paleorivers; (2) the idiosyncratic effects of island boundaries and the extent of paleorivers on diversification in each system; (3) ecology of the paleoenvironments (Heaney 2007) and the (re)emerged paleorivers that were running through the land bridges of Sundaland through the Pleistocene (Heaney 1992; Bird et al. 2005; Slik et al. 2011); (4) the geomorphology of emerged and currently submerged Sundaland land masses, creating indistinct borders among paleoriver systems; and (5) different evolutionary responses of different groups with specific life history traits. Furthermore, it has been suggested recently that the subsidence of Sundaland may have triggered the merge of Sundaland landmasses during glacial times only 400,000 years ago, implying that eustatic fluctuations prior to that period only marginally impacted the extent of emerged lands (Husson et al. 2020). Although, not in line with the initial framework used here for testing the Paleoriver Hypothesis, our observations concerning the significant effect of pre-Pleistocene geological settlement and paleorivers arrangement on biodiversity are in agreement with this recent finding. Finally, our findings propose new perspectives on the biogeography of Sundaland freshwater biotas and open new questions about the interplay between geology and paleoecology during dispersal and colonization of Sundaland islands. Riverine organisms have constrained dispersal routes, whereas terrestrial organisms do not. As such, our studies warrant new researches on the biogeography of terrestrial rainforest biotas of Sundaland.

ACKNOWLEDGEMENTS

This paper is part of the doctoral study of AS, funded by a full scholarship of the Indonesian Endowment Fund for Education (LPDP) from the Indonesian Ministry of Finance. The authors thank Hari Sutrisno and Cahyo Rahmadi as well as the staff of the Zoology division of the Research Centre for Biology (Indonesian Institute of Sciences) for hosting AS during the first year of her PhD. We thank David J. Lohman and one anonymous reviewer for the relevant comments on earlier versions of the manuscript. This publication is ISEM publication number 2020-230 SUD.

References

- Adamson E.A.S., Hurwood D.A., Mather P.B. 2010. A reappraisal of the evolution of Asian snakehead fishes (Pisces, Channidae) using molecular data from multiple genes and fossil calibration. *Mol. Phylogenet. Evol.* 56:707–717.
- Alonso D., Etienne R.S., McKane A.J. 2006. The merits of neutral theory. *Trends Ecol. Evol.* 21:451–457.
- Alter S.E., Munshi-South J., Stiassny M.L.J. 2017. Genomewide SNP data reveal cryptic phylogeographic structure and microallopatric divergence in a rapids-adapted clade of cichlids from the Congo River. *Mol. Ecol.* 26:1401–1419.
- April J., Hanner R.H., Dion-Côté A.M., Bernatchez L. 2012. Glacial cycles as an allopatric speciation pump in North eastern american freshwater fishes. *Mol. Ecol.*
- Barluenga M., Meyer A. 2004. The midas cichlid species complex: incipient sympatric speciation in Nicaraguan cichlid fishes ? *Mol. Ecol.* 13:2061–2076.
- Barraclough T.G., Nee S. 2001. Phylogenetics and Speciation. *Trends Ecol. Evol.* 16:391–399.
- Bates M., Hackett, S., J., Cracraft, J. J. 1998. Area-relationships in the neotropical lowlands: an hypothesis based on raw distributions of passerine birds. *J. Biogeogr.* 25:783–793.

847 Beck S. V., Carvalho G.R., Barlow A., Ruber L., Hui Tan H., Nugroho E., Wowor D., Mohd
848 Nor S.A., Herder F., Muchlisin Z.A., de Bruyn M. 2017. Plio-Pleistocene
849 phylogeography of the Southeast Asian Blue Panchax killifish, *Aplocheilichthys panchax*.
850 PLoS One. 12:e0179557.

851 Berra T.M. 2001. Freshwater Fish Distribution. San Diego: Academic Press.

852 Betancur-R R., Wiley E.O., Arratia G., Acero A., Bailly N., Miya M., Lecointre G., Ortí G.
853 2017. Phylogenetic classification of bony fishes. BMC Evol. Biol. 17:162.

854 Bird M.I., Taylor D., Hunt C. 2005. Palaeoenvironments of insular Southeast Asia during the
855 Last Pleistocene. Glacial Period: a savanna corridor in Sundaland? . Quat. Sci. Rev.
856 24:2228–2242.

857 Blaxter M., Mann J., Chapman T., Thomas F., Whitton C., Floyd R., Abebe E. 2005. Defining
858 operational taxonomic units using DNA barcode data. Philos. Trans. R. Soc. B Biol. Sci.
859 360:1935–1943.

860 Bouckaert R., Heled J., Kühnert D., Vaughan T., Wu C.H., Xie D., Suchard M.A., Rambaut
861 A., Drummond A.J. 2014. BEAST 2: A Software Platform for Bayesian Evolutionary
862 Analysis. PLoS Comput. Biol. 10:1–6.

863 Brittan M.R. 1972. *Rasbora*: A revision of the Indo-Malayan freshwater fish genus Rasbora.
864 Hongkong: TFH Publications.

865 Brown R.M., Siler C.D., Oliveros C.H., Esselstyn J.A., Diesmos A.C., Hosner P.A., Linkem
866 C.W., Barley A.J., Oaks J.R., Sanguila M.B., Welton L.J., Blackburn D.C., Moyle R.G.,
867 Townsend Peterson A., Alcala A.C. 2013. Evolutionary Processes of Diversification in a
868 Model Island Archipelago. Annu. Rev. Ecol. Evol. Syst. 44:411–435.

869 de Bruyn M., Rüber L., Nylander S., Stelbrink B., Lovejoy N.R., Lavoué S., Heok Hui T.,
870 Nugroho E., Wowor D., Ng P.K.L., Siti Azizah M.N., Von Rintelen T., Hall R.,
871 Carvalho G.R. 2013. Paleo-drainage basin connectivity predicts evolutionary

relationships across three Southeast Asian biodiversity hotspots. *Syst. Biol.* 62:398–410.

de Bruyn M., Stelbrink B., Morley R.J., Hall R., Carvalho G.R., Cannon C.H., van den Bergh
G., Meijaard E., Metcalfe I., Boitani L., Maiorano L., Shoup R., von Rintelen T. 2014.
Borneo and Indochina are Major Evolutionary Hotspots for Southeast Asian
Biodiversity. *Syst Biol.* 63:23.

De Bruyn M., Wilson J.A., Mather P.B. 2004. Huxley's line demarcates extensive genetic
divergence between eastern and western forms of the giant freshwater prawn,
Macrobrachium rosenbergii. *Mol. Phylogenet. Evol.* 30:251–257.

Burbrink F.T., Chan Y.L., Myers E.A., Ruane S., Smith B.T., Hickerson M.J. 2016.
Asynchronous demographic responses to Pleistocene climate change in Eastern Nearctic
vertebrates. *Ecol. Lett.* 19:1457–1467.

Cannon C.H., Morley R.J., Bush M.B. 2009. The current refugial rainforests of Sundaland are
unrepresentative of their biogeographic past and highly vulnerable to disturbance. *Proc.*
Natl. Acad. Sci. USA. 106:11188–11193.

Clark J.R., Ree R.H., Alfaro M.E., King M.G., Wagner W.L., Roalson E.H. 2008. A
comparative study in ancestral range reconstruction methods: retracing the uncertain
histories of insular lineages. *Syst. Biol.* 57:693–707.

Clements R., Sodhi N.S., Schilthuizen M., Ng P.K.L. 2006. Limestone karsts of Southeast
Asia: imperiled arks of biodiversity. *Bioscience.* 56:733–742.

Condamine F.L., Nagalingum N.S., Marshall C.R., Morlon H. 2015a. Origin and
diversification of living cycads: a cautionary tale on the impact of the branching process
prior in Bayesian molecular dating. *BMC Evol. Biol.* 15:65.

Condamine F.L., Rolland J., Morlon H. 2013. Macroevolutionary perspectives to
environmental change. *Ecol. Lett.* 16:72–85.

Condamine F.L., Rolland J., Morlon H. 2019. Assessing the causes of diversification

897 slowdowns: temperature-dependent and diversity-dependent models receive equivalent
898 support. *Ecol. Lett.* 22:1900–1912.

899 Condamine F.L., Toussaint E.F.A., Clamens A.L., Genson G., Sperling F.A.H., Kergoat G.J.
900 2015b. Deciphering the evolution of birdwing butterflies 150 years after Alfred Russel
901 Wallace. *Sci. Rep.* 5:1–11.

902 Conte-Grand C., Britz R., Dahanukar N., Raghavan R., Pethiyagoda R., Tan H.H., Hadiaty
903 R.K., Yaakob N.S., Rüber L. 2017. Barcoding snakeheads (Teleostei, Channidae)
904 revisited: Discovering greater species diversity and resolving perpetuated taxonomic
905 confusions. *PLoS One.* 12:e0184017.

906 Currie D.J., Mittelbach G.G., Cornell H. V., Field R., Guegan J.-F., Hawkins B.A., Kaufman
907 D.M., Kerr J.T., Oberdorff T., O’Brien E., Turner J.R.G. 2004. Predictions and tests of
908 climate-based hypotheses of broad-scale variation in taxonomic richness. *Ecol. Lett.*
909 7:1121–1134.

910 Dahruddin H., Hutama A., Busson F., Sauri S., Hanner R., Keith P., Hadiaty R., Hubert N.
911 2017. Revisiting the ichthyodiversity of Java and Bali through DNA barcodes:
912 taxonomic coverage, identification accuracy, cryptic diversity and identification of
913 exotic species. *Mol. Ecol. Resour.* 17:288–299.

914 Van Dam M.H., Matzke N.J. 2016. Evaluating the influence of connectivity and distance on
915 biogeographical patterns in the south-western deserts of North America. *J. Biogeogr.*
916 43:1514–1532.

917 Darriba D., Taboada G.L., Doallo R., Posada D. 2015. Europe PMC Funders Group
918 jModelTest 2 : more models , new heuristics and high- performance computing. 9:6–9.

919 Dodson J.J., Colombani F., Ng P.K.L. 1995. Phylogeographic structure in mitochondrial
920 DNA of a South-east Asian freshwater fish, *Hemibagrus nemurus* (Sluroidei; Bagridae)
921 and Pleistocene sea-level changes on the Sunda shelf. *Mol. Ecol.* 4:331–346.

922 Downing Meisner A. 2001. Phylogenetic systematics of the viviparous halfbeak genera
 923 Dermogenys and Nomorhamphus (Teleostei: Hemiramphidae: Zenarchopterinae). Zool.
 924 J. Linn. Soc. 133:199–283.

925 Duchêne S., Lanfear R., Ho S.Y.W. 2014. The impact of calibration and clock-model choice
 926 on molecular estimates of divergence times. Mol. Phylogenet. Evol. 78:277–289.

927 Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
 928 throughput. Nucleic Acids Res. 32:1792–7.

929 Esselstyn J.A., Brown R.M. 2009. The role of repeated sea-level fluctuations in the generation
 930 of shrew (Soricidae: *Crocidura*) diversity in the Philippine Archipelago. Mol.
 931 Phylogenet. Evol. 53:171–181.

932 Esselstyn J.A., Timm R.M., Brown R.M. 2009. Do geological or climatic processes drive
 933 speciation in dynamic archipelagos? the tempo and mode of diversification in southeast
 934 asian shrews. Evolution. 63:2595–2610.

935 Etienne R.S., Haegeman B., Stadler T., Aze T., Pearson P.N., Purvis A., Phillimore A.B.
 936 2012. Diversity-dependence brings molecular phylogenies closer to agreement with the
 937 fossil record. Proc. R. Soc. B Biol. Sci. 279:1300–1309.

938 Fujisawa T., Barraclough T.G. 2013. Delimiting species using single-locus data and the
 939 generalized mixed yule coalescent approach: A revised method and evaluation on
 940 simulated data sets. Syst. Biol. 62:707–724.

941 Gathorne-Hardy F.J., Syaukani, Davies R.G., Eggleton P., Jones D.T. 2002. Quaternary
 942 rainforest refugia in south-east Asia: using termites (Isoptera) as indicators. Biol. J. Linn.
 943 Soc. 75:453–466.

944 Gavrillets S., Vose A. 2006. Dynamic patterns of adaptative radiation. Proc. Natl. Acad. Sci.
 945 USA. 102:18040–18045.

946 Giam X., Koh L.P., Tan H.H., Miettinen J., Tan H.T.W., Ng P.K.L. 2012. Global extinctions

947 of freshwater fishes follow peatland conversion in Sundaland. *Front. Ecol. Environ.*
948 10:465–470.

949 Gorog A.J., Sinaga M.H., Engstrom M.D. 2004. Vicariance or dispersal? Historical
950 biogeography of three Sunda shelf marine rodents (*Maxomys surifer*, *Leopoldamys*
951 *sabanus* and *Maxomys whiteheadi*). *Biol. J. Linn. Soc.* 81:91–109.

952 Guindon S., Gascuel O. 2003. A Simple, Fast, and Accurate Algorithm to Estimate Large
953 Phylogenies by Maximum Likelihood. *Syst. Biol.* 52:696–704.

954 Guyot J.L. 1993. Hydrogeochimie des fleuves de l'Amazonie Bolivienne. .

955 Haffer J. 1997. Alternative models of vertebrate speciation in Amazonia: an overview.
956 *Biodivers. Conserv.* 6:26.

957 Hall R. 2009. Southeast Asia's changing palaeogeography. *Blumea J. Plant Taxon. Plant*
958 *Geogr.* 54:148–161.

959 Hall R. 2013. The palaeogeography of Sundaland and Wallacea since the Late Jurassic. *J.*
960 *Limnol.* 72:1–17.

961 Hardman M., Lundberg J.G. 2006. Molecular phylogeny and chronology of diversification
962 for “phractocephaline” catfishes (Siluriformes: Pimelodidae) based on mitochondrial
963 DNA and DNA recombination activating gene 2 sequences. *Mol. Phylogenet. Evol.*
964 40:410–418.

965 Heaney L.R. 1992. A synopsis of climatic and vegetational change in Southeast Asia. *Trop.*
966 *For. Clim.*:53–61.

967 Heaney L.R. 2007. Is a new paradigm emerging for oceanic island biogeography? *J.*
968 *Biogeogr.* 34:753–757.

969 Heled J., Drummond A.J. 2010. Bayesian Inference of Species Trees from Multilocus Data.
970 *Mol. Biol. Evol.* 27:570–580.

971 Hendriks K.P., Alciatore G., Schilthuizen M., Etienne R.S. 2019. Phylogeography of Bornean

972 land snails suggests long-distance dispersal as a cause of endemism. *J. Biogeogr.*

973 Herder F., Nolte A.W., Pfaender J., Schwarzer J., Hadiaty R.K., Schliewen U.K. 2006.

974 Adaptative radiation and hybridization in Wallace's dreamponds: evidence from sailfin

975 silversides in the Malili Lakes of Sulawesi. *Proc. R. Soc. B.* 273:2209–2217.

976 Ho S.Y.W., Larson G. 2006. Molecular clocks: when times are a-changin'. *TRENDS Genet.*

977 22:79–83.

978 Hoffmann M., Hilton-Taylor C., Angulo A., Böhm M., Brooks T.M., Butchart S.H.M.,

979 Carpenter K.E., Chanson J., Collen B., Cox N.A., Darwall W.R.T., Dulvy N.K., Harrison

980 L.R., Katariya V., Pollock C.M., Quader S., Richman N.I., Rodrigues A.S.L., Tognelli

981 M.F., Vié J.C., Aguiar J.M., Allen D.J., Allen G.R., Amori G., Ananjeva N.B., Andreone

982 F., Andrew P., Ortiz A.L.A., Baillie J.E.M., Baldi R., Bell B.D., Biju S.D., Bird J.P.,

983 Black-Decima P., Blanc J.J., Bolaños F., Bolivar-G. W., Burfield I.J., Burton J.A.,

984 Capper D.R., Castro F., Catullo G., Cavanagh R.D., Channing A., Chao N.L., Chenery

985 A.M., Chiozza F., Clausnitzer V., Collar N.J., Collett L.C., Collette B.B., Cortez

986 Fernandez C.F., Craig M.T., Crosby M.J., Cumberlidge N., Cuttelod A., Derocher A.E.,

987 Diesmos A.C., Donaldson J.S., Duckworth J.W., Dutson G., Dutta S.K., Emslie R.H.,

988 Farjon A., Fowler S., Freyhof J., Garshelis D.L., Gerlach J., Gower D.J., Grant T.D.,

989 Hammerson G.A., Harris R.B., Heaney L.R., Hedges S.B., Hero J.M., Hughes B.,

990 Hussain S.A., Icochea M. J., Inger R.F., Ishii N., Iskandar D.T., Jenkins R.K.B., Kaneko

991 Y., Kottelat M., Kovacs K.M., Kuzmin S.L., La Marca E., Lamoreux J.F., Lau M.W.N.,

992 Lavilla E.O., Leus K., Lewison R.L., Lichtenstein G., Livingstone S.R., Lukoschek V.,

993 Mallon D.P., McGowan P.J.K., McIvor A., Moehlman P.D., Molur S., Alonso A.M.,

994 Musick J.A., Nowell K., Nussbaum R.A., Olech W., Orlov N.L., Papenfuss T.J., Parra-

995 Olea G., Perrin W.F., Polidoro B.A., Pourkazemi M., Racey P.A., Ragle J.S., Ram M.,

996 Rathbun G., Reynolds R.P., Rhodin A.G.J., Richards S.J., Rodríguez L.O., Ron S.R.,

997 Rondinini C., Rylands A.B., de Mitcheson Y.S., Sanciangco J.C., Sanders K.L., Santos-
 998 Barrera G., Schipper J., Self-Sullivan C., Shi Y., Shoemaker A., Short F.T., Sillero-
 999 Zubiri C., Silvano D.L., Smith K.G., Smith A.T., Snoeks J., Stattersfield A.J., Symes
 1000 A.J., Taber A.B., Talukdar B.K., Temple H.J., Timmins R., Tobias J.A., Tsytsulina K.,
 1001 Tweddle D., Ubeda C., Valenti S. V., Van Dijk P.P., Veiga L.M., Veloso A., Wege D.C.,
 1002 Wilkinson M., Williamson E.A., Xie F., Young B.E., Akçakaya H.R., Bennun L.,
 1003 Blackburn T.M., Boitani L., Dublin H.T., da Fonseca G.A.B., Gascon C., Lacher T.E.,
 1004 Mace G.M., Mainka S.A., McNeely J.A., Mittermeier R.A., Reid G.M.G., Rodriguez
 1005 J.P., Rosenberg A.A., Samways M.J., Smart J., Stein B.A., Stuart S.N. 2010. The impact
 1006 of conservation on the status of the world's vertebrates. *Science* (80-.). 330:1503–1509.
 1007 Hubbell S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton:
 1008 Princeton University Press.
 1009 Hubert N., Calcagno V., Etienne R.S., Mouquet N. 2015. Metacommunity speciation models
 1010 and their implications for diversification theory. *Ecol. Lett.* 18:864–881.
 1011 Hubert N., Lumbantobing D., Sholihah A., Dahruddin H., Delrieu-Trottin E., Busson F., Sauri
 1012 S., Hadiaty R., Keith P. 2019. Revisiting species boundaries and distribution ranges of
 1013 *Nemacheilus* spp. (Cypriniformes: Nemacheilidae) and *Rasbora* spp. (Cypriniformes:
 1014 Cyprinidae) in Java, Bali and Lombok through DNA barcodes: implications for
 1015 conservation in a biodiversity hotspot. *Conserv. Genet.* 20:517–529.
 1016 Hubert N., Renno J.F. 2006. Historical Biogeography of South American Freshwater fishes. *J.*
 1017 *Biogeogr.* 33:1414–1436.
 1018 Husson L., Boucher F.C., Sarr A.C., Sepulchre P., Cahyarini S.Y. 2020. Evidence of
 1019 Sundaland's subsidence requires revisiting its biogeography. *J. Biogeogr.* 47:843–853.
 1020 Hutama A., Dahruddin H., Busson F., Sauri S., Keith P., Hadiaty R.K., Hanner R., Suryobroto
 1021 B., Hubert N. 2017. Identifying spatially concordant evolutionary significant units across

multiple species through DNA barcodes: Application to the conservation genetics of the
freshwater fishes of Java and Bali. *Glob. Ecol. Conserv.* 12:170–187.

Jiang W., Ng H.H., Yang J., Chen X. 2011. Monophyly and phylogenetic relationships of the
catfish genus *Glyptothorax* (Teleostei: Sisoridae) inferred from nuclear and
mitochondrial gene sequences. *Mol. Phylogenet. Evol.* 61:278–289.

Kadarusman, Hubert N., Hadiaty R.K., Sudarto, Paradis E., Pouyaud L. 2012. Cryptic
diversity in indo-australian rainbowfishes revealed by DNA barcoding: Implications for
conservation in a biodiversity hotspot candidate. *PLoS One.* 7.

Kekkonen M., Hebert P.D.N. 2014. DNA barcode-based delineation of putative species:
efficient start for taxonomic workflows. *Mol. Ecol. Resour.* 14:706–715.

Kekkonen M., Mutanen M., Kaila L., Nieminen M., Hebert P.D.N. 2015. Delineating Species
with DNA Barcodes: A Case of Taxon Dependent Method Performance in Moths. *PLoS
One.* 10:e0122481.

Kisel Y., McInnes L., Toomey N.H., Orme C.D.L. 2011. How diversification rates and
diversity limits combine to create large-scale species-area relationships. *Philos. Trans. R.
Soc. B Biol. Sci.* 366:2514–2525.

Knowlton N., Weight L.A., Solorzano L.A., Mills D.K., Bermingham E. 1992. Divergence of
proteins, mitochondrial DNA and reproductive compatibility across the Isthmus of
Panama. *Science* (80-.). 260:1629–1632.

Kottelat M., Whitten A.J., Kartikasari N., Wirjoatmodjo S. 1993. Freshwater Fishes of
Western Indonesia and Sulawesi. Periplus Edition (HK) Ltd.

Kumar S., Stecher G., Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* 33:1870–4.

Kusuma W.E., Ratmuangkhwang S., Kumazawa Y. 2016. Molecular phylogeny and historical
biogeography of the Indonesian freshwater fish *Rasbora lateristriata* species complex

1047 (Actinopterygii: Cyprinidae): Cryptic species and west-to-east divergences. Mol.
 1048 Phylogenet. Evol. 105:212–223.

1049 Landry L., Vincent W., Bernatchez L. 2007. Parallel evolution of lake whitefish dwarf
 1050 ecotypes in association with limnological features of their adaptive landscape. J. Evol.
 1051 Biol.

1052 Li F., Li S. 2018. Plaeocene-Eocene and Plio-Pleistocene sea-level changes as “species
 1053 pumps” in Southeast Asia: Evidence from *Althepus* spiders. Mol. Phylogenet. Evol.
 1054 127:545–555.

1055 Liao T.Y., Kullander S.O., Fang F. 2011. Phylogenetic position of rasborin cyprinids and
 1056 monophyly of major lineages among the Danioninae, based on morphological characters
 1057 (Cypriniformes: Cyprinidae). J. Zool. Syst. Evol. Res. 49:224–232.

1058 Lim H.-C., Abidin M.Z., Pulungan C.P., De Bruyn M., Mohd Nor S.A. 2016. DNA barcoding
 1059 reveals high cryptic diversity of freshwater halfbeak genus *Hemirhamphodon* from
 1060 Sundaland. PLoSONE. 11:e0163596.

1061 Lisiecki L.E., Raymo M.E. 2005. A Pliocene-Pleistocene stack of 57 globally distributed
 1062 benthic δ 18O records. Paleoclimatology. 20:1–17.

1063 Liu J., Guo X., Chen D., Li J., Yue B., Zeng X. 2019. Diversification and historical
 1064 demography of the rapid racerunner (*Eremias velox*) in relation to geological history and
 1065 Pleistocene climatic oscillations in arid Central Asia. Mol Phylogenet Evol. 130:244–
 1066 258.

1067 Lohman K., De Bruyn M., Page T., Von Rintelen K., Hall R., Ng P.K.L., Shih H.-T.,
 1068 Carvalho G.R., Von Rintelen T. 2011. Biogeography of the Indo-Australian archipelago.
 1069 Annu. Rev. Ecol. Evol. Syst. 42:205–226.

1070 Lovejoy N.R., Iranpour M., Collette B.B. 2004. Phylogeny and jaw ontogeny of beloniform
 1071 fishes. Integr. Comp. Biol. 44:366–377.

1072 Maddison W.P., Maddison D.R. 2019. Mesquite: a modular system for evolutionary analysis.
 1073 Version 3.61. <http://www.mesquiteproject.org>.

1074 Maina J.N., Maloiy G.M.O. 1986. The morphology of the respiratory organs of the African
 1075 air-breathing catfish (*Clarias mossambicus*): A light, electron and scanning microscopic
 1076 study, with morphometric observations. J. Zool. 209:421–445.

1077 Matzke N.J. 2013. Probabilistic historical biogeography: New models for founder-event
 1078 speciation, imperfect detection, and fossils allow improved accuracy and model-testing.
 1079 Front. Biogeogr. 5:242–248.

1080 Matzke N.J. 2014. Model selection in historical biogeography reveals that founder-event
 1081 speciation is a crucial process in island clades. Syst. Biol. 63:951–970.

1082 McPeck M.A. 2008. The ecological dynamics of clade diversification and community
 1083 assembly. Am. Nat. 172:e270–e284.

1084 Meisner A.D. 2001. Phylogenetic systematics of the viviparous halfbeak genera *Dermogenys*
 1085 and *Nomorhamphus* (Teleostei: Hemiramphidae: Zenarchopterinae). Zool. J. Linn. Soc.
 1086 133:199–283.

1087 Miller K.G., Kominz M.A., Browning J. V, Wright J.D., Mountain G.S., Katz M.E.,
 1088 Sugarman P.J., Cramer B.S., Christie-Blick N., Pekar S.F. 2005. The phanerozoic record
 1089 of global sea-level change. Science (80-.). 310:1293–1298.

1090 Miller M.A., Pfeiffer W., Schwartz T. 2010. Creating the CIPRES Science Gateway for
 1091 inference of large phylogenetic trees. 2010 Gatew. Comput. Environ. Work. GCE 2010.

1092 Mittelbach G.G., Schemske D.W., Cornell H. V, Allen A.P., Brown J.M., Bush M.B.,
 1093 Harrison S.P., Hurlbert A.H., Knowlton N., Lessios H.A., McCain C.M., McCune A.R.,
 1094 McDade L.A., McPeck M.A., Near T.J., Price T.D., Ricklefs R.E., Roy K., Sax D.F.,
 1095 Schluter D., Sobel J.M., Turelli M. 2007. Evolution and the latitudinal diversity gradient:
 1096 speciation, extinction and biogeography. Ecol. Lett. 10:315–331.

1097 Morlon H., Lewitus E., Condamine F.L., Manceau M., Clavel J., Drury J. 2016. RPANDA: an
1098 R package for macroevolutionary analyses on phylogenetic trees. *Methods Ecol. Evol.*
1099 7:589–597.

1100 Morlon H., Parsons T.L., Plotkin J.B. 2011. Reconciling molecular phylogenies with the
1101 fossil record. *Proc. Natl. Acad. Sci. U. S. A.* 108:16327–16332.

1102 Munshi J.S.D. 1961. The accessory respiratory organs of *Clarias batrachus* (Linn.). *J.*
1103 *Morphol.* 109:115–139.

1104 Myers N., Mittermeier R.A., Mittermeier C.G., da Fonseca G.A.B., Kent F. 2000.
1105 Biodiversity hotspots for conservation priorities. *Nature.* 403:853–858.

1106 Nagel L., Schluter D. 1998. Body size, natural selection, and speciation in sticklebacks.
1107 *Evolution.* 52:209–218.

1108 Ng H.H., Kottelat M. 2016. The *Glyptothorax* of Sundaland: A revisionary study (Teleostei:
1109 Sisoridae).

1110 Ng P.K.L., Tay J.B., Lim B.K. 1994. Diversity and conservation of blackwater fishes in
1111 Peninsular Malaysia, particularly in the North Selangor peat swamp forest.
1112 *Hydrobiologia.* 285:203–218.

1113 Nguyen T.T.T., Na-Nakorn U., Sukmanomon S., ZiMing C. 2008. A study on phylogeny and
1114 biogeography of mahseer species (Pisces: Cyprinidae) using sequences of three
1115 mitochondrial DNA gene regions. *Mol. Phylogenet. Evol.*

1116 Nolte A., Freyhof J., Stemshorn K., Tautz D. 2005. An invasive lineage of sculpins, *Cottus*
1117 sp. (Pisces, Teleostei) in the Rhine with new habitat adaptations has originated from
1118 hybridization between old phylogeographic groups. *Proc. R. Soc. London, B.* 272:2379–
1119 2387.

1120 Nores M. 1999. An alternative hypothesis for the origin of Amazonian bird diversity. *J.*
1121 *Biogeogr.* 26:475–485.

1122 Nores N. 2004. The implications of tertiary and quaternary sea level rise events for avian
1123 distribution patterns in the lowlands of northern South America. *Glob. Ecol. Biogeogr.*
1124 13:149–161.

1125 Nurul Farhana S., Muchlisin Z.A., Duong T.Y., Tanyaros S., Page L.M., Zhao Y., Adamson
1126 E.A.S., Khaironizam M.Z., de Bruyn M., Siti Azizah M.N. 2018. Exploring hidden
1127 diversity in Southeast Asia's *Dermogenys* spp. (Beloniformes: Zenarchopteridae)
1128 through DNA barcoding. *Sci. Rep.* 8:10787.

1129 Ogilvie H.A., Bouckaert R.R., Drummond A.J. 2017. StarBEAST2 brings faster species tree
1130 inference and accurate estimates of substitution rates. *Mol. Biol. Evol.* 34:2101–2114.

1131 Orti Meyer, A. G. 1997. The Radiation of Characiform Fishes and the limits of resolution of
1132 mitochondrial ribosomal DNA sequences. *Syst. Biol.* 46:75–100.

1133 Papadopoulou A., Knowles L.L. 2015a. Genomic tests of the species-pump hypothesis:
1134 Recent island connectivity cycles drive population divergence but not speciation in
1135 Caribbean crickets across the Virgin Islands. *Evolution.* 69:1501–1517.

1136 Papadopoulou A., Knowles L.L. 2015b. Species-specific responses to island connectivity
1137 cycles: Refined models for testing phylogeographic concordance across a Mediterranean
1138 Pleistocene Aggregate Island Complex. *Mol. Ecol.* 24:4252–4268.

1139 Paradis E. 2012. Analysis of phylogenetics and evolution with R. New York: Springer.

1140 Paradis E., Claude J., Strimmer K. 2004. APE: Analyses of phylogenetics and evolution in R
1141 language. *Bioinformatics.* 20:289–290.

1142 Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and
1143 evolutionary analyses in R. *Bioinformatics.* 35:526–528.

1144 Patel S., Weckstein J.D., Patane J.S., Bates J.M., Aleixo A. 2011. Temporal and spatial
1145 diversification of *Pteroglossus aracaris* (AVES: Ramphastidae) in the neotropics:
1146 constant rate of diversification does not support an increase in radiation during the

1147 Pleistocene. *Mol Phylogenet Evol.* 58:105–115.

1148 Pellissier L., Leprieur F., Parravicini V., Cowman P.F., Kulbicki M., Litsios G., Olsen S.M.,
 1149 Wisz M.S., Bellwood D.R., Mouillot D. 2014. Quaternary coral reef refugia preserved
 1150 fish diversity. *Science* (80-.). 344:1015–1019.

1151 Phillimore A.B., Price T.D. 2008. Density-dependent cladogenesis in birds. *PLoS Biol.* 6:e71.

1152 Pons J., Barraclough T.G., Gomez-Zurita J., Cardoso A., Duran D.P., Hazell S., Kamoun S.,
 1153 Sumlin W.D., Vogler A.P. 2006. Sequence-based species delimitation for the DNA
 1154 taxonomy of undescribed insects. *Syst. Biol.* 55:595–606.

1155 Pouyaud L., Sudarto, Paradis E. 2009. The phylogenetic structure of habitat shift and
 1156 morphological convergence in Asian *Clarias* (Teleostei, Siluriformes: Clariidae). *J.*
 1157 *Zool. Syst. Evol. Res.* 47:344–356.

1158 Puillandre N., Lambert A., Brouillet S., Achaz G. 2012. ABGD, Automatic Barcode Gap
 1159 Discovery for primary species delimitation. *Mol. Ecol.* 21:1864–1877.

1160 Rabosky D.L., Lovette I.J. 2008. Explosive evolutionary radiations: Decreasing speciation or
 1161 increasing extinction through time? *Evolution.* 62:1866–1875.

1162 Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior summarization
 1163 in Bayesian phylogenetics using Tracer 1.7. *Syst. Biol.* 67:901–904.

1164 Ratnasingham S., Hebert P.D.N. 2007. BOLD: The Barcode of Life Data System
 1165 (www.barcodinglife.org). *Mol. Ecol. Notes.* 7:355–364.

1166 Ratnasingham S., Hebert P.D.N. 2013. A DNA-Based Registry for All Animal Species: The
 1167 Barcode Index Number (BIN) System. *PLoS One.* 8.

1168 Read C.I., Bellwood D.R., Herwerden van L. 2006. Ancient origins of Indo-Pacific coral reef
 1169 fish biodiversity: a case study of the Leopard wrasses (Labridae: *Macropharyngodon*).
 1170 *Mol. Phylogenet. Evol.* 38:808–819.

1171 Ree R.H., Sanmartín I. 2018. Conceptual and statistical problems with the DEC+ J model of

1172 founder-event speciation and its comparison with DEC via model selection. *J. Biogeogr.*
 1173 45:741–749.

1174 Ruane S., Bryson R.W., Pyron R.A., Burbrink F.T. 2014. Coalescent species delimitation in
 1175 Milksnakes (Genus *Lampropeltis*) and impacts on phylogenetic comparative analyses.
 1176 *Syst. Biol.* 63:231–250.

1177 Rüber L., Tan H.H., Britz R. 2019. Snakehead (Teleostei: Channidae) diversity and the
 1178 Eastern Himalaya biodiversity hotspot. *J. Zool. Syst. Evol. Res.*:1–31.

1179 Sathiamurthy E., Voris K.H. 2006. Maps of Holocene Sea Level Transgression and
 1180 Submerged Lakes on the Sunda Shelf. .

1181 Schluter D. 2000. Ecological causes of adaptative radiation. *Am. Nat.* 148:40–64.

1182 Seehausen O. 2007. Evolution and ecological theory: Chance, historical contingency and
 1183 ecological determinism jointly determine the rate of adaptive radiation. *Heredity (Edinb)*.
 1184 99:361–363.

1185 Seehausen O., Takimoto G., Roy D., Jokela J. 2008. Speciation reversal and biodiversity
 1186 dynamics with hybridization in changing environments. *Mol. Ecol.* 17:30–44.

1187 Serrao N.R., Steinke D., Hanner R.H. 2014. Calibrating snakehead diversity with DNA
 1188 barcodes: Expanding taxonomic coverage to enable identification of potential and
 1189 established invasive species. *PLoS One*. 9.

1190 Sholihah A., Delrieu-Trottin E., Sukmono T., Dahrudin H., Risdawati R., Elvyra R.,
 1191 Wibowo A., Kustiati K., Busson F., Sauri S., Nurhaman U., Dounias E., Zein M.S.A.,
 1192 Fitriana Y., Utama I.V., Hubert N. 2020. Disentangling the taxonomy of the subfamily
 1193 Rasborinae (Cypriniformes, Danionidae) in Sundaland using DNA barcodes. *Sci. Rep.*
 1194 Slik J.W.F., Aiba S.-I., Bastian M., Brearley F.Q., Cannon C.H., Eichhorn K.A.O.,
 1195 Fredriksson G., Kartawinata K., Laumonier Y., Mansor A., Marjokorpi A., Meijaard E.,
 1196 Morley R.J., Nagamasu H., Nilus R., Nurtjahya E., Payne J., Permana A., Poulsen A.D.,

1197 Raes N., Riswan S., van Schaik C.P., Sheil D., Sidiyasa K., Suzuki E., van Valkenburg
1198 J.L.C.H., Webb C.O., Wich S., Yoneda T., Zakaria R., Zweifel N. 2011. Soils on
1199 exposed Sunda Shelf shaped biogeographic patterns in the equatorial forests of Southeast
1200 Asia. *Proc. Natl. Acad. Sci. USA*. 108:12343–12347.

1201 Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
1202 large phylogenies. *Bioinformatics*. 30:1312–1313.

1203 Sullivan J., Lavoué S., Hopkins C. 2002. Discovery and phylogenetic analysis of e riverine
1204 species flock of african electric fishes (Mormyridae: Teleostei). *Evolution*. 56:597–616.

1205 Tan H.H., Lim K.K.P. 2013. Three new species of freshwater halfbeaks (Teleostei:
1206 Zenarchopteridae: *Hemirhamphodon*) from Borneo. *Raffles Bull. Zool.* 61:735–747.

1207 Tan M., Armbruster J.W. 2018. Phylogenetic classification of extant genera of fishes of the
1208 order Cypriniformes (Teleostei: Ostariophysi). *Zootaxa*. 4476:6–39.

1209 Tan M.P., Jamsari A.F.J., Siti Azizah M.N. 2012. Phylogeographic Pattern of the Striped
1210 Snakehead, *Channa striata* in Sundaland: Ancient River Connectivity, Geographical and
1211 Anthropogenic Singnatures. *PLoS One*. 7:1–11.

1212 Verheyen E., Salzburger W., Snoeks J., Meyer A. 2003. Origin of the superflock of cichlid
1213 fishes from Lake Victoria, East Africa. *Science* (80-.). 300:325–329.

1214 Voris H.K. 2000. Maps of Pleistocene sea levels in Southeast Asia: Shorelines, river systems
1215 and time durations. *J. Biogeogr.* 27:1153–1167.

1216 Walker J.D., Geissman J.W., Bowring S.A., Babcock L.E. 2018. GSA Geologic Time Scale v.
1217 5.0. .

1218 Weigelt P., Steinbauer M.J., Cabral J.S., Kreft H. 2016. Late quaternary climate change
1219 shapes island biodiversity. *Nature*. 532:99–102.

1220 Wiens J.J., Donoghue M.J. 2004. Historical biogeography, ecology and species richness.
1221 *Trends Ecol. Evol.* 19:639–644.

1222 Woodruff D.S. 2010. Biogeography and conservation in Southeast Asia: How 2.7 million
 1223 years of repeated environmental fluctuations affect today's patterns and the future of the
 1224 remaining refugial-phase biodiversity. *Biodivers. Conserv.* 19:919–941.

1225 Wurster C.M., Rifai H., Zhou B., Haig J., Bird M.I. 2019. Savanna in equatorial Borneo
 1226 during the late Pleistocene. *Sci. Rep.* 9:1–7.

1227 Zhang J., Kapli P., Pavlidis P., Stamatakis A. 2013. A general species delimitation method
 1228 with applications to phylogenetic placements. *Bioinformatics.* 29:2869–2876.

1229

1230 **FIGURE AND TABLE CAPTIONS**

1231 Figure 1. Paleogeographic maps of Sundaland in the last 20 million years Pleistocene sea
 1232 levels and associated paleorivers. Reconstructions of historical land and sea distribution
 1233 during the Neogene, depicting Sundaland during the early Miocene (A), late Miocene (B) and
 1234 early Pliocene (C) (modified from Lohman et al. 2011). The Paleoriver systems in Southeast
 1235 Asia are shown (D), depicting East Sunda (1), North Sunda (2), Malacca strait (3), Siam (4)
 1236 river systems of Sundaland and the Mekong river system of IndoBurma. Lands exposed
 1237 during -60 and -120m sea level decreases are illustrated in D. The cumulative number of
 1238 sampling locations for each group within the paleoriver systems is provided (E) with color
 1239 codes as follows: orange for *Clarias*, green for *Glyptothorax*, light blue for *Zenarchopteridae*
 1240 and rose for *Channa*; and grey circles indicate that no sample was obtained from those areas.

1241

1242 Figure 2. Maximum likelihood trees and species delimitation for (A) *Clarias*, (B)
 1243 *Glyptothorax*, (C) *Zenarchopteridae* and (D-E) *Channa*. For each clade, MOTUs delimited
 1244 according to GMYC, PTP, BIN and ABGD (yellow), and the 50 percent consensus scheme of
 1245 delimitation (red) are shown. Node circles represent bootstrap proportions and nodes

numbering corresponds with the calibration nodes in Table 1. Major clades are labeled on the right.

Figure 3. Diversification through time of Sundaland freshwater fishes. Panel A shows the lineages through time (LTT) plot with confidence intervals collected from 1000 randomly sampled trees along the StarBEAST2 MCMC, superimposed with the total number of nominal species and MOTUs through time (0.5 Myr class). Panel B shows the speciation rate through time for each group, based on the most likely model of diversification (constant for *Clarias* and *Glyptothorax*, time-exponential for Zenarchopteridae, sea level-dependent for *Channa*). Panel C shows the sea level fluctuations over the last 15 Myr (adapted from Miller et al. 2005).

Figure 4. Panels a to d show the Bayesian maximum credibility trees for (A) *Clarias*, (B) *Glyptothorax*, (C) Zenarchopteridae and (D) *Channa* as well as the ancestral area reconstructions for each group, based on islands (left) and paleorivers (right). Panels E to F show exemplar specimens for (E) *Clarias*, (F) *Glyptothorax*, (G) Zenarchopteridae and (H) *Channa* with their relative size (scale bar = 1 cm).

Figure 5. Geographic pattern of speciation in Southeast Asian freshwater fishes. The plots show the numbers of speciation events associated to the four geographic patterns of speciation as follows: between islands and within paleorivers, between islands and between paleorivers, within islands and within paleorivers, and within islands and between paleorivers. Each bar shows the number of speciation events for each category for (A) all groups (cumulative), (B) Zenarchopteridae, (C) *Channa*, (D) *Clarias* and (E) *Glyptothorax*.

1271 Table 1. Reference nodes and associated time calibrations for each group. Node numbering is
1272 as in Figure 2. Age estimates result from the Bayesian StarBEAST2 analyses and parameter
1273 distributions represent median age, 5 percent and 95 percent quantiles for each reference
1274 nodes

1275

1276 Table 2. Summary statistics of geographic patterns of speciation events for *Clarias*,
1277 *Glyptothorax*, Zenarchopteridae and *Channa* inferred using ancestral area estimation by
1278 BioGeoBEARS.

1279

1280 **SUPPLEMENTARY MATERIALS**

1281 Figure S1. Maximum Likelihood trees for *Clarias* (A), *Glyptothorax* (B), *Beloniformes* (C)
1282 and *Channa* (D) including tip labels as in Table S2 (online supplementary material)

1283

1284 Figure S2. StarBEAST2 species trees for *Clarias* (A), *Glyptothorax* (B), *Beloniformes* (C)
1285 and *Channa* (D) including tip labels as in Table S2 (online supplementary material)

1286

1287 Figure S3. Lineage-through time plots (A) and species age distributions (B) according to
1288 varying substitution rate hypotheses (clock rates ranging from 0.4 to 1.8 % divergence per
1289 million years) for *Clarias*, *Glyptothorax*, *Beloniformes* and *Channa*

1290

1291 Table S1. Description of the 17 diversification models tested for *Clarias*, *Glyptothorax*,
1292 *Beloniformes* and *Channa*, including acronym, rate variation, number of parameters and
1293 parameters involved (see material and methods section for details)

1294

1295 Table S2. List of the DNA sequences compiled in the present study including systematic
1296 (order, family, genus, species and updated species names), specimen number assigned,
1297 locality, geographic distribution (locality, island-based and palaeodrainage-based), GenBank
1298 accession numbers, DNA-based species delimitation (GMYC, PTP, BIN, ABGD), OTU
1299 numbers and phylogenetic status (ingroup, outgroup)
1300

1301 Table S3. Most likely substitution models for each molecular marker within each group,
1302 selected using jModelTest
1303

1304 Table S4. Likelihood scores for each diversification model for each group. Description of
1305 each model can be found in Table S1

1306 Table S5. Likelihood scores for each diversification model for each of the seven clock rate
1307 hypotheses (0.4% to 1.6% per million years) for *Channa*.
1308

1309 Table S6. Log likelihood scores and Akaike Information Criterion values for DEC, DEC+J,
1310 DIV, DIV+J, BAY and BAY+J as calculated using BioGeoBEARS for island-based and
1311 paleoriver-based geographic areas

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5