



Investigating relationships between technological variability and ecology in the Middle Gravettian (ca. 32–28 ky cal. BP) in France

Anaïs Vignoles, William E. Banks, Laurent Klaric, Masa Kageyama, Marlon E Cobos, Daniel Romero-Alvarez

► To cite this version:

Anaïs Vignoles, William E. Banks, Laurent Klaric, Masa Kageyama, Marlon E Cobos, et al.. Investigating relationships between technological variability and ecology in the Middle Gravettian (ca. 32–28 ky cal. BP) in France. *Quaternary Science Reviews*, 2021, 253, pp.106766. 10.1016/j.quascirev.2020.106766 . hal-03385329

HAL Id: hal-03385329

<https://hal.science/hal-03385329>

Submitted on 19 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Investigating relationships between technological variability and ecology in the Middle Gravettian (*ca.* 32-28 ky cal. BP) in France

Anaïs Vignoles^{*a}, William E. Banks^{a,d}, Laurent Klaric^b, Masa Kageyama^c, Marlon E. Cobos^d, Daniel Romero-Alvarez^d

^aUMR-5199 PACEA, Université de Bordeaux, Bâtiment B2, Allée Geoffroy Saint-Hilaire, CS 50023, 33615 Pessac Cedex (France) – ^{*}anaïs.vignoles@u-bordeaux.fr (corresponding author) ; william.banks@u-bordeaux.fr

^bUMR-7055 Préhistoire et Technologie, Université Paris X-Nanterre, 21 allée de l'Université, 92023 Nanterre Cedex (France) – laurent.klaric@cnrs.fr

^cUMR-8212 LSCE, Université Paris-Saclay, Bâtiment 714, Site de l'Orme des Merisiers, Chemin Saint-Aubin – RD 128, F-91191 Gif-sur-Yvette Cedex (France) – masa.kageyama@lsce.ipsl.fr

^dBiodiversity Institute and Department of Ecology & Evolutionary Biology, University of Kansas, 1345 Jayhawk Blvd., Lawrence, KS 66045 (United States of America) – marloncobos@ku.edu ; daromero88@gmail.com

Abstract

The French Middle Gravettian represents an interesting case study for attempting to identify mechanisms behind the typo-technological variability observed in the archaeological record. Associated with the relatively cold and dry environments of GS.5.2 and 5.1, this phase of the Gravettian is characterized by two lithic typo-technical entities (*faciès* in French): the Noaillian (defined by the presence of Noailles burins) and the Rayssian (identified by the Raysse method of bladelet production).

The two *faciès* have partially overlapping geographic distributions, with the Rayssian having a more northern and restricted geographic extension than the Noaillian. Their chronological relationship, however, is still unclear, and interpretations of their dual presence at many sites within the region of overlap are not yet consensual. Nonetheless, the absence of the Raysse method south of the Garonne River suggests that this valley may have separated two different cultural trajectories for which the Rayssian represents an adaptation to environmental conditions different from those associated with Noaillian assemblages south of the Garonne River. The aim of this study is to test this hypothesis quantitatively using ecological niche modeling (ENM) methods. First, we critically evaluate published data to construct inventories of Noaillian and Rayssian archaeological sites. Next, using ENM methods, we estimate the ecological niches associated with the Middle Gravettian north (Noaillian + Rayssian) and south (Pyrenees Noaillian) of the Garonne River, which are then quantitatively evaluated and compared. Results demonstrate that, despite a relatively large degree of similarity, the niches differ significantly from one another in both geographic and environmental dimensions and that the niche associated with the northern Middle Gravettian is broader than that of the Pyrenees Noaillian. We propose that this pattern reflects different technological, subsistence and mobility strategies linked to the development of the Raysse method in the North, which was likely more advantageous in its environmental contexts than technologies employed by contemporaneous populations in the Pyrenees.

Keywords: Middle Gravettian, France, Noaillian, Rayssian, ecological niche modeling, culture-environment relationships

48 **1 Introduction**

49 The Gravettian is an Upper Paleolithic techno-complex (*sensu* Clarke, 1968) that has
50 been the subject of extensive research since its recognition (e.g. Klima, 1976; Roebroeks *et al.*,
51 1999; Rigaud, 2008, 2009; de las Heras *et al.*, 2013; Otte, 2013; Wojtal *et al.*, 2015). Spanning
52 *ca.* 34,000–26,000 calibrated years before present (y. cal. BP), its main unifying characteristics
53 are Gravette-style backed blades and bladelets (Pesesse, 2013), diagnostic graphic expressions
54 (Féreglio *et al.*, 2011), as well as a high frequency of burials (Henry-Gambier, 2008) compared
55 to preceding and subsequent archaeological cultures. These common characteristics are
56 observed in sites across Europe, from Portugal to the Don Valley in western Russia (Otte, 2013).
57 However, the term “Gravettian” groups together a wide variety of cultural traditions, especially
58 concerning lithic and osseous technology (de la Peña-Alonso, 2009; Pesesse, 2013; Noiret,
59 2013; Goutas, 2013a). This diversity is challenging to explain since it is characterized by
60 disparate data, many of which were obtained with non-modern excavation methods decades
61 ago or differing analytical approaches (de la Peña-Alonso, 2011; Pesesse, 2017). Moreover,
62 sites that date to the same chronological interval but that lack typical Gravettian features (i.e.
63 Gravette-style points) serve to challenge the definition of this techno-complex (e.g. Morala,
64 2011; Klaric *et al.*, 2011, 2018). In Western Europe, various hypotheses have been proposed to
65 explain this diversity, such as differences in site activities (e.g. Laville & Rigaud, 1973; Rigaud,
66 1988), the nature of our archaeological definitions (e.g. Touzé, 2013; Pesesse, 2017), regionally
67 differentiated populations that did not share the same technological knowledge or traditions
68 (e.g. Klaric *et al.*, 2009), or differential environmental influences (e.g. David, 1985; Djindjian
69 *et al.*, 1999). Efforts to identify and evaluate the mechanisms—defined as “a constellation of
70 factors and components that through the process of their interaction with one another stimulates
71 the trajectory of a system” (d’Errico & Banks, 2013, p. 374)—that influenced these cultural
72 traditions can aid in assessing these various hypotheses.

1.1. The French Middle Gravettian

In France, the Middle Gravettian occurs between *ca.* 32-28.5 ky cal. BP and is defined by two *faciès*, termed the “Noaillian” and the “Rayssian” that are characterized principally on the basis of their lithic industries (Touzé, 2013). The term “*faciès*” (in French) is used to describe an archaeological entity according to “the nature of the considered remains and the method employed to study them” (Touzé, 2013, p. 397). This neutral term is especially useful in the case of the Middle Gravettian, since the two *faciès* are not defined equally. The Noaillian is a typological *faciès*, defined solely by the presence of Noailles burins (Bardon & Bouyssonies, 1903; Tixier, 1958) and sometimes by the presence of bone or antler points called “Isturitz points”, although this point type’s precise chrono-cultural status remains uncertain (Goutas, 2013b). On the other hand, the Rayssian is a typo-technological *faciès*, defined by a reduction method aimed at removing, from Raysse burins, bladelets that were used as implements on projectiles (Figure 1, Movius & David, 1970; Klaric *et al.*, 2002; Lucas, 2002; Pottier, 2006; Klaric, 2017). Furthermore, Rayssian bladelet and blade reduction sequences show strong conceptual parallels and technical similarities (Klaric, 2003; Klaric, 2008).




Figure 1. Diagnostic Middle Gravettian artifact types. **a.** Noailles burin from Fourneau du Diable, Dordogne, France (drawing: A.Vignoles). **b.** Technological position of a “Picardie” bladelet on a Raysse burin-core from La Picardie, Indre-et-Loire, France (from Klaric, 2008)

The chronological relationship between the Noaillian and the Rayssian has yet to be determined with precision. This is due to the fact that very few contextually reliable ^{14}C ages are associated with these two *faciès*, and for regions north of the Garonne River the low number of available ages renders any chronological comparison between the two phases uninformative at present (Banks *et al.*, 2019). This situation is complicated by the fact that Noailles burins and the Raysse method are frequently found together within the same archaeological layer in the region of overlap. Past and on-going studies suggest that, at many sites, this association is not culturally meaningful due to imprecise excavation methods and/or disturbed stratigraphic contexts (e.g. Klaric, 2003, 2007; Vignoles *et al.*, 2019). However, in a few stratified contexts, the development of the Raysse method is always stratigraphically younger than the Noaillian (e.g., Abri Pataud and Flageolet I sites; David, 1985; Rigaud, 1982; Klaric, 2003). An exception to this statement is found at Les Jambes, a site where Noailles burins are described as being stratigraphically *above* the Raysse burins (Célérier, 1967). This configuration, though, remains to be verified. First, the two levels identified by Célérier have been described as part of a slope deposit which raises doubts as to the integrity of the levels. Moreover, stratigraphic projections of artifacts show that the levels defined by Célérier correspond to a single archaeological layer.

Finally, an ongoing reexamination (conducted by A.V) of the site's assemblage demonstrates that most of the "Noailles burins" do not correspond to the classic typological definition. In fact, only one artifact can be considered a typical Noailles burin, while all the others are highly atypical. Technological characterization of the blade/bladelet reduction sequences may provide new data with which to discuss the presence of Gravette points—a class of tool traditionally associated with the Noaillian rather than the Rayssian—at Les Jambes.

Numerous hypotheses have been proposed to explain the co-occurrence of Noaillian and Rayssian materials in the same archaeological level, such as a gradual replacement of the Noaillian by the Rayssian (David, 1985; Pottier, 2005), differing site activities (Laville & Rigaud, 1973; Rigaud, 2008, 2011), the use of different typo-technological traditions within a broad regional population (Touzé, 2013) or the result of post-depositional mixing or the inability of old excavation methods to differentiate between discrete occupations (Klaric, 2003, 2007; Vignoles *et al.*, 2019). Unfortunately, taphonomic evaluations of individual sites are not yet sufficiently numerous to evaluate these hypotheses adequately (Klaric, 2003, 2007; Pottier, 2005; Agsous, 2008; Michel, 2010; Gottardi, 2011).

With respect to geography, these two archaeological traditions have only partially overlapping distributions (Figure 2). The Noaillian is observed in regions south of the Loire River, as well as a very isolated presence in the Vosges region, with extensions into Cantabrian Spain and the Italian Peninsula. The Rayssian is restricted to a smaller geographic area situated between the Garonne River and the southern portion of the Paris Basin, with extensions into Burgundy and Brittany (Klaric, 2003, 2017; Touzé, 2013). Despite mentions of the presence of Raysse burins at the sites of La Carane-3, Isturitz and Tuto de Camalhot by David (1985), the presence of the Raysse method south of the Garonne River has not been unequivocally demonstrated: none of these artifacts have been described or pictured, nor do recent technological studies (e.g. Simonet, 2009a) mention them. The demonstration of the Raysse

method relies on precise technical criteria and more specifically on the identification of the bladelet component associated with Raysse burin-cores, and David (1985) did not use such criteria to describe the Late Noaillian (i.e. Rayssian) in these southern contexts. It is also important to point out that look-alike artifacts (*faux-amis*) have been described at Brassempouy (Klaric, 2006). The examples reported by David could therefore be misleading in the same way.

The absence of the Raysse method south of the Garonne River suggests that this valley may have played a role in the separation of the two different technological trajectories. This is also paired with the fact that the Noaillian in the Pyrenees appears to have lasted as long as the entire Middle Gravettian phase (Noaillian and subsequent Rayssian) present north of the Garonne River (Touzé, 2013; Klaric, 2017; Banks *et al.*, 2019). This pattern suggests that the environment may have played a role in the development of the cultural adaptation that serve to define the Rayssian *faciès* (David, 1985; Djindjian *et al.*, 1999).

1.2. Research question and approach

Here, we test the hypothesis that the typo-technological differences observed on either side of the Garonne River valley during the Middle Gravettian reflect the exploitation of different environmental conditions via different technological (i.e. cultural) adaptations. The application of Ecological Niche Modeling (ENM) methods to the archaeological record is one means with which to conduct such a test (Banks, 2017; d'Errico & Banks, 2013). ENM (the terminology employed in this study, cf. Peterson & Soberón, 2012; Warren, 2012) provides a means for estimating the ecological niches of past hunter-gatherer populations, employing archaeological sites as occurrence data and environmental variables derived from high-resolution paleoclimatic simulations. These data are then used by predictive modeling algorithms to identify sets of environmental parameters associated with known archaeological sites and create an estimation of the presence of suitable environmental conditions (i.e. the ecological niche) across the study area. Niche estimations can be compared with one another in order to

characterize and evaluate potential differences between niches (e.g. Warren *et al.*, 2008). The use of these tools has been demonstrated to be a valuable approach for assessing culture-environment relationships of past hunter-gatherer populations, both synchronically and diachronically (e.g. Banks *et al.*, 2008, 2009, 2011, 2013; d’Errico *et al.*, 2017). Research focused on material culture variability observed in the archaeological record is often polarized between two sets of explanations, cultural vs. functional, mirroring the well-known Bordes-Binford debate concerning Mousterian lithic industries. An advantage of ENM is that it allows one to approach evaluations of such variability from a different angle by taking into account the ecological niches exploited by archaeological hunter-gatherer populations. For example, comparisons of the ecological niches associated with two archaeological cultures or lithic industries allow one to evaluate to what extent cultural (i.e. technological) differences have an ecological basis. In such scenario, one would expect their ecological niches to differ from one another significantly due to the fact that the different typo-techno complexes represent adaptations to distinct ecological niches more than purely social factors. If their niches are highly similar, one could interpret technological variability to be more the result of internal cultural dynamics that led to the development of two cultural trajectories. As stated earlier, multiple hypotheses have been proposed to explain the typo-technological variability observed within the French Middle Gravettian and no consensus has been reached. The objective of this study is to provide new data on the ecological niches of these hunter-gatherer populations in order to evaluate to what extent ecological factors may have played a role in the observed material culture variability.




Figure 2. Sites where Noailles burins and/or Rayssian burins have been found (after Touzé, 2013; Klaric, 2017; the present study). Sites are labeled according to their number presented in Table 1 (cf. 2.2.1). Topographic background: <http://www.natureearthdata.com>

2. Materials and Methods

2.1. Conceptual framework of ecological niche modelling

The conceptual framework of ENM is based on Hutchinson's (1957) definition of the fundamental niche (N_F): an n -dimensional hypervolume whose dimensions are the non-interactive environmental variables (i.e. scenopoetic variables) necessary for a species to maintain populations indefinitely without immigrational subsidy. Following Peterson and Soberón (2005, 2012), we consider the Biotic-Abiotic-Mobility framework (**BAM**, Figure 3) to describe factors constraining geographic distribution of species. The projection of N_F in geographic space (**G**), i.e. the geographic localities corresponding to N_F , identifies areas with conditions favorable to the species (**A**). However, the geographic distribution of a species can

be constrained by at least two other types of factors: biotic interactions (**B**), i.e. the species' positive or negative interactions with other species or resources that are present, and the areas that have been physically accessible to the species over a relevant period of time (**M**). The intersection of **A** and **B** is the potential distributional area (G_P), which is the geographic expression of the realized niche (N_R) defined by Hutchinson (1957). In this study, we focus on N_F , defined solely on the basis of non-interactive variables, following the Eltonian-noise hypothesis, which argues that biotic interactions may often be manifested at fine spatial resolutions and thus may not have a significant or limiting effect on a species' distribution at broad geographic scales (Soberón, 2007). Finally, the intersection of G_P with **M** defines the occupied distributional area (G_O). In environmental space, the intersection between N_F and the environments associated with **M** define the existing fundamental niche (N_F^*), which is the portion of the fundamental niche that is actually observable in nature (Peterson & Soberón, 2012).

When applied to the archaeological record, the goal is to identify the sets of environmental conditions associated with a cultural trait or with a techno-complex, and evaluate their eventual co-variability through time (Banks, 2017). Furthermore, with respect to examinations of culture-environment relationships and cultural adaptations, it is pertinent to evaluate to what extent an archaeological typo-technological complex (archaeological culture) occupied its existing niche (i.e. G_O vs. observed distribution).




Figure 3. BAM diagram representing the factors that constrain the geographic distribution of a species at broad geographic scales, if the Eltonian-noise hypothesis holds true (after Soberón & Peterson, 2005, 2012; modified). Circles represent the different factors and black dots represent the observed distribution of the species. **G**: geographic space; **A**: non-interactive variables; **B**: biotic interactions; **M**: areas accessible to the species; **G_O**: occupied distributional area.

2.2. Data

2.2.1. Archaeological data

Occurrence data consist of the geographic coordinates of archaeological sites where Noaillian and/or Rayssian material culture assemblages have been identified (Figure 4). These data were assembled through a critical examination of the literature, although one must keep in mind that this approach has certain limitations. First, most sites were excavated and studied in the late 19th and first half of the 20th century, sometimes in an expeditious manner. Due to the fact that excavated sediments were rarely sieved and often only large, diagnostic tools were kept, many assemblages are biased and do not necessarily contain artifacts that allow the two typo-technological *faciès* to be reliably recognized, since their diagnostic artifacts are of small

size (e.g., Raysse and Picardie bladelets, some Noailles burins, Noailles burins spalls), and/or correspond to flint knapping by-products (e.g., Raysse bladelets, blades with oblique lateralized faceted platforms). As a result, there are numerous sites where one or both of these Middle Gravettian *faciès* was not initially recognized (e.g., Fourneau du Diable, Laussel or Combe-Saunière I; Klaric, 2017; Vignoles *et al.*, 2019) and this is likely the case for many others. Therefore, the corpus of sites associated with these two archaeological *faciès* should be considered incomplete at present. Furthermore, many assemblages, even those that were rather well-excavated (piece-plotted artefacts, sieved archaeological sediments, collection of unretouched artifacts) often have not been subjected to recent contextual examinations or typotechnological re-evaluations (e.g., Les Jambes, Le Facteur). This is especially a problem for the bibliographic identification of the Raysse method. Although the Raysse burin type was first described in the 1950s (Pradel, 1953; Couchard & de Sonnevile-Bordes, 1960; Movius & David, 1970), its function as a core for producing standardized projectile point implements was only demonstrated in the early 2000s (Klaric *et al.*, 2002; Lucas, 2002). It is therefore necessary to reconsider, from a technological standpoint, all previously identified Raysse burins and to identify the presence of the associated bladelet component in order to avoid attributing archaeological levels to this *faciès* on the basis of look-alike (*faux-amis*) artifacts (Klaric, 2003, 2006).

Another problem is the unequal definition of the Noaillian (i.e. sole presence of Noailles burins in an assemblage) and the Rayssian (Touzé, 2013). The latter's technical system is relatively well-described across its area of expression (Lucas, 2000, 2002; Klaric, 2003, 2017; Pottier, 2005, 2006; Guillermin, 2006; Touzé, 2011, 2013; Gottardi, 2011; Sarrazin, 2017, 2018). Variability in the use of the Raysse method has been attributed mainly to blank selection, the availability of suitable lithic raw materials, levels of technological expertise and contingencies of the reduction sequence (Klaric *et al.*, 2009; Klaric, 2017, 2018). To the

260 contrary, the technical system associated with Noailles burins has only been the subject of
261 isolated studies in the Landes (e.g. Klaric, 2003; Simonet, 2009a, 2011a; Lacarrière *et al.*,
262 2011), the Pyrenees piedmont and plateau (e.g. Foucher, 2004; Simonet, 2009a), and to lesser
263 extents the Perigord region (Lucas, 2000; Pottier, 2005) and the southern Paris Basin (Kildea
264 & Lang, 2011), thus rendering evaluations of its homogeneity difficult. Typo-technological
265 studies conducted on assemblages from Cantabrian Spain (e.g. de la Peña-Alonso, 2011), the
266 French Mediterranean coast (e.g. Onoratini, 1982; Santaniello, 2016), and the Italian peninsula
267 (e.g. Onoratini, 1982; Aranguren *et al.*, 2006, 2015; Simonet, 2010; Santaniello, 2016;
268 Santaniello & Grimaldi, 2019) have employed different methodological approaches for
269 characterizing reduction sequences (i.e. *chaînes opératoires*) and typologies, with a few
270 exceptions (e.g. Simonet, 2010), thus making comparisons to sites studied via the methods
271 traditionally used in France difficult. An additional obstacle is related to this record's
272 chronology. Noailles burins have been recovered from Cantabrian contexts that are
273 contemporaneous with the Late Aurignacian in southwestern France, as well as contexts that
274 extend into the Solutrean (de la Peña-Alonso, 2011, p. 681). Noailles burin contexts in Italy are
275 also interpreted as being younger than those in France (Touzé, 2013), but the majority of their
276 associated radiocarbon ages were produced decades ago (non-AMS) and evaluations of their
277 archaeological association are lacking, thus rendering any comparison to the French
278 archaeological record unreliable at present.




Figure 4. Noaillian and Rayssian site locations used for calibration and the study's defined accessible area (**M**) hypothesis. Topographic background: <http://www.naturalearthdata.com>; Coastlines: Siddall *et al.*, 2003; Glacier extent: Ehlers & Gibbard, 2004; Periglacial cold deserts: Bertran *et al.*, 2013; Bosq *et al.*, 2018

Taking into account these limitations, we constrain our analysis to the comparison of two adjacent regions that are thought to represent coherent territories during the Middle Gravettian: 1) the Pyrenees piedmont and plateau, based on assemblages that are regionally coherent with respect of lithic typo-technology (i.e. contexts that contain Noailles burins and in which the Raysse method is absent) and raw material circulation for the entirety of the Middle Gravettian (Foucher *et al.*, 2008; Foucher, 2013; Simonet, 2009a, 2017; Banks *et al.*, 2019), and 2) the

area constrained along its southern margin by the Garonne River Valley and by the most northerly sites where the “Raysse method” has been observed (Klaric, 2017). The latter territory is characterized by both the Noaillian and the Rayssian *faciès*, and the lack of precision, at present, concerning their chronological relationship required that we group them together, mirroring the approach employed by Banks *et al.* (2019).

In an effort to retain only sites for which a reliable techno-typological attribution could be made, published studies were carefully evaluated with respect to the pertinence of the data they contained (see Appendix). A source was considered pertinent if it provided precise counts of tool types based on the most recent typo-technological definitions (e.g. for the “Noailles burin” type: Bardon & Bouyssonies, 1903; Tixier, 1958; for “Picardie bladelet” type and “Raysse burin-core”: Klaric *et al.*, 2002; Klaric, 2003; Klaric, 2017), preferably supplemented with artifact drawings or photos. We also included personal observations made during the course of on-going and yet-to-be-published studies (L. Klaric and A. Vignoles). We, thus, did not retain sites for which one or both *faciès* were only suspected to be present in order to avoid potentially aberrant attributions based on look-alike artifacts. For example, “Raysse burins” reported in Brassempouy or Le Gratadis have been shown to not be actual Raysse burin-cores aimed at the production of La Picardie bladelets (Klaric, 2003, 2006). Bearing this in mind, we did not retain sites where Raysse burins were only mentioned with no demonstration of the presence of the Raysse method (e.g. Abri André Ragout; David, 1985). Also, sites where Raysse burins are depicted in published drawings (Klaric, 2003) but that have not been confirmed by us via direct observation and where the presence of the associated bladelet component has not yet been evaluated, such as the site of Roc de Gavaudun in Lot-et-Garonne or Lespoux shelter in Gironde (Monméjean *et al.*, 1964; Krtolitz & Lenoir, 1998), were not included. Finally, we also eliminated sites for which the presence of Raysse burin cores is not consensual, such as Chamvres in Yonne (Klaric, 2003 vs. Sarrazin, 2018). In the same vein, we did not retain sites

where reported “Noailles burins” have been demonstrated to be look-alike artifacts, such as the site of Les Jambes (ongoing examination by A.V.). Moreover, sites where Noailles burins are all atypical, such as Peutille in Lot-et-Garonne (Morala, 1984), or sites where only one or two Noailles burins are reported and their presence is not supported by published drawings, such as La Verpillière I cave in Saône-et-Loire (Floss *et al.*, 2013), were eliminated from consideration. In the end, we retained 74 sites for our analysis. North of the Garonne River, there are 9 Rayssian sites, 40 Noaillian sites, as well as 13 additional sites that yield both *faciès*. In the Pyrenees piedmont and plateau, 12 Noaillian sites were retained (Figure 4; Table 1).

Table 1. Sites retained or rejected based on a critical review of the literature, along with their geographic coordinates and references. A site was assigned to a specific data set or rejected based on criteria detailed in the Appendix. Geographic coordinates correspond to the commune (i.e. town administrative territory) in which the site is localized. Although these coordinates do not necessarily correspond to the site itself, their resolution is more than adequate considering the 11.5 km-grid resolution of the environmental data.

N°	Site	Lat. (N)	Long. (E)	Noailles burins	Rayssé method	Data set	References
1	Plasenn-al-Lomm	-3,00	48,85	absence	presence	Northern Middle Gravettian	Le Mignot, 2000; Klaric, 2003; Sarrazin 2018
2	La Martinière	-0,86	47,36	presence	suspected	Northern Middle Gravettian	Allard, 1986
3	Roc-en-Pail	2,29	48,86	insufficient	absence	<i>rejected</i>	Allard & Gruet, 1976; Gruet, 1984; Hinguant & Monnier, 2013
4	Le Taillis des Coteaux	-0,77	46,62	absence	presence	Northern Middle Gravettian	Klaric, 2017
5	La Croix-de-Bagneux	1,33	47,29	presence	absence	Northern Middle Gravettian	Kildea & Lang, 2011, 2013
6	La Picardie	0,93	46,86	absence	presence	Northern Middle Gravettian	Klaric, 2003; Klaric <i>et al.</i> , 2011; Klaric <i>et al.</i> , 2018
7	Abri Charbonnier	1,04	46,68	absence	presence	Northern Middle Gravettian	Aubry <i>et al.</i> , 2013; Klaric, pers. obs.
8	Abri Laraux	0,73	46,40	presence	suspected	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
9	Chamvres	3,36	47,96	absence	controversial	<i>rejected</i>	Klaric, 2003, 2013; Sarrazin, 2018
10	Grotte de la Verpillière I	4,73	48,81	insufficient	absence	<i>rejected</i>	Floss <i>et al.</i> , 2013
11	Grotte du Renne	3,76	47,60	absence	presence	Northern Middle Gravettian	Klaric, 2003

Table 1 - continued

N°	Site	Lat. (N)	Long. (E)	Noailles burins	Raysse method	Data set	References
12	Grotte du Trilobite	3,76	47,60	absence	presence	Northern Middle Gravettian	Klaric, 2003; David, 1985
13	Hautmougey	6,26	48,00	presence	absence	Northern Middle Gravettian	Hans, 1997
14	Abri André Ragout	0,42	45,68	presence	insufficient	Northern Middle Gravettian	Tixier, 1958; David, 1985
15	Abri du Chasseur	0,42	45,68	presence	absence	Northern Middle Gravettian	Tixier, 1958; David, 1985
16	Les Vachons	0,12	45,51	presence	suspected	Northern Middle Gravettian	David, 1985; Fontaine, 2006
17	Le Petit-Puyrousseau	0,72	45,19	presence	insufficient	Northern Middle Gravettian	de Sonneville-Bordes, 1960; Daniel, 1967; David, 1985
18	Abri Durand-Ruel	0,65	45,37	presence	suspected	Northern Middle Gravettian	de Sonneville-Bordes, 1960; Daniel & Schmider, 1972; David, 1985
19	Gisement de la Chèvre	0,59	45,32	presence	insufficient	Northern Middle Gravettian	David, 1985; Arambourou & Jude, 1964
20	Combe Saunière	0,87	45,24	presence	presence	Northern Middle Gravettian	Klaric, 2017; Klaric, pers. obs.
21	Le Fourneau du Diable	0,59	45,32	presence	presence	Northern Middle Gravettian	David, 1985; Klaric, 2003, 2017; Vignoles <i>et al.</i> , 2019
22	Les Jambes	0,72	45,19	unlikely	presence	Northern Middle Gravettian	Célérier, 1967; Vignoles, pers.obs.
23	Solvieux	0,39	45,06	presence	presence	Northern Middle Gravettian	David, 1985; Sackett, 1999; Klaric, 2003
24	Abri Sous-le-Roc	1,09	45,01	insufficient	absence	<i>rejected</i>	de Sonneville-Bordes, 1960; David, 1985
25	Abri de Fongal	1,08	44,99	insufficient	insufficient	<i>rejected</i>	de Sonneville-Bordes, 1960, p.97; David, 1985
26	Abri des Merveilles	1,11	45,00	suspected	absence	<i>rejected</i>	Delage, 1936; de Sonneville-Bordes, 1960; David, 1985
27	La Rochette	1,09	45,01	presence	suspected	Northern Middle Gravettian	de Sonneville-Bordes, 1960; Schmider, 1969; David, 1985; Klaric, 2003, p.220
28	Abri Labattut	1,11	45,00	presence	insufficient	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
29	Masnaigre	1,14	44,94	presence	suspected	Northern Middle Gravettian	David, 1985
30	Abri Pagès	1,04	44,97	presence	absence	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
31	Abri du Poisson	1,01	44,94	presence	absence	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
32	Grotte d'Oreille d'Enfer	1,01	44,94	presence	absence	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985; Pradel, 1959

Table 1 - continued

N°	Site	Lat. (N)	Long. (E)	Noailles burins	Rayssé method	Data set	References
33	La Ferrassie	0,95	44,97	presence	absence	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
34	Abri Pataud	1,01	44,94	presence	presence	Northern Middle Gravettian	David, 1985; Bricker (dir.), 1995; Pottier, 2005, 2006; Nespoulet, 2008
35	La Roque Saint-Christophe	1,08	44,99	presence	presence	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985; Vignoles, pers. obs.
36	Abri du Facteur	1,04	44,97	presence	presence	Northern Middle Gravettian	Delporte, 1968; David, 1985; Vignoles, pers. obs.
37	Grand-abri de Laussel	1,14	44,94	presence	presence	Northern Middle Gravettian	Roussot, 1985; David, 1985; Klaric, 2017; Klaric, pers. obs.
38	Grotte Maldidier	1,18	44,83	presence	presence	Northern Middle Gravettian	Klaric, 2017; Caux <i>in</i> Boudadi-Maligne, 2012
39	Le Flageolet I	1,09	44,84	presence	presence	Northern Middle Gravettian	Rigaud, 1982; David, 1985; Lucas, 2000; Gottardi, 2011
40	Cantelouve	1,28	44,84	insufficient	absence	<i>rejected</i>	Tixier, 1958; David, 1985
41	Grotte de Péchialet	1,29	44,82	suspected	absence	<i>rejected</i>	David, 1985; Breuil, 1927
42	Grotte du Bos-del-Ser	1,54	45,16	insufficient	insufficient	<i>rejected</i>	David, 1985
43	Grotte "Chez Serre"	1,53	45,10	presence	absence	Northern Middle Gravettian	David, 1985
44	Roc de Combe	1,40	45,04	presence	absence	Northern Middle Gravettian	David, 1985
45	Grotte Lacoste	1,54	45,16	presence	absence	Northern Middle Gravettian	David, 1985
46	Grotte de Champ	1,53	45,16	presence	insufficient	Northern Middle Gravettian	David, 1985; Daniel, 1969
47	Grotte Thévenard	1,54	45,16	presence	insufficient	Northern Middle Gravettian	David, 1985
48	La Font-Robert	1,54	45,16	presence	insufficient	Northern Middle Gravettian	de Sonneville-Bordes, 1960; David, 1985
49	Pré-Aubert	1,54	45,16	presence	suspected	Northern Middle Gravettian	David, 1985; Demars, 1977
50	Grotte Bouysonie	1,54	45,16	presence	presence	Northern Middle Gravettian	Touzé, 2011; Klaric, 2017; Klaric, pers. obs.
51	Bassaler-Nord	1,54	45,16	presence	presence	Northern Middle Gravettian	David, 1985; Touzé, 2011
52	Abri du Rayssé	1,54	45,16	presence	presence	Northern Middle Gravettian	David, 1985; Touzé, 2011
53	Les Morts	1,54	45,16	presence	presence	Northern Middle Gravettian	David, 1985; Sarrazin 2017
54	Les Fieux	1,71	44,85	absence	presence	Northern Middle Gravettian	Guillermin, 2006, 2008
55	Les Artigaux	-0,27	44,79	unlikely	presence	Northern Middle Gravettian	Lenoir, 1977; Klaric, 2003; Vignoles, pers. obs.

Table 1 - *continued*

N°	Site	Lat. (N)	Long. (E)	Noailles burins	Raysse method	Data set	References
56	Abri Lespaux	-0,29	44,82	presence	suspected	Northern Middle Gravettian	David, 1985; Krtoliza & Lenoir, 1998
57	Le Caillou	0,45	44,78	presence	absence	Northern Middle Gravettian	Boyer <i>et al.</i> , 1984
58	Roc de Combe-Capelle	0,82	44,77	insufficient	absence	<i>rejected</i>	de Sonneville-Bordes, 1960; David, 1985
59	Grotte du Roc de Vézac	2,52	44,89	insufficient	absence	<i>rejected</i>	Rigaud, 1982, p.262; Aujoulat <i>in</i> Leroi-Gourhan, 1984
60	Abri des Peyrugues	1,67	44,54	absence	insufficient	<i>rejected</i>	Guillermin, 2011
61	Abri de la Bergerie	1,58	44,48	presence	absence	Northern Middle Gravettian	Clottes <i>et al.</i> , 1990
62	Les Battuts	1,73	44,08	presence	suspected	Northern Middle Gravettian	Alaux, 1967; Alaux 1971; David, 1985
63	Grotte de Rouzet	1,69	44,00	presence	absence	Northern Middle Gravettian	Foucher <i>et al.</i> , 2008
64	Abri du Couvert	1,02	44,49	presence	absence	Northern Middle Gravettian	Morala, 1984
65	Abri Peyrony	0,89	44,56	presence	absence	Northern Middle Gravettian	Le Tensorer, 1981; David, 1985
66	Guiraudel	0,95	44,54	presence	absence	Northern Middle Gravettian	Morala, 1984
67	Las Pélénos	0,93	44,50	presence	absence	Northern Middle Gravettian	Morala, 1984
68	Le Callan	0,97	44,60	presence	absence	Northern Middle Gravettian	Morala, 2011
69	Le Roc de Cavart	1,07	44,54	presence	absence	Northern Middle Gravettian	Le Tensorer, 1981; David, 1985
70	Station du Fresquet	0,94	44,47	presence	absence	Northern Middle Gravettian	Morala, 1984
71	Gisement du château	1,01	44,59	suspected	absence	<i>rejected</i>	Le Tensorer, 1974, p.467
72	Peutille	0,97	44,58	suspected	absence	<i>rejected</i>	Morala, 1984
73	Termo-Pialat	0,82	44,77	suspected	absence	<i>rejected</i>	de Sonneville-Bordes, 1960; David, 1985
74	Métayer	0,89	44,56	presence	insufficient	Northern Middle Gravettian	Le Tensorer, 1981; David, 1985
75	Plateau Baillard	0,89	44,56	presence	absence	Northern Middle Gravettian	Le Tensorer, 1981; David, 1985
76	Roquecave	0,89	44,56	presence	absence	Northern Middle Gravettian	Le Tensorer, 1974; Le Tensorer, 1981
77	Le Roc de Gavaudun	0,89	44,56	presence	suspected	Northern Middle Gravettian	de Sonneville-Bordes, 1960; Monméjean <i>et al.</i> , 1964; Le Tensorer, 1981; David, 1985
78	Hin-de-Diou	-0,33	43,86	presence	absence	Pyrenees Noaillian	Briand <i>et al.</i> , 2010
79	Brassempouy	-0,69	43,63	presence	absence	Pyrenees Noaillian	Klaric, 2003; Foucher <i>et al.</i> , 2008; Simonet, 2009a

Table 1 - continued

N°	Site	Lat. (N)	Long. (E)	Noailles burins	Raysse method	Data set	References
80	Isturitz	-1,20	43,35	presence	insufficient	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a; Lacarrière <i>et al.</i> , 2011
81	Lézia	-1,58	43,31	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a
82	Gatzarria	-0,92	43,14	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a
83	Grotte des Rideaux	0,67	43,23	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a
84	Gargas	0,52	43,07	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008, 2012
85	Bois de Touaa	0,83	43,07	presence	absence	Pyrenees Noaillian	Foucher <i>et al.</i> , 2008; Clottes, 1985, p. 346
86	Tuto de Camalhot	1,13	43,01	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a
87	Grotte d'Enlène	1,20	43,02	presence	absence	Pyrenees Noaillian	Foucher <i>et al.</i> , 2008; Simonet, 2009a
88	Tarté	0,99	43,12	presence	absence	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a
89	La Carane-3	1,61	42,96	presence	insufficient	Pyrenees Noaillian	David, 1985; Foucher <i>et al.</i> , 2008; Simonet, 2009a

2.2.2. Environmental predictors

In order to employ the appropriate environmental data, it is paramount to determine accurately the precise chronology of the target archaeological culture so that it can be correlated to the appropriate climatic event or events (Banks, 2015; Banks *et al.*, 2019). Based on the results presented by Banks *et al.*, (2019), the Middle Gravettian begins between the latter part of Greenland Interstadial 5.2 (GI 5.2) and Greenland Stadial 5.2 (GS 5.2) and ends between the latter part of GS 5.1 and GI 4. It was thus present during GS 5.2 and 5.1, during which occurred Heinrich Event 3 (HE3), as well as GI 5.1. Considering the lack of chronological resolution facing this portion of the Upper Paleolithic, two important assumptions were made for the choice of the best possible environmental data. First, since most Middle Gravettian occupations have not been dated or are not associated with reliable dates, we must assume that all sites date

to some point between *ca.* 32 and 28.5 ky cal. BP (after Banks *et al.*, 2019). Second, the chronological resolution of individual radiocarbon measurements does not allow us to determine whether a specific dated occupation is associated with a specific Interstadial or Stadial (GI 5.1 or GS 5.1 for example). This limitation makes it impossible to create separate occurrence data sets such that correspond to specific Stadial or Interstadial events. Therefore, we choose environmental data such that correspond to stadial conditions since those were present during the majority of the period concerned by this study (*ca.* 2.5 ky for GS 5.2 and 5.1 combined vs. *ca.* 0.5 ky for GI 5.1).

We employed as environmental background climatic variables derived from a high-resolution paleoclimatic simulation obtained with the Atmospheric Global Circulation Model (AGCM) LMDZ5A (Hourdin *et al.*, 2013). It was run with a zoomed grid permitting a spatial resolution of *ca.* 50 km over Europe, following Sima *et al.* (2009, 2013). We used the same coastlines and ice-sheet configuration as Sima *et al.* (2009, 2013), i.e. those corresponding to the Last Glacial Maximum (LGM; Figure 4). Atmospheric greenhouse gas concentrations were also left at their LGM values (CO₂ = 185 ppm, CH₄ = 350 ppb, N₂O = 200 ppb). The orbital parameters are set to 32 ky cal. BP (Berger *et al.*, 1978), which correspond to the beginning of GS 5.2. Initial boundary conditions for prescribed sea surface temperatures (SSTs) and sea ice cover were computed with the coupled atmosphere-ocean general circulation model IPSLCM4 (Marti *et al.*, 2010). This model was obtained by setting forcing and boundary conditions of the PMIP3 LGM experiment described in Alkama *et al.* (2008) and Kageyama *et al.* (2013) to 32 ky cal. BP, thus producing an enhanced seasonal cycle of incoming insolation and surface temperatures in the Northern hemisphere.

Results were further downscaled to a spatial resolution of 11.5 km via the spline interpolation available in ArcMap 10.7.1. The simulated variables employed in this analysis are mean annual precipitation, warmest month temperature and coldest month temperature. We did

not use elevation as a covariate in this process. Elevation is usually considered to be a proxy for temperature, because these two variables are highly correlated. Moreover, elevation is also taken into account during the paleoclimatic simulation process. These reasons make it a redundant and non-informative variable for the niche analyses presented here.

2.3. Ecological niche modeling

2.3.1. Modeling preparation

Prior to estimating niches, we analyzed and filtered the occurrence datasets to reduce potential spatial biases. First, we trimmed duplicate site occurrences from each grid-cell, so that a grid-cell contained only a single occurrence point, thus ensuring that the training and testing occurrence datasets would be spatially unique (i.e., no shared occurrences). Next, we thinned the occurrence datasets to eliminate clusters of sites, thereby preventing oversampling of environmental conditions from certain areas (e.g., the northern Aquitaine area) and reducing potential spatial autocorrelation (Anderson & Gonzalez, 2011; Boria *et al.*, 2014). This consisted of eliminating occurrences such that the minimum distance between any pair of occurrence points was twice the grid resolution, i.e. *ca.* 23 km. This step was performed manually using the “Measure line” tool in QGIS 2.18.28. The final datasets consisted of 10 occurrence points for the Noaillian in the Pyrenees piedmont and plateau, and 20 occurrences for the Middle Gravettian north of the Garonne River (Noaillian and Rayssian combined).

The definition of a calibration area (**M**) relies on biogeographic assumptions (Peterson *et al.*, 2011 p. 135; Barve *et al.*, 2011). To define (**M**) for Middle Gravettian hunter-gatherers in our region of study (Figure 4), we assume that these populations could not live in regions in close proximity to ice sheets, and that they could have occupied areas exposed by the period’s lower sea levels. We therefore masked the environmental variables with coast lines 90 m below

present day sea level (Waelbroeck *et al.*, 2002; Siddall *et al.*, 2003), as well as with LGM ice sheet coverage reconstructions in the Alps, Pyrenees and the Massif Central (Ehlers & Gibbard, 2004). While these reconstructions slightly over-estimate ice coverage for *ca.* 32 ky cal. BP, they still serve as an adequate proxy since the areas in question would have been characterized by cold and dry, if not periglacial, conditions during the corresponding stadials and HE. Furthermore, based on the nature of raw material (flint) circulation in the Pyrenees region (Foucher *et al.*, 2008; Simonet, 2017), we allowed the predictive modeling architecture to extrapolate into the regions of Cantabria and Catalonia. Finally, the Rhône River Valley in the East may have served to limit the movements of hunter-gatherer populations (no raw material circulation across the valley, cf. Santaniello, 2016). Furthermore, recent geomorphological studies show that the lower and middle Rhône Valley, as well as the Mediterranean continental shelf, likely consisted of a desert, with deflation-related landforms (e.g., yardangs, pans, desert pavements) and sand deposits (dunes, sand ramps) surrounded by loess accumulations during the coldest events of the Last Glacial period (Bosq *et al.*, 2018). However, the presence of Noailles burins and Gravette points to the east indicates that they were permeable barriers. Thus, we included in our (**M**) coastal regions of Liguria, Tuscany, Lazio and Campania in present-day Italy where Noailles burins are observed in the archaeological record (Palma di Cesnola, 1991; Touzé, 2013). This step was performed with QGIS 2.18.28.

2.3.2. Model calibration and selection

Model calibration and selection were performed using the *kuenm* R package (Cobos *et al.*, 2019a), which employs Maxent 3.4 (Phillips *et al.*, 2006, 2017). In R 3.6.1 (R Core Team, 2019), we created a total of 448 candidate models for each occurrence dataset using distinct parameter settings resulting from the combination of 16 regularization multiplier values, 7 response types representing all possible combinations of the three feature classes (linear, quadratic and product), and four sets of environmental predictors derived from all possible

combinations of the three paleoclimatic variables, following Cobos *et al.* (2019b). Threshold and hinge feature classes were not used in order to reduce model complexity and overfitting. The kuenm package allows the evaluation of statistical significance via partial ROC measures (Peterson *et al.*, 2008), omission rates based on a maximum allowed error ($E = 5\%$, user defined; Anderson *et al.*, 2003; Peterson *et al.*, 2008), and model complexity by means of the Akaike information criterion corrected for small sample sizes (AICc; Warren & Seifert, 2011). For the retention of the best performing models among those that were statistically significant, we selected those with omission rates lower than E and of those we retained only the models with ΔAICc values lower than two (i.e. $\Delta\text{AICc} = \text{AICc}_i - \text{AICc}_{\min}$, where AICc_i is the AICc of the i th model and $\text{AICc}_{\min}$ is the lowest AICc among all significant models for which omission rates are below 5%).

2.3.3. Creation of final models and model comparisons

Using kuenm, we created final models within (**M**) using the parameter settings selected after model calibration (Table 2). As more than one “best parameter” setting was used to create the final models for the Pyrenees Noaillian, we created a consensus model of all results across all parameterizations by calculating a median model, following Cobos *et al.* (2019c). We also calculated the range from these final models as a variability index.

439 **Table 2.** Models calibration results. Only one set of parameters was retained for the northern Middle
440 Gravettian model, whereas 45 sets of parameters were relevant to the Pyrenees Noaillian model.
441 Parameters corresponding to the models compared via the background similarity and identity tests are
442 indicated in bold. L: linear; Q: quadratic; P: product; ctemp: temperature of the coldest month; wtemp:
443 temperature of the warmest month; mprec: mean annual precipitation.

Parameters			Evaluation results				
Regularization multiplier	Features	Environmental variables	Partial ROC	Omission rates below 5%	AICc	Δ_{AICc}	Number of parameters
<i>Northern Middle Gravettian</i>							
0.3	LQP	All	0	0	311.258	0	4
<i>Pyrenees Noaillian</i>							
0.9	QP	ctemp, mprec	0	0	170.583	0	1
1	QP	ctemp, mprec	0	0	170.680	0.098	1
2	Q	ctemp, mprec	0	0	170.680	0.098	1
3	Q	ctemp, mprec	0	0	171.206	0.623	1
3	LQ	ctemp, mprec	0	0	171.206	0.623	1
2	QP	ctemp, mprec	0	0	171.758	1.175	1
2	LQP	ctemp, mprec	0	0	171.758	1.175	1
4	Q	ctemp, mprec	0	0	171.758	1.175	1
4	LQ	ctemp, mprec	0	0	171.758	1.175	1
0.1	Q	ctemp, wtemp	0	0	172.045	1.463	2
0.2	Q	ctemp, wtemp	0	0	172.056	1.473	2
0.3	Q	ctemp, wtemp	0	0	172.072	1.489	2
0.2	QP	ctemp, wtemp	0	0	172.093	1.510	2
0.4	Q	ctemp, wtemp	0	0	172.093	1.510	2
0.4	Q	All	0	0	172.093	1.510	2
0.5	Q	ctemp, wtemp	0	0	172.119	1.536	2
0.5	Q	All	0	0	172.119	1.536	2
0.3	QP	ctemp, wtemp	0	0	172.148	1.566	2
0.6	Q	ctemp, wtemp	0	0	172.148	1.566	2
0.6	Q	All	0	0	172.148	1.566	2
0.5	LQ	ctemp, mprec	0	0	172.153	1.570	2
0.7	Q	ctemp, wtemp	0	0	172.183	1.600	2
0.7	Q	All	0	0	172.183	1.600	2
0.3	LQP	ctemp, mprec	0	0	172.197	1.614	2
0.6	LQ	ctemp, mprec	0	0	172.197	1.614	2
0.4	QP	ctemp, wtemp	0	0	172.220	1.638	2
0.8	Q	All	0	0	172.220	1.638	2
0.7	LQ	ctemp, mprec	0	0	172.247	1.664	2
0.9	Q	ctemp, wtemp	0	0	172.262	1.679	2
0.9	Q	All	0	0	172.262	1.679	2

Table 2. continued

Regularization multiplier	Features	Environmental variables	Partial ROC	Omission rates below 5%	AICc	$\Delta AICc$	Number of parameters
0.4	LQP	ctemp, mprec	0	0	172.303	1.721	2
0.8	LQ	ctemp, mprec	0	0	172.303	1.721	2
0.5	QP	ctemp, wtemp	0	0	172.306	1.724	2
1	Q	All	0	0	172.306	1.724	2
1	Q	ctemp, wtemp	0	0	172.306	1.724	2
5	Q	ctemp, mprec	0	0	172.311	1.729	1
5	LQ	ctemp, mprec	0	0	172.311	1.729	1
0.9	LQ	ctemp, mprec	0	0	172.366	1.783	2
0.6	QP	ctemp, wtemp	0	0	172.404	1.822	2
0.6	LQP	ctemp, wtemp	0	0	172.404	1.822	2
0.5	LQP	ctemp, mprec	0	0	172.433	1.851	2
1	LQ	ctemp, mprec	0	0	172.433	1.851	2
0.7	QP	ctemp, wtemp	0	0	172.513	1.930	2
0.7	LQP	ctemp, wtemp	0	0	172.513	1.930	2

Consensus niche estimations were thresholded by reclassifying as non-suitable all grid cells with suitability scores that fell within the bottom 5% of all values from grid cells that contained an occurrence point (Peterson *et al.*, 2008). Subsequently, the remaining range of suitability scores were classified as low, medium or high suitability using equal intervals in QGIS 2.18.28.

Model comparisons were performed in both geographic and environmental dimensions. To compare the geographic projections of the niches, we used the background similarity and identity tests (Warren *et al.*, 2008, 2017), which quantify the similarity / identity between two predictions by measuring their geographic overlap and then comparing the result to a null distribution. The two metrics recommended for such comparisons are Schoener (1968)'s statistic *D* for niche overlap and the similarity statistic *I*. They range from 0 (no similarity / not identical) to 1 (total similarity / identical) and are obtained via the comparison of the two empirical niche predictions. These measured metrics are then compared to those obtained from 1000 sets of null niche predictions produced using occurrences randomly sampled from the environmental background available in the calibration area. The difference between the two

predictions is deemed to be significant if the empirical value falls below the 5% limits of the null distribution. To compare niches in environmental space, we used NicheA (Qiao *et al.*, 2016). We employed a three-dimensional environmental background using the three paleoclimatic variables to develop minimum volume ellipsoid (MVE) niche estimations, with a precision of 0.01, for the Pyrenees Noaillian and the northern Middle Gravettian (Figure 4). We measured the volume of each ellipsoid and calculated the level of environmental overlap between them using the Jaccard index (Qiao *et al.*, 2016; Qiao *et al.*, 2017).

3. Results

3.1. Niche predictions

The northern Middle Gravettian niche estimation (Figure 5a) displays high suitability scores in the northern Aquitaine Basin, the southern Paris Basin, and low-to-medium suitability for the western Italian coast, northwestern Alps piedmont, southern Landes and Western Pyrenees, Brittany, and southern Britain. The site of Hautmougey has the lowest suitability score of all the occurrence points, and this is likely because it is situated in the northeastern portion of the study region and is isolated from the other sites. This is likely the result of less intensive archaeological research in this region relative to others (e.g., Southwestern France, southern Paris Basin; Angevin *et al.*, 2018). The Garonne River Valley is characterized by a low level of suitability with the exception of a narrow corridor that connects the high suitability areas in northern Aquitaine to the regions of medium suitability situated in the southern Landes and Western Pyrenees. Finally, the geographic extent of the northern Middle Gravettian niche prediction corresponds closely to the geographic distribution of the occurrence data.

The Pyrenees Noaillian niche prediction (Figure 5b) is geographically extensive and is significantly larger than the distribution of archaeological occurrence data. The estimated niche

displays continuous high suitability scores from the coast of Cantabrian Spain up to southern Britain. The lowest predicted occurrence point is that of the site of La Carane-3, and all other sites from the central Pyrenees are located in areas with low to medium suitability scores.

Variability maps indicate that the northern Middle Gravettian model is probably more reliable than that of the Pyrenees Noaillian (Figure 5c and 5d). The northern Middle Gravettian variability map displays few and relatively small areas with a range higher than 0.3. The highest suitability ranges occur in Brittany and southern Britain, the Landes platform area, the Massif Central as well as the northern Alps. The Pyrenees Noaillian variability map, however, shows variability ranges higher than 0.3 in areas along the Mediterranean coasts, the Eastern side of the Pyrenees, the Massif Central, as well as along the Atlantic coast and in the Landes periglacial desert area. The suitability estimates in these areas should therefore be considered less reliable since they can vary substantially between models.

3.2. Niche comparisons

The Pyrenees Noaillian and the northern Middle Gravettian niches are highly similar in geographic space (Figure 6). However, their differences are significant and they are significantly less similar than would be expected by chance (D metric: North vs. Pyrenees: $p = 0.03$; Pyrenees vs. North: $p = 0.006$; I metric: North vs. Pyrenees: $p = 0.029$; Pyrenees vs. North: $p = 0.007$; for a statistical significance achieved if $p < 0.05$; Figure 6). With respect to environmental dimensions, our analysis shows that the niches overlap significantly (MVE overlap volume = 1.110, which corresponds to 97% of the Pyrenees Noaillian ellipsoid volume and 7% of the Northern Middle Gravettian ellipsoid volume; Figure 7). Comparisons demonstrate that the Pyrenees Noaillian niche is smaller and less broad than that of the northern Middle Gravettian, and is primarily contained within the latter (Figure 7). It is worth noting, however, that a small portion of the Pyrenees Noaillian ellipsoid falls outside of the northern

Middle Gravettian ellipsoid, thus occupying a subset of environmental conditions not present in the latter's niche estimation.




Figure 5. Geographic projections of Maxent-produced median niche estimations. **a.** Northern Middle Gravettian. **b.** Pyrenees Noaillian. **c.** Variability map for the northern Middle Gravettian prediction showing the suitability range of 10 models produced with the same data. **d.** Variability map for the Pyrenees Noaillian prediction showing the suitability range of all 45 significant models.




Figure 6. Background similarity and identity tests results for comparisons between the northern Middle Gravettian vs. the Pyrenees Noaillian (**a.** and **c.**), and the Pyrenees Noaillian vs. the northern Middle Gravettian (**b.** and **d.**). Dashed lines represent measures between the empirical models and the histograms depict measures from 1000 background-derived comparisons. The colored areas represent the non-significance range above the 5th percentile of the distribution

4. Discussion

The fact that the northern Middle Gravettian niche is significantly broader than that of the Pyrenees Noaillian suggests that the development of the Raysse method is linked to the exploitation of a significantly expanded niche composed of colder, drier conditions that correspond to more open landscapes and associated large mammal prey species. Available archaeological data support this hypothesis.




Figure 7. NicheA Minimum Volume Ellipsoids (MVE) for the northern Middle Gravettian (blue) and the Pyrenees Noaillian (red) in environmental space during GS.5 (black points) according to different orientations of the axes. The environmental dimensions are temperature of the coldest month (ctemp), temperature of the warmest month (wtemp) and mean annual precipitation (mprec). The MVE volumes are displayed in the corresponding colored boxes.

The Raysse method is a highly standardized reduction process that can be applied to a wide variety of blanks, from blades to thick flakes produced during various stages of the blade reduction sequences or other less standardized reduction sequences (e.g., production of flakes or blade-like flakes). The bladelets produced with the Raysse method did not require further intensive modification due to their standardized morphology. Furthermore, if a problem was encountered during the final stages of their production, only minimal investment was needed to produce a new or replacement bladelet (Klaric *et al.*, 2002, 2009; Klaric, 2003, 2008, 2017). This highly standardized bladelet production system served to create end-products that could be easily transformed into *armatures* (a French term commonly used in technological and functional studies to designate either the lithic point hafted on the tip/head of a projectile or the lithic implements hafted laterally on a projectile's foreshaft) that were likely part of a composite and readily maintainable hunting weaponry toolkit. Such a technological adaptation is commonly observed in highly mobile hunter-gatherers that operate in landscapes where access to resources needed to maintain weaponry is unpredictable (Binford 1977; Bleed, 1986). The importance placed on producing highly standardized components for a maintainable and curated hunting toolkit is further supported by the emphasis that appears to have been placed on transmitting and maintaining this technique (Klaric, 2017, 2018). This has been inferred from the numerous technical details that one must take into account when using the Raysse method to replicate La Picardie bladelet blanks, and this is further supported by the frequent presence of unretouched flint knapping debris and retouched objects that appear to have been produced by apprentices or individuals who did not completely master the intricacies of the method (Klaric, 2017, 2018).

The opposite appears to have been the case in the Pyrenees, where *armature* forms are more diverse: Gravette points, microgravettes points, bi-truncated backed bladelets, and simple backed implements made from blades or bladelets (Klaric, 2003; Simonet, 2009a, 2011b, 2017).

While these *armature* types were made from straight blade or bladelet blanks, the latter do not need to be highly standardized and can be obtained from a variety of reduction methods. Intensive retouch is all that is necessary to transform an initial blank into one of these *armature* types (e.g., Gravettes points broken during fabrication indicate that the blank's width can be reduced up to 50%; Klaric, 2003, p. 257; Simonet, 2009a, fig. 21). The higher typological diversity of *armatures* in the Pyrenees may reflect a higher degree of variability in how these points were integrated into weapon systems (e.g., axial points vs. laterally mounted elements) used by these populations. These different technological strategies for producing hunting equipment between the northern Middle Gravettian and the Pyrenees Noaillian archaeological records are likely related to differences in targeted medium to large prey species and, in turn, the subsistence strategies and mobility patterns employed to exploit them.

Faunal data indicate that populations in the Pyrenees hunted a variety of animals, such as reindeer, bovids, horse, chamois, bison, deer and fox, whereas northern groups relied principally on reindeer (Lacarrière, 2015). This fact does not necessarily indicate that the availability of prey species was different between these two regions (Lacarrière, 2015, p. 347), but it is worth noting that the smaller Pyrenees niche is associated with a more diverse spectrum of prey species. Thus, it would appear that the Pyrenees piedmont plateau and plains, with its more reduced range of environmental conditions, contained a wide variety of prey species that, as indicated by seasonality data (Lacarrière, 2015), were present throughout the year. These data are consistent with inferred Gravettian occupation of the region, which is dominated by small, specialized sites (e.g., Tercis, Gatzarria) situated some distance from larger aggregation sites (e.g., Isturitz, Brassempouy), as well as an exploitation of predominantly local lithic raw materials (Simonet, 2017). Pyrenees populations, thus, were likely logistically mobile with a well-organized exploitation of resources within a relatively restricted region and narrow ecological niche. The high prey species diversity could therefore be the result of a more

generalized hunting strategy within a reduced territory. In this framework, *armature* diversity could be interpreted as an adaptation to the exploitation of a more diverse prey species spectrum within a smaller and more rugged territory, allowing hunters to easily adapt to different situations by changing their hunting weaponry (e.g. prey type, topography etc.).

To the north, in contrast, the wider range of environmental conditions (i.e., broader ecological niche) exploited by the northern Middle Gravettian populations would suggest a higher degree of mobility than is observed in the Pyrenees since the main prey species identified in northern archaeological assemblages is reindeer (Lacarrière, 2015). Higher mobility is supported by technological data. For example, the Grotte du Renne site is located some distance from high quality lithic raw material sources (35 to 120 km according to Klaric *et al.*, 2009), and its Middle Gravettian assemblage shows a high degree of curation, in contrast to other sites that are situated at or near raw material sources, such as the site of La Picardie (although a small percentage of artifacts at La Picardie are made from raw materials coming from sources located in the Charente region, some 200 km away; Delvigne *et al.*, 2020). With respect to curation, Raysse burins (i.e. bladelet cores) from the Grotte du Renne often have double bladelet production platforms, are smaller than those recovered from La Picardie, and the blanks selected to make La Picardie bladelets are generally smaller and are not always typical: many bladelets are more twisted and do not always have a *pan-revers* and/or a pointed distal end (Klaric, 2006, 2017; Klaric *et al.*, 2009). In this scenario, the Raysse method could reflect a technological specialization directed at producing *armatures* for hunting reindeer. It does not mean that reindeer was the only available species in the environment; in fact, other species are always present to a lesser extent in faunal assemblages (Lacarrière, 2015). This could therefore be the result of a cultural choice of northern populations focusing on reindeer (Klaric *et al.*, in press)—a choice that would require higher mobility. Nevertheless, one must be careful to not generalize this pattern, since it relies on only a handful of studies, and numerous sites have

assemblages that contain both by-products of the Raysse method and Gravette points, two conceptually different *chaînes opératoires* (i.e. reduction sequences), and thus different kinds of composite hunting weapons. However, we must bear in mind that occurrences of Picardie bladelets and Gravette points together are observed almost exclusively within cave or rock shelter deposits, where archaeological levels represent palimpsests of multiple occupations that often were subjected to complex post-depositional processes that can homogenize initially distinct occupations. The co-occurrence of the two *armature* types is also observed at two open-air sites: the site of Solvieux in Dordogne (Sackett, 1999), where the stratigraphic setting is extremely complex, and Sackett's analysis has raised doubts concerning the integrity of its archaeological assemblages; and the site of Les Jambes in Dordogne (Célérier, 1967), where slope depositional processes influenced the site's formation (Klaric, 2003, p. 222), thus raising doubts as to whether the distinct archaeological levels defined by Célérier are valid. Whether these two types of *armatures*—la Picardie bladelets and Gravette/microgravette points—were associated with the same hunting tool-kit or whether their association in the same archaeological level is the result of post-depositional processes or palimpsest deposits, is a subject that requires further study.

At present, the site of Callan represents the only known location of specialized occupations or activities north of the Garonne River (Morala, 2011). Its lithic assemblage is dominated by Noailles burins and no *armatures* have been recovered. This general absence of specialized sites suggests that groups in these higher latitude regions had a higher level of residential mobility than those in the Pyrenees. Such a pattern of highly mobile groups using highly standardized and curated toolkits to exploit large territories via a residential system of mobility is in sharp contradiction with the pattern observed during the same period in the Pyrenees. This pattern, though, may be influenced by the lack of chronological resolution for this time-period, which is the result of reduced stratigraphic resolution due to imprecise

excavation methods, to post-depositional mixing of levels, palimpsest deposits and the standard errors associated with radiocarbon ages for this period. Such factors make determinations of discrete activity episodes difficult, if not impossible, and the potential homogenization of archaeological levels renders making inferences of how human activities were organized across the landscape difficult.

With respect to the geographic expressions of the estimated niches and the technological differentiation observed between the two regions, what factors potentially influenced these patterns? We propose that the cold desert of the Landes region (Bertran *et al.*, 2013) and the Garonne River Valley corridor (Bruxelles & Jarry, 2011, 2012) served as an ecological barrier that played a role in the territories exploited by Middle Gravettian hunter-gatherer populations. The presence of this barrier would have been a factor in limiting the diffusion of the Raysse method to the Pyrenees area. This idea is supported by the fact that this region is associated with low suitability measures (Figure 5), and these suitability scores are even likely to be overestimated since the climatic variables used for this study insufficiently capture the particular conditions of a cold desert whose existence was the result of specific geographic and geomorphic factors (Bertran *et al.*, 2013). The presence of this ecological barrier likely influenced the territories exploited by different Middle Gravettian hunter-gatherer populations and thus in term served to create a cultural barrier between the Pyrenees and regions to the north. This hypothesis is supported by the fact that the Pyrenees Noaillian's existing niche is more geographically extensive than the distribution of sites used to reconstruct it, whereas the northern Middle Gravettian niche more closely corresponds to its occurrence data. Thus, the presence of populations in these northern habitats that were characterized by different technological systems and mobility strategies, from which one could infer other cultural differences, could have prevented the Pyrenees populations from occupying their entire existing fundamental niche.

Another observation concerns the northern model's predictions for regions beyond the borders of present-day France—regions in which Noailles burins are present in archaeological assemblages (Touzé, 2013). There exists a small area of low suitability along the western Italian coast for the northern Middle Gravettian prediction, but the region between this area and the main suitable area in France is predicted as unsuitable. As for the niche associated with the Pyrenees Noaillian, regions east of the French Massif Central are predicted as unsuitable. One possible interpretation is that the Rhône River Valley functioned as a barrier during GS.5. This interpretation, however, is contradicted by the presence of Noailles burins in the lower Rhône River Valley and along the Italian Mediterranean coast (Palma di Cesnola, 1993; Onoradini *et al.*, 2010; Touzé, 2013). Moreover, these areas are characterized by a high degree of variability depending on model parametrization, thus indicating that their corresponding suitability estimates are less reliable (Figure 5d). Additional niche predictions, comparisons, and tests that take into account Italian sites, in conjunction with detailed examinations of their lithic industries and comparisons to assemblages to the west, would be necessary to further evaluate this issue. To the west, the Cantabrian region is characterized by high suitability scores for the Pyrenees Noaillian niche, whereas it is not suitable in the northern Middle Gravettian niche prediction, except for a small portion along the Western Pyrenees. This pattern supports the idea that the Cantabrian region was part of the Pyrenees Noaillian territory—a pattern supported by the presence of Noailles burins in this region (Foucher *et al.*, 2008; Simonet, 2009a, 2017; de la Peña-Alonso, 2011). This pattern should be further evaluated with detailed studies of the archaeological assemblages and new niche predictions that include the presence of these sites. It is also paramount to couple such analyses with critical evaluations of existing chronological data (cf. Banks *et al.*, 2019), as well as with efforts to obtain new ¹⁴C ages from reliable archaeological contexts.

5. Conclusions

Comparisons of the reconstructed niches for the two Middle Gravettian technological *faciès*, the Noaillian and the Rayssian, in both geographic and environmental dimensions, indicate that their respective niches were significantly different, despite a large degree of overlap in environmental space, due to the fact that the northern Middle Gravettian niche was significantly broader than that of the Pyrenees Noaillian. We interpret this pattern as indicating that the appearance of the Raysse method is related to the exploitation of a significantly broader ecological niche. As opposed to what is observed in areas south of the Garonne River, the Raysse method appears to be associated with particular mobility and settlement strategies contained within a larger exploited territory or territories. Furthermore, this observed pattern suggests that La Picardie bladelets (products of the Raysse method) represented a technological specialization directly associated with a hunting strategy focused on reindeer. Picardie bladelets may have been considered—at least by some hunter-gatherer populations—to be more appropriate within this context than backed points and abrupt backed bladelet *armatures* (such as Gravette and microgravette points), although this hypothesis warrants further testing in the archaeological record. Furthermore, the Raysse method would have been advantageous in such contexts because it would have been more easily maintainable and adapted to hunting activities organized within territories such that access to raw material resources was less predictable or available. Conversely, we interpret *armature* diversity of the Pyrenees Noaillian as reflecting an adaptation to the organization of subsistence activities within smaller territories and the use of specialized weapon systems adapted to a variety of prey species and/or situations (e.g. topography). Furthermore, the need for a highly maintainable hunting toolkit was probably not as paramount, since access to raw material resources would have been more predictable or available.

709 These niche results further support the hypothesis that the Landes cold desert and
710 Garonne River Valley corridor served to limit the homogenization of technological traditions
711 between the Pyrenees and regions to the north. The nature of this barrier (i.e. environmental
712 and/or cultural) warrants further evaluation incorporating other potentially relevant variables
713 (distribution of cold deserts etc.) in future ecological niche modeling analyses that target the
714 concerned archaeological populations. Furthermore, continued investigations centered on
715 identifying Middle Gravettian sites and the typo-technological attributes of their archeological
716 assemblages are needed. In turn, the niche modeling results presented here will provide
717 important context and details for research addressing issues of chronology, settlement and
718 subsistence strategies, lithic raw material exploitation, lithic and bone technologies, and site
719 function pertaining to the Middle Gravettian archaeological record.

721 **Supplementary material**

722 Codes used in R to perform the steps described in sections 2.3.2 and 2.3.3 and models
723 calibration results are available online at
724 https://osf.io/35pb4/?view_only=4f84e08e6ba84754b2090e349bc2ebf4. Attached to the article
725 is an additional appendix containing a table of criteria used to identify reliable identifications
726 of the Rayssian and the Noaillian in the literature.

728 **Acknowledgements**

729 We thank A. Townsend Peterson for hosting a collaborative trip to the University of Kansas'
730 Biodiversity Institute. The climate simulations were run on the Mercure supercomputer
731 maintained by the Commissariat à l'Energie Atomique's Très Grand Centre de Calculs. We
732 thank A. Townsend Peterson, Mathieu Bosq and Pascal Bertran for their input during the

writing of the article. Finally, we thank Andreas Maier and João Marreiros for their critical reviews as part of the Peer Community In open-access reviewing process (<https://doi.org/10.24072/pci.archaeo.100003>), as well as the two anonymous reviewers for their critics and suggestions that served to improve the paper.

Funding

This research was financed by the LabEx project IMPACT (W. Banks, dir.), and the Nouvelle-Aquitaine region project “GravettoNiches” (W. Banks, dir.). The collaborative trip to the University of Kansas’ Biodiversity Institute was co-funded by the University of Bordeaux’s “Sciences et Environnements” Doctoral School.

References

Agsous, S. (2008). *Stratigraphie et sédimentologie de l’abri Pataud (Les Eyzies-de-Tayac, Dordogne, France). Contribution à l’étude des paléoenvironnements des cultures du Paléolithique supérieur ancien du Périgord*. [doctoral thesis]. Paris: Muséum National d’Histoire Naturelle.

Alaux, J.-F. (1967). Burins du type “Le Raysse” de l’abri des Battuts (Tarn). *Bulletin de la Société préhistorique française. Comptes rendus des séances mensuelles*, 64(8), 242–247. <https://doi.org/10.3406/bspf.1967.8976>

Alaux, J.-F. (1971). Burins de Noailles inachevés et burins « de technique Noailles » de l’Abri des Battuts, commune de Penne (Tarn). *Bulletin de la Société préhistorique française*, 68(8), 239–242. <https://doi.org/10.3406/bspf.1971.4332>

755 Alkama, R., Kageyama, M., Ramstein, G., Marti, O., Ribstein, P., & Swingedouw, D.
756 (2008). Impact of a realistic river routing in coupled ocean–atmosphere simulations of the
757 Last Glacial Maximum climate. *Climate Dynamics*, 30(7–8), 855–869.
758 <https://doi.org/10.1007/s00382-007-0330-1>

759 Allard, M. (1986). Le gisement préhistorique de La Martinière à La Pommeraye (M.-
760 et-L.). Mise en évidence de périgordien supérieur à burins de Noailles. *Bulletin de la Société*
761 *préhistorique française*, 83, 78–82. <https://doi.org/10.3406/bspf.1986.8738>

762 Allard, M., & Gruet, M. (1976). Les civilisations du Paléolithique supérieur dans les
763 Pays de la Loire. In H. de Lumley (dir.) *La Préhistoire française. Tome I : civilisations*
764 *paléolithiques et mésolithiques* (Vol. 2, pp. 1305–1310). Paris : Editions du CNRS.

765 Anderson, R. P., Lew, D., & Peterson, A. T. (2003). Evaluating predictive models of
766 species' distributions: criteria for selecting optimal models. *Ecological Modelling*, 162(3),
767 211–232. [https://doi.org/10.1016/S0304-3800\(02\)00349-6](https://doi.org/10.1016/S0304-3800(02)00349-6)

768 Anderson, R. P., & Gonzalez, I. J. (2011). Species-specific tuning increases robustness
769 to sampling bias in models of species distributions: an implementation with Maxent.
770 *Ecological Modelling*, 222, 279–2811. <https://doi.org/10.1016/j.ecolmodel.2011.04.011>

771 Angevin, R., Kildéa, F., Djemmali, N., Alilaire, P., Delvigne, V., Depont, J., Lafarge,
772 A., Mevel, L., Schemmama, V., Surmely, F., & Verjux, C. (2018). Le Paléolithique supérieur
773 dans le sud du Bassin parisien à la lumière des découvertes récentes : faits attendus, faits
774 nouveaux. In O. Troubat (Ed.) *Préhistoire de la France centrale: actes du colloque,*
775 *Montluçon, 19 novembre 2016* (pp. 49–71). Montluçon: Cercle archéologique de Montluçon
776 et de la région.

777 Arambourou, R., & Jude, P. E. (1964). *Le Gisement de La Chèvre à Bourdeilles*,
778 *Dordogne*. Bordeaux: R. & M. Magne.

779 Aranguren, B. M., Longo, L., Pallecchi, P., & Revedin, A. (2006). Eléments de la
780 chaîne opératoire-fonctionnelle des burins de Noailles de Bilancino (Florence, Italie). In
781 Lebrun-Ricalens F. (Ed.) *Burins préhistoriques : formes, fonctionnements, fonctions* (n°2, pp.
782 143–162). Luxembourg: Archéologiques.

783 Aranguren, B., Cavulli, F., D’Orazio, M., Grimaldi, S., Longo, L., Revedin, A., &
784 Santaniello, F. (2015). Territorial exploitation in the Tyrrhenian Gravettian Italy: The case-
785 study of Bilancino (Tuscany). *Quaternary International*, 359–360, 442–451.
786 <https://doi.org/10.1016/j.quaint.2014.07.009>

787 Aubry, T., Almeida, M., Candela, P., Chauvière, F.-X., Dimuccio, L., Fontana, L.,
788 Liard, M., Marquet, J.-C., João Neves, M., Peyrouse, J.-B., & Walter, B. (2013). Le
789 Paléolithique supérieur ancien dans le sud-ouest du Bassin parisien : du Châtelperronien au
790 Gravettien dans les vallées de la Creuse et de la Claise. In Bodu P., Chehmana L., Klaric L.,
791 Mevel L., Soriano S., Teyssandier N. (dir.) *Le Paléolithique supérieur ancien de l’Europe du*
792 *Nord-Ouest* (Mémoire 56, pp. 299–318). Paris: Société préhistorique française.

793 Banks, W. E. (2015). Constructing chronologies for the late Middle Paleolithic and
794 Upper Paleolithic: limitations and means to overcome them. *World Archaeology*, 47(4), 585–
795 600. <https://doi.org/10.1080/00438243.2015.1052544>

796 Banks, W. E. (2017). The application of Ecological Niche Modeling methods to
797 archaeological data in order to examine Culture-Environment relationships and cultural
798 trajectories. *Quaternaire*, 28(2), 271–276. <https://doi.org/10.4000/quaternaire.7966>

799 Banks, W. E., d’Errico, F., Peterson, A. T., Vanhaeren, M., Kageyama, M., Sepulchre,
800 P., Ramstein, G., Jost, A., & Lunt, D. (2008). Human ecological niches and ranges during the
801 LGM in Europe derived from an application of eco-cultural niche modeling. *Journal of*
802 *Archaeological Science*, 35(2), 481–491. <https://doi.org/10.1016/j.jas.2007.05.011>

803 Banks, W. E., Zilhão, J., d’Errico, F., Kageyama, M., Sima, A., & Ronchitelli, A.
804 (2009). Investigating links between ecology and bifacial tool types in Western Europe during
805 the Last Glacial Maximum. *Journal of Archaeological Science*, 36(12), 2853–2867.
806 <https://doi.org/10.1016/j.jas.2009.09.014>

807 Banks, W. E., Aubry, T., d’Errico, F., Zilhão, J., Lira-Noriega, A., & Townsend
808 Peterson, A. (2011). Eco-cultural niches of the Badegoulian: Unraveling links between
809 cultural adaptation and ecology during the Last Glacial Maximum in France. *Journal of*
810 *Anthropological Archaeology*, 30(3), 359–374. <https://doi.org/10.1016/j.jaa.2011.05.003>

811 Banks, W. E., d’Errico, F., & Zilhão, J. (2013). Human–climate interaction during the
812 Early Upper Paleolithic: testing the hypothesis of an adaptive shift between the Proto-
813 Aurignacian and the Early Aurignacian. *Journal of Human Evolution*, 64(1), 39–55.
814 <https://doi.org/10.1016/j.jhevol.2012.10.001>

815 Banks, W. E., Bertran, P., Ducasse, S., Klaric, L., Lanos, P., Renard, C., & Mesa, M.
816 (2019). An application of hierarchical Bayesian modeling to better constrain the chronologies
817 of Upper Paleolithic archaeological cultures in France between ca. 32,000–21,000 calibrated
818 years before present. *Quaternary Science Reviews*, 220, 188–214.
819 <https://doi.org/10.1016/j.quascirev.2019.07.025>

820 Bardon, L., Bouyssonie, J., & Bouyssonie, A. (1903). Un nouveau type de burin.
821 *Revue de l’Ecole d’Anthropologie*, 13, 165–168.

822 Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S. P., Peterson,
823 A. T., Soberón, J., & Villalobos, F. (2011). The Crucial Role of the Accessible Area in
824 Ecological Niche Modeling and Species Distribution Modeling. *Ecological Modelling*,
825 222(11), 1810–19. <https://doi.org/10.1016/j.ecolmodel.2011.02.011>.

826 Berger, A. (1978). Long-term variations of daily insolation and quaternary climatic
827 changes. *Journal of Atmospheric Sciences*, 35, 2362–2367. [https://doi.org/10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2)

829 Bertran, P., Sitzia, L., Banks, W. E., Bateman, M. D., Demars, P.-Y., Hernandez, M.,
830 Lenoir, M., Mercier, N., & Prodeo, F. (2013). The Landes de Gascogne (southwest France):
831 periglacial desert and cultural frontier during the Palaeolithic. *Journal of Archaeological*
832 *Science*, 40(5), 2274–2285. <https://doi.org/10.1016/j.jas.2013.01.012>

833 Binford, L. R. (1977). Forty-seven trips. A case study in the character of
834 archaeological formation processes. In R.V.S. Wright (Ed.) *Stone tools as cultural markers: change, evolution and complexity* (pp. 24–36). Canberra: Australian Institute of Aboriginal
835 Studies.
836 Studies.

837 Bleed, P. (1986). The Optimal Design of Hunting Weapons: Maintainability or
838 Reliability. *American Antiquity*, 51(4), 737–747. <https://doi.org/10.2307/280862>

839 Boria, R. A., Olson, L. E., Goodman, S. M., & Anderson, R. P. (2014). Spatial
840 filtering to reduce sampling bias can improve the performance of ecological niche models.
841 *Ecological Modelling*, 275, 73–77. <https://doi.org/10.1016/j.ecolmodel.2013.12.012>

842 Bosq, M., Bertran, P., Degeai, J.-P., Kreutzer, S., Queffelec, A., Moine, O., & Morin,
843 E. (2018). Last Glacial aeolian landforms and deposits in the Rhône Valley (SE France):

844 Spatial distribution and grain-size characterization. *Geomorphology*, 318, 250–269.
845 <https://doi.org/10.1016/j.geomorph.2018.06.010>

846 Boyer, M., Geneste, J.-M., & Rigaud, J. P. (1984). Le Périgordien supérieur du site de
847 plein air du Caillou, Rouffignac-de-Sigoulès (Dordogne). *Bulletin de la Société préhistorique*
848 *française*, 81(81), 302–310. <https://doi.org/10.3406/bspf.1984.8593>

849 Breuil, H. (1927). Oeuvres d’art paléolithique inédites du Périgord et art oriental
850 d’Espagne. *Revue anthropologique*, 4–6, 101–108.

851 Briand, T., Remicourt, M., Tallet, P., Fernandes, P., Rué, M., Simonet, A., Pasquini,
852 A., Fedoroff, N., Picavet, R., & Bernad, C. (2010). *Les occupations gravettiennes de Hin-de-*
853 *Diou, à Pujo-le-Plan. Opération d’archéologie préventive du 05 octobre au 18 décembre*
854 *2009*. Villard-de-Lans: Paléotime – Service régional de l’archéologie Aquitaine.

855 Bricker, H. M. (1995). *Le Paléolithique supérieur de l’abri Pataud, Dordogne : les*
856 *fouilles de H. L. Movius Jr. : Suivi d’un inventaire analytique des sites aurignaciens et*
857 *périgorgiens de Dordogne*. Paris: Ed. de la Maison des sciences de l’Homme.

858 Bruxelles, L., & Jarry, M. (2011). Climatic conditions, settlement patterns and cultures
859 in the Paleolithic: The example of the Garonne Valley (southwest France). *Journal of Human*
860 *Evolution*, 61(5), 538–548. <https://doi.org/10.1016/j.jhevol.2011.05.009>

861 Bruxelles, L., & Jarry, M. (2012). Climats et cultures paléolithiques : quand la vallée
862 devient frontière.... *Archeopages, Hors-série 3*, 72–85.
863 <https://doi.org/10.4000/archeopages.469>

864 Caux, S. (2012). *Etude de l’industrie lithique*. In Boudadi-Maligne, M. (dir.) *Grotte de*
865 *Malvidier (La Roque-Gageac, Dordogne), Rapport d’opération de fouille programmée* (pp.

866 111–118). Bordeaux: Service régional de l’archéologie, Direction régionale des affaire
867 culturelles Aquitaine.

868 Célérier, G. (1967). Le gisement périgordien supérieur des “Jambes”. Commune de
869 Périgueux (Dordogne). *Bulletin de la Société préhistorique française*, 64(1), 53–68.

870 Clarke, D. L. (1968). *Analytical Archaeology*. London: Methuen & Co Ltd.

871 Clottes, Jean. (1985). Circonscription des Pyrénées. *Gallia Préhistoire*, 28(2), 331–
872 371

873 Clottes, J., Lorblanchet, M., Fau, M.-F., & Peyre, G. (1990). L’abri périgordien de la
874 Bergerie, à Saint-Géry (Lot). *Bulletin de la Société préhistorique française*, 87(10), 342–357.
875 <https://doi.org/10.3406/bspf.1990.9920>

876 Cobos, M. E., Peterson, A. T., Barve, N., & Osorio-Olvera, L. (2019a). Kuenm: An R
877 Package for Detailed Development of Ecological Niche Models Using Maxent. *PeerJ*, 7,
878 e6281. <https://doi.org/10.7717/peerj.6281>

879 Cobos, M. E., Peterson, A. T., Osorio-Olvera, L. and Jiménez-García, D. (2019b). An
880 Exhaustive Analysis of Heuristic Methods for Variable Selection in Ecological Niche
881 Modeling and Species Distribution Modeling. *Ecological Informatics*, 53, 100983.
882 <https://doi.org/10.1016/j.ecoinf.2019.100983>.

883 Cobos, M. E., Osorio-Olvera, L., & Peterson, A. T. (2019c). Assessment and
884 representation of variability in ecological niche model predictions. *BioRxiv*, 603100.
885 <https://doi.org/10.1101/603100>

886 Couchard, J., & de Sonneville-Bordes, D. (1960). La grotte de Bassaler-Nord, près de
887 Brive et la question du Périgordien II en Corrèze. *L’Anthropologie*, 64(5–6), 415–437.

888 d'Errico, F., & Banks, W. E. (2013). Identifying Mechanisms behind Middle
889 Paleolithic and Middle Stone Age Cultural Trajectories. *Current Anthropology*, 54(8), 371–
890 387. <https://doi.org/10.1086/673388>

891 d'Errico, F., Banks, W. E., Warren, D. L., Sgubin, G., Niekerk, K. van, Henshilwood,
892 C., Daniau, A.-L., & Goñi, M. F. S. (2017). Identifying early modern human ecological niche
893 expansions and associated cultural dynamics in the South African Middle Stone Age.
894 *Proceedings of the National Academy of Sciences*, 114(30), 7869–7876.
895 <https://doi.org/10.1073/pnas.1620752114>

896 Daniel, R. (1967). Le Périgordien supérieur de Petit-Puyrousseau, commune de
897 Périgueux (Dordogne). *Bulletin de la Société préhistorique française*, 64(8), 236–241.
898 <https://doi.org/10.3406/bspf.1967.8975>

899 Daniel, R. (1969). Les burins de Noailles du Fourneau-du-Diable ; Commune de
900 Bourdeilles (Dordogne). *Bulletin de la Société préhistorique française*, 66(1), 16–18.
901 <https://doi.org/10.3406/bspf.1969.10325>

902 Daniel, R., & Schmider, B. (1972). L'abri Durand-Ruel, près Brantôme (Dordogne) et
903 ses problèmes stratigraphiques. *Gallia préhistoire*, 15(2), 323–337.
904 <https://doi.org/10.3406/galip.1972.1419>

905 David, N. (1985). *The Noaillian (level 4) assemblages and the Noaillian Culture in*
906 *Western Europe*. Cambridge: Peabody Museum of Archaeology and Ethnology.

907 de la Peña Alonso, P. (2009). Revisión crítica de los conjuntos líticos gravetienses y su
908 contexto arqueológico en la Península Ibérica. *Complutum*, 20(1), 29–53.

909 de la Peña Alonso, P. (2011). *Sobre la unidad tecnologica del Gravetiense en la*
910 *peninsula Ibérica : implicaciones para el conocimiento del Paleolitico superior inicial.*
911 [doctoral thesis]. Madrid: Universidad complutense de Madrid.

912 de las Heras, C., Lasheras, J. A., Arrizabalaga, Á., & de la Rasilla, M. (Eds.) (2013).
913 *Pensando el Gravetiense: Nuevos datos para la region cantabrica en su contexto peninsular y*
914 *pyrenaico.* Museo Nacional y Centro de Investigacion de Altamira (monografias 23).

915 de Sonnevile-Bordes, D. (1960). *Recherches sur le Paléolithique supérieur en*
916 *Périgord* [doctoral thesis]. Paris: Université de Paris.

917 Delage, F. (1936). L’abri des Merveilles à Castel-Merle (Sergeac, Dordogne). *Compte-*
918 *rendu de la XIIIe session*, 578–608.

919 Delporte, H. (1968). L’abri du Facteur à Tursac (Dordogne). *Gallia Préhistoire*, 11(1),
920 1–112. <https://doi.org/10.3406/galip.1968.1307>

921 Delvigne, V., Fernandes, P., Tuffery, C., Raynal, J.-P., & Klaric, L. (2020).
922 Taphonomic methods and a database to establish the origin of sedimentary silicified rocks
923 from the Middle-recent Gravettian open-air site of La Picardie (Indre-et-Loire, France).
924 *Journal of Archaeological Science: Reports*, 32, 102369.
925 <https://doi.org/10.1016/j.jasrep.2020.102369>

926 Demars, P.-Y. (1977). Les industries du Périgordien supérieur des grottes de Pré-
927 Aubert et des Morts près Brive (Corrèze). *Bulletin de la Société préhistorique française.*
928 *Comptes rendus des séances mensuelles*, 74(4), 103–111.
929 <https://doi.org/10.3406/bspf.1977.8490>

930 Djindjian, F., Koslowski, J., & Otte, M. (1999). *Le Paléolithique supérieur en Europe*.
931 Malakoff: Armand Colin.

932 Ehlers, J., & Gibbard, P. (2004). *Quaternary Glaciations - Extent and Chronology*,
933 *Part I: Europe* (Vol. 2a). Amsterdam: Elsevier.

934 Féruglio, V., Aujoulat, N., & Jaubert, J. (2011). L'art pariétal gravettien, ce qu'il
935 révèle de la société en complément de la culture matérielle. In Goutas N., Klaric L., Pesesse
936 D., Guillermin P. (dir.) *À la recherche des identités gravettiennes. Actualités,*
937 *questionnements, perspectives* (Mémoire 52, pp. 243–255). Paris: Société préhistorique
938 française.

939 Floss, H., Dutkiewicz, E., Frick, J., & Hoyer, C. (2013). Le Paléolithique supérieur
940 ancien en Bourgogne du sud. In Bodu P., Chehmana L., Klaric L., Mevel L., Soriano S.,
941 Teyssandier N. (dir.) *Le Paléolithique supérieur ancien de l'Europe du Nord-Ouest* (Mémoire
942 56, pp. 331–350). Paris: Société préhistorique française.

943 Fontaine, A. (2006). *Le site des Vachons (commune de Voulgézac, Charente): étude*
944 *typologique des couches 3 et 3 bis (précédée d'un avant-propos rectificatif)* [master thesis].
945 Paris: Muséum National d'Histoire Naturelle.

946 Foucher, P. (2004). *Les industries lithiques du complexe Gravettien-Solutréen dans les*
947 *Pyrénées : techno-typologie et circulation des matières siliceuses de part et d'autre de l'axe*
948 *Pyrénées-Cantabres*. [doctoral thesis]. Toulouse: Université Toulouse Le Mirail - Toulouse II.

949 Foucher, P. (2013). Synthèse chrono-culturelle sur le Gravettien des Pyrénées : constat
950 et réflexions sur la stabilité régionale des traditions techniques. In de la Heras C., Lasheras
951 J.A., Arrizalaga A., de la Rasilla M. (dir.). *Pensando el Gravetiense : nuevos datos para la*

952 *region cantabrica en su contexto peninsular y pirenaico* (monografias 23, pp. 142–159).
 953 Santander, Madrid: Museo Nacional y Centro de Investigacion de Altamira.

954 Foucher, P., San Juan-Foucher, C., Sacchi, D., & Arrizabalaga, Á. (2008). Le
 955 Gravettien des Pyrénées. *Paléo*, 20, 331–356.

956 Foucher, P., Juan-Foucher, C. S., Vercoutère, C., & Ferrier, C. (2012). La grotte de
 957 Gargas (Hautes-Pyrénées, France): l’apport du contexte archéologique à l’interprétation de
 958 l’art pariétal. In Clottes J. (Ed.) *L’art pléistocène dans le monde / Pleistocen art of the world /*
 959 *Arte pleistoceno en el mundo* (pp. 209–225). Foix: Société préhistorique ariégeoise.

960 Gottardi, G. G. (2011). *La question des faciès au Gravettien : fonctions, traditions ou*
 961 *chronologie ?* [master thesis]. Bordeaux: Université de Bordeaux I.

962 Goutas, N. (2013a). De Brassempouy à Kostienki : l’exploitation technique des
 963 ressources animales dans l’Europe gravettienne. In M. Otte (dir.) *Les Gravettiens* (pp. 105–
 964 160). Paris: Errance.

965 Goutas, N. (2013b). Nouvelles données sur l’industrie osseuse des grottes du Renne et
 966 du Trilobite à Arcy-sur-Cure (Yonne, France). In Bodu P., Chehmana L., Klaric L., Mevel L.,
 967 Soriano S., Teyssandier N. (dir.) *Le Paléolithique supérieur ancien de l’Europe du Nord-*
 968 *Ouest* (Mémoire 56, pp. 89–116). Paris: Société préhistorique française.

969 Gruet, M. (1984). L’apport de deux sites angevins à la chronologie des terrasses
 970 fluviales : Roc-en-Pail en Chalonnnes sur Loire et Port-Launay sur la Sarthe. *Bulletin de*
 971 *l’Association française pour l’étude du quaternaire*, 21(1), 13–18.
 972 <https://doi.org/10.3406/quate.1984.1473>

973 Guillermin, P. (2006). Les Fieux : une occupation gravettienne du Causse quercinois.
974 *Paléo*, 18, 69–94.

975 Guillermin, P. (2008). Les « Périgordiens » en Quercy : l'exemple du gisement des
976 Fieux. *Paléo*, 20, 357–372.

977 Guillermin, P. (2011). La fin du Gravettien dans le sud-ouest de la France : à la
978 recherche de l'identité protomagdalénienne. In Goutas N., Klaric L., Pesesse D., Guillermin
979 P. (dir.) *À la recherche des identités gravettiennes. Actualités, questionnements, perspectives*
980 (Mémoire 52, pp. 129–146). Paris: Société préhistorique française.

981 Hans, J.-M. (1997). Périgordien à burins de Noailles : le site de Hautmougey (canton
982 de Bains-les-Bains, Vosges). *Bulletin de la Société Préhistorique Luxembourgeoise*, 19, 55–
983 66.

984 Henry-Gambier, D. (2008). Comportement des populations d'Europe au Gravettien :
985 pratiques funéraires et interprétations. *Paléo*, 20, 399–438.

986 Hinguant, S., & Monnier, J.-L. (2013). Le Paléolithique supérieur ancien dans le
987 Massif armoricain : un état de la question. In Bodu P., Chehmana L., Klaric L., Mevel L.,
988 Soriano S., Teyssandier N. (dir.) *Le Paléolithique supérieur ancien de l'Europe du Nord-*
989 *Ouest* (Mémoire 56, pp. 229–238). Paris: Société préhistorique française.

990 Hourdin, F., Foujols, M.-A., Codron, F., Guemas, V., Dufresne, J.-L., Bony, S.,
991 Denvil, S., Guez, L., Lott, F., Ghattas, J., Braconnot, P., Marti, O., Meurdesoif, Y., & Bopp,
992 L. (2013). Impact of the LMDZ atmospheric grid configuration on the climate and sensitivity
993 of the IPSL-CM5A coupled model. *Climate Dynamics*, 40(9–10), 2167–2192.
994 <https://doi.org/10.1007/s00382-012-1411-3>

995 Hutchinson, G. E. (1957). Population studies: Animal ecology and demography.
 996 *Bulletin of Mathematical Biology*, 53(1–2), 193–213.

997 Kageyama, M., Braconnot, P., Bopp, L., Caubel, A., Foujols, M.-A., Guilyardi, E.,
 998 Khodri, M., Lloyd, J., Lombard, F., Mariotti, V., Marti, O., Roy, T., & Woillez, M.-N. (2013).
 999 Mid-Holocene and Last Glacial Maximum climate simulations with the IPSL model—part I:
 1000 comparing IPSL_CM5A to IPSL_CM4. *Climate Dynamics*, 40(9–10), 2447–2468.
 1001 <https://doi.org/10.1007/s00382-012-1488-8>

1002 Kildea, F., & Lang, L. (2011). Le Gravettien de la vallée du Cher : le site de la Croix-
 1003 de-Bagneux à Mareuil-sur-Cher (Loir-et-Cher, France). In Goutas N., Klaric L., Pesesse D.,
 1004 Guillermin P. (dir.) *À la recherche des identités gravettiennes. Actualités, questionnements,*
 1005 *perspectives* (Mémoire 52, pp. 273–290). Paris: Société préhistorique française.

1006 Kildea, F., Griselin, S., Lang, L., & Souffi, B. (2013). Le Paléolithique supérieur
 1007 ancien aux marges méridionales du Bassin parisien : le site de la Croix-de-Bagneux à
 1008 Mareuil-sur-Cher (Loir-et-Cher). In Bodu P., Chehmana L., Klaric L., Mevel L., Soriano S.,
 1009 Teyssandier N. (dir.) *Le Paléolithique supérieur ancien de l'Europe du Nord-Ouest* (Mémoire
 1010 56, pp. 317–330). Paris: Société préhistorique française.

1011 Klaric, L. (2003). *Technical unity of Raysse burins industries in their diachronical*
 1012 *context. Thoughts about cultural variability during Gravettian, based on the case of La*
 1013 *Picardie, Arcy-sur-Cure, Brassempouy and Le Cirque de la Patrie.* [doctoral thesis]. Paris:
 1014 Université Panthéon-Sorbonne - Paris I.

1015 Klaric, L. (2006). Des armatures aux burins : des critères de distinction techniques et
 1016 culturels à partir des productions lamellaires de quelques sites du Gravettien moyen et récent

1017 (France). In Lebrun-Ricalens F. (Ed.) *Burins préhistoriques : formes, fonctionnements,*
1018 *fonctions* (n°2, pp. 199–223). Luxembourg: Archéologiques.

1019 Klaric, L. (2007). Regional groups in the European middle Gravettian : a
1020 reconsideration of the Rayssian technology. *Antiquity*.
1021 <https://doi.org/10.1017/S0003598X00094928>

1022 Klaric, L. (2008). Anciennes et nouvelles hypothèses d’interprétation du gravettien
1023 moyen en France : la question de la place des industries à burins du Raysse au sein de la
1024 mosaïque gravettienne. *Paléo*, 20, 257–276.

1025 Klaric, L. (2017). « La réussite d’une production repose sur l’attention prêtée aux
1026 détails »: l’exemple des débitages lamellaires par méthode du Raysse (Gravettien moyen,
1027 France). *Journal of Lithic Studies*, 4(2). <https://doi.org/10.2218/jls.v4i2.2547>

1028 Klaric, L. (2018). Levels of flintknapping expertise and apprenticeship during the
1029 Mid-Upper Palaeolithic: several illustrative examples from the Early and Late Aurignacian
1030 and Middle Gravettian. In Klaric (dir.) *The Prehistoric apprentice. Investigating*
1031 *apprenticeship and expertise in prehistoric technologies* (pp. 49–116). Brno, CZ: Dolní
1032 Věstonice Studies.

1033 Klaric, L., Aubry, T., & Walter, B. (2002). Un nouveau type d’armature en contexte
1034 gravettien et son mode de production sur les burins du Raysse (la Picardie, commune de
1035 Preuilly-sur-Claise, Indre-et-Loire). *Bulletin de la Société préhistorique française*, 99(4),
1036 751–764. <https://doi.org/10.3406/bspf.2002.12754>

1037 Klaric, L., Guillermin, P., & Aubry, T. (2009). Des armatures variées et des modes de
1038 productions variables. Réflexions à partir de quelques exemples issus du Gravettien d’Europe

- occidentale (France, Portugal, Allemagne). *Gallia préhistoire*, 51(1), 113–154.
<https://doi.org/10.3406/galip.2009.2476>
- Klaric, L., Liard, M., Bertran, P., Dumarçay, G., de Araujo Igreja, M., Aubry, T., & Walter, B. (2011). La Picardie (Preuilly-sur-Claise, Indre-et-Loire) : neuf ans de fouille sur un gisement rayssien finalement pas si mal conservé ! In Goutas N., Klaric L., Pesesse D., Guillermin P. (dir.) *À la recherche des identités gravettiennes. Actualités, questionnements, perspectives* (Mémoire 52, pp. 291–310). Paris: Société préhistorique française.
- Klaric, L., Bertran, P., Dumarçay, G., & Liard, M. (2018). A long and winding road: Towards a palethnographic interpretation of the middle-Gravettian site of la Picardie (Indre-et-Loire, France). *Quaternary International*, 498, 51–68.
<https://doi.org/10.1016/j.quaint.2018.09.035>
- Klaric L., Goutas N., Lacarrière J. & Banks W. (in press). Rayssien ? Vous avez dit Rayssien ? In O. Touzé, N. Goutas, P. Noiret, H. Salomon (dir.) *NW Europe during the Gravettian: Contributions of recent research to the understanding of the societies and their environments*, Proceedings of Liège symposium, 13-14 April 2018.
- Klima, B. (Dir.) (1983). *Aurignacien, Périgordien, Gravettien—Actes des colloques XV et XVI du IXe congrès de l'UISPP, Nice, 1976* (ERAUL, Vol. 1).
- Krtolitz, Y., & Lenoir, M. (1998). Un gisement du Gravettien à burins de Noailles en Gironde, l'abri Lespoux à Saint-Quentin-de-Baron. *Revue Archéologique de Bordeaux*, 89, 47–68.
- Lacarrière, J. (2015). *Les ressources cynégétiques au Gravettien en France : acquisition et modalités d'exploitation des animaux durant la phase d'instabilité climatique*

1061 *précédant le dernier maximum glaciaire* [doctoral thesis]. Toulouse: Université Toulouse le
 1062 Mirail - Toulouse II.

1063 Lacarrière, J., Goutas, N., Normand, C., & Simonet, A. (2011). Vers une redéfinition
 1064 des occupations gravettiennes de la grotte d’Isturitz (Pyrénées-Atlantiques, France) : révision
 1065 critique des collections “anciennes” par l’approche intégrée des données lithiques, fauniques
 1066 et de l’industrie osseuse. In Goutas N., Klaric L., Pesesse D., Guillermin P. (dir.) *À la*
 1067 *recherche des identités gravettiennes. Actualités, questionnements, perspectives* (Mémoire 52,
 1068 pp. 67–84). Paris: Société préhistorique française.

1069 Laville, H., & Rigaud, J.-P. (1973). The Perigordian V industries in Périgord :
 1070 typological variations, stratigraphy and relative chronology. *World Archaeology*, 4(3), 330–
 1071 338.

1072 Le Mignot, Y. (2000). La question de la production d’armatures sur le site gravettien
 1073 de Plasenn-al-Lomm (Ile de Bréhat, Côtes d’Armor). *Revue archéologique de l’ouest*, 17(1),
 1074 7–24. <https://doi.org/10.3406/rao.2000.1099>

1075 Le Tensorer, J.-M. (1974). Le Périgordien supérieur à burins de Noailles de l’Abri
 1076 Peyrony à Gavaudun (Lot-et-Garonne). *Bulletin de la Société préhistorique française*, 71(2),
 1077 459–468. <https://doi.org/10.3406/bspf.1974.8279>

1078 Le Tensorer, Jean-Marie. (1981). *Le Paléolithique de l’Agenais*. Editions du CNRS.

1079 Lenoir, M. (1977). Un gisement de plein air du Périgordien supérieur en Gironde : les
 1080 Artigaux à Camiac-et-Saint-Denis. *Bulletin de la Société Préhistorique Française*, 74(2),
 1081 518–530. <https://doi.org/10.3406/bspf.1977.8465>

1082 Leroi-Gourhan, A. (1984). *L'art des cavernes: atlas des grottes ornées paléolithiques*
1083 *françaises*. Paris: Imprimerie nationale.

1084 Lucas, G. (2000). *Les industries lithiques du Flageolet I (Dordogne) : Approche*
1085 *économique, technologique, fonctionnelle et analyse spatiale* [doctoral thesis]. Bordeaux:
1086 Université de Bordeaux I.

1087 Lucas, G. (2002). À propos des burins du Raysse du Flageolet I (Dordogne, France).
1088 *Paléo*, 14, 63–76.

1089 Marti, O., Braconnot, P., Dufresne, J.-L., Bellier, J., Benshila, R., Bony, S.,
1090 Brockmann, P., Cadule, P., Caubel, A., Codron, F., De Noblet, N., Denvil, S., Fairhead, L.,
1091 Fichet, T., Foujols, M.-A., Friedlingstein, P., Goosse, H., Grandpeix, J.-Y., Guilyardi, E., ...
1092 Talandier, C. (2010). Key features of the IPSL ocean atmosphere model and its sensitivity to
1093 atmospheric resolution. *Climate Dynamics*, 34, 1–26. [https://doi.org/10.1007/s00382-009-](https://doi.org/10.1007/s00382-009-0640-6)
1094 [0640-6](https://doi.org/10.1007/s00382-009-0640-6)

1095 Michel, A. (2010). *L'Aurignacien récent (post-ancien) dans le Sud-Ouest de la*
1096 *France : variabilité des productions lithiques. Révision taphonomique et techno-économique*
1097 *des sites de Caminade-Est, abri Pataud, Roc-de-Combe, Le Flageolet I, La Ferrassie et*
1098 *Combemenué*. [doctoral thesis]. Bordeaux: Université de Bordeaux I.

1099 Monméjean, E., Bordes, F., & de Sonneville-Bordes, D. (1964). Le Périgordien
1100 supérieur à burins de Noailles du Roc de Gavaudun (Lot-et-Garonne). *L'anthropologie*, 68(3–
1101 4), 253–316.

1102 Morala, A. (1984). *Périgordien et Aurignacien en Haut-Agenais : étude d'ensembles*
1103 *lithiques*. Paris: Ecole des Hautes Etudes en Sciences Sociales.

1104 Morala, A. (2011). La spécialisation des activités : concept de l'archéologue et réalité
1105 archéologique ; les données du site gravettien moyen du Callan (Lot-et-Garonne). In Goutas
1106 N., Klaric L., Pesesse D., Guillermin P. (dir.) *À la recherche des identités gravettiennes.*
1107 *Actualités, questionnements, perspectives* (Mémoire 52, pp. 343–358). Paris: Société
1108 préhistorique française.

1109 Movius, H. Jr., & David, N. (1970). Burins avec modification tertiaire du biseau,
1110 burins-pointe et burins du Raysse à l'Abri Pataud, Les Eyzies (Dordogne). *Bulletin de la*
1111 *Société préhistorique française*, 67(2), 445–455. <https://doi.org/10.3406/bspf.1970.4209>

1112 Nespoulet, R. (2008). Le Gravettien de l'abri Pataud. Bilan et perspectives. *Paléo*, 20,
1113 373–380.

1114 Noiret, P. (2013). De quoi Gravettien est-il le nom ? In M. Otte (dir.) *Les Gravettiens*
1115 (pp. 28–64). Paris: Errance

1116 Onoratini, G. (1982). *Préhistoire, sédiments, climats du Würm III à l'Holocène dans le*
1117 *Sud-Est de la France* [doctoral thesis]. Aix-en-Provence: Université de droit, d'économie et
1118 des sciences d'Aix-Marseille.

1119 Onoratini, G., Cauche, D., Celiberti, V., & Simon, P. (2010). Le Noaillien dans le
1120 complexe Gravettien du sud-est de la France. *Bulletin du Musée d'Anthropologie*
1121 *préhistorique de Monaco*, 50, 39–53.

1122 Otte, M. (Ed.) (2013). *Les Gravettiens*. Paris: Errance.

1123 Palma di Cesnola, A. (1993). *Il Paleolitico Superiore in Italia*. Firenze: Garlati e
1124 Razzai.

1125 Pesesse, D. (2013). Le Gravettien existe-t-il ? Le prisme du système technique
1126 lithiques. In M. Otte (dir.) *Les Gravettiens* (pp. 66–104). Paris: Errance.

1127 Pesesse, D. (2017). Is it still appropriate to talk about the Gravettian? Data from lithic
1128 industries in Western Europe. *Quartär*, 64, 107–128.

1129 Peterson, A. T., Papeş, M., & Soberón, J. (2008). Rethinking receiver operating
1130 characteristic analysis applications in ecological niche modeling. *Ecological Modelling*,
1131 213(1), 63–72. <https://doi.org/10.1016/j.ecolmodel.2007.11.008>

1132 Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martinez-Meyer, E.,
1133 Nakamura, M., & Araújo, M. B. (2011). *Ecological niches and geographic distributions*.
1134 Princeton, NJ: Princeton University Press.

1135 Peterson, A. T., & Soberón, J. (2012). Species distribution modeling and ecological
1136 niche modeling: getting the concepts right. *Natureza & Conservação*, 10(2), 102–107.
1137 <https://doi.org/10.4322/natcon.2012.0>

1138 Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling
1139 of species geographic distributions. *Ecological Modelling*, 190(3–4), 231–259.
1140 <https://doi.org/10.1016/j.ecolmodel.2005.03.026>

1141 Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., & Blair, M. E. (2017).
1142 Opening the black box: an open-source release of Maxent. *Ecography*, 40(7), 887–893.
1143 <https://doi.org/10.1111/ecog.03049>

1144 Pottier, C. (2005). *Le Gravettien moyen de l’abri Pataud (Dordogne, France): le*
1145 *niveau 4 et l’éboulis 3/4. Etude technologique et typologique de l’industrie lithique*. [doctoral
1146 thesis]. Paris: Muséum d’Histoire Naturelle de Paris.

1147 Pottier, C. (2006). Productions lamellaires et burins du Raysse du Gravettien Moyen
 1148 de l'Abri Pataud (Dordogne, France). In Lebrun-Ricalens F. (Ed.) *Burins préhistoriques :
 1149 formes, fonctionnements, fonctions* (n°2, pp. 121–140). Luxembourg: Archéologiques.

1150 Pradel, L. (1953). Précisions sur les Burins d'Angles et les Burins Plans. *Congrès
 1151 Préhistorique de France*, 545–552.

1152 Pradel, L. (1959). Le niveau de Noailles d'Oreille-d'Enfer, Les Eyzies-de-Tayac
 1153 (Dordogne). *Bulletin de la Société préhistorique de France*, 56(3), 228–235.

1154 Qiao, H., Peterson, A. T., Campbell, L. P., Soberón, J., Ji, L., & Escobar, L. E. (2016).
 1155 NicheA: creating virtual species and ecological niches in multivariate environmental
 1156 scenarios. *Ecography*, 39(8), 805–813. <https://doi.org/10.1111/ecog.01961>

1157 Qiao, H., Escobar, L. E., & Peterson, A. T. (2017). Accessible areas in ecological
 1158 niche comparisons of invasive species: Recognized but still overlooked. *Scientific Reports*,
 1159 7(1). <https://doi.org/10.1038/s41598-017-01313-2>

1160 R Core Team (2019). *R: A Language and Environment for Statistical Computing*
 1161 (version 3.6.1). R. Vienna, Austria: R Foundation for Statistical Computing.

1162 Rigaud, J.-P. (1982). *Le Paléolithique en Périgord : les données du Sud-Ouest
 1163 sarladais et leurs implications* [doctoral thesis]. Bordeaux: Université de Bordeaux I.

1164 Rigaud, J.-P. (1988). The Gravettian Peopling of Southwestern France. In Dibble H. L.
 1165 & Montet-White A. *Upper Pleistocene prehistory of Western Eurasia* (Vol. 54, pp. 387–397).
 1166 Philadelphia: University Museum Monograph.

1167 Rigaud, J.-P. (2008). Les industries lithiques du Gravettien du nord de l'Aquitaine
 1168 dans leur cadre chronologique. *Paléo*, 20, 381–398. <https://doi.org/2101-0420>

1169 Rigaud, J.-P. (2011). Révision de quelques archéoséquences de référence du
 1170 Gravettien du nord de l'Aquitaine. In Goutas N., Klaric L., Pesesse D., Guillermin P. (dir.) *À*
 1171 *la recherche des identités gravettiennes. Actualités, questionnements, perspectives* (Mémoire
 1172 52, pp. 175–183). Paris: Société préhistorique française.

1173 Roebroeks, W., Mussi, M., Svodoba, J., Fennema, K. & Hahn, J. (Eds.). (1999).
 1174 *Hunters of the golden age: The Mid Upper Palaeolithic of Eurasia 30,000 - 20,000 BP*.
 1175 University of Leiden.

1176 Roussot, A. (1985). Inventaire typologique des industries de Laussel (tableaux inédits
 1177 de Jean Bouyssonie). *Bulletin de la Société préhistorique française*, 82(1), 15–19.
 1178 <https://doi.org/10.3406/pal.1990.1412>

1179 Sackett, J. (1999). *The Archaeology of Solvieux. An Upper Paleolithic Open Air Site in*
 1180 *France*. Los Angeles (CA): Institute of Archaeology.

1181 Santaniello, F. (2016). *Il Gravettiano dell'Italia tirrenica nel contesto mediterraneo:*
 1182 *definizione delle strategie di insediamento e mobilità attraverso lo studio delle materie prime*
 1183 *e delle industrie litiche* [doctoral thesis]. Trento, IT: Università degli studi di Trento,
 1184 Université Nice Sophia Antipolis.

1185 Santaniello, F., & Grimaldi, S. (2019). Le Gravettien entre l'Italie et la France :
 1186 nouvelles données sur l'industrie lithique de l'Abri Mochi (Balzi Rossi, Italie). *Bulletin de la*
 1187 *Société Préhistorique Française*, 116(2), 255–282.

1188 Sarrazin, C. (2017). *Les burins-nucléus du Raysse de la collection des Morts (Grotte*
 1189 *des Morts, Brive-la-Gaillarde). Interprétation du schéma opératoire et analyse des séquences*.
 1190 [master thesis]. Rennes: Université Rennes 2.

1191 Sarrazin, C. (2018). *Plasenn-al-Lomm et la stabilité du système technique rayssien*
1192 [master thesis]. Rennes: Université Rennes 2.

1193 Schmider, B. (1969). Contribution à l'étude du Périgordien supérieur de La Rochette
1194 (Dordogne). *Gallia préhistoire*, 12(1), 259–271. <https://doi.org/10.3406/galip.1969.1342>

1195 Schoener, T. W. (1968). The Anolis Lizards of Bimini: Resource Partitioning in a
1196 Complex Fauna. *Ecology*, 49(4), 704–726. <https://doi.org/10.2307/1935534>

1197 Siddall, M., Rohling, E. J., Almogi-Labin, A., Hemleben, Ch., Meischner, D.,
1198 Schmelzer, I., & Smeed, D. A. (2003). Sea-level fluctuations during the last glacial cycle.
1199 *Nature*, 423, 853–858. <https://doi.org/10.1038/nature01690>

1200 Sima, Adriana, Rousseau, D.-D., Kageyama, M., Ramstein, G., Schulz, M., Balkanski,
1201 Y., Antoine, P., Dulac, F., & Hatté, C. (2009). Imprint of North-Atlantic abrupt climate
1202 changes on western European loess deposits as viewed in a dust emission model. *Quaternary*
1203 *Science Reviews*, 28, 2851–2866. <https://doi.org/10.1016/j.quascirev.2009.07.016>

1204 Sima, A., Kageyama, M., Rousseau, D.-D., Ramstein, G., Balkanski, Y., Antoine, P.,
1205 & Hatté, C. (2013). Modeling dust emission response to North Atlantic millennial-scale
1206 climate variations from the perspective of East European MIS 3 loess deposits. *Climate of the*
1207 *Past*, 9(4), 1385–1402. <https://doi.org/10.5194/cp-9-1385-2013>

1208 Simonet, A. (2009a). *Les gravettiens des Pyrénées. Des armes aux sociétés*. [doctoral
1209 thesis]. Toulouse: Université Toulouse Le Mirail - Toulouse II.

1210 Simonet, A. (2009b). A Gravettian knapping workshop at Tercis (Landes). A probable
1211 case of apprenticeship in the fabrication of lithic weapon tips. *P@lethnologie*, 1, 183–210.

1212 Simonet, A. (2010). Les armatures lithiques du Gravettien à burins de Noailles du
 1213 niveau G de la grotte des Enfants (Balzi Rossi, Ligurie, Italie): premiers éléments d'enquête.
 1214 *Bulletin du Musée d'Anthropologie préhistorique de Monaco*, 50, 55–68.

1215 Simonet, A. (2011a). La Pointe des Vachons : nouvelles approches d'un fossile
 1216 directeur controversé du Gravettien à partir des exemplaires du niveau IV de la grotte
 1217 d'Isturitz (Pyrénées-Atlantiques, France) et des niveaux 4 des abris 1 et 2 des Vachons
 1218 (Charente, France). *Paléo*, 22, 271–298.

1219 Simonet, A. (2011b). Le Gravettien du chantier I de Brassempouy (Landes, France). In
 1220 Goutas N., Klaric L., Pesesse D., Guillermin P. (dir.) *À la recherche des identités*
 1221 *gravettiennes. Actualités, questionnements, perspectives* (Mémoire 52, pp. 57–66). Paris:
 1222 Société préhistorique française.

1223 Simonet, A. (2013). Des apprentis gravettiens ont-ils confectionné des armatures
 1224 lithiques à Tercis (Landes, France) ? *Paléo*, 23, 1–29.

1225 Simonet, A. (2017). Gravettians at Brassempouy (Landes, France), 30,000 BP: a semi-
 1226 sedentary territorial organization? *World Archaeology*, 49(5), 648–665.
 1227 <https://doi.org/10.1080/00438243.2017.1359109>

1228 Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of
 1229 species. *Ecology Letters*, 10(12), 1115–1123. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2007.01107.x)
 1230 [0248.2007.01107.x](https://doi.org/10.1111/j.1461-0248.2007.01107.x)

1231 Soberón, J., & Peterson, A. T. (2005). Interpretation of Models of Fundamental
 1232 Ecological Niches and Species' Distributional Areas. *Biodiversity Informatics*, 2(0).
 1233 <https://doi.org/10.17161/bi.v2i0.4>

1234 Tixier, J. (1958). Les burins de Noailles de l'Abri André Ragout, Bois-du-Roc,
 1235 Vilhonneur (Charente). *Bulletin de la Société préhistorique française*, 55(10), 628–644.
 1236 <https://doi.org/10.3406/bspf.1958.3707>

1237 Touzé, O. (2011). Caractérisation de la “méthode du Raysse” à Bassaler-nord et au
 1238 Raysse (Corrèze, France). *Archéo-situla*, 31, 3–27.

1239 Touzé, O. (2013). De la signification du Noaillien et du Rayssien. In de la Heras C.,
 1240 Lasheras J.A., Arrizalaga A., de la Rasilla M. (dir.). *Pensando el Gravetiense : nuevos datos*
 1241 *para la region cantabrica en su contexto peninsular y pirenaico* (monografias 23, pp. 383–
 1242 400). Santander, Madrid: Museo Nacional y Centro de Investigacion de Altamira.

1243 Vignoles, A., Klaric, L., Banks, W. E., & Baumann, M. (2019). Le Gravettien du
 1244 Fourneau du Diable (Bordeilles, Dordogne) : révision chronoculturelle des ensembles
 1245 lithiques de la “Terrasse inférieure”. *Bulletin de la Société Préhistorique Française*, 116(3),
 1246 455–478.

1247 Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J.-C., McManus, J. F., Lambeck,
 1248 K., Balbon, E., & Labracherie, M. (2002). Sea-level and deep water temperature changes
 1249 derived from benthic foraminifera isotopic records. *Quaternary Science Reviews*, 21, 295–
 1250 305. [https://doi.org/10.1016/S0277-3791\(01\)00101-9](https://doi.org/10.1016/S0277-3791(01)00101-9)

1251 Warren, D. L. (2012). In defense of ‘niche modeling.’ *Trends in Ecology & Evolution*,
 1252 27(9), 497–500. <https://doi.org/10.1016/j.tree.2012.03.010>

1253 Warren, D. L., Glor, R. E., & Turelli, M. (2008). Environmental Niche Equivalency
 1254 versus Conservatism: Quantitative Approaches to Niche Evolution. *Evolution*, 62(11), 2868–
 1255 2883. <https://doi.org/10.1111/j.1558-5646.2008.00482.x>

1256 Warren, D. L., & Seifert, S. N. (2011). Ecological niche modeling in Maxent: the
1257 importance of model complexity and the performance of model selection criteria. *Ecological*
1258 *Applications*, 21(2), 335–342. <https://doi.org/10.1890/10-1171.1>

1259 [Warren, D. L., Matzke, N., Cardillo M. & Dinnage, R. \(2017\). ENMTools: Analysis of](#)
1260 [niche evolution using niche and distribution models. R. package version 0.2.](#)

1261 Wojtal, P., Haynes, G., & Wilczynski, J. (Eds.). (2015). *World of Gravettian Hunters*
1262 (Quaternary International, Vols. 359–360).

1263