Out-of-phase Late Pleistocene glacial maxima in the Western Alps reflect past changes in North Atlantic atmospheric circulation

Natacha Gribenski, Pierre G Valla, Frank Preusser, Thibault Roattino,
Christian Crouzet, Jean-François Buoncristiani

► To cite this version:

Natacha Gribenski, Pierre G Valla, Frank Preusser, Thibault Roattino, Christian Crouzet, et al.. Out-of-phase Late Pleistocene glacial maxima in the Western Alps reflect past changes in North Atlantic atmospheric circulation. *Geology*, 2021, 49 (9), pp.1096-1101. 10.1130/G48688.1 . hal-03388270

HAL Id: hal-03388270

<https://hal.science/hal-03388270>

Submitted on 20 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Out-of-phase Late Pleistocene glacial maxima in the Western Alps reflect past changes in North Atlantic atmospheric circulation

Natacha Gribenski¹, Pierre G. Valla^{2,1}, Frank Preusser³, Thibault Roattino², Christian Crouzet², Jean-François Buoncristiani⁴

¹*Institute of Geological Sciences, University of Bern, Bern, Switzerland*

²*ISTerre, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, IRD, IFSTAR, Grenoble, France*

³*Institute of Earth and Environmental Sciences, University of Freiburg, Freiburg, Germany*

⁴*UMR 6282 Biogéosciences, CNRS/Univ Bourgogne Franche-Comté, Dijon, France*

ABSTRACT

Paleoglacier reconstructions in the northern and southern forelands of the European Alps indicate a synchronous Late Pleistocene glacial maximum during Marine Isotope Stage (MIS) 2, in phase with global ice volume records. However, strong controversy remains in the western foreland, where scarce and indirect dating as well as modelling studies suggest out-of-phase glacial maxima with the rest of the Alps. New luminescence dating brings the first direct Late Pleistocene glacial chronology for the western Alpine foreland and reveals two major glacier advances of similar maximum extent, at ca. 75-60 and ca. 40-30 ka, coinciding with MIS 4 and late MIS 3. We propose that asynchrony in glacial maxima between the western and the northern/southern Alpine forelands results from a progressive spatial reorganization of the

atmospheric circulation over the North Atlantic, in response to Northern Hemisphere ice-sheet fluctuations. While such feedback mechanism has emerged from general circulation models, our Late Pleistocene paleoglacial reconstruction permits to track the spatio-temporal evolution of moisture advection patterns over Western Europe.

INTRODUCTION

Ice-core and marine paleoclimate archives show two main periods of global temperature minima during the Late Pleistocene (LP; the last ca. 130 kyr; NGRIP, 2004), coinciding with Marine Isotope Stage (MIS) 4 and 2 (71-57 and 29-14 ka; Lisiecki and Raymo, 2005). Marine records indicate the main peak in global ice volume, also largely reported from terrestrial glacier archives, during the Last Glacial Maximum (LGM; ca. 27-19 ka; Clark et al., 2009). However, local glacial maxima well before the LGM are increasingly recorded in polar and mountain regions (Hughes et al., 2013; Batchelor et al., 2019). The emerging picture of variable ice expansion in both time and space likely reflects the regional effect of dynamic atmospheric circulations (Löfverström et al., 2014). Accurate glacial reconstructions across the globe are thus crucial to constrain past atmospheric circulation and paleo-precipitation patterns (e.g. Kuhlemann et al., 2008).

In the European Alps, major LP glaciations are recorded in prominent morainic lobes spreading over several tens of kilometers in the forelands (Ehlers et al., 2011; Fig. 1). In the northern and southern forelands, synchronous LP glacial maxima at ca. 26-23 ka have been documented (Monegato et al., 2017). On the western side, the most extensive LP glaciation occurred in the Lyon area (Fig. 1a), where an ice lobe draining the Arve (western Mont Blanc massif) and the Isère catchments developed ~50 km westward into the piedmont (Coutterand et al., 2009). In contrast to other Alpine forelands, scarce and indirect radiocarbon data in this area suggest older

LP ice-maxima than the global LGM (Mandier et al., 2003). Such asynchronous paleoglacial activity is supported by numerical simulations suggesting transgressive glacier dynamics across the Alps, with the observed maximum ice extent in the Western Alps only reached when considering paleo-precipitation partitioning or older (i.e. MIS 4) glacier advance (Becker et al., 2016; Seguinot et al., 2018).

To further examine the hypothesis of asynchronous paleoglacial dynamics across the Alps, a robust chronology in the Western Alps is now critical. Radiocarbon and cosmogenic nuclide exposure dating are challenging in this area due to the scarcity of organic material incorporated within glacial deposits and in-place moraine boulders, respectively. Here, we present the first direct chronology of LP ice-maximum extent in the Lyon area, using optically stimulated luminescence (OSL) dating of glaciofluvial sediments. These results not only enlighten the debate around the paleoglacial history of the Western Alps, but also have major implications in terms of past atmospheric circulation over the North Atlantic and Western Europe.

SETTING AND METHODS

Two main moraine complexes, separated by ~25 km, define the Middle ($\geq$ MIS 6; External Moraine Complex, EMC) and Late (Internal Moraine Complex, IMC) Pleistocene ice-maximum extents in the Lyon area (Buoncristiani and Campy, 2011; Fig. 1a). While the EMC presents discrete, highly-degraded and discontinuous moraine crests, the IMC is characterized by a series of well-defined frontal moraine ridges (Fig. 1a).

Glaciofluvial valleys (i.e. inactive today) expand over several tens of kilometers westward from the IMC (Fig. 1a), containing large sediment infills organized in step terraces. In the western and southern sectors, the IMC lays on a bedrock topographic step rising abruptly >200 m above the present-day lowland (Fig. 1b). This topographic configuration allows to unambiguously relate

glaciofluvial sediment deposition and terraces formation to major ice-extent periods (i.e. IMC ridges), when ice and meltwater could overtop this escarpment. Previous geomorphic and sedimentological studies identified at least three main terrace levels which, from high (T1/T2) to low (T4), directly connect to external (M2) to internal (M4) moraine ridges from the IMC (Mandier, 1988; Roattino et al., 2021; Fig 1a), and thus were formed by aggradation during periods of glacial advances (Marren, 2005). Typical coarse braided-river facies exposed along the terraces do not permit to argue for major discontinuity in terrace construction. Discrete outermost IMC ridges (M1) are visible only in the NW sector, suggesting an additional glacial stage hardly preserved through subsequent glacial periods (overridden moraine patterns; Fig. DR1 in the GSA Data repository), and with possible remnants in the east and south sectors of the IMC indistinguishable from the M2 deposits. In the absence of direct numerical dating, it remains challenging to quantitatively relate the observed glacial stages to distinct glaciations or short-term glacier oscillations.

We targeted the three main terrace levels (T1/T2 to T4; Fig. 1a), with six samples collected from well-sorted sand lenses exposed in quarry sections (Table 1, Appendix DR1). Samples were taken as close as possible to the terrace surfaces to capture the last depositional period related to a glacial stage. Two additional samples were collected from a glaciofluvial unit underlying till from an M1 ridge in the NW sector (PIZ site; Fig. 1a, Table 1), providing maximum age constraints of the LP outermost ice extent.

Luminescence signals and equivalent doses (D_e) were measured from quartz (green OSL) and feldspar (infra-red stimulation at 50°C and 225°C, IR₅₀ and post-IRIR₂₂₅) single grains (SG, 200-250 µm fraction). Methodological details are given in the GSA Data repository (DRII). Total burial dose since sample deposition was determined using the Central Age Model (CAM D_e) or Finite Mixture statistical model (FMM D_e) in case of partial bleaching diagnosis (Galbraith and

Roberts, 2012). Feldspar burial doses were corrected for anomalous fading (20-30% increase, Table DR3) and final ages were calculated using sample-specific dose rate (2-3 Gy ka⁻¹ range, Tables DR4,5), derived from element concentrations measured by laboratory gamma spectrometry.

RESULTS

SG quartz OSL data could only be acquired for three samples (EYP1, OYT1, SPC5; Table 1), for which ~2% of measured grains emit a suitable luminescence signal. This is consistent with low luminescence sensitivity typically observed in quartz from crystalline bedrock and/or with limited transport (Sakawuchi et al., 2011). All eight samples reveal good feldspar luminescence characteristics (20-40% suitable grains).

SG quartz D_e distributions exhibit normal-like to moderately-skewed shapes, while SG feldspar D_e distributions are widely spread, asymmetric and multimodal, with a significant fraction of saturated grain for post-IRIR₂₂₅ (Figs. 2b, DR6,7). In addition, feldspar CAM D_e are significantly larger (>50%) than for quartz. Both complex SG D_e distributions and CAM D_e differences between feldspar and quartz are strong indicators for limited pre-depositional exposure to light, typical for proximal glaciofluvial sediments (Duller, 2006). Partial bleaching, leading to age overestimate, was diagnosed for most of the samples and especially for feldspar signals. We thus applied statistical techniques (i.e. FMM model) to extract D_e from the best-bleached grain sub-populations, giving more accurate age determination. For final age calculation, quartz OSL and feldspar IR₅₀ were selected, based on their best bleaching potential.

Quartz OSL and feldspar IR₅₀ ages obtained from the highest terrace in the central part of the study area (T1/T2: SPC5; Fig. 1a) and from the intermediate (T3: ART2, OYT1, EYP1) to lower (T4: ART1) terrace levels overlap between ca. 42 and 39 ka (Fig. 3b; Table 1). Where both

available, quartz OSL and feldspar IR_{50} ages agree within uncertainties, bringing further confidence in the statistical treatment applied to complex feldspar IR_{50} D_e distributions. The southern highest terrace level (PEN1) resulted in a distinctly older IR_{50} age of 63.0 ± 11.4 ka, coinciding with two IR_{50} ages of 60.8 ± 12.3 and 76.4 ± 15.1 ka obtained for the glaciofluvial unit underlying the north-western M1 ridge (PIZ1 and PIZ2; Table 1).

DISCUSSION

Luminescence dating indicates two periods of glaciofluvial aggradation beyond the IMC: between ca. 60-75 ka and ca. 30-40 ka, coinciding with MIS 4 and late MIS 3 respectively. This result strongly suggests that western Alpine glaciers reached the IMC at least twice during the LP, with similar maximum ice-extent configuration. The main terrace-moraine sequence (T2/T3/T4 - M2/M3/M4) observed in the area would hence result from glacier margin oscillations during a single glaciation period (i.e. late MIS 3), whereas scarce and outer remnant deposits (T1 - M1) were preserved from an earlier MIS 4 glacial maxima.

Our new MIS 4 chronology further corroborates earlier studies based on scarce chronological data in the Swiss Alpine foreland (Gaar et al., 2019 and references therein), and in the nearby Pyrenees (Delmas et al., 2011), as well as results from numerical simulations (Fig. 3a; Seguinot et al., 2018), all of which proposed significant glaciations in the Western European mountain ranges at that time. The late MIS 3 glaciation clearly predates the last glacial maxima recorded in the northern and southern Alpine forelands during MIS 2 (Monegato et al., 2017). However, an extended MIS 3 ice advance has been suggested in the southwestern Alpine foreland (e.g. Ivy-Ochs et al., 2018), while valleys and forelands in the northern and central Alps were reported to be ice-free before 30 ka (Preusser et al., 2011; Barret et al., 2017). The absence of MIS 2 ages in our dataset, in particular for the lowest (T4) terrace level (ART 1, 33.8 ± 7.3 ka), suggests that no

glacier advance reached the IMC after late MIS 3. This is further supported by radiocarbon ages (Fig. 1a; Mandier et al., 2003) of ca. 27 to >35 ka from a paleosol above till (i.e. minimum ages) in the NW sector of the IMC (Balan) and of 28-30 ka from glacio-lacustrine deposits 10-20 km upstream of the IMC (Moras and Malville), which suggest shorter glacier extent in the area during MIS 2. It is unlikely that the time lag (ca. 10 kyr) observed for the last major advance between the Lyon and northern/southern central Alpine forelands is caused by methodological uncertainties, as the applied dating approaches have shown age consistency on such timescales in similar glacial settings (e.g. Smedley et al., 2016; Gribenski et al., 2018).

In addition to multiple and similar LP glacial maxima in the western Alpine foreland predating the global LGM, our data indicates spatially and temporarily variable paleo-glacier extent patterns across the European Alps, which likely reflect differential precipitation distribution combined with periods of global cooling. In contrast to the modern dominance of north-west atmospheric circulation, it has been proposed that south-dominated moisture advection prevailed across the Alps during the LGM (Florineth and Schlüchter, 2000; Luetscher et al., 2015; Monegato et al., 2017). The southward latitudinal shift of the North Atlantic storm track, as far as 40°N, was forced by the development of large ice masses and sea ice in the North Hemisphere (NH; Fig. 3c), as indicated by atmospheric general circulation models (e.g. Hofer et al., 2012). The extensive MIS 4 glaciation in the western Alpine foreland coincides with a major cooling period observed at the global and regional scales (NGRIP, 2004; Helmens, 2014; Moseley et al., 2020). Further elaborating on the above circulation model, we propose that the moderate expansion of the NH ice sheets during MIS 4, as indicated by deep-marine sediment records (Lisiecki and Raymo, 2005), could have similarly initiated a more discrete storm-track migration southward (Löffverström et al., 2014). Moisture was thus dominantly brought from the west and promoted high precipitation in the west-facing massifs of the Western Alps, enabling favorable

conditions (i.e. wet and cold) for large piedmont glaciers. While cold MIS 4 conditions are expected across the entire Alps, limited precipitation in the Central and Eastern Alps may have resulted in more restricted ice extents in the main northern/southern central forelands, with glacial deposits likely eroded during the greater MIS 2 extent.

During MIS 3, global ice records indicate a decrease of temperatures already at 40 ka before reaching minima at 25-20 ka (NGRIP, 2004). Few existing long-term regional records suggest similar trends in SW Europe, even with a potentially earlier cooling onset at 45 ka (Moreno et al., 2014; Moseley et al., 2020). After significant shrinkage during early MIS 3, NH ice sheets started to expand again from ca. 40 ka (Batchelor et al., 2019). Similar to our proposed MIS 4 scenario, dominant western moisture advection would have been maintained throughout mid/late MIS 3 with a progressive southward shift of the polar storm track, again favoring glacier extent in the Western Alps. This configuration changed with the ongoing growth of NH ice sheets towards the LGM, which further pushed the polar storm track southward and led shortly after 30 ka to the main southerly moisture advection over the European Alps (Luetscher et al., 2015). Interestingly, the lacustrine record at Les Echets (Fig. 1a) also points towards a major change in lake productivity at ca. 28-30 ka, with a climatic transition from highly-oscillating to relatively stable, dry and cold conditions (Veres et al., 2009). We propose that this change in atmospheric circulation resulted in significant moisture decrease over the Western Alps, while enabling maximum glacier extent in the southern and northern Alpine forelands during MIS 2.

CONCLUSIONS

Our luminescence chronology constitutes the first direct evidence of two major Late Pleistocene glaciations reaching similar maximum extents in the western Alpine foreland, during MIS 4 (76-60 ka) and late MIS 3 (42-29 ka). These multiple and early glacial maxima revealed in the

western foreland contrast with MIS 2 maximum ice extent recorded in the northern and southern sides of the European Alps, in phase with the global LGM. Such spatial variability in glacier fluctuations has been observed elsewhere around the globe and likely translates swaying atmospheric circulation and precipitation partitioning during the last glacial cycle. We propose that such configuration may be related to the gradual migration of the polar storm track, forced by the progressive development of NH ice sheets. This study further highlights the importance of regional paleo-glacier reconstructions and numerical chronologies for quantifying past changes in atmospheric circulation patterns.

ACKNOWLEDGMENTS

This work was supported by the SNSF (PP00P2_170559), INSU and USMB (AAP Montagne). We thank Gachet TP, Eiffage, Cemex Granulat Rhône Méditerranée, Nord Isère Matériaux and Vicat Granulat for providing access to gravel pits. We thank the four anonymous reviewers for their valuable suggestions.

REFERENCES

- Barrett, S.J., Starnberger, R., Tjallingii, R., Brauer, A., and Spötl, C., 2017, The sedimentary history of the inner-alpine Inn Valley (Austria): extending the Baumkirchen type section further back in time with new drilling: *Journal of Quaternary Science*, v. 32, p. 63–79, <https://doi.org/10.1002/jqs.2924>
- Batchelor, C.L., Margold, M., Krapp, M., Murton, D.K., Dalton, A.S., Stokes, C.R., Murton, J.B., and Manica, A., 2019, The configuration of Northern Hemisphere ice sheets through

the Quaternary: *Nature Communications*, v. 10, p. 3713, <https://doi.org/10.1038/s41467-019-11601-2>

Becker, P., Seguinot, J., Jouvet, G., and Funk, M., 2016, Last Glacial Maximum precipitation pattern in the Alps inferred from glacier modelling: *Geographica Helvetica*, v. 71, p. 173–187, <https://doi.org/10.5194/gh-71-173-2016>

Buoncrisiani, J.-F. and Campy, M., 2011, Quaternary glaciations in the French Alps and Jura: *Developments in Quaternary Sciences*, v. 15, p. 117–126, <https://doi.org/10.1016/B978-0-444-53447-7.00010-6>

Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X., Hostetler, S.W. and McCabe, A.M., 2009, The Last Glacial Maximum: *Science*, v. 325, p. 710–714, <https://doi.org/10.1126/science.1172873>

Coutterand, S., Schoeneich, P., Nicoud, G., 2009, Le lobe glaciaire lyonnais au maximum würmien : glacier du Rhône ou/et glaciers savoyard?. Philip Deline, Ludovic Ravanel. *Neige et glace de montagne : Reconstitution, dynamique, pratiques*, Collection EDYTEM - Cahiers de Géographie, v. 8, p. 11–22, <halshs-00389085>

Delmas, M., Calvet, M., Gunnell, Y., Braucher, and R., Bourlès, D., 2011, Paleogeography and ¹⁰Be exposure-age chronology of Middle and Late Pleistocene glacier systems in the northern Pyrenees: implications for reconstructing regional paleoclimates: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 305, p.109–122, <https://doi.org/10.1016/j.palaeo.2011.02.025>

Duller, G.A.T., 2006, Single grain optical dating of glacial deposits: *Quaternary Geochronology*, v. 1, p. 296–304, <https://doi.org/10.1016/j.quageo.2006.05.018>

Ehlers, J., Gibbard, P.L., and Hughes, P.D., eds., 2011, Quaternary Glaciations - Extent and
 Chronology: A Closer Look: Developments in Quaternary Science, v. 15. Elsevier,
 Amsterdam, 1108 p.

Florineth, D., and Schlüchter, C., 2000, Alpine evidence for atmospheric circulation patterns in
 Europe during the Last Glacial Maximum: Quaternary Research, v. 54, p. 295–308,
<https://doi.org/10.1006/qres.2000.2169>

Gaar, D., Graf, H.R., and Preusser, F., 2019, New chronological constraints on the timing of Late
 Pleistocene glacier advances in northern Switzerland, E & G Quaternary Science Journal,
 v. 68, p. 53–73, <https://doi.org/10.5194/egqsj-68-53-2019>

Galbraith, R.F., and Roberts, R.G., 2012, Statistical aspects of equivalent dose and error
 calculation and display in OSL dating: An overview and some recommendations:
 Quaternary Geochronology, v. 11, p. 1–27, <https://doi.org/10.1016/j.quageo.2012.04.020>

Gribenski, N., Jansson, K.N., Preusser, F., Harbor, J.M., Stroeve, A.P., Trauerstein, M.,
 Blomdin, R., Heyman, J., Caffee, M.W., Lifton, N., and Zhang, W., 2018, Re-evaluation
 of MIS 3 glaciation using cosmogenic radionuclide and single grain luminescence ages,
 Kanas Valley, Chinese Altai: Journal of Quaternary Science, v. 33, p. 55–67,
<https://doi.org/10.1002/jqs.2998>

Helmens, K.F., 2014, The Last Interglacial Glacial cycle (MIS 5-2) re-examined based on long
 proxy records from central and northern Europe: Quaternary Science Reviews, v. 86, p.
 115–143, <http://dx.doi.org/10.1016/j.quascirev.2013.12.012>

Hofer, D., Raible, C.C., Merz, N., Dehnert, A., and Kuhlemann, J., 2012, Simulated winter
 circulation types in the North Atlantic and European region for preindustrial and glacial
 conditions: Geophysical Research Letters, v. 39, L15805,
<http://dx.doi.org/10.1029/2012GL052296>

260 Hughes, P.D., Gibbard, P.L., Ehlers, J., 2013, Timing of glaciation during the last glacial cycle:
 261 evaluating the concept of a global ‘Last Glacial Maximum’ (LGM): *Earth-Science*
 262 *Reviews*, v. 125, p. 171–198, <https://doi.org/10.1016/j.earscirev.2013.07.003>
 263 Ivy-Ochs, S., Lucchesi, S., Baggio, P., Fioraso, G., Gianotti, F., Monegati, G., Graf, A.A., Akçar,
 264 N., Christl, M., Carraro, F., Gabrialla Forno, M., and Schlüchter, C., 2018, New
 265 geomorphological and chronological constraints for glacial deposits in the Rivoli
 266 Avigliana end-moraine system and the lower Susa Valley (Western Alps, NW Italy):
 267 *Journal of Quaternary Science*, v. 33, p. 550–562, <https://doi.org/10.1002/jqs.3034>
 268 Kuhlemann, J., Rohling, E.J., Krumrei, I., Kubik, P., Ivy-Ochs, S. and Kucera, M., 2008,
 269 Regional synthesis of Mediterranean atmospheric circulation during the Last Glacial
 270 Maximum: *Science*, v. 321, p. 1338–1340, <https://doi.org/10.1126/science.1157638>
 271 Lisiecki, L.E. and Raymo, M.E., 2005, A Pliocene-Pleistocene stack of 57 globally distributed
 272 benthic ^{18}O records: *Paleoceanography*, v. 20, p. 1–17,
 273 <https://doi.org/10.1029/2004PA001071>
 274 Löffverström, M., Caballero, R., Nilsson, J., and Kleman, J., 2014, Evolution of the large-scale
 275 atmospheric circulation in response to changing ice sheets over the last glacial cycle:
 276 *Climate of the Past*, v. 10, p. 1453–1471, <https://doi.org/10.5194/cp-10-1453-2014>
 277 Luetscher, M., Boch, R., Sodemann, H., Spötl, C., Cheng, H., Edwards, R. L., Frisia, S., Hof, F.,
 278 and Müller, W., 2015, North Atlantic storm track changes during the Last Glacial
 279 Maximum recorded by Alpine speleothems: *Nature Communications*, v. 6, p. 6344,
 280 <https://doi.org/10.1038/ncomms7344>
 281 Mandier, P., 1988, Le relief de la moyenne vallée du Rhône au Tertiaire et au Quaternaire: essai
 282 de synthèse paléogéographique. Bureau de Recherches Géologiques et Minières, Orléans,
 283 865 pp.

284 Mandier, P., Evin, J., Argant, J., and Petiot, R., 2003, Chronostratigraphie des accumulations
 285 würmiennes dans la moyenne vallée du Rhône. L'apport des dates radiocarbone:
 286 Quaternaire, v. 14/2, p. 113–127, <https://doi.org/10.3406/quate.2003.1735>
 287 Marren, P., 2005, Magnitude and frequency in proglacial rivers: a geomorphological and
 288 sedimentological perspective: Earth Science Reviews, v. 70, p. 203–251,
 289 <https://doi.org/10.1016/j.earscirev.2004.12.002>
 290 Moreno, A., Svensson, A., Brooks, S.J., Connor, S., Engels, S., Fletcher, W., Genty, D., Heiri,
 291 O., Labuhn, I., Persiou, A., Peyron, O., Sadori, L., Valero-Garces, B., Wulf, S., and
 292 Zanchetta, G., 2014, A compilation of Western European terrestrial records 60-8 ka BP:
 293 towards an understanding of latitudinal climatic gradients: Quaternary Science Reviews,
 294 v. 106, p. 167–185. <https://doi.org/10.1016/j.quascirev.2014.06.030>
 295 Monegato, G., Scardia, G., Hajdas, I., Rizzini, F., and Piccin, A., 2017, The Alpine LGM in the
 296 boreal ice-sheets game: Nature Scientific reports, v. 7, p. 2078,
 297 <https://doi.org/10.1038/s41598-017-02148-7>
 298 Moseley, G.E., Spötl, C., Brandstätter, S., Erhardt, T., Luetscher, M., and Edwards, R.L., 2020,
 299 NALPS19: sub-orbital-scale climate variability recorded in northern Alpine speleothems
 300 during the last glacial period: Climate of the Past, v. 16, p. 29–50,
 301 <https://doi.org/10.5194/cp-16-29-2020>
 302 NGRIP members, 2004, High-resolution record of Northern Hemisphere climate extending into
 303 the last interglacial period: Nature, v. 43, p. 147–151,
 304 <https://doi.org/10.1038/nature02805>
 305 Preusser, F., Graf, H. R., Keller, O., Krayss, E., and Schlüchter, C., 2011, Quaternary glaciation
 306 history of northern Switzerland: E&G Quaternary Science Journal, v. 60, p. 21,
 307 <https://doi.org/10.3285/eg.60.2-3.06>

- Roattino, T., Crouzet, C., Buoncristiani, J-F., Tissoux, H., 2021, Geometry of glaciofluvial deposits and dynamics of the Lyonnais lobe ice front during the last glacial period (France, Northern Alps): *Earth Science Bulletin BSGF*, *Accepted*
- Sakawuchi, A.O., Blair, M.W., DeWitt, R., Faleiros, F.M., Hyppolito, T., and Guedes, C.C.F., 2011, Thermal history versus sedimentary history: OSL sensitivity of quartz grains extracted from rocks and sediments: *Quaternary Geochronology*, v. 6, p. 261–272, <https://doi.org/10.1016/j.quageo.2010.11.002>
- Seguinot, J., Ivy-Ochs, S., Jouvet, G., Huss, M., Funk, M., and Preusser, F., 2018, Modelling last glacial cycle ice dynamics in the Alps: *The Cryosphere*, v. 12, p. 3265–3285, <https://doi.org/10.5194/tc-12-3265>
- Smedley, R.K., Glasser, N.F., and Duller, G.A.T., 2016, Luminescence dating of glacial advances at Lago Buenos Aires (~46°S), Patagonia: *Quaternary Science Reviews*, v. 134, p. 59–73, <http://dx.doi.org/10.1016/j.quascirev.2015.12.010>
- Veres, D., Lallier-Vergès, E., Wohlfarth, B., Lacourse, T., Kéravis, D., Björck, S., Preusser, F., Andrieu-Ponel, V., and Ampel, L., 2009, Climate-driven changes in lake conditions during late MIS 3 and MIS 2: a high-resolution geochemical record from Les Echets, France: *Boreas*, v. 38, p. 230–243, <https://doi.org/10.1111/j.1502-3885.2008.00066.x>
- Wirsig, C., Zasadni, J., Christl, M., Akçar, N., and Ivy-Ochs, S., 2016, Dating the onset of LGM ice surface lowering in the High Alps: *Quaternary Science Reviews*, v. 143, p. 37–50, <https://doi.org/10.1016/j.quascirev.2016.05.001>

TABLE & FIGURE CAPTIONS

Table 1: Sample locations and luminescence ages.

Figure 1: (a) Glacial geomorphology of the study area (Mandier, 1988; Roattino et al., 2021). Inset: geomorphic limits of the LP ice maximum in the Alps (Ehlers et al., 2011) with main LGM ice flows (light blue arrows; Wirsig et al., 2016), and location of the study area (black box) with main ice-drainage catchments (dark blue limit, AR: Arve, IS: Isère). (b) Longitudinal topographic profile (LP1) showing bedrock escarpement underlying the IMC ridges and downstream glaciofluvial sediments (undifferentiated), when the glacier reached its LP maximum extent. (c) Topographic cross-profile (ART site; CP1) showing the main three terrace levels. The bedrock/sediment contact is extrapolated from borehole data (Roattino et al., 2021).

Figure 2: Quartz normal-like and feldspar (no fading correction) complex single-grain burial age distributions for a representative sample (SPC5). Consistent quartz and feldspar sample burial ages are obtained using the CAM and FMM models respectively, based on the different partial bleaching diagnosis.

Figure 3: (a) Late Pleistocene Northern Alpine speleothems $\delta^{18}\text{O}$ (purple; Moseley et al., 2020) and global marine sediments $\delta^{18}\text{O}$ (grey curve with MIS in shaded bars; Lisiecki and Raymo, 2005) records, with modelled Alpine ice volume (blue, GRIP paleoclimate forcing; Seguinot et al., 2018). (b) Luminescence ages showing two main groups around MIS 4 (green) and late MIS 3 (orange), consistent with existing radiocarbon ages from the NW sector (green) and upstream (red) of the IMC (Mandier et al., 2003). (c) Conceptual sketch illustrating the southward shift of the North Atlantic storm track forced by gradual NH ice sheet growth (LGM extent in blue), with impact on the paleo-atmospheric circulation (coloured arrows).

354 ¹GSA Data Repository item 201Xxxx, [Details of samples' geomorphological and
355 sedimentological context, and luminescence dating], is available online at
356 www.geosociety.org/pubs/ft20XX.htm, or on request from editing@geosociety.org.

Figure 1

Figure 2

a)

Marine Isotope Stages (MIS)

b)

c)

Gribenski et al., 2021—Table

Table 1. Sample locations and luminescence ages

Sample ID	Lat./ Long. (°N/°E)	Elevation (m a.s.l.)	Depth (m below surface)	Geomorphic unit	Quartz (OSL) age (ka)	Feldspar (IR ₅₀) age (ka)
EYP1	45.496/5.009	262	5	T3	32.9±2.3	29.4±2.8
OYT1	45.575/5.030	265	3.2	T3	36.4±3.8	32.4±4.3
SPC5	45.666/5.015	250	5	T1/T2	38.6±4.5	41.6±7.9
ART1	45.537/5.182	430	3	T4	-	33.8±7.3
ART2	45.536/5.182	420	12	T3 or T4	-	40.6±4.2
PEN1	45.363/5.186	313	12	T1/T2	-	63.0±11.4
PIZ1	45.876/5.088	250	13	underlying M1	-	60.8±12.3
PIZ2	45.876/5.088	256	2	underlying M1	-	76.4±15.1