Tick and tick-borne disease circulation in a changing marine ecosystem

Karen D. McCoy

► To cite this version:

Karen D. McCoy. Tick and tick-borne disease circulation in a changing marine ecosystem. *Climate, Ticks and Disease*, 2021. hal-03388523

HAL Id: hal-03388523

<https://hal.science/hal-03388523>

Submitted on 20 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ch 77. Tick and tick-borne disease circulation in a changing marine ecosystem

Karen D. McCoy

CNRS, France

1. Expert opinion

One may wonder how ticks and tick-borne disease could be present in marine ecosystems. It is true that ticks do not live underwater (although they can survive immersion) and do not exploit fish. However, ticks do feed on other vertebrates that are marine animals, in particular those that spend part of their lives on land, predominantly seabirds. Indeed, at least 30 tick species occur regularly in this ecosystem, and can be found across the globe, including the most extreme areas of both hemispheres (Table 1).

As in terrestrial systems, high infestations by ticks can reduce seabird body condition and lower reproductive success, which in turn can affect seabird population dynamics (Boulinier and Danchin, 1996; Gauthier-Clerc *et al.*, 1998; Mangin *et al.*, 2003). The impact of tick exploitation can be especially important under marine conditions, where thermal stress and the physiological demands of foraging can be arduous. Hyper-infestations by *Ixodes uriae*, resulting in blood loss and hypothermia, was, for example, implicated as a direct cause of death of breeding adult King penguins, *Aptenodytes patagonicus* (Gauthier-Clerc *et al.*, 1998). According to attachment site preferences, heavy infestations can also directly impair locomotion, leading to starvation or predation (Fig. 1a). Even lighter tick infestations have been associated with lower reproductive success and colony abandonment (Bergstrom *et al.*, 1999; Monticelli *et al.*, 2008; Sanz-Aguilar *et al.*, 2020); this can be due to direct impacts on young birds or irritation that perturbs normal feeding and vigilance behaviours in adults. Tropical seabirds are reported to shift colony locations as a means of avoiding the buildup of large numbers of ticks, such as *Ornithodoros capensis* (Fig. 1b; Duffy, 1983; Feare, 1976).

In addition to these direct effects, general surveys of marine ticks have shown that a range of infectious agents are present and circulating in these systems, including diverse bacteria, viruses, and protozoan parasites (Table 1). Some of these agents are known pathogens for humans and/or

their domestic animals, such as the bacterial agents of Lyme disease, carried by *Ixodes uriae*, or of African tick-bite fever (*Rickettsia africae*) found in *Amblyomma loculosum*. The impact of these agents on seabirds (or on ticks, for that matter) remain largely unknown. However, it is likely that the presence of these organisms is of some consequence for seabird health and population ecology. For example, the Soldado virus (Hughes serogroup) has been isolated from sick birds, and experimentally shown to kill chickens (Converse *et al.*, 1975). Infection by *Babesia peircei* has been associated with higher mortality rates in certain seabirds brought to rehabilitation centres, particularly penguins (Parsons *et al.*, 2018, 2017). More generally, quantifying the impact of infection on seabirds, as in other wild populations, is extremely difficult when pathogenicity is chronic in nature and acts in concert with other sources of environmental stress (e.g., food availability, predators, etc).

It is evident that current global changes will alter interactions between ticks and their marine hosts, with subsequent consequences for the circulation of infectious agents. This may occur via several possible mechanisms:

(i) Marine ticks can only exist where their hosts breed. The presence of breeding colonies is dependent on both suitable nesting habitat and adequate food resources for nestlings. The distribution of seabird species has been dynamic over evolutionary time, shifting with changes in climate. For example, the formation of the Antarctic Polar Front during the Miocene led to the formation of a region of extremely high biodiversity, favouring its use by a large diversity of seabirds (Bost *et al.*, 2009). The discovery of fossilized nests on Bermuda showed that albatross bred within the North Atlantic up until approximately 400,000 years ago, likely going extinct during an interglacial period when sea water levels and storm surge were maximised (Olson and Hearty, 2003). Climate-driven changes in gyres where seabirds feed are regularly noted and known to affect adult survival (Fluhr *et al.*, 2017). Current rapid changes will therefore certainly result in the displacement or extinction of certain colonies. Indeed, as warming waters cause thermal expansion and ice sheets to melt, sea levels will rise and will eliminate low-lying colonies. Increases in water column stratification, deoxygenation, and salinity, along with shifts in ocean circulation patterns, will accompany increasing sea surface temperatures and affect the availability of local food sources for birds (Hoegh-Guldberg *et al.*, 2018). Increasing distances between established nesting sites and food sources will put hatchlings at risk (Jenouvrier *et al.*, 2015) and may lead to colony abandonment. Some seabird species may respond rapidly to such changes, particularly if replacement zones for reproduction and foraging can be found. For example, warmer conditions in high latitudes may allow some species to expand their ranges poleward (Péron *et al.*, 2010). However, some species will not have this opportunity and will need to adapt to a warmer world. Given the 'slow pace' of life of

seabirds and the relative speed of current changes, many of these species will likely be driven to extinction, along with their parasites and pathogens (Grose *et al.*, 2020).

(ii) Climate changes may directly impact the presence and abundance of ticks in coastal marine environments. Marine ticks are only found on their hosts for the time of the bloodmeal and have adapted to a range of climates, with only the very coldest and driest polar regions seemingly unoccupied at present. Increasing temperatures and modified humidity regimes will change this distribution. Indeed, studies have shown that many terrestrial tick species are currently expanding their distributions northward due to milder winters that enable tick establishment (see references in Léger *et al.*, 2013). Surveys now suggest the same pattern holds true for marine ticks. For example, *Ixodes uriae* seems to be increasingly present in high Arctic seabird colonies due to milder temperatures (Descamps, 2013). However, as the current distribution of marine ticks is still only partially known, such changes will be difficult to quantify. More importantly, *if ticks expand in some ranges, will they actually retract in others? How plastic are ticks to changing conditions, and is the current rate of change too fast for adaptation to follow?* We know that marine ticks can survive under extreme conditions. For example, *I. uriae* has the largest temperature tolerance range known in ticks, from -30 °C to +30 °C, but is almost absent from seabird colonies in latitudes between 43 °N and 45 °S. Is this due to temperature/humidity interactions (Benoit *et al.*, 2007) or to some other environmental or ecological factor (e.g., competition)? Ticks seem to have large, highly repetitive genomes (Ullmann *et al.*, 2005). This may enable rapid adaptation to changing environmental conditions and shifts in host species use, ensuring their survival. Such a mechanism has been hypothesized to explain the worldwide distribution and broad host range of *I. uriae* (Dietrich *et al.*, 2014b).

(iii) The impact of ticks and their infectious agents can be exacerbated by the concurrent influence of other environmental factors, such as extreme weather conditions, reduced food resources, pollution, and the presence of predators or coinfecting parasites/pathogens. It is the combination of these different individual factors which may have the greatest impact on marine vertebrate populations (Dias *et al.*, 2019), and as a consequence on their interactions, with ticks and tick-borne infectious agents. These additional sources of stress are augmenting rapidly. For example, by-catch and competition with fisheries for depleting stocks have increased over time, leading to seabird population declines (Grémillet *et al.*, 2018). Pollutants are ubiquitous in the marine environment and have reached even the most untouched zones of the planet via ocean currents (Cózar *et al.*, 2017). Indeed, massive accumulations of plastics have formed, destroying foraging and breeding habitats for marine animals and bioaccumulating along trophic food chains with potential chronic toxicity effects (Auta *et al.*, 2017). Climate change is rapidly modifying access to habitats and

food resources used by marine animals, and destabilising populations via an increasing number of extreme weather events (Jenouvrier *et al.*, 2015).

Although seabirds are among the most vagile creatures on earth, most show strong migratory and breeding site fidelity, limiting their movements to within well-defined limits. Exploratory movements only tend to occur when individuals prospect for new locations to breed. This is seen in young birds during the period between fledgling and first reproduction, but also in adult birds that have failed a reproductive attempt. As the proportion of local nest success declines within a colony, the probability that a given breeding pair will leave to prospect increases (Boulinier *et al.*, 2008). Failed birds in colonies with widespread failure will thus show reduced site fidelity the following year, and will have a higher probability of changing breeding partners. This behaviour sets-up a feedback loop in which breeding failures (caused by a combination of potential factors), leads to unstable spatio-temporal population dynamics and a higher probability of dispersal. This, in turn, increases the probability of dispersing parasites (ticks) and infectious agents harboured by the birds, allowing the invasion of new locations and novel hosts. Therefore, the combined effect of different environmental stresses can alter the nature and outcome of host-parasite interactions and result in large-scale changes to the ecological and evolutionary pathways taken by these species. As global changes increase pressure on marine birds, this phenomenon may become more frequent, resulting in the large-scale homogenization of ticks and infectious agents across ranges of suitable ecological conditions.

2. Future predictions

So can we make any concrete predictions? This is a difficult question to answer in a complex and rapidly changing marine ecosystem. However, I expect that certain opportunistic tick species, such as *A. loculosum* and *O. capensis* *sl*, will expand their distributions and explode in numbers. Other, more specialist species, and particularly polar species such as *Ixodes signatus* and *I. kerguelensis*, will likely follow the dynamics of their hosts, and decline in prevalence and abundance. Still other, more plastic, species like *I. uriae*, may be able to maintain the status quo, shifting their distributions to the limits of their physiological capacities and adapting to those host species available locally. What is sure is that these changes will interplay with other environmental factors that affect seabird community dynamics, patterns of colony use, and dispersal. This, in turn, will feedback on the interaction to alter the spread of ticks and the infectious agents these ticks and their host birds carry. As competition for limiting marine resources continue to intensify, contact between humans and wildlife are likely to increase, opening new links between marine and terrestrial tick and tick-borne disease systems. In this light, a better understanding of what limits the spatial distribution of

marine ticks, along with more basic information on the infectious agents they carry, seem essential first steps for predicting future impacts of ticks and tick-borne disease on marine ecosystems and beyond.

References

- Auta, H.S., Emenike, C., and Fauziah, S. (2017) Distribution and importance of microplastics in the marine environment: a review of the sources, fate, effects, and potential solutions. *Environment International*, 102, 165–176.
- Benoit, J.B., Yoder, J.A., Lopez-Martinez, G., Elnitsky, M.A., Lee, R.E. and Denlinger, D.L. (2007) Habitat requirements of the seabird tick, *Ixodes uriae* (Acari : Ixodidae), from the Antarctic Peninsula in relation to water balance characteristics of eggs, nonfed and engorged stages. *Journal of Comparative Physiology B*, 177, 205–215.
- Bergstrom, S., Haemig, P.D. and Olsen, B. (1999) Increased mortality of black-browed albatross chicks at a colony heavily infested with the tick *Ixodes uriae*. *International Journal for Parasitology*, 29, 1359–1361.
- Bost, C.A. *et al.*, 2009. The importance of oceanographic fronts to marine birds and mammals of the southern oceans. *Journal of Marine Systems*, 78, 363–376.
- Boulinier, T. and Danchin, E. (1996) Population trends in Kittiwake *Rissa tridactyla* colonies in relation to tick infestation. *Ibis*, 138, 326–334.
- Boulinier, T., McCoy, K.D., Yoccoz, N.G., Gasparini, J. and Tveraa, T. (2008) Public information affects breeding dispersal in a colonial bird: kittiwakes cue on neighbours. *Biology Letters*, 4, 538–540.
- Converse, J.D., Hoogstraal, H., Moussa, M.I., Feare, C.J. and Kaiser, M.N. (1975) Soldado virus (Hughes group) from *Ornithodoros (Alectorobius) capensis* (Ixodoidea - Argasidae) infesting sooty tern colonies in Seychelles, Indian Ocean. *American Journal of Tropical Medicine and Hygiene*, 24, 1010–1018.
- Coulson, S.J., Lorentzen, E., Strom, H. and Gabrielsen, G.W. (2009) The parasitic tick *Ixodes uriae* (Acari: Ixodidae) on seabirds from Spitsbergen, Svalbard. *Polar Research*, 28, 399–402.
- Cózar, A. *et al.* (2017) The Arctic Ocean as a dead end for floating plastics in the North Atlantic branch of the Thermohaline Circulation. *Science Advances*, 3, e1600582.
- Descamps, S. (2013) Winter temperature affects the prevalence of ticks in an Arctic seabird. *PLoS One*, 8, e65374.
- Dias, M.P. *et al.*, 2019. Threats to seabirds: A global assessment. *Biological Conservation*, 237, 525–537.

Dietrich, M., Gomez-Diaz, E. and McCoy, K.D. (2011) Worldwide distribution and diversity of seabird ticks: Implications for the ecology and epidemiology of tick-borne pathogens. *Vector-Borne and Zoonotic Diseases*, 11, 453-470.

Dietrich, M. *et al.* (2014a) *Rickettsia* spp. in seabird ticks from Western Indian Ocean islands, 2011-2012. *Emerging Infectious Diseases*, 20, 838-842.

Dietrich, M., Lobato, E., Boulinier, T. and McCoy, K.D. (2014b) An experimental test of host specialization in a ubiquitous polar ectoparasite: a role for adaptation? *Journal of Animal Ecology*, 83, 576-587.

Duffy, D.C. (1983) The ecology of tick parasitism on densely nesting Peruvian seabirds. *Ecology*, 64, 110-119.

Dupraz, M. *et al.* (2017) Population structure of the soft tick *Ornithodoros maritimus* and its associated infectious agents within a colony of its seabird host *Larus michahellis*. *International Journal for Parasitology: Parasites and Wildlife*, 6, 122-130.

Feare, C.J. (1976) Desertion and abnormal development in a colony of Sooty Terns *Sterna fuscata* infested by virus-infected ticks. *Ibis*, 118, 112-115.

Fluhr, J., Strøm, H., Pradel, R., Duriez, O., Beaugrand, G. and Descamps, S. (2017) Weakening of the subpolar gyre as a key driver of North Atlantic seabird demography: a case study with Brünnich's guillemots in Svalbard. *Marine Ecology Progress Series*, 563, 1-11.

Gauthier-Clerc, M., Clerquin, Y. and Handrich, Y. (1998) Hyperinfestation by ticks *Ixodes uriae*: a possible cause of death in adult king penguins, a long-lived seabird. *Colonial Waterbirds*, 21, 229-233.

Gomard, Y. *et al.* (2020). Changes in bacterial diversity, composition and interactions during the development of the seabird tick *Ornithodoros maritimus* (Argasidae). *Microbial Ecology*, 1-14.

Grémillet, D., Ponchon, A., Paleczny, M., Palomares, M.-L.D., Karpouzi, V. and Pauly, D. (2018) Persisting worldwide seabird-fishery competition despite seabird community decline. *Current Biology*, 28, 4009-4013.e2.

Grose, S.O., Pendleton, L., Leathers, A., Cornish, A. and Waitai, S. (2020) Climate change will re-draw the map for marine megafauna and the people who depend on them. *Frontiers in Marine Science*, 7, 547.

Hoegh-Guldberg, O. *et al.* (2018) Impacts of 1.5°C global warming on natural and human systems. *An IPCC Special Report*.

Jenouvrier, S., Péron, C. and Weimerskirch, H. (2015) Extreme climate events and individual heterogeneity shape life-history traits and population dynamics. *Ecological Monographs*, 85, 605-624.

Kwak, M.L., Kuo, C.-C. and Chu, H.-T. (2020) First record of the sea snake tick *Amblyomma nitidum* Hirst and Hirst, 1910 (Acari: Ixodidae) from Taiwan. *Ticks and Tick-borne Diseases*, 11, 101383.

Léger, E., Vourc'h, G., Vial, L., Chevillon, C. and McCoy, K.D. (2013) Changing distributions of ticks: causes and consequences. *Experimental and Applied Acarology*, 59, 219–244.

Llanos-Soto, S., Muñoz-Leal, S. and González-Acuña, D. (2019) Hyperparasitism in the seabird tick *Ornithodoros amblus* (Acari: Argasidae). *Systematic and Applied Acarology*, 24, 525–528.

Mangin, S., Gauthier-Clerc, M., Frenot, Y., Gendner, J.-P. and Le Maho, Y. (2003) Ticks *Ixodes uriae* and the breeding performance of a colonial seabird, king penguin *Aptenodytes patagonicus*. *Journal of Avian Biology*, 34, 30–34.

Munro, H.J. et al. (2017) Evidence for *Borrelia bavariensis* infections of *Ixodes uriae* within seabird colonies of the North Atlantic Ocean. *Applied and Environmental Microbiology*, 83, 20.

Monticelli, D., Ramos, J.A., Hines, J.E., Nichols, J.D. and Spendelow, J.A. (2008) Juvenile survival in a tropical population of roseate terns: interannual variation and effect of tick parasitism. *Marine Ecology-Progress Series*, 365, 277–287.

Olson, S.L. and Hearty, P.J. (2003) Probable extirpation of a breeding colony of Short-tailed Albatross (*Phoebastria albatrus*) on Bermuda by Pleistocene sea-level rise. *Proceedings of the National Academy of Sciences, USA*, 100, 12825–12829.

Parsons, N.J. et al. (2018) Novel vagrant records and occurrence of vector-borne pathogens in King Penguins (*Aptenodytes patagonicus*) in South Africa. *Polar Biology*, 41, 79–86.

Parsons, N.J., Voogt, N.M., Schaefer, A.M., Peirce, M.A. and Vanstreels, R.E.T. (2017) Occurrence of blood parasites in seabirds admitted for rehabilitation in the Western Cape, South Africa, 2001-2013. *Veterinary Parasitology*, 233, 52–61.

Péron, C., Authier, M., Barbraud, C., Delord, K., Besson, D. and Weimerskirch, H. (2010) Interdecadal changes in at-sea distribution and abundance of subantarctic seabirds along a latitudinal gradient in the Southern Indian Ocean. *Global Change Biology*, 16, 1895–1909.

Sanz-Aguilar, A. et al. 2020. Infestation of small seabirds by *Ornithodoros maritimus* ticks: Effects on chick body condition, reproduction and associated infectious agents. *Ticks and Tick-borne Diseases*, 11, 101281.

Schulze, P., 1936. The Norwegian zoological expedition to the Galapagos Island. *Die Ixodoidea der Galapagos-Inseln. Meddelelser fra det Zoologiske Museum*, 45, 155–162.

Ullmann, A.J., Lima, C.M.R., Guerrero, F.D., Piesman, J. and Black, W.C. (2005) Genome size and organization in the blacklegged tick, *Ixodes scapularis* and the Southern cattle tick, *Boophilus microplus*. *Insect Molecular Biology*, 14, 217–222.

Table 1. Known tick species from the marine ecosystem and associated infectious agents. Updated from Dietrich *et al.* (2011) with only new references included.

Tick genera	Species	Recorded Hosts	Known distribution	Infectious agents
Argasidae				
<i>Argas</i>	<i>cucumerinus</i>	<i>Pelecanus thagus</i>	Peru	unknown
	<i>macrostigmatus</i>	<i>Phalacrocorax aristotelis</i>	Corsica, Crimea	unknown
	<i>monolakensis</i>	<i>Larus californicus</i>	California	Mono Lake virus (Orbivirus, Kemerovo serogroup)
<i>Ornithodoros</i> (<i>Carios/</i> <i>Alectorobius</i>)	<i>amblus</i>	< 10; Suliformes, Charadriiformes, Spheniiformes ¹	Peru, Chile	Huacho virus (Orbivirus, Kemerovo serogroup), Punta salinas virus (Nairovirus, Hughes serogroup)
	<i>capensis</i>	> 35; diverse Procellariiformes, Suliformes, Charadriiformes, Spheniiformes, Phaethoniformes, Pelecaniformes	Globally distributed across tropical/subtropical regions	Soldado, Farallon viruses (Nairovirus, Hughes serogroup); Upolu, Aransas Bay viruses (Thogotovirus), Saumarez Reef virus (Flavivirus), Midway virus (Nyamivirus), Johnson Atoll virus (Quarantavirus); Adenoviruses (group B); <i>Rickettsia bellii</i> ; <i>R. hoogstraalii</i> ²
	<i>cheikhi</i>	<i>Sterna</i> sp.	Mauritania	Unknown
	<i>denmarki</i>	> 10; diverse Procellariiformes, Suliformes, Charadriiformes, Spheniiformes	Hawaii islands, Peru, Mexico, Jamaica, Martinique, Florida, Australia, New Zealand	Soldado virus (Nairovirus, Hughes serogroup), Farallon-like (Nairovirus, Hughes serogroup), Frazer Point and Raza viruses; Midway virus (Nyamivirus), Johnson Atoll virus (Quarantavirus)
	<i>maritimus</i>	> 35; diverse Procellariiformes, Suliformes,	British Isles, France, Spain, Portugal, Morocco	Essaouira, Kala Iris, Baku, Moellez viruses (Orbivirus, Kemerovo serogroup); Soldado,

		Charadriiformes, Phaethontiformes, Pelecaniformes, Spheniformes		Great Saltee, Pettico Wick, Puffin Island viruses (Nairovirus, Hughes serogroup); West Nile, Caspiy, Meaban viruses (Flavivirus), <i>Rickettsia</i> spp., <i>Borrelia</i> spp, <i>Babesia</i> spp. ³
	<i>muesebecki</i>	< 10; Charadriiformes, Suliformes	Arabian sea	Zirqa virus (Nairovirus, Hughes serogroup)
	<i>sawaii</i>	2 species; Procellariiformes	Japan	<i>Rickettsia</i> sp., <i>Borrelia</i> sp.
	<i>spheniscus</i>	<i>Spheniscus humboldti</i>	Peru	Unknown
	<i>yunkerii</i>	< 10; certain Procellariiformes, Suliformes, Charadriiformes, Spheniformes	Galapagos	Unknown
<hr/>				
Ixodidae				
<i>Ixodes</i>	<i>amersoni</i>	> 5 species: Procellariiformes (Diomedidae), Suliformes (Sulidae), Charadriiformes (Laridae)	Central Pacific Ocean	unknown
	<i>auritulus</i>	>35 species : diverse Procellariiformes, Suliformes, Charadriiformes, Spheniformes	Southern circumpolar region	unknown
	<i>diomedea</i>	1 species: Procellariiformes (Diomedidae)	Tristan da Cunha	unknown

<i>eudyptidus</i>	>10 species : diverse Procellariiformes, Suliformes, Charadriiformes, Spheniiformes	Australia, New Zealand	Saumarez Reef virus (Flavivirus)
<i>jacksoni</i>	1 species: Suliformes	Australia, New Zealand	unknown
<i>keruelensis</i>	>10 species : diverse Procellariiformes (Diomedidae, Procellariidae)	Southern circumpolar region	unknown
<i>kohli</i>	>5 species : Procellariiformes, Suliformes, Charadriiformes, Spheniiformes	Australia	unknown
<i>layanensis</i>	10 species : Procellariiformes, Suliformes, Charadriiformes, Phaethontiformes	Central Pacific ocean	unknown
<i>murreleti</i>	1 species: <i>Synthliboramphus</i> <i>hypoleucus</i>	Mexico	unknown
<i>percavatus</i>	2 species: Procellariiformes (Diomedidae), Spheniiformes	Tristan da Cunha	unknown
<i>philipi</i>	2 species: Procellariiformes	Japan	unknown
<i>rothschildi</i>	>10 species : diverse Procellariiformes, Suliformes, Charadriiformes	North-eastern Atlantic	Phlebovirus (Uukuniemi group)

	<i>signatus</i>	~4 species : Suliformes, Charadriiformes	North Pacific Ocean	Orbivirus (Kemerovo group); Orthonairovirus (Sakhalin group), Phlebovirus (Uukuniemi group)
	<i>unicavatus</i>	2 <i>Phalacrocorax</i> spp.	Eastern north Atlantic, Baltic sea	unknown
	<i>uriae</i>	> 60 species: diverse Procellariiformes, Suliformes, Charadriiformes, Spheniiformes	Circumpolar (both hemispheres)	Orbivirus (Kemerovo group); Orthonairovirus (Sakhalin and Hughes groups); Phlebovirus: (Uukuniemi group); Flaviviruses; <i>Borrelia burgdorferi</i> sl (<i>B. garinii</i> , <i>B. bavariensis</i> , <i>B. burgdorferi</i> ss, <i>B. lusitaniae</i>) ⁴
<i>Amblyomma</i>	<i>darwini</i>	1 : <i>Amblyrhynchus</i> <i>cristatus</i> (iguana) ⁵	Galapagos	unknown
	<i>loculosum</i>	>10 species : diverse Procellariiformes, Suliformes, Charadriiformes, Phaethontiformes	Australia, Indian Ocean	Soldado virus (Nairovirus, Hughes serogroup) <i>Rickettsia africae</i> . ²
	<i>nitidum</i>	Sea snakes (<i>Laticauda</i>) ⁶	South western Pacific Ocean	unknown

¹ Llanos-Soto *et al.*, 2019

² Dietrich *et al.*, 2014a

³ Dupraz *et al.*, 2017; Gomard *et al.*, 2020; Sanz-Aguilar *et al.*, 2020

⁴ Munro *et al.*, 2017

⁵ Schulze, 1936

⁶ Kwak *et al.*, 2020

Fig. 1. Marine ticks can pose significant problems for breeding seabirds in cases of heavy infestations and via the transmission of infectious agents. While infestations by ixodid ticks are relatively easy to quantify, as seen by this massively infested Atlantic puffin *Fratercula arctica* (a), those by argasid ticks are much more difficult. Indeed, soft ticks can be present in significant numbers within tropical and temperate seabird colonies, such as in this Sooty tern, *Onychoprion fuscatus*, colony on the Ile de Lys, Iles Eparses (b), but only leave shelters in the vegetation for their short bloodmeal during the night (insets: sampling of *Ornithodoros capensis*). Photo credits: (a) D. Playle used with permission from A. Spellman (www.mullbirds.com); (b) K.D. McCoy.