Manipulation of worker size diversity does not affect colony fitness under natural conditions in the ant *Temnothorax nylanderi*

Romain Honorio¹, Claudie Doums^{2,3} & Mathieu Molet¹

¹ Sorbonne Université, Université Paris Est Créteil, Université Paris Diderot, CNRS, INRAE, IRD, Institute of Ecology and Environmental Sciences-Paris, iEES-Paris, F-75005 Paris, France

² Institut de Systématique Évolution Biodiversité, ISYEB, F-75005 Paris, Sorbonne Université, MNHN, CNRS, EPHE, Université des Antilles, France

³ EPHE, PSL University, F-75014 Paris, France

Correspondence: R. Honorio, Sorbonne Université, Université Paris Est Créteil, Université Paris Diderot, CNRS, INRAE, IRD, Institute of Ecology and Environmental Sciences-Paris, iEES-Paris, F-75005 Paris, France

E-mail address: romain.honorio@sorbonne-universite.fr

Abstract

In social insects, within-colony worker diversity has long been thought to improve colony fitness. Recent studies start to question this assertion in species with moderate diversity. However, demonstrating an absence of effect is difficult as this absence could be due to the measurement of only few life history traits or to artificial conditions. In order to circumvent these limitations, we experimentally decreased worker size diversity within colonies of the ant *Temnothorax nylanderi*, with or without affecting mean worker size and we reintroduced them in the field for four months (spring and early summer). We then thoroughly measured their fitness based on survival, growth and reproductive success. Our results show that our

manipulation did not affect colony fitness. In addition, colonies did not restore diversity to its initial level, further suggesting that worker size diversity is not a key parameter to them. We found the classically observed positive relationship between colony size, colony growth and reproductive success. Overall, our results confirm that worker size diversity within colony is not necessarily adaptive in species where it is moderate. We discuss the alternative mechanisms that could explain the evolutionary persistence of moderate worker size diversity.

Significance statement

Organisms that live in groups can greatly benefit from the emergence of novel group-level traits. For instance, social insects show significant variability in worker size within colonies. This size diversity increases the division of labour among workers and improves colony fitness. However, in species where size diversity is moderate, this relationship may not always be verified. Here, we manipulated both worker size diversity and mean worker size within colonies of the ant *Temnothorax nylanderii*, we reintroduced them in the field, and we extensively measured colony fitness after four months. We found no impact on colony survival, growth and reproduction. We discuss how social life and its associated cooperation and conflicts could cause variation in worker size, without any positive effect on colony fitness.

Key-words: colony size, social insects, mean body size, survival, growth, reproductive success

Introduction

Animal societies can exhibit genetic, behavioral, physiological and morphological diversity among individuals of the same group and this is thought to provide benefits to the group (Sendova-Franks and Franks 1999; Robinson 2009). One of the most obvious and ecologically important traits in social insects is size diversity of individuals within colonies (bees: Brand & Chapuisat 2012; Linksvayer et al. 2011 - wasps: Kovacs et al. 2010 - ants: Molet et al. 2012). Worker size diversity improves division of labour and colony efficiency in many species (e.g. bumble bee: Peat et al. 2005; stingless bee: Ramalho et al. 1998). Among social insects, ants are the most morphologically diverse group and the one where the highest level of morphological diversity within colonies can be observed (Wheeler 1991). In ant species with high worker polymorphism (including discrete worker and soldier castes), studies support a general positive relationship between worker size diversity and colony performance (Davidson 1978; Wilson 1984; Yang et al. 2004; Evison et al. 2008; Powell 2009; Arnan et al. 2011 - but see: Rissing and Polloek 1984; Calabi and Traniello 1989). However, most species have moderate worker size diversity (with an intra-colonial coefficient of variation lower than 0.2 (Fjerdingstad and Crozier (2006))), and in this case the link between worker size diversity and colony performance is less clear. (Beshers and Traniello 1994, 1996) suggest that the adaptiveness of size variation in moderately polymorphic ants is unrelated to division of labor, and size could be driven by other pressures which increase survival or colony growth. Some studies show a positive relationship (Porter and Tschinkel 1985a, 1986; Heinze et al. 2003; Billick and Carter 2007; Modlmeier et al. 2013), others do not (Billick 2002; Modlmeier and Foitzik 2011; Westling et al. 2014; Yilmaz et al. 2019). In ant species with moderate worker size diversity, within group morphological diversity could therefore be non-adaptive at the colony level.

However, it is difficult to reliably rule out any adaptive value of worker size diversity as the presence or the lack of positive effects could always result from the limited number of

traits investigated or the rearing setup under artificial laboratory conditions. Indeed, some studies focus on a single trait, such as foraging (Billick and Carter 2007; Westling et al. 2014), brood production (Porter and Tschinkel 1985a, 1986; Billick 2002; Modlmeier and Foitzik 2011), thermal tolerance (Heinze et al. 2003; Yilmaz et al. 2019) or resistance to starvation (Modlmeier et al. 2013). Accordingly, trade-offs with other traits that could be negatively affected by size diversity may have been overlooked, resulting in a biased assessment of fitness. Moreover, some of these studies are correlative, so they do not prove a causal link between size diversity and colony efficiency (e.g. Modlmeier and Foitzik 2011; Westling et al. 2014). Finally, all the experimental studies were performed in the laboratory where realistic environmental pressures are lacking (e.g. Porter and Tschinkel 1985a; Colin et al. 2017). Indeed the distinct environmental conditions (photoperiod, temperature, food availability, social environment) of laboratory and field experiments cause variation in the behavior of both vertebrates and invertebrates (Campbell et al. 2009).

Size diversity and mean worker size increase with colony size (respectively: Ferguson-Gow et al. 2014; Porter and Tschinkel 1985b). Only colonies with sufficient resources can produce large workers and enhance worker size diversity. It is important to disentangle these two traits in order to properly conclude about the adaptive value of size diversity and mean worker size within colonies of social insects. Indeed, a recent study on bumblebee colonies highlighted a positive correlation between colony performance and mean worker size, but no link with worker size diversity (Herrmann et al. 2018).

In this study, we aimed at quantifying the impact of within-colony worker size diversity and worker mean size on colony fitness using a long-lasting field-based experiment with extensive measures of fitness (survival, growth and reproductive success). We focused on an ant species lacking discrete worker subcastes and exhibiting moderate worker size diversity, *Temnothorax nylanderii* (see Appendix A1). In *Temnothorax* species, worker size

does not determine worker specialization (Pinter-Wollman et al. 2012), task-allocation is highly flexible (Robinson, Feinerman, & Franks 2009) and it relies on physiological parameters (Robinson et al. 2012). Colin et al. (2017) manipulated worker size diversity within *T. nylanderii* colonies and challenged colonies in the laboratory with various tasks (nest emigration, foraging, hygienic behavior, cold shock, etc.). They did not find any differences in performance between colonies with reduced size diversity and control colonies with unmanipulated diversity. Moreover, neither mean size nor size diversity correlate with per-capita productivity (Modlmeier and Foitzik 2011). This suggests that worker size diversity is not adaptive in this species, although larger individuals survive longer to cold temperature (Heinze et al. 2003) and starvation (Modlmeier et al. 2013). We tested whether this holds true in the field by manipulating the size of workers in colonies in the laboratory. Importantly, we manipulated both within colony worker size diversity (decrease) and mean worker size (increase, decrease or no change) in order to disentangle their effects. We subsequently re-implanted colonies in the field for four months to expose them to natural biotic and abiotic selective pressures, and we subsequently measured colony fitness extensively. In *Temnothorax* species, brood production is highly synchronized (Headley 1943; Kipyatkov 1993) so, in addition to colony survival, we were able to quantify two additional fitness parameters, namely colony growth (based on the production of new workers) and reproductive success (based on the production of sexuals). In addition, because workers control larval development (Linksvayer et al. 2011), we expected colonies to compensate for the size of the removed individuals and restore diversity by producing workers of the very same size if size diversity is somewhat important for colony fitness.

Material and methods

125 *Ants*

126 Colonies of *Temnothorax nylanderi* nest in twigs or acorns and are thus easy to pick up from
127 the forest leaf litter. One hundred and forty-six colonies of *T. nylanderi* were collected in
128 February 2019 in the 'Bois de Vincennes' forest (Paris, France, 48° 50' 22.14" N, 2°26'
129 51.96" E). In the laboratory, each colony was installed in an artificial nest consisting of two
130 microscope slides separated by a 1-mm auto-adhesive plastic foam with three chambers. The
131 nest was placed in a plastic box (11.5 × 11.5 × 5.5 cm) providing a foraging area (see
132 Appendix A2). During their two-week stay in the laboratory, colonies were kept in a climatic
133 chamber at 10-12°C with a natural photoperiod mimicking the natural conditions at this
134 period. Water was provided *ad libitum* in a tube plugged with cotton. Colonies were fed once
135 with a freshly killed mealworm (*Tenebrio molitor*). Workers and brood were counted.

136

137 *Manipulation of worker size*

138 Out of the 146 colonies, 80 colonies were selected for the experiment based on the presence
139 of one queen and at least 70 workers. Four treatments were performed and, in all treatments,
140 50% of workers were removed from the 20 colonies. In treatment 25S25L, we decreased
141 worker diversity without changing mean worker size by removing the 25% smallest workers
142 and the 25% largest workers. In the two other treatments, we decreased worker diversity but
143 also either increased mean worker size by removing the 50% smallest workers (treatment
144 50S) or decreased mean worker size by removing the 50% largest workers (treatment 50L).
145 The last treatment was the control (treatment 50R) as we removed 50% of workers randomly.
146 The worker removal protocols were similar to Colin et al. (2017) and based on the apparent
147 global body size: the sorting of large and small workers was done by eye under a stereoscopic
148 microscope (Zeiss®, x50 magnification) whereas the removal of random workers was done
149 without a microscope to make sure that worker size could not be evaluated. This method was

proven to be reliable (Colin et al. 2017). Colonies were assigned to the four treatments based on their number of workers in order to keep a similar distribution of colony sizes among the four treatments. Workers remaining in the colonies after manipulation constituted the initial workforce of the colonies ($67.8 \text{ workers} \pm 22.5$). Colonies initially contained $91.6 \text{ larvae} \pm 27.7$.

Colony rearing in the field

We manufactured artificial nests to make them match natural nests as closely as possible, so that colonies could easily live in them after reintroduction in nature. For that purpose, we used 2.5x 2cm truncated cone corks. We drilled a 4, 5 or 6mm-wide chamber from the larger side of the cone cork, and plugged this side with a glue gun to seal the chamber. Then, a 1mm-wide entrance gallery was pierced from the smaller side of the cone cork using a pointed plier in order to connect the chamber to the outside. In the laboratory, six corks (two of each size) were placed inside each plastic box, and we induced the emigration of colonies into the corks of their choice by removing the cover glass of the original nest. Then, the six corks from each plastic box (one containing the colony) were reintroduced in the 'réserve ornithologique du Bois de Vincennes' in a semi-buried (10cm deep) 40 x 35cm bucket with a pierced bottom (for water draining) and containing local soil (see Appendix A3). The bucket lid was cut off into a ring shape and the bottom side was covered with fluon®, a slippery coating, to prevent ant escape while retaining a wide entrance. The six corks were placed randomly. This allowed colonies to relocate to the nest of their choice (size and location) inside the bucket whenever they wanted to. Indeed, *T. nylanderi* colonies readily switch nest depending on environmental conditions (Foitzik et al. 2003). Because the 80 buckets had been in place for a year, the soil and litter in the buckets were very similar to those of the surrounding forest; many arthropods and soil organisms could come in and out of the buckets. A preliminary food-supplementation

experiment showed that adding food into the buckets did not increase colony survival and growth, so the 40cm-wide foraging area provided enough food for the colonies. Colonies were left in the buckets from March 26th to July 16th, and subsequently collected and brought back to the laboratory. Corks were collected in the early morning, when workers were inactive because of cool temperature, in order to make sure that complete colonies were collected. Cork nests were destroyed and colonies were forced to move to artificial microscope-slide nests. Workers, sexual and worker pupae and larvae were counted. There were no sexual adults yet. Brood production is highly synchronized during Summer (July-August) in *Temnothorax* (Headley 1943; Kipyatkov 1993; Modlmeier and Foitzik 2011), therefore the individuals that we recovered as pupae most likely reflected the total production of the colonies. We allowed pupae to molt into adults in order to assess the sex of the brood. To do so, colonies were kept in the laboratory at room temperature ($21^{\circ}\text{C} \pm 2^{\circ}\text{C}$) with a natural photoperiod. Water was provided *ad libitum* and colonies were fed once a week with half a freshly killed mealworm. Emerging adults were killed by freezing and then stored in 96° ethanol. The workers, males and gynes counted as ‘newly produced’ by colonies were individuals that emerged in the laboratory after developing into pupae in the field. All adults had emerged by August 16th and all remaining individuals were killed.

We considered that colonies had survived if they were recovered in July and if their queen was still alive. This loose definition of survival is an over-estimate relative to the proper life-history trait ‘colony survival’, as some colonies may actually have escaped from the bucket (entirely or only a fragment containing the queen, although this is highly unlikely because the bottom side of the lid was covered with a slippery coating – see Methods), and some colonies could have recovered from orphanage by later adopting a new fertilized queen (Foitzik and Heinze 2000). Queenless colonies were excluded from the analyses. Mortality rate of adult workers was computed as $1 - (\text{number of workers retrieved on July 16th} / \text{initial$

workforce). Indeed, it is very unlikely that any workers were produced during this period (Headley 1943; Kipyatkov 1993; Modlmeier and Foitzik 2011). We also counted and measured the workers and sexuals (gynes and males) produced out of the pupae brought back to the laboratory. Traits related to colony growth were the mortality rate of adult workers, the number of workers produced, the mean size of the workers produced and its coefficient of variation (CV). Traits related to reproductive success were the number of produced sexuals (gynes and males), the mean size of the sexuals, and the sexual investment (number of sexuals / number of workers produced). All three main fitness components of colonies were thus measured.

Morphological measurements

The workers and sexuals that emerged in the laboratory were measured. Heads were separated from bodies. Heads were stuck on double-sided tape and photographed under a Discovery.V12 Zeiss stereomicroscope connected to a computer via a Zeiss AxioCam ICc 5 video camera. Head width, a standard measurement of body size in ants (Tschinkel et al. 2003), was measured with ImageJ 1.8 (Abràmoff et al. 2004) and measurements were blind to treatment.

Statistical analysis

We investigated the effect of treatments (predictor variable, four levels) on colony survival, traits related to colony growth (mortality rate of adult workers, number of new workers produced, size of the produced workers and CV of the size of produced workers) and reproductive success (number of produced sexuals, size of the sexuals and its CV, sexual investment). The mortality rate of adult workers and the number of produced individuals were also used as predictor variables in some models (Tables 1 and 2). So, in addition to

treatments, we considered four other predictor variables: the initial workforce (number of workers in a colony after the experimental removal of 50% of workers), the initial number of larvae in March, the mortality rate of adult workers and the number of produced workers and sexuals. All first-order interactions between treatment and these variables were also included.

All statistical analyses were carried out with R v3.6.1 (www.r-project.org). All plots were generated using *ggplot2* (Wickham 2009). First, we tested the effect of predictor variables on colony-level dependent variables (mortality rate of adult workers, number of produced workers and males, sexual investment, and size and CV of the produced individuals) using generalized linear models (GLMs). Counting data (number of produced individuals and sexual investment) were analyzed using quasipoisson distribution. Mortality rate of adult workers and CV were analyzed using Gaussian distribution. Finally, we tested the effect of predictor variables on the head size of emerged individuals, an individual-level dependent variable, using linear mixed models (LMMs, package *nlme*; Pinheiro et al 2016), with colony as random factor. All models are summarized in Table 1 and 2.

For each analysis, the minimum adequate model was selected using a backward stepwise approach where predictor variables were removed one by one from a full model based on a log likelihood ratio test. We used log likelihood ratio tests, i.e. F-tests for Gaussian distribution or χ^2 tests for quasi-poisson distribution, to obtain the *P*-values for each predictor variable by comparing the minimum adequate model with a model excluding or including the variable of interest (according to whether the variable was present in or absent from the minimal adequate model respectively). Normality of the residuals and homogeneity of variances were checked visually following Pinheiro and Bates (2000); no transformation of the data was necessary.

Results

Effect of treatments on colony survival

Only five of the 80 manipulated colonies could not be recovered during colony collection in July (three colonies 50S, one colony 50L and one colony 25L25S). Six of the 75 recovered colonies had lost their queen (three colonies 50R and three colonies 50L). These queenless colonies were excluded from the statistical analysis. Our final dataset consisted of 69 queenright colonies.

Effect of treatments on growth

The mortality rate of adult workers depended on the initial number of larvae in interaction with the treatment (treatment x initial larvae interaction, $F_{57:60} = 2.95$, $P = 0.04$). The mortality rate of adult workers decreased with the initial number of larvae with a steeper slope in treatment 50S (figure 2a). The mortality rate of adult workers did not differ among the treatments ($F_{57:66} = 1.94$, $P = 0.063$; figure 1a). Predictors that had a marginal effect ($0.05 < P\text{-value} < 0.07$) and could therefore slightly affect the dependent variable were retained in the minimum adequate model.

A total of 4 341 workers emerged in the laboratory from 66 queenright colonies (3 colonies did not produce any worker). The number of produced workers was positively correlated with the initial workforce ($F_{66:67} = 51.81$, $P < 0.001$; figure 2b) and negatively correlated with the mortality rate of adult workers ($F_{66:67} = 4.44$, $P = 0.04$), without significant interaction with the treatments (see Table 1). The number of produced workers did not differ among the treatments ($F_{66:63} = 0.38$, $P = 0.8$; figure 1b). The coefficient of variation of the size of produced workers did not differ among the treatments ($F_{63:60} = 0.39$, $P = 0.8$),

and it was not explained by any other predictor (see Table 1). The size of produced workers increased with the number of produced workers (mixed effect model: $\chi^2_1 = 38.51$, $P < 0.001$; figure 3a) and with the initial number of larvae ($\chi^2_1 = 3.79$, $P = 0.05$). The treatments had no effect on the size of produced workers ($\chi^2_3 = 3.25$, $P = 0.3$; figure 1d). Detailed statistics are presented in Table 1.

Effect of treatments on reproductive success

No gynes were produced except for three in one colony of treatment 50S. However, 57 queenright colonies produced 809 males. The number of males was positively correlated with the initial workforce ($F_{67:68} = 20.11$, $P < 0.001$; figure 2c) but was not affected by the treatment ($F_{67:64} = 1.22$, $P = 0.3$). The size of produced males was positively correlated with the number of males ($\chi^2_1 = 4.03$, $P = 0.04$; figure 3b). The sexual investment (number of sexuals / number of produced workers) and the CV of the size of produced males were not explained by any predictor. Detailed statistics are presented in Table 2.

Discussion

In this study, we aimed at quantifying the impact of worker size and size diversity within colony on colony fitness using a laboratory manipulation followed by a reintroduction in the field. Because brood production is synchronized in *Temnothorax nylanderi* (Headley 1943; Kipyatkov 1993), we were able to measure both colony growth and sexual investment in addition to survival, and thus to extensively quantify fitness. In agreement with a previous laboratory study (Colin et al. 2017), reducing worker size diversity did not affect colony fitness. Colony survival, reproductive success (the number of males and the investment in

300 sexuals) and colony growth (the number of produced workers) did not differ among the four
301 treatments (removal of large, small, large and small, or random workers). Surprisingly,
302 treatments that reduced or increased mean worker size did not affect fitness either. We also
303 found that colonies did not compensate for the size of the removed workers by producing
304 workers of the same size, which further suggests that worker size diversity has no strong
305 effect on colony fitness.

306 The initial workforce of colonies had a major impact on colony fitness in our
307 experiment, which is not the case in other species of the *Temnothorax* genus (Dornhaus et al.
308 2008; Bengston and Dornhaus 2013). Indeed, we found a positive relationship between the
309 initial workforce and the number of produced workers and males. This confirms that in *T.*
310 *nylanderi*, large groups grow and reproduce better (Foitzik and Heinze 2000; Foitzik et al.
311 2003). Only one colony produced gynes. This might be a consequence of the experimental
312 removal of half of the workers from each colony, which resulted in a loss of workforce and a
313 redirection of resources towards workers and males, as already known in *T. nylanderi* (Foitzik
314 and Heinze 2000; Foitzik et al. 2003).

315 Our results are in line with Colin et al. (2017), who found no impact of reducing
316 worker size diversity on colony performance in the laboratory in the same species, and
317 Modlmeier & Foitzik (2011) who found no correlation between worker size diversity and per
318 capita productivity in the field in *T. longispinosus*. Modlmeier & Foitzik (2011) also found no
319 effect of the mean worker size on colony productivity. Our experimental manipulation of
320 mean worker size confirms that it does not affect colony fitness. Moreover, colonies do not
321 compensate for the size of the removed workers, reinforcing the idea that worker size
322 diversity is not of primary value to them. Billick (2002) found a similar pattern in *Formica*
323 *neorufibarbis*. He collected 45 workers from field colonies, and removed the 15 largest
324 workers (7% of colony size). He followed colony development for 35 days and found no

consequence on the number of pupae produced per worker. In addition, although a positive relationship between initial workforce and both mean worker size and worker size diversity is usually found in ants (Porter and Tschinkel 1985a; Ferguson-Gow et al. 2014), we did not find such a correlation. Our results hence confirm that this trait is not of primary importance for colonies. For the first time, our study proves the absence of effect of both mean worker size and worker size diversity in the field, over a long period of time, and with an extensive measure of colony fitness (three main life-history traits), making it a highly reliable result. We discuss three non-mutually exclusive hypotheses that could explain why within-colony worker size diversity still occurs.

First, worker size diversity could only bring advantages in the context of harsher environmental conditions. In *T. nylanderii*, large workers resist better to starvation (Modlmeier et al. 2013). In the closely related *Leptothorax acervorum*, larger individuals survive longer in cold temperature (Heinze et al. 2003). Based on our results, such individual-level response (individual survival) does not seem to translate to the group level (colony survival). Further experiments are needed to fully explore the role of size diversity in the context of harsher environmental conditions (e.g. during overwintering) or over a full life cycle.

Second, size diversity could result from relaxed selection. Group living can buffer selective pressures on individual development because (1) phenotypes can diverge and still be viable under the protection of the social environment (Molet et al. 2012) and (2) workers themselves can control the environment of developing larvae and adjust their development if needed (Smith et al. 2008; Segers et al. 2015). This could thus decrease the level of canalization (the ability of development to remain unchanged despite environmental or genetic variation; Debat and David 2001), and thereby increase worker size diversity. The cost of maintaining developmental canalization (Van Buskirk and Steiner 2009) could be outbalanced by the lack of benefits that it brings in a social context. Accordingly, size

diversity, in species where it is moderate, could be not adaptive *per se* but just economical to avoid the costs of canalization (Hunt et al. 2011; Colin et al. 2017).

Third, larvae could exhibit selfish strategies (Amor et al. 2011, 2016). Larvae may seek to receive more resources in order to become larger individuals, and increase their direct fitness by laying male eggs (Stroeymeyt et al. 2007). Heinze & Oberstadt (1999) showed that large workers of the ant *Leptothorax gredleri* (closely related to *T. nylanderi*) have higher social ranks than small ones, so they are more likely to be the ones that lay male eggs. It would be interesting to compare larval development in the presence or absence of workers to shed light on this potential conflict. In *Ectatomma tuberculatum*, larval begging does not fully reflect the nutritional status, suggesting that larvae do cheat to receive more food (Sauvaget 2017).

The ecological success of *T. nylanderi* (up to 10 colonies per m²: Heinze et al. 1996 - with a wide distribution throughout Western European forests, woods and parks: Csoz et al. 2015; Khimoun et al. 2020), probably originates from unassessed traits other than mean worker size or worker size diversity, such as behavioral diversity. Indeed, *Temnothorax* ants demonstrate highly flexible and context-dependent task specialization, (Pinter-Wollman et al. 2012) and they can quickly respond to changes in the required tasks (Robinson et al. 2009). Colonies with high within-colony behavioral diversity are more productive than colonies with less variation (Modlmeier et al. 2012). This may apply more generally to social insects. Indeed, in a ponerine ant lacking size diversity, age-associated division of labour is highly flexible. Foragers can revert to nursing and egg laying (Bernadou et al. 2015). Moreover, the functioning of honeybee societies is based on the coexistence of physiologically specialized individuals that perform fixed functions such as nectar processing, and physiologically plastic individuals that are also more flexible behaviorally (Johnson 2003). To summarize, the ecological success of social insects with high size-diversity could come from their sharp

division of labour in relation with workers morphology (with the exception of *Veromessor pergandei* and *Pheidole dentata*, respectively Rissing and Polloek 1984; Calabi and Traniello 1989), whereas the success of species with moderate size diversity could result from their behavioral ability to quickly adapt to specific contexts.

Acknowledgments

We thank Romain Péronnet for technical assistance and ant collection in Vincennes. We are grateful towards Michel Neff and the Division du Bois de Vincennes de la Ville de Paris for letting us use an area of wood land of the Réserve Ornithologique du Bois de Vincennes as an experimental plot. We thank the two anonymous reviewers for their helpful comments on the manuscript.

Author's contributions

RH designed the study, collected and reared ants, performed the experiment and statistical analyses, and wrote the manuscript. CD designed the study, contributed to statistical analyses and wrote the manuscript. MM designed the study, wrote the manuscript and supervised the project. All authors read and approved the final manuscript.

Funding

This work was funded by the Institute of Ecology and Environmental Sciences - Paris and the Institut de la Transition Environnementale de Sorbonne Université.

Data availability

The dataset analyzed during the current study is available from the corresponding author on reasonable request.

Compliance with ethical standards

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

All applicable international, national and/or institutional guidelines for the care and use of animals were followed.

References

- Abràmoff MD, Magalhães PJ, Ram SJ (2004) Image processing with imageJ. *Biophotonics Int* 11:36–41.
- Amor F, Ortega P, Jowers MJ, et al (2011) The evolution of worker-queen polymorphism in *Cataglyphis* ants: Interplay between individual-and colony-level selections. *Behav Ecol Sociobiol* 65:1473–1482.
- Amor F, Villalta I, Doums C, et al (2016) Nutritional versus genetic correlates of caste differentiation in a desert ant. *Ecol Entomol* 41:660–667.
- Arnan X, Ferrandiz-Rovira M, Pladevall C, Rodrigo A (2011) Worker size-related task partitioning in the foraging strategy of a seed-harvesting ant species. *Behav Ecol Sociobiol* 65:1881–1890.
- Bengston SE, Dornhaus A (2013) Colony size does not predict foraging distance in the ant *Temnothorax rugatulus*: A puzzle for standard scaling models. *Insectes Soc* 60:93–96.
- Bernadou A, Busch J, Heinze J (2015) Diversity in identity: behavioral flexibility, dominance, and age polyethism in a clonal ant. *Behav Ecol Sociobiol* 69:1365–1375.
- Beshers SN, Traniello JFA (1994) The Adaptiveness of Worker Demography in the Attine Ant *Trachymyrmex septentrionalis*. *Ecology* 75:763–775.
- Beshers SN, Traniello JFA (1996) Polyethism and the adaptiveness of worker size variation in the attine ant *Trachymyrmex septentrionalis*. *J Insect Behav* 9:61–83.
- Billick I (2002) The relationship between the distribution of worker sizes and new worker production in the ant *Formica neorufibarbis*. *Oecologia* 132:244–249.
- Billick I, Carter C (2007) Testing the importance of the distribution of worker sizes to colony

432 performance in the ant species *Formica obscuripes* forel. *Insectes Soc* 54:113–117.
 433 Brand N, Chapuisat M (2012) Born to be bee, fed to be worker? The caste system of a primitively
 434 eusocial insect. *Front Zool* 9:35.
 435 Calabi P, Traniello JFA (1989) Social organization in the ant *Pheidole dentata* - Physical and temporal
 436 caste ratios lack ecological correlates. *Behav Ecol Sociobiol* 24:69–78.
 437 Campbell DLM, Weiner SA, Starks PT, Hauber ME (2009) Context and control: behavioural ecology
 438 experiments in the laboratory. *Ann Zool Fennici* 46:112–123.
 439 Colin T, Doums C, Péronnet R, Molet M (2017) Decreasing worker size diversity does not affect
 440 colony performance during laboratory challenges in the ant *Temnothorax nylanderi*. *Behav Ecol*
 441 *Sociobiol* 71:92.
 442 Csoz S, Heinze J, Mikó I (2015) Taxonomic synopsis of the ponto-mediterranean ants of
 443 *temnothorax nylanderi* species-group. *PLoS One* 10(11):e0140000.
 444 Davidson DW (1978) Size Variability in the Worker Caste of a Social Insect (*Veromessor pergandei*
 445 Mayr) as a Function of the Competitive Environment. *Am Nat* 112:523–532.
 446 Debat V, David P (2001) Mapping phenotypes: canalization, plasticity and developmental stability.
 447 *Trends Ecol Evol* 16:555–561.
 448 Dornhaus A, Holley JA, Pook VG, et al (2008) Why do not all workers work? Colony size and
 449 workload during emigrations in the ant *Temnothorax albipennis*. *Behav Ecol Sociobiol* 63:43–
 450 51.
 451 Evison SEF, Hart AG, Jackson DE (2008) Minor workers have a major role in the maintenance of
 452 leafcutter ant pheromone trails. *Anim Behav* 75:963–969.
 453 Ferguson-Gow H, Sumner S, Bourke AFG, Jones KE (2014) Colony size predicts division of labour in
 454 attine ants. *Proc R Soc B Biol Sci* 281:20141411.
 455 Fjerdingstad EJ, Crozier RH (2006) The evolution of worker caste diversity in social insects. *Am Nat*
 456 167:390–400.
 457 Foitzik S, Heinze J (2000) Intraspecific parasitism and split sex ratios in a monogynous and
 458 monandrous ant (*Leptothorax nylanderi*). *Behav Ecol Sociobiol* 47:424–431.
 459 Foitzik S, Strätz M, Heinze J (2003) Ecology, life history and resource allocation in the ant,
 460 *Leptothorax nylanderi*. *J Evol Biol* 16:670–680.
 461 Headley AE (1943) Population studies of two species of ants, *Leptothorax longispinosus* Roger and
 462 *Leptothorax curvispinosus* Mayr. *Ann Entomol Soc Am* 36:743–753.
 463 Heinze J, Foitzik S, Fischer B, et al (2003) The significance of latitudinal variation in body size in a
 464 holarctic ant, *Leptothorax acervorum*. *Ecography* 26:349–355.
 465 Heinze J, Foitzik S, Hippert A, Hölldobler B (1996) Apparent dear-enemy phenomenon and
 466 environment-based recognition cues in the ant *Leptothorax nylanderi*. *Ethology* 102:510–522.
 467 Heinze J, Oberstadt B (1999) Worker age, size and social status in queenless colonies of the ant
 468 *Leptothorax gredleri*. *Anim Behav* 58:751–759.

469 Herrmann JD, Haddad NM, Levey DJ (2018) Mean body size predicts colony performance in the
 470 common eastern bumble bee (*Bombus impatiens*). *Ecol Entomol* 43:458–462.

471 Hunt BG, Ometto L, Wurm Y, et al (2011) Relaxed selection is a precursor to the evolution of
 472 phenotypic plasticity. *Proc Natl Acad Sci U S A* 108:15936–15941.

473 Johnson BR (2003) Organization of work in the honeybee: a compromise between division of labour
 474 and behavioural flexibility. *Proc R Soc B Biol Sci* 270:147–152.

475 Khimoun A, Doums C, Molet M, et al (2020) Urbanization without isolation: the absence of genetic
 476 structure among cities and forests in the tiny acorn ant *Temnothorax nylanderi*. *Biol Lett*
 477 16:20190741.

478 Kipyatkov VE (1993) Annual cycles of development in ants: diversity, evolution, regulation. *Proc.*
 479 *Colloq. Soc. Insects* 2:25–48.

480 Kovacs JL, Hoffman EA, Marriner SM, et al (2010) Environmental and genetic influences on queen
 481 and worker body size in the social wasp *Vespula maculifrons*. *Insectes Soc* 57:53–65.

482 Linksvayer TA, Kaftanoglu O, Akyol E, et al (2011) Larval and nurse worker control of
 483 developmental plasticity and the evolution of honey bee queen–worker dimorphism. *J Evol Biol*
 484 24:1939–1948.

485 Modlmeier AP, Foitzik S (2011) Productivity increases with variation in aggression among group
 486 members in *Temnothorax* ants. *Behav Ecol* 22:1026–1032.

487 Modlmeier AP, Foitzik S, Scharf I (2013) Starvation endurance in the ant *Temnothorax nylanderi*
 488 depends on group size, body size and access to larvae. *Physiol Entomol* 38:89–94.

489 Modlmeier AP, Liebmann JE, Foitzik S (2012) Diverse societies are more productive: A lesson from
 490 ants. *Proc R Soc B Biol Sci* 279:2142–2150.

491 Molet M, Wheeler DE, Peeters C (2012) Evolution of novel mosaic castes in ants: modularity,
 492 phenotypic plasticity, and colonial buffering. *Am Nat* 180:328–341.

493 Peat J, Tucker J, Goulson D (2005) Does intraspecific size variation in bumblebees allow colonies to
 494 efficiently exploit different flowers? *Ecol Entomol* 30:176–181.

495 Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC (2016) Nlme: linear and nonlinear mixed effects
 496 models. R Package version 3.1-128. <http://CRAN.R-project.org/package=nlme>.

497 Pinheiro JC, Bates DM (2000) Mixed-effects models in S and S-PLUS, Statistics and computing. In:
 498 *Statistics and Computing*. Springer.

499 Pinter-Wollman N, Hubler J, Holley JA, et al (2012) How is activity distributed among and within
 500 tasks in *Temnothorax* ants? *Behav Ecol Sociobiol* 66:1407–1420.

501 Porter SD, Tschinkel WR (1985a) Fire ant polymorphism: the ergonomics of brood production. *Behav*
 502 *Ecol Sociobiol* 16:323–336.

503 Porter SD, Tschinkel WR (1985b) Fire ant polymorphism (Hymenoptera: Formicidae): factors
 504 affecting worker size. *Ann Entomol Soc Am* 78:381–386.

505 Porter SD, Tschinkel WR (1986) Adaptive value of nanitic workers in newly founded red imported

506 fire ant colonies (Hymenoptera: Formicidae). *Ann Entomol Soc Am* 79:723–726.

507 Powell S (2009) How ecology shapes caste evolution: Linking resource use, morphology, performance
508 and fitness in a superorganism. *J Evol Biol* 22:1004–1013.

509 Ramalho M, Imperatriz-Fonseca VL, Giannini TC (1998) Within-colony size variation of foragers and
510 pollen load capacity in the stingless bee *Melipona quadrifasciata* anthidioides Lepeletier
511 (Apidae, Hymenoptera). *Apidologie* 29:221–228.

512 Rissing SW, Polloek GB (1984) Worker size variability and foraging efficiency in *Veromessor*
513 *pergandei* (Hymenoptera: Formicidae). *Behav Ecol Sociobiol* 15:121–126.

514 Robinson EJJ (2009) Physiology as a caste-defining feature. *Insectes Soc* 56:1–6.

515 Robinson EJJ, Feinerman O, Franks NR (2012) Experience, corpulence and decision making in ant
516 foraging. *J Exp Biol* 215:2653–2659.

517 Robinson EJJ, Feinerman O, Franks NR (2009) Flexible task allocation and the organization of work
518 in ants. *Proc R Soc B Biol Sci* 276:4373–4380.

519 Sauvaget M (2017) Les relations jeunes-adultes chez les fourmis : sollicitations alimentaires des larves
520 de l'espèce *Ectatomma tuberculatum* [Doctoral dissertation]. Villetaneuse, France: Université
521 Paris XIII.

522 Segers FHID, Menezes C, Vollet-neto A, et al (2015) Soldier production in a stingless bee depends on
523 rearing location and nurse behaviour. *Behav Ecol Sociobiol* 69:613–623.

524 Sendova-Franks AB, Franks NR (1999) Self-assembly, self-organization and division of labour. *Philos*
525 *Trans R Soc B Biol Sci* 354:1395–1405.

526 Smith CR, Anderson KE, Tillberg C V., et al (2008) Caste determination in a polymorphic social
527 insect: nutritional, social, and genetic factors. *Am Nat* 172:497–507.

528 Stroeymeyt N, Brunner E, Heinze J (2007) “Selfish worker policing” controls reproduction in a
529 *Temnothorax* ant. *Behav Ecol Sociobiol* 61:1449–1457.

530 Tschinkel WR, Mikheyev AS, Storz SR (2003) Allometry of workers of the fire ant, *Solenopsis*
531 *invicta*. *J Insect Sci* 3:1–11.

532 Van Buskirk J, Steiner UK (2009) The fitness costs of developmental canalization and plasticity. *J*
533 *Evol Biol* 22:852–860.

534 Westling JN, Harrington K, Bengston S, Dornhaus A (2014) Morphological differences between
535 extranidal and intranidal workers in the ant *Temnothorax rugatulus*, but no effect of body size on
536 foraging distance. *Insectes Soc* 61:367–369.

537 Wheeler DE (1991) The developmental basis of worker caste polymorphism in ants. *Am Nat*
538 138:1218–1238.

539 Wickham H (2009) *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New-York.

540 Wilson EO (1984) The relation between caste ratios and division of labor in the ant genus *Pheidole*
541 (Hymenoptera: Formicidae). *Behav Ecol Sociobiol* 16:89–98.

542 Yang AS, Martin CH, Nijhout HF (2004) Geographic variation of caste structure among ant

543 populations. *Curr Biol* 14:514–519.

544 Yilmaz AR, Chick LD, Perez A, et al (2019) Remarkable insensitivity of acorn ant morphology to
545 temperature decouples the evolution of physiological tolerance from body size under urban heat
546 islands. *J Therm Biol* 85:102426.

548

Figure captions

Figure 1: Boxplots comparing the effects of the four treatments (50R: 50% of randomly removed workers; 25S25L: the 25% smallest and the 25% largest removed workers; 50L: the 50% largest removed; 50S: the 50% smallest removed) on four life history traits (a) mortality rate of adult workers, (b) number of produced workers, (c) number of produced males and (d) head size of produced workers (mm). Boxes show median, quartiles and extremes. Statistics are presented in Table 1 for workers and Table 2 for males.

Figure 2: Correlation between the initial larvae and workforce with three different traits. When the initial larvae increased, (a) the mortality rate of adult workers (50S: solid line type, estimate = $-2.141e^{-3} \pm 1.14e^{-3}$; 50R: dotdash, estimate = $-1.072e^{-3} \pm 1.08e^{-3}$; 50L: dotted, estimate = $-1.074e^{-3} \pm 1.16e^{-3}$; 25L25S: longdash, estimate = $-0.701e^{-3} \pm 1.03e^{-3}$). When the initial workforce increased, (b) the number of produced workers increased similarly for all treatments, and (c) the number of produced males increased similarly for all treatments. The regression lines are drawn from the coefficients of the different models. Statistics are presented in Table 1 for workers and Table 2 for males.

Figure 3: Correlation between the mean head size (mm) and the number of (a) produced workers and (b) produced males. The regression lines are drawn from the coefficients of the models and did not differ among treatments. Statistics are presented in Table 1 and Table 2.

Table 1: Models and statistics related to colony growth. ‘Minimum model’ means that the predictor was retained in the minimum model selected. A mixed effect model with colony as random factor was used for the size of produced workers in order to control for the colony of origin of workers.

575 Table 2: Models and statistics related to reproductive success. ‘Minimum model’ means that
576 the predictor was retained in the minimum model selected. A mixed effect model with colony
577 as random factor was used for the size of produced males in order to control for the colony of
578 origin of males.

Predictors	Response variables			
	Mortality rate of adult workers	Produced workers	CV of produced workers	Size of produced workers
Treatment	Minimum model $F_{57:66} = 1.94, P = 0.063$	$F_{66:63} = 0.38, P = 0.8$	$F_{63:60} = 0.39, P = 0.8$	$\chi^2_3 = 3.25, P = 0.3$
Initial workforce	Minimum model $F_{57:61} = 2.31, P = 0.069$	Minimum model $F_{66:67} = 51.81, P < 0.001$	$F_{63:62} = 0.02, P = 0.9$	$\chi^2_1 = 0.002, P = 0.9$
Treatment - initial workforce interaction	Minimum model $F_{57:60} = 2.52, P = 0.067$	$F_{66:63} = 0.50, P = 0.7$	$F_{63:59} = 0.43, P = 0.8$	$\chi^2_4 = 4.24, P = 0.4$
Initial larvae	Minimum model $F_{57:61} = 2.28, P = 0.07$	$F_{66:65} = 1.94, P = 0.2$	$F_{63:62} = 0.22, P = 0.6$	Minimum model $\chi^2_1 = 3.79, P = 0.05$
Treatment - initial larvae interaction	Minimum model $F_{57:60} = 2.95, P = 0.04$	$F_{66:62} = 0.55, P = 0.7$	$F_{63:59} = 0.54, P = 0.7$	$\chi^2_3 = 4.42, P = 0.22$
Mortality rate of adult workers		Minimum model $F_{66:67} = 4.44, P = 0.04$	$F_{63:62} = 0.05, P = 0.8$	$\chi^2_1 = 0.49, P = 0.5$
Treatment - mortality rate of adult workers interaction		$F_{66:63} = 0.40, P = 0.7$	$F_{63:59} = 0.33, P = 0.8$	$\chi^2_4 = 4.95, P = 0.3$
Produced workers			$F_{63:62} = 0.26, P = 0.6$	Minimum model $\chi^2_1 = 38.51, P < 0.001$
Treatment - produced workers interaction			$F_{63:59} = 1.01, P = 0.4$	$\chi^2_3 = 3.30, P = 0.3$

Predictors	Response variables			
	Males produced	Sexual investment	CV of produced males	Size of produced males
Treatment	$F_{67:64} = 1.22$, $P = 0.3$	$F_{65:62} = 0.23$, $P = 0.9$	$F_{45:42} = 0.32$, $P = 0.8$	$\chi^2_3 = 0.22$, $P = 0.9$
Initial workforce	Minimum model $F_{67:68} = \mathbf{20.11}$, $P < \mathbf{0.001}$	$F_{65:64} < 0.001$, $P = 0.3$	$F_{45:44} = 0.20$, $P = 0.6$	$\chi^2_1 = 0.38$, $P = 0.5$
Treatment - initial workforce interaction	$F_{67:64} = 1.11$, $P = 0.3$	$F_{65:61} = 0.39$, $P = 0.9$	$F_{45:41} = 0.37$, $P = 0.8$	$\chi^2_4 = 1.15$, $P = 0.9$
Initial larvae	$F_{67:66} = 0.19$, $P = 0.7$	$F_{65:64} < 0.001$, $P = 0.9$	$F_{45:44} = 2.54$, $P = 0.1$	$\chi^2_1 = 0.02$, $P = 0.9$
Treatment - initial larvae interaction	$F_{67:63} = 1.28$, $P = 0.3$	$F_{65:61} = 0.24$, $P = 0.9$	$F_{45:41} = 1.08$, $P = 0.4$	$\chi^2_4 = 0.46$, $P = 0.9$
Adult worker mortality rate	$F_{67:66} = 0.07$, $P = 0.8$	$F_{65:64} = 0.36$, $P = 0.3$	$F_{45:44} = 0.08$, $P = 0.8$	$\chi^2_1 = 0.72$, $P = 0.4$
Treatment - adult worker mortality rate interaction	$F_{67:63} = 0.51$, $P = 0.7$	$F_{65:61} = 0.73$, $P = 0.7$	$F_{45:41} = 0.08$, $P = 0.9$	$\chi^2_4 = 1.78$, $P = 0.8$
Males produced			$F_{45:44} = 1.32$, $P = 0.3$	Minimum model $\chi^2_1 = \mathbf{4.03}$, $P = \mathbf{0.04}$
Treatment - males produced interaction			$F_{45:41} = 0.43$, $P = 0.8$	$\chi^2_3 = 0.12$, $P = 0.9$

587 **Appendix captions**

588

589 Appendix A1: Worker size diversity within one colony of *Temnothorax nylanderi*. In our
590 experiment, the head width of the smallest worker produced was 0.384mm, that of the largest
591 was 0.661mm.

592

593 Appendix A2: Laboratory rearing setup. The nest consisted of two microscope slides
594 separated by a 1-mm auto-adhesive plastic foam with three chambers, covered with black
595 paper for darkness. It was placed in a plastic box (foraging area) with water (tube plugged
596 with cotton) and food (mealworm).

597

598 Appendix A3: Field rearing setup. Eighty semi-buried buckets were installed in the ‘réserve
599 ornithologique du Bois de Vincennes’. Each bucket contained local soil and was pierced at
600 the bottom for water draining. Six nests (drilled corks) were available for the colony to
601 choose. The foraging area was restricted to the bucket. Cut-off lids are not present on the
602 photos.