Structural insights into photosystem II assembly

Jure Zabret, Stefan Bohn, Sandra K Schuller, Oliver Arnolds, Madeline Möller, Jakob Meier-Credo, Pasqual Liauw, Aaron Chan, Emad Tajkhorshid, Julian D Langer, et al.

► To cite this version:

Jure Zabret, Stefan Bohn, Sandra K Schuller, Oliver Arnolds, Madeline Möller, et al.. Structural insights into photosystem II assembly. *Nature Plants*, 2021, 7 (4), pp.524-538. 10.1038/s41477-021-00895-0 . hal-03406233

HAL Id: hal-03406233

<https://hal.science/hal-03406233>

Submitted on 27 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Structural insights into photosystem II assembly

Jure Zabret^{1#}, Stefan Bohn^{2#}, Sandra K. Schuller^{3,4}, Oliver Arnolds⁵, Madeline Möller¹, Jakob Meier-Credo⁶, Pasqual Liauw¹, Aaron Chan⁷, Emad Tajkhorshid⁷, Julian D. Langer^{6,8}, Raphael Stoll⁵, Anja Krieger-Liszkay⁹, Benjamin D. Engel^{2,10,11}, Till Rudack^{12,13,*}, Jan M. Schuller^{3,4,*}, Marc M. Nowaczyk^{1,*}

Affiliations:

¹Department of Plant Biochemistry, Faculty of Biology & Biotechnology, Ruhr University Bochum, 44780 Bochum, Germany.

²Department of Molecular Structural Biology, Max Planck Institute of Biochemistry, 82152 Martinsried, Germany.

³Department of Structural Cell Biology, Max Planck Institute of Biochemistry, 82152 Martinsried, Germany.

⁴CryoEM of Molecular Machines, SYNMIKRO Research Center and Department of Chemistry, Philipps University of Marburg, 35043 Marburg, Germany.

⁵Biomolecular Spectroscopy and RUBiospek|NMR, Faculty of Chemistry and Biochemistry, Ruhr-University Bochum, 44780 Bochum, Germany

⁶Department of Molecular Membrane Biology, Max Planck Institute of Biophysics, 60438 Frankfurt/Main, Germany.

⁷NIH Center for Macromolecular Modeling and Bioinformatics, Beckman Institute for Advanced Science and Technology, Department of Biochemistry, and Center for Biophysics and Quantitative Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois

⁸Max Planck Institute for Brain Research, Max von Laue Strasse 4, 60438 Frankfurt/Main, Germany.

⁹Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France

¹⁰Helmholtz Pioneer Campus, Helmholtz Zentrum München, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany.

¹¹Department of Chemistry, Technical University of Munich, Lichtenbergstraße 4, 85748 Garching, Germany.

¹²Biospectroscopy, Center for Protein Diagnostics (ProDi), Ruhr University Bochum, 44801 Bochum, Germany.

¹³Department of Biophysics, Faculty of Biology & Biotechnology, Ruhr University Bochum, 44780 Bochum, Germany.

#Equal contribution

*corresponding authors: till.rudack@rub.de, jan.schuller@synmikro.uni-marburg.de, marc.m.nowaczyk@rub.de

Keywords

photosynthesis, photosystem II biogenesis, assembly factors, bicarbonate binding, reactive oxygen species, protection mechanisms, oxygen evolving complex, photoactivation, cryo-EM

Abstract

Biogenesis of photosystem II (PSII), nature's water splitting catalyst, is assisted by auxiliary proteins that form transient complexes with PSII components to facilitate stepwise assembly events. Using cryo-electron microscopy, we solved the structure of such a PSII assembly intermediate from *Thermosynechococcus elongatus* at 2.94 Å resolution. It contains three assembly factors (Psb27, Psb28, Psb34) and provides detailed insights into their molecular function. Binding of Psb28 induces large conformational changes at the PSII acceptor side, which distort the binding pocket of the mobile quinone (Q_B) and replace the bicarbonate ligand of non-heme iron with glutamate, a structural motif found in reaction centers of non-oxygenic photosynthetic bacteria. These results reveal novel mechanisms that protect PSII from damage during biogenesis until water splitting is activated. Our structure further demonstrates how the PSII active site is prepared for the incorporation of the Mn₄CaO₅ cluster, which performs the unique water splitting reaction.

1 Introduction

Photosystem II (PSII) is the only enzyme that catalyzes the light-driven oxidation of water, a thermodynamically demanding reaction that drives photosynthesis, sustaining life on our planet¹⁻³. This multi-subunit membrane protein complex is located in the thylakoid membranes of cyanobacteria, algae and plants. PSII strips electrons from water and injects them into the photosynthetic electron transport chain (PET). It forms a homodimer with a molecular mass of ~500 kDa⁴, with each monomer composed of at least 20 protein subunits and numerous cofactors, including chlorophylls, quinones, carotenoids, lipids, bicarbonate and the unique Mn₄CaO₅ cluster⁵⁻⁷. The two core proteins D1 and D2 form a central, membrane-intrinsic heterodimer, which binds all important redox cofactors involved in internal electron transfer⁸. Light-excitation leads to a charge-separated state in which an electron is transferred from the reaction center chlorophylls P₆₈₀⁹ to the nearby pheophytin¹⁰. Subsequently, the electron is passed to the bound plastoquinone (Q_A) and then to the mobile plastoquinone molecule (Q_B), which leaves the complex after accepting two electrons and two protons¹¹. The electron hole at P₆₈₀ is filled by oxidation of an adjacent tyrosine residue (Tyr_Z)¹² and finally by the oxygen

evolving complex (OEC), which contains the Mn_4CaO_5 cluster. In cyanobacteria, the cluster is shielded on the luminal side by the three extrinsic proteins, PsbO, PsbU and PsbV, which regulate access to the OEC by forming an intricate network of channels for different substrates and products¹³. Light energy is collected and funneled towards P_{680} by the two membrane-intrinsic antenna proteins CP43 and CP47. These proteins bind most of the chlorophyll molecules and are located on opposite sides of the D1/D2 heterodimer¹⁴. Moreover, at least twelve small transmembrane subunits with one or two transmembrane helices have been identified in PSII¹⁵, including cytochrome- b_{559} ¹⁶.

Structural and spectroscopic investigations have revealed these aforementioned comprehensive insights into PSII function¹⁷⁻²¹, but we are far from understanding PSII biogenesis with molecular detail. How nature facilitates the assembly of a multi-subunit, multi-cofactor membrane protein complex is a fundamental unsolved question. The biogenesis of PSII is even more challenging, as the mature complex performs sophisticated and extreme redox chemistry to catalyze the light-driven oxidation of water. This can easily lead to the formation of reactive oxygen species (e.g., singlet oxygen is produced by triplet chlorophyll in the PSII reaction center) and subsequent loss of function due to damaged proteins and cofactors^{22,23}. Biogenesis intermediates with only partially functional fragments of the redox chain are particularly prone to damage, thus demanding specialized protection mechanisms for the assembly process²⁴. Therefore, PSII biogenesis is not a spontaneous process but rather must be tightly regulated by the action of assembly factors. Thus far, more than 20 auxiliary proteins have been identified that guide the stepwise assembly of PSII subunits and cofactors via intermediate modules, which are assembled independently and then joined together to produce mature PSII²⁵⁻²⁷. In cyanobacteria, PSII biogenesis begins with the formation of the D1/D2 heterodimer reaction center (RC) complex from the D1 precursor protein (pD1) and the D2 protein. This is assisted by the PSII assembly factor Ycf48 after partial processing of the D1 C-terminal extension by the D1 specific peptidase CtpA^{28,29}. In the next step, the assembly factor Psb28 helps CP47 join the RC complex to form the RC47 complex, in which iD1 is further processed to its mature form by CtpA^{30,31}. Almost all ligands of the Mn_4CaO_5 cluster are already present at this stage, except for those provided by CP43, which comes pre-constructed with assembly factor Psb27 and several small subunits (together called the CP43 module)³². Psb28 is released as CP43 binds, and the resulting Psb27-PSII monomer is activated by maturation of the OEC and the binding of the extrinsic proteins PsbO, PsbU and PsbV³³⁻³⁵. OEC assembly is a multistep process that requires a functional upstream redox chain for the oxidation of Mn^{2+} to build up the cluster's μ -oxo bridges between the manganese atoms^{36,37}. The mechanistic and structural

50 details of this photoactivation process are not yet understood. In the consensus ‘two quantum
51 model’³⁸⁻⁴⁰, a single Mn^{2+} ion bound to the high-affinity site (HAS)⁴¹ is oxidized to Mn^{3+} . This
52 initiating light-dependent step is followed by a slow light-independent phase and further fast
53 light-dependent steps in which the remaining Mn^{2+} ions are oxidized and incorporated.
54 Understanding the light-independent slow phase is key to unraveling the mechanism of
55 photoactivation. Finally, PSII biogenesis completes with dimerization of two fully assembled
56 monomers and attachment of the soluble phycobilisome antenna complexes. Interestingly,
57 deletion of *psbJ*, which encodes a small single transmembrane helix protein at the entrance of
58 the PSII plastoquinone channel, decreases PSII-mediated oxygen evolution and increases the
59 lifetime of the reduced primary acceptor (Q_A^-) in cyanobacterial and tobacco chloroplasts⁴².
60 These effects are caused by a massive accumulation of an intermediate monomeric PSII
61 complex, which contains both assembly factors Psb27 and Psb28⁴³. Physiological studies of
62 Psb27 and Psb28 deletion strains point towards multifaceted functions. Cyanobacterial mutants
63 lacking Psb28 exhibited slower autotrophic growth, particularly under stress conditions^{31,44}, and
64 limited synthesis of Chl-binding proteins but without decreased PSII functionality³¹. The Psb28
65 mutant also exhibited an overall increase in PSII repair and faster recovery from
66 photodamage³¹. Chemical cross-linking combined with mass spectrometry revealed that Psb28
67 binds to the cytosolic side of CP47 close to cytochrome-*b*₅₅₉ and the Q_B binding site. Based on
68 this, researchers postulated a protective role for Psb28, where it blocks electron transport to the
69 acceptor side of PSII to shield the RC47 complex from excess photodamage during the
70 assembly process⁴⁵. This hypothesis is strengthened by the observation that Psb28 is also found
71 in PSII repair complexes⁴⁶. The luminal PSII assembly factor Psb27 has been similarly well
72 investigated. This lipoprotein is predominantly associated with inactive PSII fractions involved
73 in assembly or repair^{32,34,46-50}, stabilizing the CP43 luminal domain and presumably facilitating
74 the assembly of the OEC.

75 Our current knowledge of PSII biogenesis mainly describes the order of events and protein
76 composition of each intermediate, as well as the general roles of PSII assembly factors.
77 However, the precise molecular functions of these intermediate complexes and the involved
78 assembly factors are still elusive due to their low abundance and intrinsic instability. High-
79 resolution structural information is of vital importance to gain a deeper understanding into the
80 molecular action of PSII assembly factors, as they are proposed to alter the structures of their
81 associated PSII proteins to provide protection or facilitate specific biogenesis transitions.

82 Here, we use cryo-EM single particle analysis to describe the first molecular structure of a PSII
83 assembly intermediate. This structure represents one of the key transitions in PSII biogenesis:

the attachment of the CP43 module to the pre-assembled RC47 reaction center complex, which precedes incorporation and activation of the Mn_4CaO_5 cluster. We complement this structural data with spectroscopic analysis, revealing the first detailed insights into the molecular mechanisms of PSII assembly. Our study provides mechanistic answers to three long-standing questions: i) How do assembly factors modulate the structures of PSII subunits to assist biogenesis? ii) How is PSII protected from photodamage during assembly? iii) How is the PSII active site prepared for incorporation of the Mn_4CaO_5 cluster?

Results

Stable PSII intermediates were purified from the *T. elongatus* ΔpsbJ mutant⁴³ by affinity chromatography using a twin-strep-tag fused to the C-terminus of the CP43 subunit and subsequent ion exchange chromatography (Fig. S1a). The main peak of the IEC profile corresponds primarily to monomeric PSII, which lacks the extrinsic subunits PsbO, PsbU and PsbV that are indicative for water splitting activity (Fig. S1b and c). Single particle cryo-EM analysis of this PSII fraction resulted in three different high-resolution maps that allowed model building with high confidence and excellent statistics (Fig. S2, Table S1). In addition to the protein subunits, we also faithfully assigned all essential non-protein cofactors, including chlorophylls, quinones, carotenoids and lipids, which are also present in the mature PSII complex (Fig. S3). Consistent with previous biochemical studies^{34,43,48}, the EM density corresponding to the fully assembled, active Mn_4CaO_5 cluster is missing in the purified biogenesis intermediates. The first cryo-EM map (2.94 Å), which we call PSII-I (for PSII-Intermediate), provides a snapshot of the attachment of the CP43 module to the pre-assembled RC47 reaction center complex (Fig. 1). This PSII intermediate contains three assembly factors (Psb27, Psb28 and Psb34), as well as almost all the membrane-intrinsic subunits and cofactors found in mature PSII, although CP43 is bound in an immature conformation. Psb27 and Psb28 are well-known assembly factors^{31,32,35,43,51}, whereas the additional single transmembrane helix protein (tsl0063), which we named Psb34, has not been described before. The small subunit PsbY, which is known to be loosely bound⁵², is not resolved in our structure. In addition, PsbJ is not present, as the corresponding gene was inactivated to stall PSII assembly at this specific transition⁴³.

The two additional maps serve as internal controls. PSII-I' (2.76 Å) lacks Psb27 but is otherwise comparable to PSII-I; the root mean square deviation (RMSD) of the C_α atomic positions between similar subunits of the two complexes is 0.4 Å. Most likely, Psb27 was partly lost

during sample preparation. The third cryo-EM map (2.82 Å), which we call PSII-M (for PSII-Monomer), represents a monomeric PSII complex without bound assembly factors. Comparison of our PSII-M structure with a crystal structure of monomeric PSII⁵² (PDB-ID 3KZI, 3.6 Å) reveals only minimal differences between both structures, with a Cα RMSD of 1.3 Å, which verifies that the structural changes observed in PSII-I are not caused by the deletion of PsbJ.

Psb34 binds to CP47 and is involved in the RC47 transition

Our PSII-I structure provides the first identification of the single transmembrane helix protein Psb34 bound to a PSII assembly intermediate (Fig. 2a), which we also confirmed by mass spectrometry (Fig. 2b). Psb34 was probably overlooked previously due to its hydrophobicity and small size. It has a single transmembrane helix that binds to the CP47 antenna protein in close proximity to PsbH (Fig. 2a). Its conserved long N-terminal arm is located at the side and top of the D2 subunit (Fig. 2a). In addition, we independently confirmed the interaction of Psb34 with PSII assembly intermediates by isolation of strep-tagged Psb34 complexes from otherwise unmodified cells, indicating a specific function of Psb34 in the attachment of CP43 to RC47 (Fig. 2c). In this case, two distinct PSII intermediates were isolated via pulldown of strep-tagged Psb34: the RC47 complex with bound Psb28 and the subsequent PSII intermediate after attachment of CP43 and Psb27 (Fig. 2c). This observation implies that Psb28 is usually released from the PSII intermediate after attachment of CP43, probably after incorporation of PsbJ, as this trigger is missing in the analyzed *ΔpsbJ* mutant. PSII-I, which only accumulates in the *ΔpsbJ* mutant, seems to represent a hybrid state of both PSII intermediates that might be stabilized by the incomplete binding of CP43. Psb34 shows sequence similarity to high-light inducible proteins (HLIPs), which play a role in transient chlorophyll storage and chlorophyll biosynthesis⁵³. However, the chlorophyll binding motive is missing in Psb34 (Table S2), suggesting a distinct function for this protein in PSII biogenesis.

Psb28 induces conformational changes to the D1 D-E loop

Psb28 binds on the cytosolic faces of the D1 and D2 subunits, directly above the Q_B binding site (Fig. 3a), which differs from the position that was previously predicted by mass spectrometry⁴⁵. Its binding induces the formation of an extended beta-hairpin structure that incorporates the central anti-parallel beta-sheet of Psb28, the C-terminus of CP47 and the D1 D-E loop⁵⁴ (Fig. 3a). Binding of Psb28 to the C-terminus of CP47 also imparts a directionality

to the assembly process. In the Psb28-free complex (PSII-M), the CP47 C-terminus blocks the Psb28 binding site by interacting with the D1 D-E loop, thus preventing the reverse process and perturbation of active PSII by Psb28. Using nuclear magnetic resonance (NMR) spectroscopy, we performed chemical shift perturbation (CSP) experiments with recombinant Psb28 and a synthetic peptide of the conserved CP47 C-terminus to characterize this interaction in detail and determine the dissociation constant (K_D) (Fig. 3 and Fig. S4). The CSP measurements indicated significant shifts with a chemical shift difference ($\Delta\delta$) of more than one standard deviation located at strands $\beta 3$ and $\beta 4$ as well at the C-terminal region of Psb28 (Fig. 3c and d). Upon peptide binding, resonances for several residues gradually appeared with increasing peptide concentration, which were line-broadened beyond detection for the free form of Psb28. This observation indicates a less dynamic and more rigid complex structure. This is further supported by the heteronuclear Overhauser effect (NOE) data, which show that the C-terminus of Psb28 becomes rigid from L108 to K112 upon CP47 peptide binding due to creation of an intermolecular β -sheet (Fig. 3e). 2D-lineshape analysis was performed, yielding a K_D of $53.92 \pm 0.41 \mu\text{M}$ and a dissociation rate k_{off} of $10.14 \pm 0.16 \text{ s}^{-1}$, which is consistent with the observed slow-exchange in the NMR spectra (Fig. 3b). The affinity of Psb28 for full-length CP47 and PSII might indeed be even higher due to additional contacts between Psb28 and the D-E loop of D1 (Fig 3a).

Psb28 prevents full association of CP43 and distorts the Q_B -site

Binding of Psb28—with support of Psb34—causes major structural perturbations at the PSII acceptor side (Supplementary Movies 1 and 2), which mainly involve the D-E loops of the central D1 and D2 subunits. Comparison of the CP43 structure in PSII-I with that in our PsbJ-free control PSII-M (Fig. 4a-d) or with that in mature monomeric PSII (PDB-ID 3KZI) (Fig. 4c and d) reveals several differences. The CP43 C-terminus is not resolved in PSII-I, probably due to an immature position of the last transmembrane helix of CP43 and an altered conformation of the D1 D-E loop, which may prevent binding of the CP43 C-terminus to the cytoplasmic PSII surface (Fig. 4b). This region is close to the loop between helices D and E of the D2 subunit, which is also altered by binding of Psb28, as clearly shown by movement of D2 Arg233 (Fig. 4b, Fig. S5a and b). After dissociation of Psb28, the CP43 module undergoes a rigid body rotation where it clicks into place (Fig. 4b-d, Supplementary Movie 1), whereas binding of PsbJ and the extrinsic proteins PsbO, PsbV and PsbU during further maturation has very little influence on the CP43 binding position (Fig. 4c and d). The part of PSII that

originates from RC47 shows almost no difference between PSII-I and mature PSII (Fig. 4d), except for PsbE, which binds adjacent to PsbJ (Fig. 4c).

Most importantly, the structural changes in the D1 D-E loop may have a direct functional impact on PSII electron transfer (Fig. 4e and f), as this region coordinates several important PSII cofactors. In functional PSII, after charge separation at the reaction center P_{680} , electrons are transferred via pheophytin to the bound plastoquinone (Q_A) and further to mobile plastoquinone (Q_B). In our PSII-I structure, the Q_A site is fully assembled, and a well-resolved Q_A molecule is bound (Fig. 4e and f, Fig. S5c and d). The nearby non-heme iron is also already in place in PSII-I (Fig. 4e and f, Fig. S5e and f). The Q_B binding site of the PSII-M control is comparable to mature PSII, although it is not occupied by Q_B in our preparation (Fig. S5g). In contrast, the Q_B binding site of PSII-I is immature due to the Psb28- and Psb34-induced structural changes in the D1 D-E loop (Fig. 4e and f, Fig. S5h). Notably, D1 Phe265, which coordinates the head group of Q_B in mature PSII, is clearly at a different position²¹ (Supplementary Movie 2).

Binding of Psb28 protects PSII during biogenesis

A more detailed analysis of the structural environment close to the Q_A/Q_B binding sites revealed differences in the coordination and the hydrogen-bond network of the adjacent non-heme iron, which also indicate functional consequences for PSII electron transfer and charge recombination processes. In mature PSII, the non-heme iron is coordinated by four histidine residues and bicarbonate as the fifth ligand (Fig. 5a and c), whereas in PSII-I, the bicarbonate molecule is replaced by the E241 side-chain of D2 (Fig. 5b and d, Fig. S5e and f, Supplementary Movie 3). Other residues, including D1 E244 and Y246, which bind to the bicarbonate molecule in mature PSII (Fig. 5a), are also displaced in PSII-I due to the conformational change of the D1 D-E loop (Fig. 5b, Fig. S5e and f, Supplementary Movie 3). Binding of bicarbonate is important for PSII efficiency⁵⁵, as it lowers the redox potential of (Q_A/Q_A^-) to favor forward electron transport^{56,57}. If charge recombination occurs, the lower redox potential favors indirect charge recombination via $P_{680}^{+}/Pheo^+$. This back reaction yields triplet chlorophyll and subsequently singlet oxygen⁵⁷, a highly reactive oxidizing species. Changes in the redox potential of (Q_A/Q_A^-) have been proposed to tune the efficiency of PSII depending on the availability of CO_2 as the final electron acceptor and thereby protect PSII under low CO_2 conditions⁵⁷. Therefore, we used flash-induced variable fluorescence to monitor the stability of Q_A^- in active PSII, in the PSII-I assembly intermediate and in inactivated, Mn-depleted PSII, both of which lack a functional OEC (Fig. 5e, Fig. S6a and b). The results indicate that the

unique fluorescence traces observed with PSII-I are a consequence of Q_A/Q_A^- redox tuning by an absent OEC, an immature Q_B -binding pocket and probably the replacement of bicarbonate by glutamate as a ligand of the non-heme iron (see supplementary discussion for details). To determine the functional consequences of this redox tuning in PSII-I, we measured the formation of 1O_2 by EPR spectroscopy using the spin probe TEMPD. The data clearly show that 1O_2 formation is reduced by ~30% in PSII-I compared to inactivated PSII (Fig. 5f).

Psb27 binds to loop E of CP43 at the luminal PSII surface

Psb27 binds to the luminal side of the PSII complex, adjacent to loop E of the CP43 subunit (Fig. 6a and b). We could successfully model the Psb27 N-terminus until Cys22, which is the first residue after N-terminal processing of Psb27³⁴. Psb27-Cys22 is located close to the membrane plane, but the lipid moiety is not well enough resolved to build a model. In contrast to previously proposed models^{58,59}, the binding site of Psb27 has little overlap with the binding sites of the extrinsic subunits (PsbO, PsbV and PsbU) and has at least no direct contact to the Mn_4CaO_5 cluster binding site, particularly not to the D1 C-terminus (Fig. 6a and b), which has been suggested previously^{58,60}. Instead, Psb27 is bound at a remote position that might be occupied by CyanoQ in the mature complex⁶¹. This position of Psb27 may still prevent proper binding of PsbO³⁴ by lowering its binding affinity due to the partial overlap of the binding sites, although both proteins can bind together, as PsbO-containing Psb27-PSII complexes have been isolated via his-tagged Psb27⁴⁹. Additionally, Psb27 might stabilize loop E of CP43 in the unassembled state and facilitate its binding to the D1 subunit. This is of particular importance, as loop E of CP43 provides Arg345 and Glu342 (Arg357 and Glu354 in previous publications, see figure caption), to position the Mn_4CaO_5 cluster in mature PSII (Fig. 6b, dashed box). Glu342 serves as a ligand, whereas Arg345 stabilizes the Mn_4CaO_5 cluster by coordinating Asp170 of D1 through a hydrogen bond. Moreover, in the Psb27-bound state (PSII-I), the D1 C-terminus, which is directly involved in coordination of the Mn_4CaO_5 cluster²¹, is bound away from the cluster (Fig. 6c, Fig. S7), which might be an additional consequence of Psb27-induced stabilization of the CP43 E-loop. Thus, our PSII-I structure reveals not only how the Psb27 protein binds to CP43 and thus stabilizes it, but also indicates a role for the maturation of the oxygen evolving cluster that is consistent with functional data from previous studies^{32,35,48}.

PSII-I contains a single cation at the immature OEC site

The unique Mn_4CaO_5 cluster is a key feature of PSII that splits water into oxygen and protons. However, our PSII-I complex does not show any oxygen-evolving activity, suggesting that the oxygen evolving complex (OEC) is not fully assembled. In mature PSII, the Mn_4CaO_5 cluster is submerged in the complex and additionally capped by the extrinsic subunits PsbO, PsbU and PsbV (Fig. 6a and b). In our PSII-I structure, these subunits are absent, which leaves two parts of the CP43 E-loop (residues 320-327 and 397-404) in a flexible conformation, exposing the binding site of Mn_4CaO_5 cluster to the lumen. There is no strong density feature at this position that would correspond to the fully assembled metal-redox cofactor. Thus, our PSII-I structure provides a model for an immature OEC. Comparing our structure with the high-resolution crystal structure of mature PSII²¹ (PDB-ID 3WU2) provides insights into the first-steps of OEC biogenesis (Fig. 7).

The D1 C-terminus is an essential component for formation of the of OEC, as it provides several charged residues that are responsible for coordination of the chloride ion and the Mn_4CaO_5 cluster (Fig. 7a, b and d). The density for these C-terminal residues is weak in our PSII-I map, but traceable (Fig. S7a), indicating a flexibility that confirms the absence of the OEC. Compared to the mature complex, the last 12 residues of the C-terminal tail of D1 would need to undergo significant conformational changes to bring the side chains of Glu333, His337, Asp342, and the Ala344 C-terminus into the correct position to coordinate the Mn_4CaO_5 cluster (Supplementary Movie 4).

Moreover, we identify a clearly visible density at the position of the chloride ion, which is coordinated by Lys317 (D2) and the hydrogen atom of the backbone nitrogen of Glu333 (D1) in mature PSII (Fig. 7b and e). Despite the similar position, the Cl^- is coordinated by the nitrogen atom of the ring of adjacent His332 (D1) in PSII-I (Fig. 7a and e, Fig. S7d). Surprisingly, we identified another density in the area where the Mn_4CaO_5 cluster is located in mature PSII (Fig. 7a-c and f, Fig. S7c). However, this density is not large enough to reflect the whole cluster. Based on its size and interaction partners (Fig. 7f), it corresponds to one positively charged ion. In the structural context, this ion is most likely Mn^{2+} , but it could also be Ca^{2+} or any other positively charged ion.

This ion is coordinated by the side chains of D1 Asp170, Glu189, and His332, which are already in similar positions compared to mature PSII. Glu342 and Arg345 (Glu354 and Arg357 in previous publications) of CP43, which are both involved in the second coordination sphere of the Mn_4CaO_5 cluster, are also already pre-positioned through the interaction between Arg345

with D1 Asp170 (Fig. 7g). However, there are still significant conformational changes necessary for the transition from PSII-I to mature PSII, as highlighted in Figure 7d and g, as well as in Supplementary Movie 4. The D1 C-terminal tail must bring the side chains of Glu333, His337 and Asp342, as well as the C-terminus of Ala344, into correct alignment to coordinate the Mn_4CaO_5 cluster. In addition, the C-terminal tail of D2 needs to flip towards the D1 C-terminus (Fig. 7c, Fig. S7b, Supplementary Movie 4). In summary, PSII-I is characterized by only one positive charged ion bound instead of the complete Mn_4CaO_5 cluster, resulting in significantly different conformations of the D1 and D2 C-termini compared to the structural model containing a mature Mn_4CaO_5 cluster. However, the PSII-I structure seems to be prepared to accept the Mn_4CaO_5 cluster, as indicated by the above described similarities in side chain positioning.

Discussion

PSII biogenesis is a complex process that requires the action of specific assembly factors. These auxiliary proteins are not present in the mature complex and interact only transiently with specific subunits or preassembled PSII intermediates. Although more than 20 factors have been identified and assigned to specific transitions, their precise molecular functions in PSII assembly remain elusive in almost all cases. Our study provides the first detailed molecular insights into the function of PSII assembly factors Psb27, Psb28 and Psb34, which are involved in an important transition prior to activation of the OEC. The determined binding positions of Psb27 and Psb28, which are two of the most studied PSII assembly factors, define the binding with atomic resolution, in contrast to previous models^{58,59,62-66}. This provides precise information about the molecular function of Psb27 and Psb28 in the assembly process.

Binding of Psb28 and Psb34 to the cytoplasmic side of PSII induces large conformational changes in the D1 D-E loop (Fig. 4), which has been identified previously as an important location for PSII photoinhibition and D1 degradation^{67,68}. Structural changes observed in the PSII-I Q_B binding pocket and coordination of the non-heme iron suggest a functional impact on PSII electron transfer to protect the immature complex until water splitting is activated. In particular, D2 Glu241 replacing bicarbonate as a ligand of non-heme iron suggests a regulatory role, as binding of bicarbonate was proposed to tune PSII efficiency by changing the redox potential of ($\text{Q}_\text{A}/\text{Q}_\text{A}^-$)^{56,57}. As a functional consequence, PSII-I generates less singlet oxygen compared to inactivated PSII (Fig. 5f).

Interestingly, the coordination of the non-heme iron in PSII-I resembles that in non-oxygenic bacterial reaction centers (BRCs)⁶⁹ (Fig. S6c). In BRCs, the fifth ligand of the non-heme iron is provided by E234 of the M subunit⁷⁰, and mutagenesis of this residue induces changes in the free energy gap between the $P^{•+}/Q_A^{•-}$ radical pair⁷¹. These findings indicate that the environment of the non-heme iron is important for the Q_A redox potential, which influences charge recombination and the formation of reactive oxygen species⁵⁶.

The Psb27-bound and -unbound structures do not differ substantially (Fig. 6b), suggesting a rather subtle action in PSII biogenesis. Previous work demonstrated that Psb27 is already bound to free CP43³², where it might protect free CP43 from degradation or stabilize the E-loop in a specific conformation to chaperone the subsequent association with the RC47 complex. This step is crucial for preparing the binding site of the Mn_4CaO_5 cluster, as the CP43 E-loop is involved in its second coordination sphere⁷². Moreover, a recent high-speed AFM study with spinach PSII has shown fluctuations of the C43 E-loop at room temperature after removal of the extrinsic subunits and destruction of the Mn_4CaO_5 cluster⁷³, which might suggest a role of Psb27 in restraining the scope of conformational fluctuations for further assembly of the OEC. Consistent with this model, recent spectroscopic analysis of PSII from *Synechocystis* sp. PCC 6803 revealed that Psb27 facilitates photoactivation, probably by stabilization of intermediates with increased accessibility for ions, especially Ca^{2+} , to the site of cluster assembly⁷⁴. However, our data indicate a self-organization of the CP43 E-loop, at least in *T. elongatus*. The corresponding structural models for this region are almost identical, independent of binding of Psb27 or the extrinsic proteins, whereas the positions of the D1 and D2 C-termini are clearly different between intermediate and mature PSII. Binding of PsbO seems to be the main driver for these structural differences, and its presence interferes with efficient photoactivation, probably by closing the cluster binding site and thus preventing the exchange of ions⁷⁴. The overlap of the Psb27 and PsbO binding sites, although small, might decrease the PsbO affinity and prevent its premature binding, as initially suggested^{34,37}, or Psb27 may promote binding of PsbO in a different conformation (both proteins can bind simultaneously in a stable complex⁴⁹) that might support photoactivation.

Previous structural studies aimed to obtain mechanistic insights into the dark-rearrangement during photoactivation by removing the Mn_4CaO_5 cluster from fully assembled PSII, either by depleting it directly from PSII crystals by chemical treatment⁷⁵ or by cryo-EM single particle analysis in manganese- and calcium-free buffer⁷⁶. The X-ray structure was indeed missing the Mn_4CaO_5 cluster, but the D1 C-terminus followed mostly the same trajectory as found in the mature PSII-dimer structure. The authors suggested that the D1 C-terminus might not rearrange

during Mn₄CaO₅ cluster assembly. However, the crystal structure was dimeric and still had the extrinsic subunits PsbO, PsbU, and PsbV bound. It is known that these subunits are typically not associated with Mn₄CaO₅ cluster-depleted PSII. Thus, the structure might be artificially stabilized by crystal packing forces. The cryo-EM structure, on the other hand, revealed a monomeric PSII that lacks extrinsic subunits and the Mn₄CaO₅ cluster⁷⁶. This structure is more similar to our PSII biogenesis intermediate PSII-I, as PsbY, PsbZ and PsbJ are also missing. The PsbJ subunit is surprising; it is an integral subunit of PSII and should not be easily detached, yet it is missing from this structure, as we deleted it to stabilize our PSII-I complex. These observations might indicate a more specific and regulatory role of PsbJ in PSII biogenesis. Additionally, the D1 C-terminus is disordered in this previous cryo-EM structure, and the authors suggest that the dark-rearrangement involves a transition from a disordered to an ordered state.

Our structure now reveals the fate of the D1 C-terminus with the assembly factor Psb27 bound. The D1 C-terminus follows a different trajectory compared to the mature PSII. Thus, we provide structural evidence that the slow dark-rearrangement involves a conformational change of the D1 C-terminus rather than the previously proposed disorder-to-order transition after initial photoactivation⁷⁶. Compared to mature PSII, twelve residues of the D1 C-terminal tail must undergo significant conformational changes to bridge the side chains of Glu333, His337 and Asp342, as well as to bring the C-terminus of Ala344 to the correct position to coordinate the Mn₄CaO₅ cluster (Fig. 6c and 7d, Supplementary Movie 4), which is consistent with previous models^{39,77,78}. We also identified a single positively charged ion in our PSII-I structure, coordinated by Asp170, Glu189 and His332 of D1 (Fig. 7f), at the position of the Mn₄CaO₅ cluster of mature PSII. This binding site most likely corresponds to the long-sought single high-affinity site (HAS), where the first Mn²⁺ binds prior to the first photoactivation step in OEC biogenesis⁴¹. The presence of manganese in immature PSII, due to a PrtA-facilitated pre-loading mechanism, has been previously suggested⁷⁹. However, we cannot exclude binding of Ca²⁺, which was shown to bind with a much lower affinity^{77,80}, or any other positively charged ion at this position. Nevertheless, Asp170 has been identified as the most critical residue for the HAS^{81,82}, which supports our hypothesis. Further photoactivation steps occur presumably after cooperative binding of calcium and manganese. The binding of the extrinsic subunit PsbO, potentially after release of Psb27 and maturation of the water oxidizing complex, is the next step of the PSII assembly line *in vivo*, which leads to the next unsolved question in PSII biogenesis: what triggers the release of an assembly factor? For Psb27, its detachment might be promoted by the binding of PsbO, as their binding sites partially overlap.

Membrane protein complexes play a fundamental role in bioenergetics to sustain and proliferate life on Earth. They drive the light-to-chemical energy conversion in photosynthetic organisms and are essential for energy supply in heterotrophs. These intricate molecular machines are assembled from numerous single proteins in a spatiotemporally synchronized process that is facilitated by a network of assembly factors. These auxiliary proteins are the key players of Nature's assembly lines. Our PSII-I cryo-EM structure reveals the first molecular snapshot of PSII biogenesis and, accompanied by our spectroscopic and biochemical analyses, provides clear mechanistic insights into how three assembly factors (Psb27, Psb28 and Psb34) coordinate the stepwise construction of this powerful catalyst of life.

Methods

Cultivation of *Thermosynechococcus elongatus* BP-1

Cell growth and thylakoid membrane preparation were performed as described previously⁸³. In brief, *T. elongatus* mutant strains (Δ psbJ psbC-TS and psb34-TS) were grown in BG-11 liquid medium inside a 25-litre foil fermenter (Bioengineering) at 45°C, 5% (v/v) CO₂-enriched air bubbling and 50-200 μ mol photons m⁻² s⁻¹ white light illumination (depending on the cell density). Cells were harvested at an OD₆₈₀ of ~ 2 after 5-6 days of cultivation and concentrated to ~ 0.5 l, using an Amicon DC10 LA hollow fibre system, pelleted (3500 rcf, 45 min and 25 °C) and resuspended in 150 ml of Buffer D (100 mM Tris-HCL, pH 7.5, 10 mM MgCl₂, 10 mM CaCl₂, 500 mM mannitol and 20% (w/v) glycerol). The harvested cells were flash-frozen in liquid nitrogen and stored at -80 °C until further use.

Preparation of *T. elongatus* mutant strains

Thermosynechococcus elongatus ΔpsbJ psbC-TS was generated based on the previously described strain *T. elongatus* ΔpsbJ⁴³ that was transformed with the plasmid pCP43-TS. The plasmid is based on pCP34-10His³⁴. The His-tag sequence was exchanged with TwinStrep-tag by PCR using the primers CP43TS_rev (5'CCCGATATCTTACTTCTCAAATTGCGGATGAGACCACGCAGAACCACCAGAACCACCGCCGCTGCCGCCGCCTTTTTTCGAACTGCGGGTGGCTCC 3') and NTCP43 (5' TGCTCTAGAATGAAAACCTTTGTCTTCCCAGA 3'). The resulting PCR product was ligated back into an empty pCP34-10His backbone using XbaI and EcoRV restriction endonucleases. *T. elongatus* BP-1 cells were transformed as described previously⁸⁴. Mutant colonies were selected by frequent re-plating onto agar plates with increasing antibiotic concentrations, stopping at 8 μg/ml of chloramphenicol and 80 μg/ml

of kanamycin. Complete segregation of the mutant was confirmed by PCR with the primers CTCP43DS (5' CCGCTCGAGACCATCCAAGCTTGGCAGCA 3') and NTCP43 (5' TGCTCTAGAATGAAAACCTTTGTCTTCCCAGA 3').

T. elongatus psb34-TS was generated by transformation with the plasmid pPsb34-TS. The plasmid DNA was obtained from TwistBioscience. It consisted of psb34 (tsl0063) with a C-terminal TwinStrep-tag and a kanamycin resistance cassette, flanked by tsl0063-upstream and downstream regions (900 bp each). *T. elongatus* BP-1 cells were transformed⁸⁴ and mutant selection took place³⁴. Complete segregation of the mutant was verified by PCR. The primers used were tsl0063-up-for (5' CATATGGTCTCGCAATTATTTGCCCCATGC 3') and tsl0063-down-rev (5' GGTACCCCG ACACAGTTGATCACCGC 3').

Purification of photosystem II assembly intermediates

Thawed cells were diluted in 100 ml of Buffer A (100 mM Tris-HCL, pH 7.5, 10 mM MgCl₂ and 10 mM CaCl₂) and pelleted again (21 000 rcf, 20 min and 4°C). The pellet was resuspended in 100 ml of Buffer A with 0.2% (w/v) lysozyme and dark incubated for 75-90 min at 37 °C. This was followed by cell disruption by Parr bomb (Parr Instruments Company) and pelleting (21 000 rcf, 20 min and 4°C). All following steps were performed under green illumination. The pellet was resuspended in 150 ml of Buffer A and pelleted again (21 000 rcf, 20 min and 4°C). This step was repeated three times, with the last resuspension in 80 ml of Buffer B (100 mM Tris-HCL, pH 7.5, 10 mM MgCl₂, 10 mM CaCl₂ and 500 mM mannitol). The isolated thylakoid membranes were flash frozen in liquid nitrogen and stored at -80 °C.

Strep-Tactin-affinity purification of PsbC-TS and Psb34-TS assembly intermediates were performed under green illumination. Membrane protein extraction was performed as described previously⁸³, with certain adaptations. Isolated thylakoid membranes were supplemented with 0.05% (w/v) n-Dodecyl β-maltoside (DDM) (Glycon) and pelleted (21000 rcf, 20 min and 4°C). The sample was resuspended in extraction buffer (100 mM Tris-HCL, pH 7.5, 10 mM MgCl₂, 10 mM CaCl₂, 1.2% (w/v) DDM, 0.5% (w/v) sodium-cholate and 0.01% (w/v) DNase) to a final chlorophyll concentration of 1 mg/ml and incubated for 30 min at 20 °C. The solubilized membrane proteins were ultra-centrifugated (140000 rcf, 60 min and 4 °C) and NaCl was added to the supernatant to a final concentration of 300 mM.

The supernatant was filtered through a 0.45 μm syringe filter (Sarsted AG, Germany) and applied to a 5 ml Strep-Tactin Superflow HC column (IBA Lifesciences), equilibrated in Buffer W (100 mM Tris-HCL, pH 7.5, 10 mM MgCl₂, 10 mM CaCl₂, 500 mM mannitol, 300 mM

NaCl and 0.03% (w/v) DDM) at a flowrate of 3 ml/min. The column was washed with Buffer W until a stable baseline (A_{280}) was reached. Strep-tagged protein complexes were eluted by an isocratic elution with Buffer E (100 mM Tris-HCL, pH 7.5, 10 mM $MgCl_2$, 10 mM $CaCl_2$, 500 mM mannitol, 300 mM NaCl 2.5 mM desthiobiotin and 0.03% (w/v) DDM). The captured fractions were equilibrated in Buffer F (20 mM MES, pH 6.5, 10 mM $MgCl_2$, 10 mM $CaCl_2$, 500 mM mannitol and 0.03% (w/v) DDM) with a spin concentrator (Amicon, Ultra – 15, 100000 NMWL), flash-frozen in liquid nitrogen and stored at -80 °C until analysis.

PsbC-TS containing assembly intermediates were further separated by ion exchange chromatography (IEC). Captured elution fraction from the Strep-Tactin-affinity purification were loaded onto a anion exchange column (UNO Q-6, Biorad) with a flowrate of 4 ml/min, pre-equilibrated in Buffer F. Protein complexes were eluted by a liner gradient of $MgSO_4$ (0-150 mM) using Buffer G (20 mM MES, pH 6.5, 10 mM $MgCl_2$, 10 mM $CaCl_2$, 500 mM mannitol, 150 mM $MgSO_4$ and 0.03% (w/v) DDM). Fractions containing PSII assembly intermediates were collected, concentrated to 100 – 10 μ M reaction centers, using a spin concentrator (Amicon, Ultra – 15, 100 000 NMWL), aliquoted, flash frozen in liquid nitrogen and stored at -80 °C until further analysis.

Cryo-electron microscopy

For cryo-EM sample preparation, 4.5 μ l of purified protein complexes were applied to glow discharged Quantifoil 2/1 grids, blotted for 3.5 s with force 4 in a Vitrobot Mark III (Thermo Fisher) at 100% humidity and 4°C, then plunge frozen in liquid ethane, cooled by liquid nitrogen. Cryo-EM data was acquired with a FEI Titan Krios transmission electron microscope using the SerialEM software⁸⁵. Movie frames were recorded at a nominal magnification of 22,500x using a K3 direct electron detector (Gatan), The total electron dose of ~55 electrons per $\text{\AA}^2$ was distributed over 30 frames at a calibrated physical pixel size of 1.09 $\text{\AA}$. Micrographs were recorded in a defocus range of -0.5 to -3.0 μ m.

Image processing, classification and refinement

Cryo-EM micrographs were processed on the fly using the Focus software package⁸⁶ if they passed the selection criteria (iciness < 1.05 , drift $0.4 \text{ \AA} < x < 70 \text{ \AA}$, defocus $0.5 \mu\text{m} < x < 5.5 \mu\text{m}$, estimated CTF resolution $< 6 \text{ \AA}$). Micrograph frames were aligned using MotionCorr⁸⁷ and the contrast transfer function (CTF) for aligned frames was determined using Gctf⁸⁸. Using Gautomatch (<http://www.mrc-lmb.cam.ac.uk/kzhang/>) 693,297 particles were picked template-free on 824 acquired micrographs. Particles were extracted with a pixel box size of 260 using RELION 3.1⁸⁹ and imported into Cryosparc 2.3⁹⁰. After reference-free 2D classification,

675,123 particles were used for ab initio construction of initial models and subjected to multiple rounds of 3D classification to obtain models with and without Psb28 density. Non-uniform refinement in Cryosparc resulted in models with an estimated resolution of ~3.2 Å. Particles belonging to 3D classes with and without Psb28 (150,090 and 166,411 particles, respectively) were reextracted in RELION with a pixel box size of 256 and subjected to several rounds of CTF-refinement (estimation of anisotropic magnification, fit of per-micrograph defocus and astigmatism and beam tilt estimation) and Bayesian polishing⁹¹. Both classes were refined using the previously generated starting models. 3D classification without further alignment using a mask around the Psb27 region separated particles in the Psb28-containing class into distinct classes with and without Psb27 (57,862 and 91,473 particles, respectively). Final refinement of each of the three classes (with Psb27 and Psb28 (PSII-I), with Psb28 but without Psb27 (PSII-I'), and without Psb27 and Psb28 (PSII-M)) resulted in models with global resolutions of 2.94 Å, 2.76 Å and 2.82 Å, respectively (Gold standard FSC analysis of two independent half-sets at the 0.143 cutoff). Local-resolution and 3D-FSC plots (Extended Data Fig. 2) were calculated using RELION and the “Remote 3DFSC Processing Server” web interface⁹², respectively.

Atomic model construction

The 3.6 Å resolution X-ray structure of monomeric PSII from *T. elongatus* with PDB-ID 3KZI⁵² was used as initial structural model that was docked as rigid body using Chimera⁹³ into the obtained cryo EM densities for PSII-M and PSII-I. The cofactors that had no corresponding density were removed. The subunit PsbJ was also removed, as it was deleted in the experimental design. By highlighting the still unoccupied parts of the PSII-I density map, we identified densities that lead to the structures of Psb27, Psb28, and Psb34.

The 2.4 Å resolution X-ray structures of isolated Psb28 from *T. elongatus* with PDB-ID 3ZPN⁹⁴ and the 1.6 Å resolution X-ray structure of isolated Psb27 from *T. elongatus* with PDB-ID 2Y6X⁶⁵ were docked as rigid bodies into the unoccupied densities. The 1.6 Å resolution X-ray structure of CyanoQ from *T. elongatus* with PDB-ID 3ZSU⁹⁵ does not fit into the density and was therefore not modeled.

As there was no experimentally resolved structural model of Psb34 available, we first used the sequence with UniProt-ID Q8DMP8 to predict structures using the webserver SWISS Model⁹⁶ and LOMETS⁹⁷. We also predicted the secondary structure through the meta server Bioinformatics Toolkit⁹⁸ and CCTOP⁹⁹. The results of the secondary structure prediction are summarized in Table S4. Combining these predictions together with the unassigned cryo-EM

density, we used COOT¹⁰⁰ to build an initial model of Psb34 that has one α -helix from amino acid number 28 to 55.

Model Refinement

The initial model of the complex described above was refined in real space against the cryo-EM density of PSII-I, and structural clashes were removed using molecular dynamics flexible fitting (MDFF)¹⁰¹. MDFF simulations were prepared in VMD 1.9.4a35¹⁰² using QwikMD¹⁰³ and the MDFF plugin. The simulations were carried out with NAMD 2.13¹⁰⁴ employing the CHARMM36 force field. Secondary structure, cis peptide and chirality restraints were employed during 800 steps of minimization followed by a 40 ps MDFF simulation at 300K. Due to the employed restraints, only conformational changes of side chains and subunit movements compared to the initial structure are identified during the initial MDFF run. We checked the fit to density of the structure by calculating cross-correlation values of the backbone atoms. For PSII-I, we identified residues 217 to 269 from PsbA and residues 467 to 499 from PsbB and PsbZ as main regions where the structural model was not yet in accordance with the density after the initial MDFF run. For these three regions, we employed an iterative combination of MDFF with Rosetta^{105,106}. Here, we used the optimized strategy as described for model construction of the 26S proteasome^{107,108}.

To obtain an atomic model that fit the PSII-M density, we used the initial model based on 3KZI described above, but without PsbJ, Psb27, Psb28, and Psb34. After the initial MDFF run, the cross-correlation check did not reveal any regions with significant deviation between model and density. Therefore, no further refinement was necessary. This fast convergence reflects that there are no crucial differences between the PSII-M model and the X-ray structure 3KZI.

To obtain the atomic model that fit the PSII-I' density, we used the final PSII-I model without Psb27 for MDFF. After the initial MDFF run, the cross-correlation check did not reveal any regions with significant deviation between model and density. This fast convergence reflects that there are no crucial differences between the PSII-I and PSII-I' models, except for the presence of the Psb27 subunit.

Last, the PSII-M, PSII-I, and PSII-I' models were used to initiate one final round of real-space refinement in Phenix¹⁰⁹.

Acknowledgements

We thank C. König, M. Völkel, and R. Oworah-Nkruma for excellent technical assistance, Kristin Becker for cloning of the pIVEX_{Psb28His} plasmid, Bibi Erjavec for preparation of the

scheme in Fig. 1 and Nicholas Cox for helpful discussion. J.M.S. is grateful to E. Conti for scientific independence and great mentorship and to J. M. Plitzko and W. Baumeister for access to the cryo-EM infrastructure and early career support. M.M.N. is grateful to his mentor M. Rögner for generous support. **Funding:** Financial support was provided by the Max Planck Society, the Helmholtz Zentrum München, the DFG research unit FOR2092 (EN 1194/1-1 to B.D.E., NO 836/3-2 to M.M.N.), the DFG priority program 2002 (NO 836/4-1 to M.M.N.), the grant NIH P41-GM104601 (to E.T.) and an Emmy-Noether fellowship (SCHU 3364/1-1 to J.M.S.). A.K.-L. was supported by the LabEx Saclay Plant Sciences-SPS (grant number ANR-10-LABX-0040-SPS) and the French Infrastructure for Integrated Structural Biology (FRISBI; grant number ANR-10-INSB-05). R.S. gratefully acknowledges support from the DFG (INST 213/757-1 FUGG and INST 213/843-1 FUGG). **Author contributions:** B.D.E., T.R., J.M.S. and M.M.N. conceived the research, prepared the figures, and wrote the manuscript with the contributions of all other authors. M.M.N. coordinated the activities. Preparation of mutants, PSII isolation and biochemical analysis were performed by J.Z., M.M., P.L. and M.M.N. Mass spectrometry analysis was done by J.M.-C. and J.D.L. J.M.S., S.B. and B.D.E. performed the cryo-EM analysis and T.R. built the structural model with the help of S.K.S., A.C. and E.T. Fluorescence spectroscopy was carried out by J.Z. and M.M.N. EPR experiments were conducted by A.K.-L. NMR experiments were conducted and analyzed by O.A. and R.S.. All authors approved the final version of the manuscript. **Competing interests:** The authors declare no competing interests. **Data availability:** The cryo-EM density maps will be deposited in the Electron Microscopy Data Bank, the atomic models of the cryo-EM structures in the worldwide Protein Data Bank (wwPDB) and the NMR assignments for Psb28 in the Biological Magnetic Resonance Bank (BMRB), respectively.

References

- 1 Hohmann-Marriott, M. F. & Blankenship, R. E. Evolution of photosynthesis. *Annual review of plant biology* **62**, 515-548, doi:10.1146/annurev-arplant-042110-103811 (2011).
- 2 Sanchez-Baracaldo, P. & Cardona, T. On the origin of oxygenic photosynthesis and Cyanobacteria. *New Phytol* **225**, 1440-1446, doi:10.1111/nph.16249 (2020).
- 3 Vinyard, D. J., Ananyev, G. M. & Dismukes, G. C. Photosystem II: the reaction center of oxygenic photosynthesis. *Annual review of biochemistry* **82**, 577-606, doi:10.1146/annurev-biochem-070511-100425 (2013).
- 4 Boekema, E. J. *et al.* Supramolecular structure of the photosystem II complex from green plants and cyanobacteria. *Proc Natl Acad Sci U S A* **92**, 175-179, doi:10.1073/pnas.92.1.175 (1995).

- 569 5 Cox, N., Pantazis, D. A. & Lubitz, W. Current Understanding of the Mechanism of
570 Water Oxidation in Photosystem II and Its Relation to XFEL Data. *Annual review of*
571 *biochemistry* **89**, 795-820, doi:10.1146/annurev-biochem-011520-104801 (2020).
- 572 6 Shen, J. R. The Structure of Photosystem II and the Mechanism of Water Oxidation in
573 Photosynthesis. *Annual review of plant biology* **66**, 23-48, doi:10.1146/annurev-arplant-
574 050312-120129 (2015).
- 575 7 Yano, J. *et al.* Light-dependent production of dioxygen in photosynthesis. *Met Ions Life*
576 *Sci* **15**, 13-43, doi:10.1007/978-3-319-12415-5_2 (2015).
- 577 8 Ferreira, K. N., Iverson, T. M., Maghlaoui, K., Barber, J. & Iwata, S. Architecture of
578 the photosynthetic oxygen-evolving center. *Science* **303**, 1831-1838,
579 doi:10.1126/science.1093087 (2004).
- 580 9 Cardona, T., Sedoud, A., Cox, N. & Rutherford, A. W. Charge separation in
581 photosystem II: a comparative and evolutionary overview. *Biochim Biophys Acta* **1817**,
582 26-43, doi:10.1016/j.bbabbio.2011.07.012 (2012).
- 583 10 Holzwarth, A. R. *et al.* Kinetics and mechanism of electron transfer in intact
584 photosystem II and in the isolated reaction center: pheophytin is the primary electron
585 acceptor. *Proc Natl Acad Sci U S A* **103**, 6895-6900, doi:10.1073/pnas.0505371103
586 (2006).
- 587 11 Müh, F., Glöckner, C., Hellmich, J. & Zouni, A. Light-induced quinone reduction in
588 photosystem II. *Biochim Biophys Acta* **1817**, 44-65, doi:10.1016/j.bbabbio.2011.05.021
589 (2012).
- 590 12 Faller, P. *et al.* Rapid formation of the stable tyrosyl radical in photosystem II. *Proc*
591 *Natl Acad Sci U S A* **98**, 14368-14373, doi:10.1073/pnas.251382598 (2001).
- 592 13 Roose, J. L., Frankel, L. K., Mummadisetti, M. P. & Bricker, T. M. The extrinsic
593 proteins of photosystem II: update. *Planta* **243**, 889-908, doi:10.1007/s00425-015-
594 2462-6 (2016).
- 595 14 Müh, F. & Zouni, A. Structural basis of light-harvesting in the photosystem II core
596 complex. *Protein Sci* **29**, 1090-1119, doi:10.1002/pro.3841 (2020).
- 597 15 Shi, L. X., Hall, M., Funk, C. & Schröder, W. P. Photosystem II, a growing complex:
598 updates on newly discovered components and low molecular mass proteins. *Biochim*
599 *Biophys Acta* **1817**, 13-25, doi:10.1016/j.bbabbio.2011.08.008 (2012).
- 600 16 Stewart, D. H. & Brudvig, G. W. Cytochrome b559 of photosystem II. *Biochim Biophys*
601 *Acta* **1367**, 63-87, doi:10.1016/s0005-2728(98)00139-x (1998).
- 602 17 Cox, N. *et al.* Photosynthesis. Electronic structure of the oxygen-evolving complex in
603 photosystem II prior to O-O bond formation. *Science* **345**, 804-808,
604 doi:10.1126/science.1254910 (2014).
- 605 18 Kern, J. *et al.* Structures of the intermediates of Kok's photosynthetic water oxidation
606 clock. *Nature* **563**, 421-425, doi:10.1038/s41586-018-0681-2 (2018).
- 607 19 Kupitz, C. *et al.* Serial time-resolved crystallography of photosystem II using a
608 femtosecond X-ray laser. *Nature* **513**, 261-265, doi:10.1038/nature13453 (2014).
- 609 20 Suga, M. *et al.* An oxyl/oxo mechanism for oxygen-oxygen coupling in PSII revealed
610 by an x-ray free-electron laser. *Science* **366**, 334-338, doi:10.1126/science.aax6998
611 (2019).
- 612 21 Umena, Y., Kawakami, K., Shen, J. R. & Kamiya, N. Crystal structure of oxygen-
613 evolving photosystem II at a resolution of 1.9 Å. *Nature* **473**, 55-60,
614 doi:10.1038/nature09913 (2011).
- 615 22 Krieger-Liszakay, A., Fufezan, C. & Trebst, A. Singlet oxygen production in
616 photosystem II and related protection mechanism. *Photosynth Res* **98**, 551-564,
617 doi:10.1007/s11120-008-9349-3 (2008).
- 618 23 Vass, I. Molecular mechanisms of photodamage in the Photosystem II complex.
619 *Biochim Biophys Acta* **1817**, 209-217, doi:10.1016/j.bbabbio.2011.04.014 (2012).

- 24 Shevela, D. *et al.* 'Birth defects' of photosystem II make it highly susceptible to photodamage during chloroplast biogenesis. *Physiologia plantarum* **166**, 165-180 (2019).
- 25 Heinz, S., Liauw, P., Nickelsen, J. & Nowaczyk, M. Analysis of photosystem II biogenesis in cyanobacteria. *Biochim Biophys Acta* **1857**, 274-287, doi:10.1016/j.bbabi.2015.11.007 (2016).
- 26 Nickelsen, J. & Rengstl, B. Photosystem II assembly: from cyanobacteria to plants. *Annual review of plant biology* **64**, 609-635, doi:10.1146/annurev-arplant-050312-120124 (2013).
- 27 Nixon, P. J., Michoux, F., Yu, J., Boehm, M. & Komenda, J. Recent advances in understanding the assembly and repair of photosystem II. *Ann Bot* **106**, 1-16, doi:10.1093/aob/mcq059 (2010).
- 28 Komenda, J. *et al.* Cleavage after residue Ala352 in the C-terminal extension is an early step in the maturation of the D1 subunit of Photosystem II in *Synechocystis* PCC 6803. *Biochim Biophys Acta* **1767**, 829-837, doi:10.1016/j.bbabi.2007.01.005 (2007).
- 29 Komenda, J. *et al.* The cyanobacterial homologue of HCF136/YCF48 is a component of an early photosystem II assembly complex and is important for both the efficient assembly and repair of photosystem II in *Synechocystis* sp. PCC 6803. *J Biol Chem* **283**, 22390-22399, doi:10.1074/jbc.M801917200 (2008).
- 30 Boehm, M. *et al.* Subunit composition of CP43-less photosystem II complexes of *Synechocystis* sp. PCC 6803: implications for the assembly and repair of photosystem II. *Philos Trans R Soc Lond B Biol Sci* **367**, 3444-3454, doi:10.1098/rstb.2012.0066 (2012).
- 31 Dobáková, M., Sobotka, R., Tichy, M. & Komenda, J. Psb28 protein is involved in the biogenesis of the photosystem II inner antenna CP47 (PsbB) in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol* **149**, 1076-1086, doi:10.1104/pp.108.130039 (2009).
- 32 Komenda, J. *et al.* The Psb27 assembly factor binds to the CP43 complex of photosystem II in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol* **158**, 476-486, doi:10.1104/pp.111.184184 (2012).
- 33 Mamedov, F., Nowaczyk, M. M., Thapper, A., Rögner, M. & Styring, S. Functional characterization of monomeric photosystem II core preparations from *Thermosynechococcus elongatus* with or without the Psb27 protein. *Biochemistry* **46**, 5542-5551, doi:10.1021/bi7000399 (2007).
- 34 Nowaczyk, M. M. *et al.* Psb27, a cyanobacterial lipoprotein, is involved in the repair cycle of photosystem II. *Plant Cell* **18**, 3121-3131, doi:10.1105/tpc.106.042671 (2006).
- 35 Roose, J. L. & Pakrasi, H. B. The Psb27 protein facilitates manganese cluster assembly in photosystem II. *J Biol Chem* **283**, 4044-4050, doi:10.1074/jbc.M708960200 (2008).
- 36 Bao, H. & Burnap, R. L. Photoactivation: The Light-Driven Assembly of the Water Oxidation Complex of Photosystem II. *Front Plant Sci* **7**, 578, doi:10.3389/fpls.2016.00578 (2016).
- 37 Becker, K., Cormann, K. U. & Nowaczyk, M. M. Assembly of the water-oxidizing complex in photosystem II. *J Photochem Photobiol B* **104**, 204-211, doi:10.1016/j.jphotobiol.2011.02.005 (2011).
- 38 Cheniae, G. & Martin, I. Photoactivation of the manganese catalyst of O₂ evolution. I- Biochemical and kinetic aspects. *Biochim Biophys Acta* **253**, 167-181, doi:10.1016/0005-2728(71)90242-8 (1971).
- 39 Dasgupta, J., Ananyev, G. M. & Dismukes, G. C. Photoassembly of the Water-Oxidizing Complex in Photosystem II. *Coord Chem Rev* **252**, 347-360, doi:10.1016/j.ccr.2007.08.022 (2008).

670 40 Radmer, R. & Cheniae, G. M. Photoactivation of the manganese catalyst of O₂
671 evolution. II. A two-quantum mechanism. *Biochimica et Biophysica Acta* **253**, 182-186,
672 doi: 10.1016/0005-2728(71)90243-X (1971).

673 41 Nixon, P. J. & Diner, B. A. Aspartate 170 of the photosystem II reaction center
674 polypeptide D1 is involved in the assembly of the oxygen-evolving manganese cluster.
675 *Biochemistry* **31**, 942-948, doi:10.1021/bi00118a041 (1992).

676 42 Regel, R. E. *et al.* Deregulation of electron flow within photosystem II in the absence
677 of the PsbJ protein. *Journal of Biological Chemistry* **276**, 41473-41478 (2001).

678 43 Nowaczyk, M. M. *et al.* Deletion of *psbJ* leads to accumulation of Psb27-Psb28
679 photosystem II complexes in *Thermosynechococcus elongatus*. *Biochim Biophys Acta*
680 **1817**, 1339-1345, doi:10.1016/j.bbabi.2012.02.017 (2012).

681 44 Sakata, S., Mizusawa, N., Kubota-Kawai, H., Sakurai, I. & Wada, H. Psb28 is involved
682 in recovery of photosystem II at high temperature in *Synechocystis* sp. PCC 6803.
683 *Biochim Biophys Acta* **1827**, 50-59, doi:10.1016/j.bbabi.2012.10.004 (2013).

684 45 Weisz, D. A. *et al.* Mass spectrometry-based cross-linking study shows that the Psb28
685 protein binds to cytochrome b559 in Photosystem II. *Proc Natl Acad Sci U S A* **114**,
686 2224-2229, doi:10.1073/pnas.1620360114 (2017).

687 46 Bečková, M. *et al.* Association of Psb28 and Psb27 Proteins with PSII-PSI
688 Supercomplexes upon Exposure of *Synechocystis* sp. PCC 6803 to High Light. *Mol*
689 *Plant* **10**, 62-72, doi:10.1016/j.molp.2016.08.001 (2017).

690 47 Bentley, F. K., Luo, H., Dilbeck, P., Burnap, R. L. & Eaton-Rye, J. J. Effects of
691 inactivating psbM and psbT on photodamage and assembly of photosystem II in
692 *Synechocystis* sp. PCC 6803. *Biochemistry* **47**, 11637-11646, doi:10.1021/bi800804h
693 (2008).

694 48 Grasse, N. *et al.* Role of novel dimeric Photosystem II (PSII)-Psb27 protein complex in
695 PSII repair. *J Biol Chem* **286**, 29548-29555, doi:10.1074/jbc.M111.238394 (2011).

696 49 Liu, H., Roose, J. L., Cameron, J. C. & Pakrasi, H. B. A genetically tagged Psb27 protein
697 allows purification of two consecutive photosystem II (PSII) assembly intermediates in
698 *Synechocystis* 6803, a cyanobacterium. *J Biol Chem* **286**, 24865-24871,
699 doi:10.1074/jbc.M111.246231 (2011).

700 50 Weisz, D. A. *et al.* A novel chlorophyll protein complex in the repair cycle of
701 photosystem II. *Proc Natl Acad Sci U S A* **116**, 21907-21913,
702 doi:10.1073/pnas.1909644116 (2019).

703 51 Mabbitt, P. D., Wilbanks, S. M. & Eaton-Rye, J. J. Structure and function of the
704 hydrophilic Photosystem II assembly proteins: Psb27, Psb28 and Ycf48. *Plant Physiol*
705 *Biochem* **81**, 96-107, doi:10.1016/j.plaphy.2014.02.013 (2014).

706 52 Broser, M. *et al.* Crystal structure of monomeric photosystem II from
707 *Thermosynechococcus elongatus* at 3.6-Å resolution. *J Biol Chem* **285**, 26255-26262,
708 doi:10.1074/jbc.M110.127589 (2010).

709 53 Komenda, J. & Sobotka, R. Cyanobacterial high-light-inducible proteins--Protectors of
710 chlorophyll-protein synthesis and assembly. *Biochim Biophys Acta* **1857**, 288-295,
711 doi:10.1016/j.bbabi.2015.08.011 (2016).

712 54 Mulo, P. *et al.* Mutagenesis of the D-E loop of photosystem II reaction centre protein
713 D1. Function and assembly of photosystem II. *Plant Mol Biol* **33**, 1059-1071,
714 doi:10.1023/a:1005765305956 (1997).

715 55 Eaton-Rye, J. J. & Govindjee. Electron transfer through the quinone acceptor complex
716 of photosystem II in bicarbonate-depleted spinach thylakoid membranes as a function
717 of actinic flash number and frequency. *Biochimica et Biophysica Acta (BBA)-*
718 *Bioenergetics* **935**, 237-247 (1988).

719 56 Allen, J. F. & Nield, J. Redox Tuning in Photosystem II. *Trends Plant Sci* **22**, 97-99,
720 doi:10.1016/j.tplants.2016.11.011 (2017).

721 57 Brinkert, K., De Causmaecker, S., Krieger-Liszkay, A., Fantuzzi, A. & Rutherford, A.
722 W. Bicarbonate-induced redox tuning in Photosystem II for regulation and protection.
723 *Proc Natl Acad Sci U S A* **113**, 12144-12149, doi:10.1073/pnas.1608862113 (2016).

724 58 Cormann, K. U., Möller, M. & Nowaczyk, M. M. Critical Assessment of Protein Cross-
725 Linking and Molecular Docking: An Updated Model for the Interaction Between
726 Photosystem II and Psb27. *Front Plant Sci* **7**, 157, doi:10.3389/fpls.2016.00157 (2016).

727 59 Liu, H., Huang, R. Y., Chen, J., Gross, M. L. & Pakrasi, H. B. Psb27, a transiently
728 associated protein, binds to the chlorophyll binding protein CP43 in photosystem II
729 assembly intermediates. *Proc Natl Acad Sci U S A* **108**, 18536-18541,
730 doi:10.1073/pnas.1111597108 (2011).

731 60 Wei, L. *et al.* LPA19, a Psb27 homolog in *Arabidopsis thaliana*, facilitates D1 protein
732 precursor processing during PSII biogenesis. *Journal of Biological Chemistry* **285**,
733 21391-21398 (2010).

734 61 Wei, X. *et al.* Structure of spinach photosystem II-LHCII supercomplex at 3.2 Å
735 resolution. *Nature* **534**, 69-74, doi:10.1038/nature18020 (2016).

736 62 Cormann, K. U. *et al.* Structure of Psb27 in solution: implications for transient binding
737 to photosystem II during biogenesis and repair. *Biochemistry* **48**, 8768-8770,
738 doi:10.1021/bi9012726 (2009).

739 63 Fagerlund, R. D. & Eaton-Rye, J. J. The lipoproteins of cyanobacterial photosystem II.
740 *J Photochem Photobiol B* **104**, 191-203, doi:10.1016/j.jphotobiol.2011.01.022 (2011).

741 64 Liu, H. *et al.* Mass spectrometry-based footprinting reveals structural dynamics of loop
742 E of the chlorophyll-binding protein CP43 during photosystem II assembly in the
743 cyanobacterium *Synechocystis* 6803. *J Biol Chem* **288**, 14212-14220,
744 doi:10.1074/jbc.M113.467613 (2013).

745 65 Michoux, F. *et al.* Crystal structure of the Psb27 assembly factor at 1.6 Å: implications
746 for binding to Photosystem II. *Photosynth Res* **110**, 169-175, doi:10.1007/s11120-011-
747 9712-7 (2012).

748 66 Weisz, D. A., Gross, M. L. & Pakrasi, H. B. The Use of Advanced Mass Spectrometry
749 to Dissect the Life-Cycle of Photosystem II. *Front Plant Sci* **7**, 617,
750 doi:10.3389/fpls.2016.00617 (2016).

751 67 Kettunen, R., Tyystjärvi, E. & Aro, E. M. Degradation pattern of photosystem II
752 reaction center protein D1 in intact leaves. The major photoinhibition-induced cleavage
753 site in D1 polypeptide is located amino terminally of the DE loop. *Plant Physiol* **111**,
754 1183-1190, doi:10.1104/pp.111.4.1183 (1996).

755 68 Mulo, P., Laakso, S., Maenpää, P. & Aro, E. M. Stepwise photoinhibition of
756 photosystem II. Studies with *Synechocystis* species PCC 6803 mutants with a modified
757 D-E loop of the reaction center polypeptide D1. *Plant Physiol* **117**, 483-490,
758 doi:10.1104/pp.117.2.483 (1998).

759 69 Stowell, M. H. *et al.* Light-induced structural changes in photosynthetic reaction center:
760 implications for mechanism of electron-proton transfer. *Science* **276**, 812-816,
761 doi:10.1126/science.276.5313.812 (1997).

762 70 Wang, X. *et al.* Is bicarbonate in Photosystem II the equivalent of the glutamate ligand
763 to the iron atom in bacterial reaction centers? *Biochim Biophys Acta* **1100**, 1-8,
764 doi:10.1016/0005-2728(92)90119-m (1992).

765 71 Cheap, H. *et al.* M234Glu is a component of the proton sponge in the reaction center
766 from photosynthetic bacteria. *Biochim Biophys Acta* **1787**, 1505-1515,
767 doi:10.1016/j.bbabi.2009.07.004 (2009).

768 72 Burnap, R. L. D1 protein processing and Mn cluster assembly in light of the emerging
769 Photosystem II structure. *Physical Chemistry Chemical Physics* **6**, 4803-4809 (2004).

770 73 Tokano, T., Kato, Y., Sugiyama, S., Uchihashi, T. & Noguchi, T. Structural dynamics
771 of a protein domain relevant to the water-oxidizing complex in photosystem II as

visualized by high-speed atomic force microscopy. *The Journal of Physical Chemistry B* **124**, 5847-5857 (2020).

74 Avramov, A. P., Hwang, H. J. & Burnap, R. L. The role of Ca²⁺ and protein scaffolding in the formation of nature's water oxidizing complex. *Proceedings of the National Academy of Sciences* **117**, 28036-28045 (2020).

75 Zhang, M. *et al.* Structural insights into the light-driven auto-assembly process of the water-oxidizing Mn₄CaO₅-cluster in photosystem II. *Elife* **6**, doi:10.7554/eLife.26933 (2017).

76 Gisriel, C. J. *et al.* Cryo-EM Structure of Monomeric Photosystem II from *Synechocystis* sp. PCC 6803 Lacking the Water-Oxidation Complex. *Joule* (2020).

77 Kolling, D. R., Cox, N., Ananyev, G. M., Pace, R. J. & Dismukes, G. C. What are the oxidation states of manganese required to catalyze photosynthetic water oxidation? *Biophys J* **103**, 313-322, doi:10.1016/j.bpj.2012.05.031 (2012).

78 Zaltsman, L., Ananyev, G. M., Bruntrager, E. & Dismukes, G. C. Quantitative kinetic model for photoassembly of the photosynthetic water oxidase from its inorganic constituents: requirements for manganese and calcium in the kinetically resolved steps. *Biochemistry* **36**, 8914-8922, doi:10.1021/bi970187f (1997).

79 Stengel, A. *et al.* Initial steps of photosystem II de novo assembly and preloading with manganese take place in biogenesis centers in *Synechocystis*. *The Plant Cell* **24**, 660-675 (2012).

80 Tyryshkin, A. M. *et al.* Spectroscopic evidence for Ca²⁺ involvement in the assembly of the Mn₄Ca cluster in the photosynthetic water-oxidizing complex. *Biochemistry* **45**, 12876-12889, doi:10.1021/bi061495t (2006).

81 Campbell, K. A. *et al.* Dual-mode EPR detects the initial intermediate in photoassembly of the photosystem II Mn cluster: the influence of amino acid residue 170 of the D1 polypeptide on Mn coordination. *Journal of the American Chemical Society* **122**, 3754-3761 (2000).

82 Cohen, R. O., Nixon, P. J. & Diner, B. A. Participation of the C-terminal region of the D1-polypeptide in the first steps in the assembly of the Mn₄Ca cluster of photosystem II. *J Biol Chem* **282**, 7209-7218, doi:10.1074/jbc.M606255200 (2007).

83 Kuhl, H. *et al.* Towards structural determination of the water-splitting enzyme. Purification, crystallization, and preliminary crystallographic studies of photosystem II from a thermophilic cyanobacterium. *J Biol Chem* **275**, 20652-20659, doi:10.1074/jbc.M001321200 (2000).

84 Iwai, M., Katoh, H., Katayama, M. & Ikeuchi, M. Improved genetic transformation of the thermophilic cyanobacterium, *Thermosynechococcus elongatus* BP-1. *Plant Cell Physiol* **45**, 171-175, doi:10.1093/pcp/pch015 (2004).

85 Mastronarde, D. N. Automated electron microscope tomography using robust prediction of specimen movements. *Journal of structural biology* **152**, 36-51 (2005).

86 Biyani, N. *et al.* Focus: The interface between data collection and data processing in cryo-EM. *J Struct Biol* **198**, 124-133, doi:10.1016/j.jsb.2017.03.007 (2017).

87 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193 (2017).

88 Zhang, K. Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12, doi:10.1016/j.jsb.2015.11.003 (2016).

89 Scheres, S. H. A Bayesian view on cryo-EM structure determination. *J Mol Biol* **415**, 406-418, doi:10.1016/j.jmb.2011.11.010 (2012).

90 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296, doi:10.1038/nmeth.4169 (2017).

823 91 Scheres, S. H. Beam-induced motion correction for sub-megadalton cryo-EM particles.
824 *Elife* **3**, e03665, doi:10.7554/eLife.03665 (2014).

825 92 Tan, Y. Z. *et al.* Addressing preferred specimen orientation in single-particle cryo-EM
826 through tilting. *Nat Methods* **14**, 793-796, doi:10.1038/nmeth.4347 (2017).

827 93 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research
828 and analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

829 94 Bialek, W. *et al.* Crystal structure of the Psb28 accessory factor of
830 *Thermosynechococcus elongatus* photosystem II at 2.3 Å. *Photosynth Res* **117**, 375-
831 383, doi:10.1007/s11120-013-9939-6 (2013).

832 95 Michoux, F. *et al.* Crystal structure of CyanoQ from the thermophilic cyanobacterium
833 *Thermosynechococcus elongatus* and detection in isolated photosystem II complexes.
834 *Photosynth Res* **122**, 57-67, doi:10.1007/s11120-014-0010-z (2014).

835 96 Schwede, T., Kopp, J., Guex, N. & Peitsch, M. C. SWISS-MODEL: An automated
836 protein homology-modeling server. *Nucleic Acids Res* **31**, 3381-3385,
837 doi:10.1093/nar/gkg520 (2003).

838 97 Wu, S. & Zhang, Y. LOMETS: a local meta-threading-server for protein structure
839 prediction. *Nucleic Acids Res* **35**, 3375-3382, doi:10.1093/nar/gkm251 (2007).

840 98 Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics Toolkit with
841 a New HHpred Server at its Core. *J Mol Biol* **430**, 2237-2243,
842 doi:10.1016/j.jmb.2017.12.007 (2018).

843 99 Dobson, L., Remenyi, I. & Tusnady, G. E. CCTOP: a Consensus Constrained TOPology
844 prediction web server. *Nucleic Acids Res* **43**, W408-412, doi:10.1093/nar/gkv451
845 (2015).

846 100 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of
847 Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501,
848 doi:10.1107/S0907444910007493 (2010).

849 101 Trabuco, L. G., Villa, E., Schreiner, E., Harrison, C. B. & Schulten, K. Molecular
850 dynamics flexible fitting: a practical guide to combine cryo-electron microscopy and X-
851 ray crystallography. *Methods* **49**, 174-180, doi:10.1016/j.ymeth.2009.04.005 (2009).

852 102 Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol*
853 *Graph* **14**, 33-38, doi:10.1016/0263-7855(96)00018-5 (1996).

854 103 Ribeiro, J. V. *et al.* QwikMD - Integrative Molecular Dynamics Toolkit for Novices and
855 Experts. *Sci Rep* **6**, 26536, doi:10.1038/srep26536 (2016).

856 104 Phillips, J. C. *et al.* Scalable molecular dynamics with NAMD. *J Comput Chem* **26**,
857 1781-1802, doi:10.1002/jcc.20289 (2005).

858 105 Leaver-Fay, A. *et al.* ROSETTA3: an object-oriented software suite for the simulation
859 and design of macromolecules. *Methods Enzymol* **487**, 545-574, doi:10.1016/B978-0-
860 12-381270-4.00019-6 (2011).

861 106 Lindert, S. & McCammon, J. A. Improved cryoEM-Guided Iterative Molecular
862 Dynamics--Rosetta Protein Structure Refinement Protocol for High Precision Protein
863 Structure Prediction. *J Chem Theory Comput* **11**, 1337-1346, doi:10.1021/ct500995d
864 (2015).

865 107 Guo, Q. *et al.* In Situ Structure of Neuronal C9orf72 Poly-GA Aggregates Reveals
866 Proteasome Recruitment. *Cell* **172**, 696-705 e612, doi:10.1016/j.cell.2017.12.030
867 (2018).

868 108 Wehmer, M. *et al.* Structural insights into the functional cycle of the ATPase module of
869 the 26S proteasome. *Proc Natl Acad Sci U S A* **114**, 1305-1310,
870 doi:10.1073/pnas.1621129114 (2017).

871 109 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons
872 and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-
873 877, doi:10.1107/S2059798319011471 (2019).

Fig. 1: Cryo-EM map of a PSII assembly intermediate (PSII-I) from *T. elongatus*, segmented by subunit. a, 15 PSII subunits and 3 assembly factors are colored and named (PSII subunits: D1, D2, CP43, CP47, PsbE, PsbF, PsbH, PsbI, PsbK, PsbL, PsbM, PsbT, PsbX, PsbZ and Psb30; assembly factors: Psb27, Psb28 and *tsl0063*, which we named Psb34) (front view). **b,** Parts of PSII that originate from the CP43 module (comprised of CP43, Psb27, PsbZ, Psb30 and PsbK) and the RC47 complex are indicated by dashed lines (back view). **c,** Schematic model of the PSII assembly process starting with the formation of PSII-I from the CP43 module and RC47. Small PSII subunits were omitted for simplicity.

Fig. 2: Psb34 binds to RC47 during attachment of the CP43 module. **a**, Binding site of Psb34 at CP47, close to PsbH (top view), with extended binding of the Psb34 N-terminus along the cytoplasmic PSII surface (dashed box). **b**, MALDI-ToF analysis of PSII assembly intermediates. Mass spectrum of Psb34 (tsl0063) from the PSII complex (inset) and the fragment spectrum obtained for m/z 5936.356 with annotated b- and y-ion series matching the Psb34 sequence. Observed fragmentation sites are indicated by dashes in the sequence. Mascot score: 171. **c**, Subunit composition of Psb34-PSII assembly intermediates analyzed by 2D-PAGE.

Fig. 3: The role of the CP47 C-terminus in binding of Psb28. **a**, Binding of Psb28 at the cytoplasmic/stromal PSII surface (side view, colors correspond to Fig. 1) and continuation of the central Psb28 beta-sheet by the CP47 C-terminus and the D-E loop of D1 (dashed box). For comparison, mature monomeric PSII (PDB-ID 3KZI) is shown in gray. **b**, Superimposed 2D ^1H - ^{15}N -HSQC spectra of free Psb28 (blue) and Psb28 bound to the C-terminal peptide of CP47 (magenta). Upper left inset: representation of slow exchange behavior for the proton amide resonance of T24, ranging from 126.9 ppm to 128.6 ppm in the ^{15}N dimension. **c**, CSPs of more than one SD projected onto the model representation of Psb28. **d**, Weighted $^1\text{H}/^{15}\text{N}$ chemical shift perturbations observed for

Psb28 upon binding to the CP47 peptide. Red line indicates one standard deviation (SD), residues that yield resonances only in the complex form are indicated in orange. **e**, Backbone ^{15}N $\{^1\text{H}\}$ -heteronuclear NOE of free Psb28 (*blue*) and Psb28 bound to the C-terminal region of the CP47 peptide (*magenta*). Smaller I/I_0 ratios correspond to regions that exhibit dynamics on the pico- to nanosecond timescale.

Fig. 4. Structural changes of the D1 and D2 D-E loops induced by binding of Psb28 and Psb34. **a**, Side view of the CP43 antenna protein in PSII-I (*teal*) and the PSII-M control (*light blue*). **b**, Structural changes between PSII-I and the PSII-M control in the cytoplasmic D2 D-E loop (*yellow*: PSII-I, *blue*: PSII-M) and attachment of CP43 (*teal*: PSII-I, *light blue*: PSII-M control) (top view). Details of the structural changes in the D2 loop are shown in Fig. S5A and B. **c**, Side view and **d**, top view of the PSII-I structure (*orange*) compared to the PSII-M control (*light blue*) and mature monomeric PSII (*light red*, PDB-ID 3KZI). **e**, Side view and **f**, top view of the Psb28-induced structural changes in the D1 D-E loop (*orange*) and perturbation of the Q_B binding site compared to PSII-M (*light blue*), which lacks the assembly factors. Q_A is shown in *yellow* (PSII-I) or *light blue* (PSII-M), respectively. See Fig. S5c-h for enlarged views of the Q_A and Q_B binding site and the adjacent non-heme iron.

Fig. 5: Binding of Psb28 displaces bicarbonate as a ligand of the non-heme iron and protects PSII from damage. **a**, The electron transfer from PQ_A to PQ_B is coordinated by the non-heme iron (Fe^{2+}), with the binding of bicarbonate (Bic) serving as a regulatory mechanism⁵⁷ in mature PSII (PDB-ID 3WU2). **b**, Binding of Psb28 to the PSII-I assembly intermediate induces a conformational change in the cytoplasmic D2 D-E loop, where the side

chain of Glu241 replaces bicarbonate as a ligand of the non-heme iron. The respective fits of the non-heme iron binding sites are shown in Fig. S5E and F. A similar coordination is also found in non-oxygenic bacterial reaction centers⁶⁹ (Fig. S6C). **c**, Electron transfer (purple arrows) in mature PSII. Light-induced charge separation at the reaction center chlorophylls (P_{D1} , P_{D2} , Chl_{D1} , Chl_{D2}) leads to electron transfer via pheophytin ($Pheo_{D1}$) and plastoquinone A (Q_A) towards Q_B . The electron gap at the reaction center is filled by the oxygen evolving complex (OEC). **d**, Reoxidation of Q_A^- by direct and safe charge recombination is favored in the PSII assembly intermediate, as indicated by the purple arrows. **e**, Flash-induced fluorescence decay of PSII. Blue lines represent active PSII and red lines correspond to PSII-I. Black and grey lines represent PSII control samples without a functional OEC (Apo-PSII: hydroxylamine treated PSII; PSII (-OUV): extrinsic proteins removed). Dotted corridors depict SD ($n = 3$). **f**, The protective role of Psb28 binding was further confirmed by EPR spectroscopy using the spin probe TEMPd, which is specific for 1O_2 , the major reactive oxygen species in PSII generated by triplet chlorophyll (3P).

Fig. 6: The role of Psb27 in Mn_4CaO_5 cluster assembly. **a**, Bottom view of the luminal PSII surface for PSII-I (orange), the PSII-M control (light blue) and mature monomeric PSII (PDB-ID 3KZI) (light red). **b**, Side view of CP43 (teal) and Psb27 (brown) in PSII-I, as well as of CP43 (blue) and PsbO (purple) in mature monomeric PSII (PDB-ID 3KZI). Dashed box: CP43 E loop with residues Arg345 and Glu342 (shown as sticks), which form the second coordination sphere of the Mn_4CaO_5 cluster. We changed the numbering of CP43 residues due to a corrected N-terminal sequence (www.UniProt.org). The residues correspond to Arg357 and Glu354 in previous publications. The high-resolution structure of the Mn_4CaO_5 cluster is taken from Umena et al. 2011 (PDB-ID 3WU2). **c**, Position of the D1 C-terminus in PSII-I (orange) and mature monomeric PSII (PDB-ID 3KZI) (light red).

Figure 7: Conformational changes within the active site of the Mn_4CaO_5 cluster. The Mn_4CaO_5 cluster performs PSII's unique water-splitting reaction. **a**, The active site of the Mn_4CaO_5 cluster is resolved within our PSII-I structural model but is not yet oxygen-evolving. **b**, Crystal structure of the oxygen-evolving, mature PSII (PDB-ID 3WU2, resolution 1.9 Å). **c**, Overlay of both structures, illustrating significant differences in the backbone conformation of the D1 and D2 C-terminal tails. **d**, Accompanying side chain rearrangements of the D1 C-terminus. The Cl^- (**e**), Ion^+ (**f**) and Mn_4CaO_5 (**g**) cluster coordination partners are compared in detailed. The validation of the fit to density for the structural details shown here is provided in Figure S7.