Physical activity and aerobic fitness in children with inherited cardiac diseases

Aymeric Boisson, Gregoire de La Villeon, Helena Huguet, Hamouda Abassi, Jean Luc Pasquié, Kathleen Lavastre, Stefan Matecki, Sophie Guillaumont, Anne Requirand, Johanna Calderon, et al.

► To cite this version:

Aymeric Boisson, Gregoire de La Villeon, Helena Huguet, Hamouda Abassi, Jean Luc Pasquié, et al.. Physical activity and aerobic fitness in children with inherited cardiac diseases. Archives of cardiovascular diseases, 2021, 10.1016/j.acvd.2021.07.004 . hal-03411440

HAL Id: hal-03411440

<https://hal.science/hal-03411440>

Submitted on 2 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Physical activity and aerobic fitness in children with inherited cardiac diseases[☆]

Activité physique chez l'enfant atteint de maladie cardiaque héréditaire

Aymeric Boisson^a, Gregoire De La Villeon^{a,b},
Helena Huguet^c, Hamouda Abassi^{a,d},
Jean-Luc Pasquie^{a,d}, Kathleen Lavastre^a,
Stefan Matecki^{d,e}, Sophie Guillaumont^{a,b},
Anne Requirand^a, Johanna Calderon^{a,d},
Pascal Amedro^{f,g,*}

^a Paediatric and Congenital Cardiology Department, M3C Regional Reference Centre, University Hospital, 34295 Montpellier, France

^b Paediatric Cardiology and Rehabilitation Unit, Saint-Pierre Institute, 34250 Palavas-Les-Flots, France

^c Clinical Research and Epidemiology Unit, CHU de Montpellier, University of Montpellier, 34295 Montpellier, France

^d Inserm, PhyMedExp, CNRS, University of Montpellier, 34295 Montpellier, France

^e Department of Physiology, CHU de Montpellier, 34295 Montpellier, France

^f Department of Paediatric and Adult Congenital Cardiology, M3C National Reference Centre, Bordeaux University Hospital, 33604 Pessac, France

^g Inserm, U1045, IHU Liryc, Bordeaux Cardio-Thoracic Research Centre, Electrophysiology and Heart Modelling Institute, University of Bordeaux, Fondation Bordeaux Université, 33604 Pessac, France

KEYWORDS

Physical activity;
Inherited cardiac
arrhythmia;

Summary

Background. — Because of sports and exercise restrictions, children with inherited cardiac disease are at risk of physical deconditioning. Guidelines on sports participation in cardiovascular disease have become less restrictive over time, but their real-life application and behavioural impact have seldom been evaluated in children.

Abbreviations: CHD, congenital heart disease; CPET, cardiopulmonary exercise test; ESC, European Society of Cardiology; RER, respiratory exchange ratio; VO₂, oxygen uptake; WHO, World Health Organization.

[☆] Tweet: currently, most children with inherited cardiac arrhythmia or inherited cardiomyopathy have a good aerobic physical fitness and comply with the 2020 European Society of Cardiology guidelines on sports cardiology and exercise in cardiovascular disease.

* Corresponding author.

E-mail address: pascal.amedro@chu-bordeaux.fr (P. Amedro).

Aims. — We aimed to evaluate adherence to the 2020 European Society of Cardiology guidelines on sports and exercise in children with inherited cardiac arrhythmia and inherited cardiomyopathy; we also sought to evaluate their aerobic fitness, and the behavioural impact of inherited cardiac diseases on physical activity in children.

Methods. — Children aged 6–18 years with inherited cardiomyopathy or inherited cardiac arrhythmia were eligible for this cross-sectional study. Clinical, demographic and qualitative data were analysed.

Results. — A total of 32 children were included in the study (mean age 12.7 ± 3.5 years). Most children (81.3%) complied with the 2020 European Society of Cardiology guidelines; they were physically active and had good overall aerobic fitness, with a mean peak oxygen uptake (VO_2) value of $36.5 \pm 8.0 \text{ mL/kg/min}$ ($84.0 \pm 17.2\%$ of theoretical value). As a result of personal or parental behaviour, some children at risk of sudden cardiac death did not comply with the recommended upper limit of physical activity intensity, whereas others at low risk did not comply with the lower limit.

Conclusion. — Most children with inherited cardiac arrhythmia or inherited cardiomyopathy complied with current 2020 European Society of Cardiology guidelines on sports cardiology and exercise in cardiovascular disease.

MOTS CLÉS

Activité physique ;
Arythmie cardiaque
héréditaire ;
Cardiomyopathie
d'origine génétique ;
Pédiatrie ;
Recommandations

Résumé

Contexte. — En raison des restrictions sur l'activité physique, les enfants souffrant de cardiopathies héréditaires sont à risque de déconditionnement physique. Les recommandations sur la pratique sportive dans les cardiopathies sont plus permissives avec le temps, mais leurs applications dans la vie réelle et leurs impacts comportementaux ont été peu évalués chez l'enfant.

Objectifs. — Évaluer l'adhésion aux recommandations européennes (ESC 2020) sur l'activité physique et sportive chez l'enfant avec arythmie cardiaque héréditaire ou cardiomyopathie d'origine génétique, ainsi que leur aptitude physique aérobie et l'impact comportemental des cardiopathies héréditaires sur l'activité physique chez l'enfant.

Méthodes. — Les enfants âgés de 6 à 18 ans ayant une maladie rythmique héréditaire ou une cardiomyopathie d'origine génétique étaient éligibles pour cette étude. Les données cliniques, démographiques et qualitatives ont été analysées.

Résultats. — Trente-deux enfants ont été inclus dans l'étude (âge moyen $12,7 \pm 3,5$ ans). La plupart des enfants (81,3 %) suivaient les recommandations ESC 2020. Ils avaient une bonne condition aérobie avec une valeur moyenne de pic de VO_2 de $36,5 \pm 8,0 \text{ mL/kg/min}$ ($84,0 \pm 17,2\%$ des valeurs théoriques). Certains enfants à haut risque de mort subite ne respectaient pas la limite supérieure d'intensité recommandée, tandis que certains à bas risque ne respectaient pas la limite inférieure.

Conclusion. — La majorité des enfants ayant une maladie rythmique héréditaire ou une cardiomyopathie d'origine génétique respectent les recommandations 2020 de l'ESC.

Background

Physical activity is known for its important benefits in terms of physical and mental health [1,2]. Based on the World Health Organization (WHO) guidelines, children and adolescents aged 5–17 years should undertake at least 60 minutes of moderate to vigorous physical activity daily [3]. Indeed, repeated physical activity and exercise help to prevent obesity, the prevalence of which has increased dramatically in the young [4]. A sedentary lifestyle and its consequences in

the paediatric population currently represent a major public health issue, especially in children with pre-existing cardiac conditions [5–7].

Traditionally, congenital cardiac disorders are divided into three groups: congenital heart disease (CHD) for structural cardiac anomalies; inherited cardiac arrhythmias; and inherited cardiomyopathies. In the CHD population, early onset of physical deconditioning during childhood has been reported [8], even after successful cardiac surgical repair, and despite the absence of any sports restriction [9].

Parental and social barriers to physical activity in young people with CHD have been observed [10,11], resulting in a rapid decrease in aerobic fitness and a lower health-related quality of life [12,13]. Therefore, most children with CHD are currently encouraged to follow the WHO guidelines on physical activity or, for severe conditions, to undergo cardiac rehabilitation programmes [14].

Data on the physical activity of children with inherited cardiac arrhythmia and inherited cardiomyopathy are scarce. Moreover, the risk of sudden cardiac death during exercise in these two specific groups often makes the balance between a precautionary approach and the promotion of physical activity difficult [15–19]. Nevertheless, the guidelines on sports participation in inherited cardiac arrhythmia or inherited cardiomyopathy have become progressively less restrictive over time [20–27]. In 2020, guidelines on sports participation and exercise in patients with cardiovascular disease from the European Society of Cardiology (ESC) supported the concept of shared decision making between patients, families and physicians in the promotion of physical activity in patients with inherited cardiac disorders [28].

However, these guidelines are mostly dedicated to adult patients, and their application in the paediatric population has not been evaluated. Importantly, few data are currently available on actual physical activity in children with inherited cardiac arrhythmia or inherited cardiomyopathy, in terms of both intensity capacity and aerobic fitness.

The primary aim of this study was to assess the level of adherence to the 2020 ESC guidelines on sports participation and exercise in children with inherited cardiomyopathy or inherited cardiac arrhythmia; we also sought to evaluate their aerobic fitness, and the behavioural impact of the disease on physical activity in daily life.

Methods

Study design and population

This observational cross-sectional study was carried out from May to December 2020 in two paediatric cardiology tertiary care hospitals, labeled as rare cardiac disease expert centres by national health authorities www.filiere-cardiogen.fr: the Paediatric and Congenital Cardiology Department of Montpellier University Hospital; and the Paediatric Cardiology and Rehabilitation Unit of the St-Pierre Institute in Palavas-Les-Flots, France. Patients were screened consecutively during their annual follow-up. All children aged from 6 to 18 years with inherited cardiomyopathy (e.g. hypertrophic cardiomyopathy, dilated cardiomyopathy or arrhythmogenic right ventricular cardiomyopathy) or inherited cardiac arrhythmia (e.g. long QT syndrome, Brugada syndrome or catecholaminergic polymorphic ventricular tachycardia), who underwent their routine paediatric follow-up during the study period, were eligible to participate in the study. Children with severe neurocognitive impairments, preventing them from understanding and/or answering the questions during the interview, were not included in the study. Children were ineligible if they had absolute contraindications for cardiopulmonary exercise test: fever; uncontrolled

asthma; respiratory failure; acute myocarditis or pericarditis; uncontrolled arrhythmia; uncontrolled heart failure; acute pulmonary embolus or pulmonary infarction; and mental impairment leading to inability to cooperate.

The study complies with the declaration of Helsinki. Ethical approval was granted from Montpellier ethics committee (IRB-MPT_2020_03_202000385) before any data were collected. The study is registered on ClinicalTrials.gov (Identifier: NCT04650009). Informed consent was required from all parents or legal guardians, as well as the children's assent.

Physical activity outcomes

Patients' exercise intensity levels were categorized as low, medium or high, according to the 2020 ESC guidelines on sports participation and exercise in patients with cardiovascular disease [28]. Children who practiced less than 60 minutes per day of moderate-to-intense physical activity were considered to be physically inactive, as defined by the WHO guidelines [3]. The physical activity profile of each patient was assessed using the physical activity self-questionnaire from Ricci & Gagnon [29], and was categorized as inactive (total score < 18), active (total score $\geq$ 18 and $\leq$ 35) or very active (total score > 35). This questionnaire has three summary scores: sedentary behaviour (one item); recreational physical activities (four items); and daily physical activities (four items). Each item uses a 5-point scale from 1 to 5.

All interviewed children completed the questionnaires under the supervision of a paediatric cardiology research fellow (A. B.). The interview of the child and at least one parent or legal guardian lasted approximately 30 minutes, and was conducted in a quiet room, immediately after the annual paediatric cardiology consultation. If needed, the interview could also be conducted by videoconference. The clinical research team drafted a semistructured interview guide, with open-ended and closed-ended questions on physical activity (Table 1). The qualitative part of the interview focused on children's experiences, motivations and feelings regarding physical activity in the context of their cardiac condition. Once the child had answered the questionnaire, the parents were asked to answer the same questions.

Aerobic fitness outcomes

To assess aerobic fitness, children underwent a cardiopulmonary exercise test (CPET), using the same methodology as in our previous paediatric CPET studies [8,30,31]. The following variables were measured: maximum oxygen uptake ($\text{VO}_{2\text{max}}$); ventilator anaerobic threshold; ventilatory efficiency (VE/VCO_2 slope); oxygen uptake efficiency slope; maximum load; peak heart rate; and peak respiratory exchange ratio (RER). Peak VO_2 , ventilator anaerobic threshold and peak heart rate were normalized as percentage of predicted peak using normal values from Wasserman and Cooper [1,32,33]. The two investigating centres used the same CPET cycle ergometer protocol, in order to obtain a homogeneous incremental overall duration of between 8 and 12 minutes: a 1-minute rest; a 3-minute warm-up (10–20 watts) in increments of 10, 15 or 20 watts each minute, depending on the patient; a pedalling rate of 60–80

Table 1 Physical activity interview guide.

Do you practice any sports, exercise or physical activity? If yes, name them
Do you practice competitive sports? ^a
Do you know if exercise limitations exist in your cardiac condition? ^a
How did you react when you discovered that you were limited in your physical activity?
Have you ever been in a situation where you would have liked to participate in a physical activity, but could not due to your cardiac condition? ^a
What did you feel in that moment?
Do you have extracurricular activities apart from physical activities?
If yes, describe your feeling and emotions regarding these activities
Do you have a medical sports certificate? Describe it if possible
Who gave you this medical sports certificate?
Do you have any current treatment for your cardiac condition?
What impact does your heart condition have on your physical activity?
Have you ever received advice or guidelines regarding therapeutic education during a consultation? (Drugs and medications to avoid, warm-up before physical activity, life hygiene)
Would you be interested in a dedicated consultation on this subject?
Have you ever experienced any cardiac symptoms related to your cardiac condition? ^a
If yes, what did you feel at that moment?
Can you describe your feelings and emotion related to physical activity?

^a Closed-ended questions.

revolutions per minute; a 3-minute active recovery (20 watts); and a 2-minute rest. The CPET was considered as maximal when the following criteria were reached: RER ($\text{VCO}_2/\text{VO}_2 \geq 1$; limit of the patient's tolerance despite verbal encouragement; plateau of VO_2 ($\text{VO}_{2\text{max}}$) despite increasing exercise intensity; and patient's inability to provide a minimum pedaling frequency of 60 revolutions per minute despite verbal encouragement. If the $\text{VO}_{2\text{max}}$ did not reach the criteria for maximal effort, the peak VO_2 was recorded, as is usual in paediatric CPET studies [34,35].

Clinical outcomes

The following clinical data were collected: demographic data (age, sex, time since diagnosis); anthropometric data; clinical cardiac diagnosis; age at diagnosis; time since diagnosis; circumstance of diagnosis; current treatment; presence of a cardiovascular event in the last 24 months; functional status (New York Heart Association classification); left ventricular ejection fraction; and presence and type of genetic mutation, as reported in medical records.

Quantitative and qualitative data analyses

Based on the interview data, patients were classified into two groups, according to level of adherence to the 2020 ESC guidelines on sports participation and exercise in patients with cardiovascular disease [28]: full adherence (e.g. strict compliance with the recommendations); or no adherence (e.g. practice of a non-recommended physical activity, in terms of type and/or intensity of exercise). Children from the "full-adherence" group were divided into three subgroups according to their level of maximum intensity exercise practice: (1) "maximum permitted level" group, if the child practiced exercise at the maximum intensity level authorized by the guidelines; (2) "lower level than

permitted" group, if the child practiced exercise one intensity level down compared with the maximum intensity level authorized; and (3) "much lower level than permitted" group, if the child practiced exercise more than one intensity level down compared with the maximum intensity level authorized. In order to provide a double independent assessment, a blinded classification of patients in groups and subgroups was carried out by a senior sports cardiologist, who did not participate in the interview (G. D. L. V.).

Emotions and discourse of children and their parents were classified after the interview into three groups: positive (including joy, well-being, optimism and resilience); neutral (if participants expressed neutral discourse); and negative (including fear, sadness, frustration, disappointment and anger). Field notes were made during the interview. The examiner's notes and transcribed information were included for analysis. All data were anonymized by the principal investigator. Data were processed according to the thematic analysis method [36], and were collected in Excel spreadsheets.

Statistical analyses

The study population is described using means and standard deviations for quantitative variables and frequencies for qualitative variables. Continuous variable distributions were tested using the Shapiro–Wilk test. Quantitative variables were compared using Student's *t*-test when the distribution was Gaussian, and with the Mann–Whitney test otherwise. For qualitative variables, groups were compared using the Chi² test or Fisher's exact test. For comparisons according to intensity of adherence to ESC guidelines, Holm's correction was applied for pairwise comparisons. Statistical significance was set at 0.05, and analyses were performed using Statistical Analysis Systems Enterprise Guide, version 4.3 (SAS Institute, Cary, NC, USA).

Table 2 Population characteristics.

Inherited cardiac disease	
Long QT syndrome	18 (56.3)
Brugada syndrome	1 (3.1)
ARVC	2 (6.2)
Hypertrophic cardiomyopathy	6 (18.8)
Dilated cardiomyopathy	2 (6.2)
CPVT	3 (9.4)
Sex	
Male	21 (65.6)
Female	11 (34.4)
Age (years)	12.7 ± 3.5
Age at diagnosis (years)	6.6 ± 5.2
Body mass index (kg/m ²)	19.2 ± 3.9
Left ventricular ejection fraction (%)	67.3 ± 6.6
Cardiovascular event in the last 24 months ^a	3 (9.4)
Identification of a genetic mutation	29 (90.6)
Current treatment	
Beta-blockers	29 (90.6)
Implantable cardioverter defibrillator	3 (9.4)
Circumstance of diagnosis	
Familial genetic screening	17 (53.1)
Symptoms	14 (43.8)
Fortuitous diagnosis	1 (3.1)
Physical inactivity (WHO guidelines)	7 (21.9)

ARVC: arrhythmogenic right ventricular cardiomyopathy; CPVT: catecholaminergic polymorphic ventricular tachycardia; WHO: World Health Organization. Data are expressed as number (%) or mean ± standard deviation.

^a Cardiac arrhythmia, cardiac syncope, sudden cardiac arrest, chest pain at exertion.

Results

Patient characteristics

During the study period, 33 consecutive eligible patients were asked to participate after their regular consultation. After gaining consent from the parent or legal guardian, a total of 32 children were included in the study (mean age 12.7 ± 3.5 years; 65.6% male patients). One child and his family refused to participate because of a lack of time for the interview. Inclusions were distributed evenly between the two centres. Long QT syndrome represented the predominant cardiac disorder ($n=18$). Most of the population was treated with beta-blockers ($n=29$), whereas few had an implantable cardioverter defibrillator ($n=3$). Regarding the circumstances of diagnosis, distribution was equitable between those diagnosed through familial genetic screening and those diagnosed after presenting cardiac symptoms related to their disease. Fortuitous diagnosis was exceptional. Few ($n=3$) cardiac events during exercise were reported (e.g. non-severe moderate chest pain), and no sudden cardiac death or cardiac arrhythmia occurred in the last 24 months. No patients presented with heart failure symptoms at enrolment. Detailed population characteristics are presented in [Table 2](#).

Quantitative data on physical activity

Most children with inherited cardiac arrhythmia or inherited cardiomyopathy enrolled in the study ($n=26$, 81.3%)

complied with the current 2020 ESC guidelines on sports cardiology and exercise in patients with cardiovascular disease. From the physical activity self-questionnaire, the patients included in the study presented an active ($n=28$) or inactive ($n=4$) profile, with a mean overall score of 24.8 ± 5.3. In addition, in terms of amount of time being physically active, most enrolled children (78.1%) complied with the WHO guidelines and were therefore not considered to be physically inactive. Children who complied with the WHO guidelines had a significantly higher score on the physical activity self-questionnaire than physically inactive children (26.2 ± 4.6 vs. 19.7 ± 4.8; $P<0.01$) ([Table 3](#)).

In comparison with children complying with the 2020 ESC guidelines, children who did not follow them ($n=6$; mainly patients with catecholaminergic polymorphic ventricular tachycardia and arrhythmogenic right ventricular cardiomyopathy) showed a significantly lower maximum heart rate (125.6 ± 36.4 vs. 152.9 ± 24.0; $P=0.03$) and percentage of predicted heart rate (61.2 ± 16.9 vs. 73.5 ± 11.6; $P=0.03$) during exercise test. No other significant differences were observed between these two groups.

From the CPET results, aerobic fitness in children with inherited cardiomyopathy or inherited cardiac arrhythmia was good overall, with a mean peak VO_2 value of 36.5 ± 8.0 mL/kg/min (84.0 ± 17.2% of predicted peak VO_2), and no evidence of physical deconditioning in the ventilatory variables ([Table 3](#)).

Among patients who followed the 2020 ESC guidelines ($n=26$), 15 children practiced exercise at the maximum intensity level authorized, seven children practiced exercise

Table 3 Physical activity and aerobic fitness.

Variable	Total (n = 32)	2020 ESC guidelines			WHO guidelines		
		Adherence	No adherence	P	Adherence	No adherence	P
		(n = 26)	(n = 6)		(n = 25)	(n = 7)	
Physical activity level	24.8 ± 5.3	24.9 ± 5.4	24.5 ± 5.8	0.88	26.2 ± 4.6	19.7 ± 4.8	< 0.01 ^a
Peak VO ₂ (mL/kg/min)	36.5 ± 8.0	36.2 ± 7.1	38.0 ± 13.2	0.69	36.3 ± 8.5	37.5 ± 5.9	0.79
% predicted peak VO ₂	84.0 ± 17.2	84.1 ± 15.4	83.3 ± 28.3	0.93	84.3 ± 18.1	82.0 ± 14.1	0.81
VAT (mL/kg/min)	26.3 ± 6.8	26.5 ± 6.3	25.5 ± 10.2	0.80	26.1 ± 6.8	27.8 ± 7.7	0.66
% predicted VAT	60.6 ± 13.4	61.1 ± 12.2	57.5 ± 20.8	0.63	60.4 ± 12.8	61.5 ± 18.6	0.88
VE/VCO ₂ slope	30.8 ± 5.9	30.1 ± 5.8	35.3 ± 6.1	0.16	30.9 ± 6.2	30.3 ± 5.0	0.86
OUES	1427 ± 853	1415 ± 796	1549 ± 1766	0.84	1398 ± 844	1605 ± 1079	0.71
OUES/kg	32.8 ± 15.8	32.2 ± 14.7	38.3 ± 31.8	0.62	34.0 ± 16.4	25.5 ± 10.6	0.40
Peak RER	1.14 ± 0.10	1.14 ± 0.10	1.11 ± 0.08	0.56	1.14 ± 0.09	1.16 ± 0.15	0.43
Peak HR (beats/min)	147.8 ± 28.0	152.9 ± 24.0	125.6 ± 36.4	0.03 ^a	148.2 ± 28.2	146.0 ± 30.5	0.88
% predicted HR	71.2 ± 13.3	73.5 ± 11.6	61.2 ± 16.9	0.03 ^a	71.4 ± 13.1	70.4 ± 15.6	0.85
Maximum load (watts)	118.9 ± 57.7	119.6 ± 56.5	116.0 ± 69.7	0.85	112.5 ± 54.1	147.0 ± 70.9	0.16

ESC: European Society of Cardiology; HR: heart rate; OUES: oxygen uptake efficiency slope; peak VO₂: maximum oxygen uptake; RER: respiratory exchange ratio; VAT: ventilatory anaerobic threshold; VE/VCO₂ slope: ventilatory efficiency; WHO: World Health Organization. Data are expressed as mean ± standard deviation.

^a Significant *P*-value.

Table 4 Physical activity characteristics.

	Adherence to 2020 ESC guidelines			P
	Intensity at the maximum level permitted	Intensity at a lower level than permitted	Intensity at a much lower level than permitted	
	(n = 15)	(n = 7)	(n = 4)	
Physical activity level	27.3 ± 4.5	23.7 ± 3.5	17.8 ± 4.3	< 0.01 ^a
Peak VO ₂ (mL/kg/min)	35.7 ± 4.5	40.0 ± 8.4	31.8 ± 9.9	0.20
% predicted peak VO ₂	83.0 ± 12.8	93.3 ± 16.3	73.3 ± 16.2	0.12
VAT (mL/kg/min)	25.8 ± 4.3	29.8 ± 7.3	23.3 ± 8.8	0.25
% predicted VAT	60.5 ± 11.4	66.7 ± 8.2	54.8 ± 17.9	0.32
VE/VCO ₂ slope	30.9 ± 6.2	29.3 ± 7.1	29.0 ± 2.3	0.93
OUES	1493 ± 737	1466 ± 793	1156 ± 1104	0.67
OUES/kg	34.9 ± 13.4	36.8 ± 16.5	19.9 ± 11.5	0.17
Peak RER	1.15 ± 0.08	1.14 ± 0.15	1.14 ± 0.10	0.99
Peak HR (beats per minute)	147.2 ± 24.2	165.3 ± 21.3	151.3 ± 26.3	0.33
% predicted HR	70.8 ± 11.3	79.2 ± 10.6	73.0 ± 14.0	0.37
Maximum load (watts)	111.7 ± 48.9	119.2 ± 54.8	143.8 ± 86.7	0.83

ESC: European Society of Cardiology; HR: heart rate; OUES: oxygen uptake efficiency slope; peak VO₂: maximum oxygen uptake; RER: respiratory exchange ratio; VAT: ventilatory anaerobic threshold; VE/VCO₂ slope: ventilatory efficiency. Data are expressed as mean ± standard deviation.

^a Significant *P*-value.

at a lower level than authorized and four children practiced exercise at a much lower level than authorized. A total of 11 children (34.4%) complied with the 2020 ESC guidelines but could perform physical activity at a higher intensity level. The three subgroups of actual over authorized exercise levels were significantly different in terms of physical activity self-questionnaire scores (Table 4).

Qualitative data on physical activity

Detailed data on patients' physical activities are presented in Table 5. Half of the interviewed children (n = 16) expressed positive emotions regarding physical activity in their daily life; for them, exercise was perceived as a source of pleasure and a factor of good global health, despite their

Table 5 Patients' detailed physical activities.

	Skill	Power	Mixed	Endurance
Low intensity	Golf (<i>n</i> = 1)	Shot putting ^a	Soccer ^a (<i>n</i> = 10)	Jogging
	Table tennis (<i>n</i> = 2)	Discus ^a	Basketball ^a (<i>n</i> = 3)	Long-distance walking
	Shooting	Alpine skiing ^a (<i>n</i> = 1)	Handball ^a (<i>n</i> = 5)	Swimming ^a (<i>n</i> = 7)
Medium intensity	Sailing	Short-distance running (<i>n</i> = 3)	Volleyball (<i>n</i> = 1)	Cycling ^a (<i>n</i> = 10) Speed walking
	Yachting	Shot putting	Tennis (doubles)	Mid-/long-distance running
	Equestrian (<i>n</i> = 1)	Discus Alpine skiing Judo/karate (<i>n</i> = 2)		Gymnastics (<i>n</i> = 2)
High intensity		Weight lifting (<i>n</i> = 2)	Ice-hockey	Cycling (road)
		Wrestling	Hockey	Mid-/long-distance swimming
		Boxing (<i>n</i> = 3)	Climbing (<i>n</i> = 1)	Long-distance skating
			Rugby (<i>n</i> = 2)	Pentathlon
			Fencing	Athletics (<i>n</i> = 1)
			Tennis (singles) (<i>n</i> = 1)	Rowing
			Water polo	Canoeing
			Soccer (competition) (<i>n</i> = 1)	Cross-country skiing
			Basketball (competition)	Biathlon
			Handball (competition)	Triathlon

n indicates the physical activity for the patients enrolled in the study. Multiple choices per patient were possible.

^a Recreational activity with adapted intensity.

heart disease. These patients stressed the social aspect of physical activity, as it allowed them to interact with peers during activities, even if they reported having to adapt the intensity of exercise ("I can still play football with my friends, as a goalkeeper for example"). Resilience skills were frequently observed when analysing the discourse of those children ("it was difficult at first, but now that I understand why it's important, it does not bother me anymore"). One third of the interviewed children practiced non-exercise related activities, such as music lessons or circus arts; they all expressed strong positive feelings about those activities, and placed them at the same level as exercise and sports in terms of a source of wellbeing, especially for children with strict sports restrictions ("music is all my life now, I like it even more than sports").

Negative emotions were reported by 34% of children, who were mainly frustrated by limitations in terms of intensity of exercise, especially those who did not respect the ESC guidelines on sports cardiology and exercise in cardiovascular disease ("I can't exercise as much as I want to, I feel very limited"). However, a feeling of fear related to physical activity in the context of their cardiac condition was not reported by any of the interviewed children ("I'm not afraid of practicing sports").

When considering parents' reports, more than half (59%) reported negative emotions regarding physical activity in the context of their child's cardiac condition. Fear was the predominant emotion reported by parents ("I'm always in fear that something may happen when I'm not there"); they

mainly expressed the need to supervise their child when practicing a physical activity, out of fear of sudden cardiac death or another serious clinical event. Most reported having set physical activity limits for their child, even if the guidelines and their physician were not as restrictive ("with such limitations, I feel safer"). Only 25% of the interviewed parents reported positive emotions regarding their child's physical activity; according to them, controlled and supervised physical activity was still perceived as something beneficial for their children. Few children and parents (*n* = 5) expressed neutral emotions when reporting little to no impact on their daily life ("exercise and sports are not important to me, I am happy without it").

A small number of patients (*n* = 5) and parents (*n* = 1) were not aware of exercise and sports restriction with regard to their child's cardiac disease, although they all respected the 2020 ESC guidelines "by chance".

Most families had received medical advice related to contraindicated drugs and medications or life hygiene. Half of the patients and families expressed the wish to receive structured patient therapeutic education dedicated to those topics.

Discussion

This study has reported, for the first time, the actual degree of implementation of the recent 2020 ESC guidelines on sports cardiology and exercise in a population of children

with cardiovascular diseases. We found that a large majority of children with inherited cardiac arrhythmia or inherited cardiomyopathy complied with the 2020 ESC guidelines, and that only a minority was physically inactive. Furthermore, no serious cardiac events occurred in the last 24 months among enrolled children, which is consistent with recent studies suggesting that exercise-related events are rare in appropriately managed children with long QT syndrome and hypertrophic cardiomyopathy [37,38]. As this study was carried out concomitantly with the publication of the 2020 ESC guidelines, we may hypothesize that paediatric cardiologists had anticipated the recommendations by promoting physical activity at a safe level in children with cardiac conditions. Despite the absence of specific paediatric guidelines, these results are in line with the evolution of guidelines in sport cardiology and exercise in patients with cardiovascular disease in recent decades [20–27]. Nevertheless, our results contrast with recent public health data reporting that half of boys and more than two thirds of girls from the French general paediatric population did not follow the WHO guidelines regarding daily physical activity, and were considered to have a sedentary lifestyle [39]. In our study, analyses from the interviews suggest the existence of resilience skills in children with inherited cardiac disease, as classically described in many chronic diseases [40]. Added to changes in physician practice, this form of resilience may have contributed to promote physical activity in this population.

Of note, even if most children complied with the 2020 ESC guidelines on sports cardiology and exercise, those who did not follow them were among those at higher risk of sudden cardiac death during exercise. Indeed, all interviewed patients with catecholaminergic polymorphic ventricular tachycardia or arrhythmogenic right ventricular cardiomyopathy were found to practice high intensity exercise, even after their physicians advised them not to. This result may be explained by the absence of real physical disability during exercise, combined with strict exercise restrictions, in this group of patients. Yet, according to recent studies, approximately 30% of patients with catecholaminergic polymorphic ventricular tachycardia have a family history of sudden cardiac death before the age of 40 years [41,42], and the mortality rate rises to 50% in severely affected and untreated patients [43]. In our study, the interviews showed that children who did not respect the 2020 ESC guidelines were mainly frustrated by the imposed intensity limitations for physical activities and were aware of the risks. To avoid any serious cardiac events in this selected population, it is necessary to build structured paediatric therapeutic education programmes promoting safe and adapted physical activity. Moreover, many children reported that non-exercise-related activities provided as much well-being as sports used to, which suggests good resilience skills. Indeed, mental health care to support the anticipation and management of their condition should be integrated into educational therapeutic programmes for children with inherited cardiac disorders.

In our study, children with inherited cardiomyopathy and inherited cardiac arrhythmia had good overall aerobic fitness, with no evidence of physical deconditioning. Data from exercise tests corroborated the results from the physical activity questionnaires and interviews, and the enrolled children did practice exercise and sports-related

activities. No significant differences in aerobic fitness were reported when comparing children who complied with the 2020 ESC guidelines and those who did not, except for a lower maximum heart rate at exercise in children who did not follow the recommendations (i.e. those with catecholaminergic polymorphic ventricular tachycardia). Higher dosages of beta-blockers are used in these patients, which might explain this difference [42,44].

Conversely, almost half of children who followed the 2020 ESC guidelines practiced physical activity at a lower level of intensity than permitted. From the interviews, parental barriers to physical activity were predominant in this group, as previously reported in children with CHD [11]. Future patient education programmes in children with inherited cardiac disorders should involve parental psychological follow-up to support adaptive coping strategies with regard to their child's condition. Finally, these children with inherited cardiac disorders are at risk of physical deconditioning, impaired health-related quality of life and additional cardiovascular burden in adulthood, as observed in CHD [8,12,13,45]. Thus, there is an urgent need to develop and implement holistic rehabilitation and educational programmes for children with inherited cardiac disorders and their families.

Study limitations

This study was performed in tertiary care centres and may not represent the overall population of children with inherited cardiac disease. However, in France, children with these cardiac disorders are usually followed in rare disease expert centres labelled by national health authorities (www.filiere.cardiogen.fr).

These preliminary results are limited by the study sample size; in addition, data on physical activity came from interviews and questionnaires, and no fitness tracker was used. Moreover, given the number of tests performed, marginal statistically significant differences should be interpreted with caution. Nevertheless, this pilot study was necessary to draft the current on-going QUALIMYOTYM trial, a controlled multicentre prospective study on health-related quality of life, physical activity and aerobic fitness in children with inherited cardiac arrhythmia or inherited cardiomyopathy [46].

Conclusions

This observational cross-sectional study has highlighted that most children with inherited cardiac arrhythmia or inherited cardiomyopathy comply with the current 2020 ESC guidelines on sports cardiology and exercise in patients with cardiovascular disease. Our results showed that these children were physically active and had good aerobic fitness. Nevertheless, because of personal or parental behaviour, some children at risk of sudden cardiac death did not comply with the upper limit of physical intensity, whereas others at low risk did not comply with the lower limit.

Funding

This study was supported by the paediatric and congenital clinical research unit of Montpellier University Hospital and funded by a research grant from Cardiogen www.filiere-cardiogen.fr.

Acknowledgments

We sincerely thank the families for their generous participation.

Disclosure of interest

The authors declare that they have no competing interest.

References

- [1] Physical activity and cardiovascular health. NIH consensus development panel on physical activity and cardiovascular health. *JAMA* 1996;276:241–6.
- [2] Fletcher GF, Balady G, Blair SN, et al. Statement on exercise: benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the committee on exercise and cardiac rehabilitation of the council on clinical cardiology, American Heart Association. *Circulation* 1996;94:857–62.
- [3] Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020;54:1451–62.
- [4] Praznoczy C, Lambert C, Pascal C. État des lieux de l'activité physique et de la sédentarité en France – Édition. Observatoire national de l'activité physique et de la sédentarité – Rapport; 2017. p. 130 [Available at: <https://onaps.fr/actualisation-de-letat-des-lieux-de-lactivite-physique-et-de-la-sedentarite-en-france-edition-2017/>].
- [5] Karsenty C, Maury P, Blot-Souletie N, et al. The medical history of adults with complex congenital heart disease affects their social development and professional activity. *Arch Cardiovasc Dis* 2015;108:589–97.
- [6] Papasavvas T, Alhashemi M, Micklewright D. Association between depressive symptoms and exercise capacity in patients with heart disease: a meta-analysis. *J Cardiopulm Rehabil Prev* 2017;37:239–49.
- [7] Tamayo C, Manlhiot C, Patterson K, Lalani S, McCrindle BW. Longitudinal evaluation of the prevalence of overweight/obesity in children with congenital heart disease. *Can J Cardiol* 2015;31:117–23.
- [8] Amedro P, Gavotto A, Guillaumont S, et al. Cardiopulmonary fitness in children with congenital heart diseases versus healthy children. *Heart* 2018;104:1026–36.
- [9] Longmuir PE, Brothers JA, de Ferranti SD, et al. Promotion of physical activity for children and adults with congenital heart disease: a scientific statement from the American Heart Association. *Circulation* 2013;127:2147–59.
- [10] Berg AE, Meyers LL, Dent KM, Rothwell EW, Everitt MD. Psychological impact of sports restriction in asymptomatic adolescents with hypertrophic cardiomyopathy, dilated cardiomyopathy, and long QT syndrome. *Prog Pediatr Cardiol* 2018;49:57–62.
- [11] Moola F, Fusco C, Kirsh JA. The perceptions of caregivers toward physical activity and health in youth with congenital heart disease. *Qual Health Res* 2011;21:278–91.
- [12] Amedro P, Dorka R, Moniotte S, et al. Quality of life of children with congenital heart diseases: a multicenter controlled cross-sectional study. *Pediatr Cardiol* 2015;36:1588–601.
- [13] Amedro P, Picot MC, Moniotte S, et al. Correlation between cardio-pulmonary exercise test variables and health-related quality of life among children with congenital heart diseases. *Int J Cardiol* 2016;203:1052–60.
- [14] Amedro P, Gavotto A, Legendre A, et al. Impact of a centre and home-based cardiac rehabilitation program on the quality of life of teenagers and young adults with congenital heart disease: the QUALI-REHAB study rationale, design and methods. *Int J Cardiol* 2019;283:112–8.
- [15] Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults? *J Am Coll Cardiol* 2003;42:1959–63.
- [16] Emery MS, Kovacs RJ. Sudden cardiac death in athletes. *JACC Heart Fail* 2018;6:30–40.
- [17] Harmon KG, Asif IM, Maleszewski JJ, et al. Incidence, cause, and comparative frequency of sudden cardiac death in national collegiate athletic association athletes: a decade in review. *Circulation* 2015;132:10–9.
- [18] Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980–2006. *Circulation* 2009;119:1085–92.
- [19] Solberg EE, Gjertsen F, Haugstad E, Kolsrud L. Sudden death in sports among young adults in Norway. *Eur J Cardiovasc Prev Rehabil* 2010;17:337–41.
- [20] Ackerman MJ, Zipes DP, Kovacs RJ, Maron BJ. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 10: the cardiac channelopathies: a scientific statement from the American Heart Association and American College of Cardiology. *J Am Coll Cardiol* 2015;66:2424–8.
- [21] Hammond-Haley M, Patel RS, Providencia R, Lambiase PD. Exercise restrictions for patients with inherited cardiac conditions: current guidelines, challenges and limitations. *Int J Cardiol* 2016;209:234–41.
- [22] Maron BJ, Chaitman BR, Ackerman MJ, et al. Recommendations for physical activity and recreational sports participation for young patients with genetic cardiovascular diseases. *Circulation* 2004;109:2807–16.
- [23] Maron BJ, Udelson JE, Bonow RO, et al. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 3: hypertrophic cardiomyopathy, arrhythmogenic right ventricular cardiomyopathy and other cardiomyopathies, and myocarditis: a scientific statement from the American Heart Association and American College of Cardiology. *Circulation* 2015;132:e273–80.
- [24] Pelliccia A, Fagard R, Bjornstad HH, et al. Recommendations for competitive sports participation in athletes with cardiovascular disease: a consensus document from the Study Group of Sports Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005;26:1422–45.
- [25] Pelliccia A, Solberg EE, Papadakis M, et al. Recommendations for participation in competitive and leisure time sport in athletes with cardiomyopathies, myocarditis, and pericarditis: position statement of the Sport Cardiology Section of the European Association of Preventive Cardiology (EAPC). *Eur Heart J* 2019;40:19–33.
- [26] Priori SG, Blomstrom-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: the task force for the management of patients with ventricular

- arrhythmias and the prevention of sudden cardiac death of the European Society of Cardiology (ESC), endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J* 2015;36:2793–867.
- [27] Zipes DP, Ackerman MJ, Estes 3rd NA, Grant AO, Myerburg RJ, Van Hare G. Task force 7: arrhythmias. *J Am Coll Cardiol* 2005;45:1354–63.
 - [28] Pelliccia A, Sharma S, Gati S, et al. 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* 2021;42:17–96.
 - [29] Vuillemin A, Denis G, Guillemin F, Jeandel C. A review of evaluation questionnaires for physical activity. *Rev Epidemiol Sante Publique* 1998;46:49–55.
 - [30] Abassi H, Huguet H, Picot MC, et al. Health-related quality of life in children with congenital heart disease aged 5 to 7 years: a multicentre controlled cross-sectional study. *Health Qual Life Outcomes* 2020;18:366.
 - [31] Gavotto A, Vandenbergh D, Abassi H, et al. Oxygen uptake efficiency slope: a reliable surrogate parameter for exercise capacity in healthy and cardiac children? *Arch Dis Child* 2020;105:1167–74.
 - [32] Cooper DM, Berry C, Lamarra N, Wasserman K. Kinetics of oxygen uptake and heart rate at onset of exercise in children. *J Appl Physiol* 1985;59:211–7.
 - [33] Cooper DM, Weiler-Ravell D, Whipp BJ, Wasserman K. Aerobic parameters of exercise as a function of body size during growth in children. *J Appl Physiol Respir Environ Exerc Physiol* 1984;56:628–34.
 - [34] Barker AR, Williams CA, Jones AM, Armstrong N. Establishing maximal oxygen uptake in young people during a ramp cycle test to exhaustion. *Br J Sports Med* 2011;45:498–503.
 - [35] Rowland TW, Cunningham LN. Oxygen uptake plateau during maximal treadmill exercise in children. *Chest* 1992;101:485–9.
 - [36] Vaismoradi M, Turunen H, Bondas T. Content analysis and thematic analysis: implications for conducting a qualitative descriptive study. *Nurs Health Sci* 2013;15:398–405.
 - [37] Chambers KD, Beausejour Ladouceur V, Alexander ME, et al. Cardiac events during competitive, recreational, and daily activities in children and adolescents with long QT syndrome. *J Am Heart Assoc* 2017;6:e005445.
 - [38] Saarel EV, Law I, Berul CI, et al. Safety of sports for young patients with implantable cardioverter-defibrillators: long-term results of the multinational ICD sports registry. *Circ Arrhythm Electrophysiol* 2018;11:e006305.
 - [39] Verdot C, Salanave B, Deschamps V. Activité physique et sédentarité dans la population française. Situation en 2014–2016 et évolution depuis 2006–2007. *Bull Epidemiol Hebd* 2020;15:296–304.
 - [40] Cal SF, de Sá LR, Glustak ME, Santiago MB. Resilience in chronic diseases: a systematic review. *Cogent Psychol* 2015;2:1024928.
 - [41] Leenhardt A, Lucet V, Denjoy I, Grau F, Ngoc DD, Coumel P. Catecholaminergic polymorphic ventricular tachycardia in children. A 7-year follow-up of 21 patients. *Circulation* 1995;91:1512–9.
 - [42] Priori SG, Napolitano C, Memmi M, et al. Clinical and molecular characterization of patients with catecholaminergic polymorphic ventricular tachycardia. *Circulation* 2002;106:69–74.
 - [43] Priori SG, Wilde AA, Horie M, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm* 2013;10:1932–63.
 - [44] Hayashi M, Denjoy I, Extramiana F, et al. Incidence and risk factors of arrhythmic events in catecholaminergic polymorphic ventricular tachycardia. *Circulation* 2009;119:2426–34.
 - [45] Saha P, Potiny P, Rigdon J, et al. Substantial cardiovascular morbidity in adults with lower-complexity congenital heart disease. *Circulation* 2019;139:1889–99.
 - [46] Amedro P, Werner O, Abassi H, et al. Health-related quality of life and physical activity in children with inherited cardiac arrhythmia or inherited cardiomyopathy: the prospective multicentre controlled QUALIMYORHYTHM study rationale, design and methods. *Health Qual Life Outcomes* 2021;19:187.