Accepted, Quaternary Science Reviews, 13/11/2020. MS No.: JQSR-D-20-00139R3

**Ocean surface and bottom water conditions, iceberg drift and sediment transport on
the North Iceland margin during MIS 3 and MIS 2**

Andrews, J.T.¹, Smik, L.², Belt, S.T.², Sicre, M-A³, McCave, I. N.⁴

1. Institute of Arctic and Alpine Research, and Department of Geological Sciences,
University of Colorado, Boulder, CO 80303

2. Biogeochemistry Research Centre, School of Geography, Earth and
Environmental Sciences, University of Plymouth, Drake Circus, Plymouth, PL4
8AA, UK.

3. LOCEAN, CNRS, Sorbonne Université, Campus Pierre et Marie Curie, 4 place
Jussieu, Paris, France.

4. Godwin Laboratory for Palaeoclimate Research, Department of Earth Sciences,
University of Cambridge, Downing Street, Cambridge CB2 3EQ, U.K.

Orcid: Andrews: 0000-0003-3169-5979

Orcid: Belt: 0000-0002-1570-2924

Orcid: Sicre: 0000-0002-5015-1400

Orcid; McCave: 0000-0002-4702-5489

corresponding: andrewsj@colorado.edu

Key words: Iceland Plateau, MIS 2 and 3, sea ice biomarkers, IP₂₅, alkenones,

sortable silt, sediment provenance

28 **Abstract**

29 Radiocarbon dates and marine tephra suggest that the upper 10 m of core MD99-2274,
30 off North Iceland, extends from ~0 to ~65 ka BP. A multi-proxy sediment and biomarker
31 study at a millennial-scale resolution is used to derive a paleoclimate scenario for this
32 area of the southwestern Nordic Seas, which during the Holocene had intermittent
33 excursions of icebergs and a seasonal cover of drifting sea ice across the site. The
34 sortable silt mean size ($\overline{SS}$) suggests a bottom current (1000 m depth) flow speed
35 maximum to minimum range of ~8 cm/s during Marine Isotope Stages 2 to 3, but the data
36 are unreliable for the Holocene. Slow-down in flow speeds may be associated with
37 massive ice and water discharges linked to the Hudson Strait ice stream (H-events) and to
38 melt of icebergs from Greenland in the Nordic seas where convection would have been
39 suppressed. Five pulses of sediment with a distinct felsic component are associated with
40 iceberg transport from E/NE Greenland. Sea ice, open water and sea surface temperature
41 (SST) biomarker proxies (i.e. IP₂₅, HBI III, brassicasterol and alkenones) all point
42 towards near-perennial sea ice cover during MIS 3 and 2, rather than seasonal sea ice or
43 open water conditions. Indeed, our biomarker and sediment data require that the seas
44 north of Iceland experienced a nearly continuous cover of sea ice, together with icebergs
45 calved from ice stream termini, which drifted southward. The cross-correlation of the
46 quartz % records between MD99-2274 and the well-dated core PS2644 in Blosseville
47 Basin indicates significant coherence in the records at a multi-millennial (~8 ky)
48 timescale. A transition to open ocean conditions is evident from the early Holocene
49 onwards, albeit with the occurrence of some drift ice and icebergs.

50

1. Introduction

1.1 Aims of study

In order to gain some understanding of the complex marine environments that prevailed during the Late Quaternary we need to employ a multi-proxy approach that not only characterizes ocean surface and bottom water conditions, but also provides direct measurement of glacial influences on sediment supply. Several studies have been reported from the North Iceland Shelf (Fig. 1) that document late glacial/Holocene records (e.g. Andrews et al., 2018; McCave and Andrews, 2019a & b; Sicre et al., 2008; Knudsen et al., 2003) but there are only limited references to conditions during Marine Isotope Stages (MIS) 2, 3 or 4. Therefore, with the primary goal of establishing a framework for environmental conditions in this sector of the Iceland Sea from MIS 2 to MIS 4, we selected a previously unstudied core, MD99-2274 (Labeyrie et al., 2003) (Fig. 1), and sampled the upper 10 m. MD99-2274 (henceforth #2274) is a 10-cm diameter 26 m Calypso core retrieved from 67.582°N and 17.073°W at 1000 m water depth (Labeyrie et al., 2003) during the IMAGES V cruise aboard the French RV *Marion Dufresne*. For further context, we note that the core site is located 200 km east of the well-studied core PS2644 (van Kreveld et al., 2000; Voelker, 1999; Voelker and Haflidason, 2015) and 163 km from core P57-7 (Sejrup et al., 1989) (Table 1, Fig. 1A). The main questions we posed were: 1) what is an appropriate depth/age model, 2) is there evidence for either pervasive sea ice or an ice shelf (Boers et al., 2018; Dokken et al., 2013; Petersen et al., 2013), which have been called for to explain D-O cycles, 3) what were sea surface temperatures (SSTs), and 4) are there substantial changes in grain-size and mineral composition that can be associated with changes in bottom current flow speed and

changes in glacial sediment provenance? Given the location of the core (Fig. 1A) we were particularly interested in whether we could discriminate between glacial sediments derived from Iceland versus those from E/NE Greenland.

1.2 Present-day oceanography

The Iceland and Greenland Seas (Fig. 1A) are key areas for the formation of dense overflow waters (Brakstad et al., 2019) that flow south through sills in the Scotland-Greenland Ridge (Fig. 1C). The North Icelandic Jet (NIJ) flows southwestward along the slope below ~1000 m (Fig. 1C) with a mean speed of $9.3 \text{ m} \pm 2.7 \text{ m/sec}$ towards Denmark Strait (Mauritzen, 1996; Pickart et al., 2005) where it exits to form a major component of North Atlantic Deep Water “....and points to the Iceland Sea as an important place for this water mass formation.” (Jonsson and Valdimarsson, 2004). The study site lies in a sensitive area with the surface flow being the East Icelandic Current (EIC), which brings cold and relatively fresh surface water as a spin-off from the East Greenland Current (EGC), whereas the North Icelandic Irminger Current (NIIC), sourced from the southern warmer and saltier waters of the North Atlantic Drift (Stefansson, 1962), continues as an eastward flow over the inner North Iceland Shelf (NIC) (Fig. 1C).

Sea ice in the form of drift ice has been noted to reach the area in modern times, although the average position of the sea ice edge (30% sea ice cover by area) lies north of our site (Divine and Dick, 2006) (Fig. 1C). Thirty years of observations on the presence of icebergs (Andrews et al., 2019), their Fig. 7A) indicate that icebergs from E/NE Greenland drift across the site.

1.3 Background to study region

Stein and colleagues (Nam et al., 1995; Stein, 2008; Stein et al., 1996) studied a comprehensive suite of cores on the Scoresby Sund Trough Mouth Fan (Fig. 1 A, TMF) and reported both ice-rafted debris (IRD) and $\delta^{18}\text{O}$ on the near-surface planktonic foraminifera *Neoglobquadrina pachyderma* (Table 1). The cores included discrete IRD peaks (counts $10\text{ cm}^3 > 500\text{ }\mu\text{m}$), which they suggested may have been coeval with the massive ice and water discharges of the Hudson Strait Heinrich (HS H-) events (Andrews and Voelker, 2018b; Heinrich, 1988; Hemming, 2004; Hesse 2016). However, whether the response of the Greenland, Iceland, and European ice sheets was synchronous or asynchronous with the Laurentide Ice Sheet collapse events still requires clarification (Dowdeswell et al., 1999; Elliot et al., 2001). Verplanck et al (2009) provided radiogenic isotope data fingerprinting sediment sources from two cores on the Scoresby Sund TMF (O'Cofaigh et al., 2002) (JR51-GC31 and -GC32) and another core (PS62/017-4) from the Blossville Basin (Milo et al., 2005) (Table 1). Stein et al. (1996) and Verplanck et al. (2009) described events in cores PS1730 and PS62/017-4 (Table 1, Fig. 1) that they considered coeval with the HS H-events. Andrews and Voelker (2018) have argued that the use of the term “Heinrich events” for locations such as the Nordic Seas is not appropriate and should be modified. For example, the IRD-rich layer in PS2644 correlated with HS H-2 (Voelker et al., 1998) is now referred to as PS2644 IRD#2 (Andrews and Voelker, 2018). In our study, events that might correlate with HS H-events will be termed #2274-IRD#.

There is no firm agreement on the extent and duration of sea ice cover in the Nordic Seas during MIS 2 and MIS 3. The CLIMAP data showed an extensive cover across the Nordic Seas (Ruddiman and McIntyre, 1981) whereas Sarnthein et al. (2003)

argue that the Nordic Seas during MIS 2 were “..largely ice free” during the summer months. The presence of an ice shelf buttressing the East Greenland ice streams has also triggered a debate especially as to a possible answer to the cause of D-O oscillations (Pettersen et al., 2013; van Kreveld et al., 2000). However, other researchers working at sites in the eastern Nordic Seas have rather focused on the role of sea ice (Dokken et al., 2013; Hoff et al., 2016) and changes in the structure of the water column, and concluded that during Greenland interstadials in MIS 3, sea ice was limited in extent and duration.

The presence of thick, pervasive sea ice could potentially limit the export of icebergs from E and NE Greenland Ice Streams (Reeh et al., 1999), although the sediment records from numerous sediment cores retrieved from the floor of the Arctic Ocean clearly document that iceberg rafting occurred throughout the Pleistocene (Clark, 1990a,b; Stein, 2008; Phillips and Grantz, 2001; Stokes et al., 2005), with some evidence that the timing of events in some cores were similar to those for HS H-events. For example, IRD peaks in cores from the Arctic Ocean were linked to the McClure Ice Stream in the NW sector of the Laurentide Ice Sheet and dated at 12.9, 15.6, ~22, and 30 ka BP (Stokes et al., 2005). Iceberg drift is primarily a function of the integrated current direction and speed over depth, plus a component associated with wind forcing on the exposed “sail” (Bigg, 2016). In many ways, sea ice protects icebergs as it inhibits wave action, which is the greatest cause of iceberg disintegration (Bigg, 2016; Venkatesh et al., 1994).

1.4 Ice sheet extent MIS 1 to MIS 3

#2274 lies only 60 km north of the LGM limit of the Iceland Ice Sheet (IIS) (Fig. 1) (Andrews and Helgadóttir, 2003; Patton et al., 2017) with the onset of retreat associated

with calibrated radiocarbon dates of between 14 and 15 ka BP, depending on the ocean reservoir correction (Andrews et al., 2018; Andrews and Helgadottir, 2003; Knudsen et al., 2003). Retreat from the maximum position was rapid (Andrews et al., 2018; Norðdahl and Ingolfsson, 2015; Patton et al., 2017), and the ice sheet was at or behind the present-day coast by the time of the deposition of the Vedde tephra ~12.2 ka BP (Lohne et al., 2013). Little detail is known about the history of this ice sheet during MIS 3 (e.g. Andrews et al., 2017). Moles et al. (2019) argued that the North Atlantic Ash Zone II (NAAZII) tephra, dated ca 54 ka BP (Austin and Hibbert, 2012), was erupted under >400 m of ice, thus indicating a reasonably extensive IIS during the Greenland ^{18}O stadial 15.2 (Moles et al., 2019; Rasmussen et al., 2014), but no specific information is currently available on the MIS 3 history of the ice sheet.

The Greenland Ice Sheet (GIS) extended to the shelf break during the LGM (Funder et al., 2011b; Vasskog et al., 2015) but little is known about its history during MIS 3 or MIS 4. Judging from the delivery of quartz-rich sediments to cores along Denmark Strait, especially PS2644 and MD99-2323 (Andrews and Vogt, 2020a), it is probable that the ice also reached a similar position at these times. Peterson et al. (2013) suggested that an ice shelf may have extended out from the East Greenland Shelf across Blosseville Basin, although the sedimentary evidence for this is scanty (Andrews and Vogt, 2020a).

1.5 Bedrock Geology and source signatures

In terms of the mineral composition of #2274 sediments, the bedrock in glacial source areas consists primarily of either mafic (basalt) or felsic (granites/gneisses/sandstones),

although finer source identification is possible (Andrews and Vogt, 2014; 2020) (Fig. 1A). Further, Andrews and Vogt (2014) demonstrated that the sediment mineral signature of sediments offshore from the Caledonian Fold Belt was dominated by high wt% of quartz, illite, and muscovite. Detrital carbonate sediments derived from the Paleozoic outcrops of N Greenland and the Canadian Arctic are also recognized by color and mineralogy. However, radiogenic isotopes (White et al., 2016; Verplanck et al., 2009) allow more age-related differentiations, which in terms of our region (Fig. 1A and B), consists of Archaean, Paleoproterozoic, Caledonian Fold Belt, and Tertiary volcanics (Henriksen, 2008).

2. Environmental proxies and age model

2.1 Data methods

The proxies used in this paper are the sea ice biomarkers IP₂₅ and HBI II (Belt et al., 2007; Belt and Müller, 2013; Belt, 2018), brassicasterol and HBI III as indicators of open water primary production (Volkman, 1986; Belt et al., 2015), alkenones (for SST) (Sicre et al., 2008a), % C37:4 alkenone to identify polar surface waters, grain-size indicators of bottom flow and deposition (McCave and Andrews, 2019a, b; McCave et al., 2017), magnetic susceptibility, and quantitative X-ray diffraction estimates of mineral wt% (Andrews et al., 2017; Andrews and Vogt, 2014). The X-ray diffraction data for #2274 are available (Andrews and Vogt, 2020b) The full details of these methods are included as Supplementary Material.

2.2 Depth/age model

The age model is based on radiocarbon dates and the occurrence of tephras (Table 2). There are significant problems associated with obtaining and interpreting calibrated ages because of the uncertainty of the ocean reservoir correction (ORC), which has varied spatially and temporally, and might be as much as 1000 yr (Andrews et al., 2018; Skinner et al., 2019; Voelker, 1999; Voelker et al., 1998). Three radiocarbon dates were obtained on the near-surface planktonic foraminifera *Neogloboquadrina pachyderma* and the other on lustrous shell fragments. Most tephras older than the Borrobol (ca 14.5 ka BP) (Lind et al., 2016; Matthews et al., 2011) are dated by reference to GIS cores, which themselves are based on a variety of assumptions and whose error increases with the estimated age (Boers et al., 2017). The qXRD data (Andrews et al., 2013; 2018) suggest the presence of high wt% of volcanic glass in two cores on the Iceland Shelf that might be coeval with the Vedde and NAAZII tephras (Brendryen et al., 2011; Lohne et al., 2013). The tephra bed at 607 cm in #2274 was identified by Haflidasson (person. commun. 2018) as being similar to FMAZ IV dated at ~47.12 ka BP (Davies et al., 2008; Rasmussen et al., 2003; Voelker and Haflidason, 2015) and that date is used in our depth/age models (see Supplementary Material). Other discrete layers of black basaltic glass were noted in the shipboard log at 99, 127.5, 717, and 740 cm (Labeyrie et al., 2003, p 477), and age estimates were obtained from our depth/age model (see later).

We used the Bayesian radiocarbon calibration program “Bacon” (Blaauw and Christen, 2005) to construct depth/age models, but we also acknowledge the many problems associated with establishing accurate depth/age models (Telford et al., 2003; Trachsel and Telford, 2017). The first model is based solely on the available ^{14}C dates and the 607 cm tephra (Table 2A and B), while the second one is based on an assumed

age estimate for the core top of 500 ± 500 (i.e. little sediment loss) and the inferred presence of the Vedde and NAAZII tephras. Given the uncertainty in the OCR, we used a $\Delta R = 0$. In practice, there is relatively little difference in the median age estimates (Fig. 2A). The average sediment accumulation rate (SAR) is 68 yr/cm or 14.7 cm/ky, thus our 10 cm sampling density permits millennial-scale evaluations, with an average spacing between samples of 0.5 cal ky. Remarkably, for MD cores of this vintage (1999), the upper part of the core shows no evidence of piston-induced stretching (Skinner and McCave, 2003). However, the spread between minimum and maximum age estimates is often considerable given the relative paucity of dated levels, and the Bayesian approach results in an age estimate for the core top of 3600 yr BP, although the estimated date of 500 ± 500 yr BP finds some support in our data. The estimated ages for the logged tephra layers are: ~11, 13.2, 53, and 56 ka BP. A possible age for the 99 cm basaltic tephra is the 10.2 ka BP Saksunarvatn tephra (Lohne et al., 2013), which is widespread on the north Iceland Shelf (Krisjansdottir et al., 2007; Eiriksson and Knudson, 2002). All our subsequent data have been converted to a common depth/age model using the data in Table 2B; thus, robust inter-proxy comparisons can be made. To ensure that we have not forced our data into an existing framework we have not tuned our model to other records (Blaauw, 2012).

We have also obtained radiocarbon dates on several *Vema* cores that were taken to the north of Iceland and #2274 (Fig. 1; Table 3) (Manley and Jennings, 1996). The calibrated radiocarbon dates range from ~13 to > 49 ka BP ($\Delta R = 0$) and were obtained on relatively large samples of *N. pachyderma* (Table 3). Several tephras were noted in the core description (Suppl. Data), thus indicating that conditions allowed for the deposition

of discrete tephras. The dates from these cores also provide additional information on the presence of significant numbers of the planktonic foraminifera *N. pachyderma* (Greco et al., 2019) and hence inferences about sea ice cover and light conditions.

3. Results

3.1 Lithology and Grain-size

The core log of core #2274 (Labeyrie et al., 2003. p. 477) described the sediment as being principally mottled silty clay with colors ranging between 2.5Y4/2 to 5Y4/1. Visible ice-rafted clasts occur but are not common. The grain-size measurements were undertaken on sample splits from the qXRD samples and only 30 samples were processed, resulting in a coarse resolution data set (on average one sample every 2300 yr). The sediments vary between a very coarse to a fine silt with average grain-sizes varying between 54.3 to 6.05 μm . Sand $> 240 \mu\text{m}$ is considered to be ice-rafted (McCave and Andrews, 2019a) and occurs in low % throughout the core, except for two notably coarser intervals with $\text{IRD}_{240} > 5\%$, (Fig. 3).

We have also undertaken an analysis of the sortable silt mean size ($\overline{SS}$) and $SS\%$ in the 63-10 μm fractions (McCave et al., 1995). The correlation coefficient between these two variables is $r = 0.804$ indicating, relative to other cores (McCave and Andrews, 2019a,b), a somewhat noisy correlation, but a generally current-sorted signal (Supplementary Fig. 1). Computation of the running correlation between $SS\%$ and $\overline{SS}$ yields high average values ($r > 0.9$) between ~ 11 and 42 ka BP but values unacceptable for flow speed inference occur in the Holocene and during brief interval ~ 57 ka BP (Fig. 3). Variations in the flow speed of bottom currents (Fig. 1C) in this region reflect changes in the vigour of the ocean overturning system because the NIJ feeds into the

Denmark Strait overflow, a key starting point for the North Atlantic western Boundary Undercurrent.

The overall range (minimum-maximum) in flow speed indicated by this record is ~ 8 cm/s. Calibration of the sortable silt proxy yields a sensitivity ($\text{cm s}^{-1}/\mu\text{m}$) rather than an absolute speed-size relationship (McCave et al., 2017). In favourable circumstances actual speeds may be estimated by matching core-top $\overline{SS}$ data to nearby current meter measurements and plotting the differences downcore. Unfortunately, because the Holocene data are unreliable as a speed record, we cannot relate this to the present nearby flow speed measurements of 9.3 cm/sec (Jonsson, 2004). Nevertheless, low speeds correspond to HS H 1 (~ 15 ka), 4 (~ 40 ka), and 6 (~ 60 ka) (Fig. 3) as expected from previous work on the impact of Heinrich and other cold events on N. Atlantic circulation (e.g. Kleiven et al., 2011), on the basis of which, speeds of < 5 cm/s are probable.

3.2 Mineral composition

On Figure 4, we plot the changes in the weight % of key minerals as determined by qXRD as well as the ratio quartz/pyroxene, which we use as a measure of felsic/mafic bedrock (as opposed to quartz/plagioclase which was used by Moros et al. (2004)). The quartz wt% in a surface grab from this site is $\sim 5\%$ (Andrews and Eberl, 2007), and the median for the whole record is 5.3 % with a maximum of 16.8 %. The magnetic susceptibility record for #2274 (Fig. 2A) is clearly inversely associated with the variations in quartz (Fig. 2B), which, together with the K-feldspars, are diamagnetic minerals (Robinson et al., 1995; Watkins and Maher, 2003). A similar inverse relationship was noted in other cores from the area (Andrews and Vogt, 2020a). Hence

the magnetic susceptibility fluctuations support our interpretation that there are substantial variations in the inputs of felsic- versus mafic-rich sediments.

The Holocene record mimics that from many sites on the North Icelandic Shelf (NIS) in showing an increase in quartz toward the present-day (Andrews et al., 2019). Quartz and pyroxene have an antiphase relationship ($r^2 = 0.47$), which in part is related to the mineral data summing to 100% (i.e. a closed array), and which provides some constraints on the interpretation (Aitchison, 1986; Chayes, 1971). There are five sustained peaks in the quartz wt % (Fig. 4), and K-feldspar (not shown, K-feldspar values track those of quartz (Andrews and Vogt, 2020a)) are therefore not included in this figure), which we interpret as indicating the influx of sediment from NE Greenland and possibly farther afield from Canada or Fennoscandia. Of these possible mechanisms, icebergs alone carry basal and englacial debris that includes all size fractions from cobbles to clay ($> 1 \mu\text{m}$). The variations in quartz are frequently matched by the sum of calcite and dolomite (carbonate) (Fig. 4) ($r^2 = 0.13$, $p < 0.0001$) although the correlations are much more significant for dolomite ($r^2 = 0.22$) than calcite ($r^2 = 0.07$). This probably represents transport of glacially derived material from the carbonate bedrock of NE and N Greenland and/or the Canadian Arctic Islands and Channels (Darby and Zimmerman, 2008; Lakeman et al., 2018; Phillips and Grantz, 2001). The estimated ages for the 5 peaks are 14.4, 31.5, 40, 54.7, and 61.8 ka BP (Fig. 4) with a possible smaller episode ~ 22.8 ka BP. These age estimates are somewhat similar to the HS H-events (Andrews and Voelker, 2018a; Heinrich, 1988; Hemming, 2004) (see Fig. 3) but their duration are longer than the < 1 ky episodes of detrital carbonate deposition associated with the HS H-events (Andrews and Voelker, 2018a).

Previous work on sediment sources in this area (Verplanck et al., 2009) provide temporally limited but critical information using radiogenic isotopes on the < or > 63 μ m fractions. Debris flow from the two Scoresby Sund TMF sites (JR51-GC31,-32, Table 1, Fig. 1B) lay along the 1.7 Ga Paleoproterozoic isochron; the samples contained abundant quartz and lesser amounts of basalt (Verplanck et al., 2009, p.53). However, the sediments in the Blossville Basin (core 17-4, Fig. 1A, Table 1), some 150 km downstream (Fig. 1A), and considered to be coeval with HS H events-1, -2, and -3, all cluster along the 0.5 Ga isochron (Calendonide bedrock, that outcrops on the eastern edge of NE Greenland north of Scoresby Sund (Fig. 1B)). The same isotopic signature characterized the non-HS H sediments in this core. Pb systematics indicate that the Holocene sediment samples at sites 907 (Table 1) and JR51-GC28 are dominated by the 0.5 Ga Calendonides (White et al., 2016). Given the sediment SedUnMix results (Fig. 5) and the reported radiogenic isotopic data (Verplanck et. al., 2009; White et al., 2016), the variations in quartz are most probably associated with sediment discharge events from glacial erosion and transport in ice streams flowing through the numerous fiords north of Scoresby Sund and primarily within the Caledonian Fold Belt outcrop (Evans et al., 2002, 2009; Stein, 2008).

The SedUnMix analysis included sediments from NE Greenland (Calendoides, ~73N; Andrews et al., 2016), E Greenland (basalt), and Iceland. The analysis of possible bedrock sources for the #2274 compositional changes indicated (as might be expected given the bedrock geology of E and NE Greenland, and Iceland) that the NE Greenland source had a granite and gneissic composition, whereas E Greenland and Iceland were linked to basalt and also dolerite (Brooks and Nielsen, 1982; Henriksen, 2008; Higgins et

al., 2008; Kristjansson et al., 1979). The results (Fig. 5) indicate that felsic-rich sediments from NE Greenland or farther afield (Arctic Canada, Fennoscandia) (Verplanck et al., 2009) were deposited in a series of events that mimic the influx of quartz to the site (Figs. 2B and 4); the correlation between the NE Greenland Calendonides source estimates in #2274 and the quartz wt% explains 79% of the variance. The average “unaccounted” or “unexplained” composition averaged 20 ± 5 % and degree of fit or average absolute bias is 2.3 ± 0.4 wt% indicating that the input mineral source regions provide a good fit to the #2274 mineral compositions. Figure 5 highlights two periods when the mineral composition indicates little deposition of sediment that could be ascribed to a felsic source centered around 20 and 57 ka BP, the latter being a time of substantial deposition of tephra at this site and also a time when glacial ice covered at least some of Iceland (Moles et al., 2019). Source estimates from E. Greenland (sites seaward of the early Tertiary basalt outcrop on the Geikie Plateau) and SW Iceland resulted in nearly identical patterns over the last ~65 ka BP (Fig. 5), but the results from considering Icelandic basal glacial marine diamictos (Dmm) as a source are different. The reasons for these two differing estimates are presently unclear.

The provenance time-series thus suggests that we can identify four episodes in the arrival of foreign sourced sediments; 1) from ~65 to 38 ka BP when distinct pulses of NE Greenland sourced sediments arrived; 2) 38 to 17 ka BP when there was an overall decrease in this source with virtually no quartz noted ~20 ka BP; 3) a large pulse of these sediments centered ~ 15 ka BP; and 4) the last 10 ka or so that shows a steady increase in this source. This latter event is also noted in MD99-2269 (Fig. 7) and is matched by changes in the sea ice biomarker IP₂₅ (Cabedo-Sanz et al., 2016).

3.3 Biomarkers

The sea ice biomarkers IP₂₅ and HBI II were absent or below the limit of detection in the majority of the sediment sections analyzed with only a few exceptions (Fig. 6). Of the two, HBI II was always more abundant, consistent with findings from previous studies from the study region and elsewhere in the Arctic (e.g. Massé et al., 2011; Xiao et al., 2013; Bai et al., 2019). In some cases, only HBI II could be identified and quantified, with IP₂₅ likely also present in such horizons but below the detection limit.

Alkenones and brassicasterol were found at very low concentrations in glacial sediments contrasting with higher abundances in Holocene sediments. Further, the open water biomarker HBI III was only detected in Holocene sediments (data not shown). While alkenone-SSTs ranged from 7 to 9°C during the Holocene, they are unexpectedly high in the glacial portion of the record, spanning from 8 to 16°C.

4. Discussion

4.1 Icebergs and IRD during MIS 3 and MIS 2

There is no general theory about the association of sea ice and icebergs and there is no observational census of the icebergs being transported in the EGC as there is for the Labrador Shelf off Newfoundland, apart from a 30-yr count of icebergs on the Iceland shelves (Jónsdóttir *in* Andrews et al., 2019). However, Cabedo-Sanz et al. (2016) and Darby et al. (2017) showed that Holocene variations in the wt% of quartz and the sea ice biomarker IP₂₅ co-varied in cores to the west and south of #2274, yet this was not the case at #2274 during MIS 2 and MIS 3 (Figs. 4 and 6). In N Greenland, semi-permanent sea-ice conditions prevail today and did so intermittently during the Holocene (Funder et

al., 2011a) and it is reasonable to assume that sea-ice would have been more extensive during MIS 2 and MIS 3 when the GIS may have reached the shelf break. However, cosmogenic dates pertaining to the extent of the Northeast Greenland Ice Stream at ~78°N (Larsen et al., 2018) have been used to argue that this ice stream was behind its present margin "...41-26 ka."

Several authors have argued for the presence of an ice shelf fringing the E/NE GIS (Boers et al., 2018; Petersen et al., 2013). However, sediments recovered from beneath ice shelves are invariably fine-grained and lack ice-rafted debris (Domack et al., 1999; Jennings et al., 2019; McKay et al., 2016), whereas the sediments from the Scoresby Sund TMF (Fig. 1) and margin contain clear IRD (Stein et al., 1996) (Fig. 5; see also Table 3). Radiocarbon dates in Stein et al., (1996a) were based on 2000 *N. pachyderma* specimens per sample, and the numerous MIS 2 and MIS 3 radiocarbon dates on *N. pachyderma* from PS2644 (Sarthein et al., 2003; Voelker, 1999; Voelker et al., 1998, 2000) were obtained on 10 mg samples of 800-2300 tests in 1-cm sediment samples. Although the complete ecology of *N. pachyderma* is not well known, a study of plankton hauls (Greco et al., 2019) indicates a relationship between sea ice cover and chlorophyll, hence suggesting that "*light or light-dependent processes might influence the ecology of this species.*" In addition, several of these cores have discrete tephra layers indicating rapid accumulation of tephras by particles falling through the water column, versus a more dispersed occurrence if the tephra was deposited on multi-year sea ice. Together these data indicate that the sea ice, at times during MIS2 and 3 and probably seasonally, must have had extensive leads and open-water areas.

397 Stein et al (1996) present detailed IRD data (counts $10 \text{ cm}^3 > 500 \mu\text{m}$) from a
 398 series of radiocarbon dated cores on the Scoresby Sund TMF (Fig. 1; PS1726 and
 399 PS1730, Fig. 1B) that reflect delivery of coarse sediments in a discrete series of episodes
 400 (data from www.Pangaea.de). Stein (2008) noted coarse sediment intervals that were
 401 attributed to iceberg-rafting at ~4-15, 16, 17-18, 20-21, and 22-23 ka BP. There are no
 402 mineral composition data for PS1730, but data exist for PS2644, which is 300 km away
 403 (Table 1, Fig. 1B) (Andrews and Vogt, 2020a; Vogt, 2017). A comparison between
 404 PS2644 and #2274 (Fig. 8A) indicates that PS2644, closer to the Scoresby Sund Ice
 405 Stream, has more quartz wt% but there are some notable corresponding peaks in both
 406 series. However, we note that the quartz wt% were obtained via two different but
 407 comparable quantitative methods (Andrews and Vogt, 2020a; Vogt, 2017; Zou, 2016). To
 408 evaluate similarities and differences between these two sites we used cross-wavelet
 409 analysis (Roesch and Schmidbauer, 2018; Hammer et al., 2001) (Fig. 8). The wavelet
 410 analysis of the two quartz records (Fig. 8A) demonstrates both important coeval events as
 411 well as obvious differences. In addition, the overall match between these sites for the
 412 earlier part of the record adds confidence to our age model, and also emphasizes the
 413 important differences between 35 and 65 ka. The reconstructed wavelets for PS2644
 414 show three major pulses of quartz at ~13, 20, and 29 ka BP, and these are matched by
 415 much lower peaks at #2274. Conversely, there are no distinct peaks during MIS 3 in
 416 PS2644 but there are in #2274. The sense of the directional arrows in Figure 8B is that
 417 PS2644 either leads or is in phase with #2274, and there is a hint of a significant shorter
 418 period ~60 ka BP with the two records being anti-phase. The cross-wavelet power
 419 spectrum (Fig. 8B) confirms the presence of a significant zone of coherence extending

from ~10-34 ka BP with the average cross-wavelet power peaking at ~8 ky (Fig. 8C); this is of course similar to the periodicity of HS H-events (see Clark et al., 2007) (e.g. Fig. 3) but lacks the diagnostic carbonate provenance indicators (Andrews and Voelker, 2018). Possibly because of our 0.6 ky sample spacing (Fig. 8A), there is no obvious D-O signal in the quartz PS2644 data, whereas it is evident in the $\delta^{18}\text{O}$ *Np* data (Suppl. Fig. 3). The difference in signals between #2274 and PS2644 during MIS 3 (Fig. 8A) suggests a change in either the delivery of quartz-rich sediments or a dampening down of sediment delivery.

The sortable silt evidence indicates that even at the glacial maximum there was flow along the slope in the precursor to the NIJ. As this presently heads toward the Denmark Strait outflow, we suggest that the Nordic Seas acted as a source of deep waters (probably formed in the east where Atlantic inflow continued (Sarnthein et al, 1994)) that overflowed to the North Atlantic where they formed a deep water mass (Howe et al., 2016; Keigwin and Swift, 2017). The classical view of Nordic Sea behaviour during cold periods is that freshwater from melting ice-sheets and -bergs suppresses convection resulting in a severe reduction or even cessation of the AMOC inflow and overflow (e.g. a recent model, including consideration of the EGC, analysing this is from Liu et al, (2018)). However an emerging view is of a slowdown (not cessation) of Nordic Sea overflows in cold periods (Howe et al., 2016; Keigwin and Swift, 2017). A very recent view is that ice discharges in the North Pacific precede Heinrich events and may be implicated as a triggering mechanism (Walczuk et al., 2020). In the Nordic Seas Atlantic water inflow persisted throughout the Pleistocene glacials over the Norwegian slope (Sarnthein et al., 1994; Newton et al., 2018). The evidence here indicates a persistent

outflow along the N Iceland Slope with reductions during HS H- events 1, 4, and 6. Flow speed decreases have been noted for both shallow and deep flows in this region during stadials and glacial intervals of the late and mid-Quaternary (Kleiven et al., 2011; McCave and Andrews, 2019b). The Younger Dryas often shows speed decreases but some cores record increased flow (McCave and Andrews, 2019b), as is seen here. These disparities remain a puzzle.

4.2 Rationalizing mineralogical and biomarker proxies for sea ice reconstruction

When detected, the concentrations of IP₂₅ and HBI II were mainly much lower than those reported previously for mid-late Holocene (ca. 6-0 cal. ka) and deglacial (ca. 15-11 ka) sites from the NIS (Cabedo-Sanz et al., 2016; Xiao et al., 2017). However, the presence and concentration of IP₂₅ at ca. 3.7 ka aligns with previous data reported from core JR51-GC35 (located 76 km SW of #2274 (Figs. 1B and 7; Table 1)) for the mid-Holocene (Cabedo-Sanz et al., 2016), consistent with the delivery of drift ice across the NIS at that time (Fig. 7). The otherwise general absence of IP₂₅ and HBI II in #2274 points towards an environment unfavorable for sea ice diatom growth, namely ice-free conditions or a setting of near-permanent ice cover. To distinguish between these two scenarios, we measured three other biomarker types indicative of open water conditions, i.e. brassicasterol, HBI III and alkenones. In the case of brassicasterol, a phytosterol characteristic of marine diatoms (Volkman, 1986), concentrations in selected sediments from #2274 were relatively high in the Holocene section and typically two orders of magnitude lower in older (>14 ka) intervals, indicative of much lower glacial primary productivity reflecting near-perennial sea ice cover. Similarly, HBI III, a biomarker

466 derived from certain open water diatoms (Belt et al., 2017), was only detected in
467 Holocene sections (data not shown). Consistent with these findings, concentrations of
468 alkenones derived from coccolithophorid blooming in mid-late summers were also
469 substantially lower in the older sections compared to those in the Holocene (Fig. 6).
470 Further, the relatively high percentage contribution of the tetra-unsaturated alkenone C_{37:4}
471 prior to the Holocene (mean value 36% compared to 6% for the Holocene) is consistent
472 with a dominance of polar waters (Sicre et al., 2002; Bendle et al., 2005) potentially
473 laden with sea ice. Alkenone-derived SST estimates for the Holocene (ca. 7–9°C) are in
474 line with those reported from other high-resolution studies from the NIS (e.g. Bendle and
475 Rosell-Melé, 2007; Sicre et al., 2008b; Kristjansdottir et al. 2016). In contrast, SST
476 estimates prior to the Holocene were somewhat higher (ca. 8–16°C; mean 11.4°C)
477 although the accuracy of such estimates might be lower than for the Holocene owing to
478 the relatively high contributions from C_{37:4} (Bendle and Rosell-Melé, 2004).
479 Anomalously warm SSTs associated with low alkenone concentrations during glacial
480 time have been reported in previous studies and attributed to advection of detrital
481 alkenones (Sicre et al., 2005; Knutz et al., 2011). Such advection by surface currents can
482 introduce significant bias in regions where there are large productivity and SST gradients,
483 thereby overprinting any local signal (Bendle and Rosell-Melé, 2004; Conte et al., 2006).
484 With extremely low alkenone production due to the presence of ice at #2274, transport of
485 allochthonous alkenones within the IC likely explains the deviation in SSTs towards
486 seemingly unrealistic warmer values. In any case, the most robust aspects of the
487 biomarker data point towards near-perennial sea ice cover prior to the Holocene, although
488 the presence of both phytosterols and alkenones (albeit at low concentrations) indicates

the occurrence of at least partial open water conditions, potentially restricted to leads or regions of partial ice melt within otherwise heavily consolidated pack ice. Such conditions would likely have led to short-term and reduced primary production during relatively short summer seasons and limited to the near-surface layer due to a strongly-stratified water column resulting from partial ice melt. Both such uppermost surface layer production conditions in leads and advection of allochthonous alkenones within the IC would account for the anomalously high glacial SSTs.

Our conclusion of near-perennial sea ice during MIS 3 and MIS 2 is broadly consistent with outcomes from a recent 120,000 yr reconstruction of sea-ice conditions for the North Atlantic (Maffezzoli et al., 2019) based on the analysis of enriched bromine (Br_{enr}) in an ice core from the Renland Ice Cap (RIC) 560 km WNW from #2274 (Figs. 1 and 7 [RIC]). Albeit at a much broader spatial resolution (i.e. 50-85° N), Maffezzoli et al. (2019) proposed that MIS 3 and MIS 2 experienced a (variable) mix of multi-year and first-year sea ice, before transitioning to mainly first-year ice and open water conditions following the termination of the LGM. Interestingly, the greater range of sea ice cover inferred from the RIC Br_{enr} record is not at all clear in our #2274 record, but is evident in a biomarker record from the eastern Nordic Seas, with extensive/near-perennial sea ice cover during stadials and H-events (i.e. comparable to #2274) but ice-free conditions during interstadials (since ca. 90 ka BP); such differences between marine sites in the western and eastern Nordic Seas presumably reflects the variable influence of warm Atlantic water, limited to the eastern Nordic Seas (Hoff et al., 2016). The most prominent signature of first-year ice in the Br_{enr} records occurred during the Younger Dryas and it is noteworthy that a transition from permanent to increasing seasonal sea ice at the NIS was

reported for this interval following a biomarker-based reconstruction of surface oceanographic conditions from core #2272 (Fig. 1; 7; Xiao et al., 2017). Further, based on relatively high concentrations of IP₂₅ in MD99-2272 during the Younger Dryas and the preceding Bølling-Allerød, Xiao et al. (2017) concluded that biomarker production was likely associated with locally formed first year ice rather than from advected drift ice, the latter being a feature of modern-day oceanography. In contrast, our new data from #2274 indicate still near-permanent sea ice cover at this time (Fig. 7). As such, we interpret the combined ice core and marine sediment core data to suggest that as climate conditions ameliorated at the end of the LGM, near-permanent sea ice cover transitioned to first-year seasonal sea ice in the southern part of the region, especially during the Bølling-Allerød and Younger Dryas, likely due to increasing influence of the IC (Xiao et al., 2017); however, the spatial extent of this area of first year ice, located southward of the near-permanent sea ice front that characterizes MIS 3 and MIS 2, remains uncertain at this point (see Fig. 7 sub-panel). Large-scale sea ice reduction then characterized the early Holocene (Fig. 7), with a marked increase in all open water primary productivity biomarker proxies (Fig. 6). Increasing drift ice subsequently became a characteristic of the NIS from the mid Holocene onwards (Fig. 7; Cabedo-Sanz et al., 2016).

Conclusions

The multi-proxy sediment data from core #2274 130 km off the north Iceland coast appears at first sight to yield conflicting interpretations depending on whether sediment mineral composition or biomarker proxy data are being considered; however, these can be resolved through a more detailed consideration of the mode(s) of iceberg drift and trajectory through largely consolidated and near-pervasive sea ice. The low- resolution

sampling for grain-size restricts detailed interpretation but the sediments are mostly moderately sorted in the silt range allowing a valid record of bottom flow speed. This shows low flow speeds during H-events 1, 4 and 6 related to decrease in Nordic Sea overflow, but not cessation, and a peak in the Younger Dryas.

The mineral composition of the < 2 mm grain-size sediment samples shows peaks with wt% of quartz values significantly higher than Holocene values. The variations in the quartz wt% are also reflected in the estimated contributions of sediment from Precambrian and Caledonian bedrock sources of NE Greenland. These data require sediment transport to the #2274 site during MIS 3 and MIS 2. If the transport is by icebergs then the sea ice cover had to allow icebergs to drift southward, as they do at present (Figs. 1C, 7). A framework of near-permanent sea ice is confirmed from ultra-low seasonal sea ice and open water biomarker concentrations. On the other hand, the occurrence of non-zero concentrations of some phytoplanktic biomarkers, and numbers of near-surface planktonic foraminifera (Table 3) points to some short-term open water conditions, either from limited sea ice melt or following the opening of leads; the presence of drifting icebergs may be significant in this respect (Fig. 7).

An underlying question for HS H-events is whether North Atlantic-wide glacial marine sediment events were triggered as a response to events in Hudson Strait or whether the events are part of a shared response to broader regional oceanographic conditions (e.g. Marcott, et al., 2011; Bassis et al., 2017; Velay-Vitow et al., 2019). Thus, were “coeval” HS H- events on the East Greenland margin (Stein et al., 1996; Andrews et al., 1998; Voelker, 1999), or lagged events (e.g. Baffin Bay: Simon et al., 2014 Jennings et al., 2018), triggered in response to events in the Hudson Strait ice

stream? If our quartz and IRD events (Figs. 3 and 8) are indeed coeval with HS H-
events, this implies that the stability of ice streams on the NE and E Greenland shelf (and
N Iceland) and Hudson Strait may all have been affected by basin-wide subsurface
warming in response to a reduction in the Atlantic meridional overturning circulation
(Shaffer et al., 2004; Clark et al., 2007; Marcott et al., 2011).

Acknowledgements

We thank Dr Anne E. Jennings for picking and providing the foraminifera for the
radiocarbon dates, and Dr Haflidi Haflidasson for the identification and geochemical
analysis of the tephra. Professor Grant Bigg provided guidance on the role of icebergs
and sea ice. We thank the Centre National de la Recherche Scientifique (CNRS) for
MAS salary and the crew of the *RV Marion Dufresne* for coring operations during the
IMAGESV cruise. Data from this study will be archived at: www.Pangaea.de, along
with other IMAGESV data for MD99-2274. We acknowledge the availability and our
use of data from PS1726, PS1730, and PS2644, which we accessed through
www.Pangaea.de. Finally, we thank three anonymous reviewers who provided critical
and helpful feedback on previous versions of the manuscript.

575 **Tables**

576 Table 1 Location of the cores referenced in this study and showing distance from MD99-
 577 2274. Cores located on Fig. 1A and B unless noted as NA. The last 5 sites are cores that
 578 specify sediment sources based on radiogenic isotopic data (Verplanck et al., 2009; White
 579 et al., 2016).

580

581 Table 2 A and B: Data for two possible depth/age models for MD99-2274 used in the
 582 Bayesian “Bacon” model—see text. cc = 0 when date derived from other sources and
 583 does not require calibration; cc = 2 when ocean reservoir correction $\Delta R = 0$ is used
 584 (marine IntelCal 13; Reimer et al., 2013).

585

586 Table 3: Depth/age data and calibrated ages for radiocarbon dates on near-surface
 587 planktonic foraminifera (see Figs. 1 and 5). Ocean reservoir correction $\Delta R = 0$.

588

589 **Suppl. Table:** Geochemistry of the tephra layer (see text). Courtesy Dr. H. Haflidasson)

590

591 **Figure Captions**

592 Figure 1: A) location of MD99-2274 and some other cores noted in the paper (Table 1)
 593 (ODV, Schlitzer, 2011). The shaded areas represent the late glacial maximum (LGM)
 594 extent of the ice sheets north of Denmark Strait; the words “basalt” and “felsic” define
 595 the primary sediment mineral sources and the arrows show probable flow paths for
 596 icebergs. BB = Blosseville Basi; TMF = Scoresby Sund Trough Mouth Fan; B)
 597 Additional cores referenced in the paper (see also Table 1). Note that “Cald” on this
 598 figure references the southern outcrop of the Greenland Caledonides (Higgins et al.,

2008). SS = Scoresby Sund; RIC = Renland Ice Cap. C) Surface and bottom currents and historical April sea-ice edge (1870-1920) (dashed white line; Divine and Dick, 2006). NIIC = North Iceland Irminger Current; EGC = East Greenland Current; EIC = East Iceland Current; Yellow lines: Bottom Currents DSOW = Denmark Strait Overflow Water; NIJ = North Iceland Jet., S = Separated East Greenland Current; OC = Iceland Sea Ocean Convection site (after Harden et al., 2016).

Figure 2: A) Downcore plot of magnetic susceptibility (SI^5) and Bayesian ((Blaauw and Christen, 2016) depth age plots for MD99-2274 (see Table 2)---the red curve is for the initial available data blue curve is for the estimated ages with the addition of an estimated core top age and the presence of the Vedde and NAAZII tephras (see text). The Marine Isotope Stage (MIS) boundaries are indicated. Location of radiocarbon dates and tephras are noted. B) Plot of the departures from the median values of magnetic susceptibility ($2.03 \times 10^{-3} SI$) and quartz wt% (5.3). Note that the quartz axis is reversed.

Figure 3: Variation in the Sortable Silt mean size (3-point 1-2-1 weighted smoothing with raw data dots) and IRD% >240 μm . Minima in $\overline{SS}$ are seen at the time of Hudson Strait H events -H6, -H4 and -H1 while -H4, -H2, early -H1 and the YD (-H0) are marked by elevated IRD %. Blue bars are regions where the data are unreliable indicators of flow speed according to the $\overline{SS}$ -SS% correlation criterion of McCave and Andrews, (2019a)

Figure 4: Plots of the variations in the weight% of minerals in MD99-2274, the quartz/pyroxene ratio, and magnetic susceptibility. The green shaded areas represent Holocene values, hence points above represent departures. Numbers 1 through 5 identify IRD quartz peaks. The vertical blue shading areas represent times when the weight% of quartz exceeds Holocene limits.

Figure 5: Plots of the sediment source percentages and the degree of fit (DOF), that is the average absolute bias in the SedUnMix calculation of (observed mineral wt% - predicted mineral weight%) for each sample. The top panel shows the location of measurable quantities of gravel, and sites of tephra layers and the radiocarbon dates on near-surface planktonic foraminifera (Table 3). Numbers on the NE Greenland panel represent the peaks in that source and the yellow bars locate areas with minimal input from that area.

Figure 6: Biomarker data (A) IP₂₅ and HBI II concentrations; (B) $\sum C_{37:3} + C_{37:2}$ alkenone and brassicasterol concentrations; C) SST° C estimates and the %C_{37:4}; and D) Weight % quartz and different coarse sediment fractions.

Figure 7: Schematic presentation of changes in sea ice and iceberg distribution. The first panel (upper left) shows core locations (see Table 1 and Fig. 1A and B) and the adjoining panel the inferred conditions during MIS 3 and 2 with pervasive sea ice and embedded icebergs. The remaining panels show the proposed evolution in the state of sea ice and iceberg supply (red triangles) during deglaciation into the Holocene (adapted from

643 Cabedo-Sanz et al., 2016; Xiao et al., 2017). SS =Scoresby Sound, RIC=Renland Ice
 644 Cap.

645

646

647 Figure 8: Analysis of the quartz wt% records from PS2644 (Vogt, 2017) and MD99-2274
 648 at a common 0.6 ky spacing. A) Original quartz data (black line) and the wavelet
 649 reconstructions for the two records; B) Cross-wavelet power spectrum of quartz wt% for
 650 PS2644 and MD99-2274. The cone of confidence indicated by the light grey areas;
 651 0.05% probability area demarcated by white line. Arrows pointing to the right mean that
 652 the two records are in phase, arrows pointing down mean that x leads y, arrows pointing
 653 to the left indicate the records are anti-phase and pointing up indicates that #2274 leads
 654 PS2644. C) Cross-wavelet (Fig. 8B) average power. The 0.05 significance period is red
 655 and delimited by the dashed slanting line. The horizontal dashed line indicates the peak
 656 periodicity (~8.5 ky).

657

658

659 Suppl. Figure 1: Data for VM30-130 (see Fig. 1 and Table 3).

660

661 Suppl. Figure 2: Showing the reduced major axis association between sortable silt mean
 662 size ($\overline{SS}$) and SS%.

663

664 Suppl. Figure 3: $\delta^{18}\text{O}$ *N. pachyderma* plots of cores from the Blosseville Basin/Scoresby
 665 Sund Trough Mouth Fan (see Fig. 1 and 8) from cores PS1730 (Stein et al., 1996a,b,
 666 and PS2644 (Voelker, 1999).

References

- Aitchison, J., 1986. The statistical analysis of compositional data. Chapman and Hall, London.
- Andrews, J.T., Bjork, A.A., Eberl, D.D., Jennings, A.E., Verplanck, E.P., 2015. Significant differences in late Quaternary bedrock erosion and transportation: East versus West Greenland $\sim 70^{\circ}\text{N}$ and the evolution of glacial landscapes. *Journal of Quaternary Science* 30, 452-463.
- Andrews, J.T., Cabedo-Sanz, P., Jennings, A.E., Olafsdottir, S., Belt, S.T., Geirsdottir, A., 2018. Sea ice, ice-rafting, and ocean climate across Denmark Strait during rapid deglaciation (similar to 16-12 cal ka BP) of the Iceland and East Greenland shelves. *Journal of Quaternary Science* 33, 112-130.
- Andrews, J.T., Cooper, T.A., Jennings, A.E., Stein, A.B., Erlenkeuser, H., 1998: Late Quaternary iceberg-rafted detritus events on the Denmark Strait–Southeast Greenland continental slope ($\sim 65^{\circ}\text{N}$): related to North Atlantic Heinrich events? *Marine Geology* 149, 211-228.
- Andrews, J.T., Dunhill, G., Vogt, C., Voelker, A.H.L., 2017. Denmark Strait during the Late Glacial Maximum and Marine Isotope Stage 3: Sediment sources and transport processes. *Marine Geology* 390, 181-198.
- Andrews, J.T., Eberl, D.D., 2007. Quantitative mineralogy of surface sediments on the Iceland shelf, and application to down-core studies of Holocene ice-rafted sediments. *Journal of Sedimentary Research* 77, 469-479.
- Andrews, J.T., Eberl, D.D., 2012. Determination of sediment provenance by unmixing the mineralogy of source-area sediments: The "SedUnMix" program. *Marine Geology* 291, 24-33.
- Andrews, J.T., Helgadottir, G., 2003. Late Quaternary ice cap extent and deglaciation of Hunaflaall, North West Iceland: Evidence from marine cores. *Arctic, Antarctic, and Alpine Research* 35, 218-232.
- Andrews, J.T., Jónsdóttir, I., Geirsdóttir, A., 2019. Tracking Holocene drift-ice limits on the NW/SW Iceland shelf: comparing proxy data with observation and historical evidence. *Arctic, Antarctic, and Alpine Research*. 51, 96-114.

- 697 Andrews, J.T., Kristjansdottir, G.B., Eberl, D.D., Jennings, A.E., 2013. A quantitative X-
 698 ray diffraction inventory of tephra and volcanic glass inputs into the Holocene
 699 marine sediment archives of Iceland: A contribution to V.A.S.T. Polar Research 1-
 700 15.
- 701 Andrews, J.T., Stein, R., Moros, M., Perner, K., 2016. Late Quaternary changes in
 702 sediment composition on the NE Greenland margin (~73 degrees N) with a focus on
 703 the fjords and shelf. *Boreas* 45, 381–397.
- 704 Andrews, J.T., Voelker, A., 2018. "Heinrich events" (& sediments): A history of
 705 terminology and recommendations for future usage. *Quaternary Science Reviews*
 706 187, 31-40.
- 707 Andrews, J.T., Vogt, C., 2014. Source to Sink: Statistical identification of regional
 708 variations in the mineralogy of surface sediments in the western Nordic Seas (58°N –
 709 75°N; 10° W -- 40°W). *Marine Geology* 357, 151-162.
- 710 Andrews, J.T., Vogt, C., 2020a. Variations in felsic- versus mafic-sources in the Western
 711 Nordic Seas during MIS 1 to MIS 4 *Marine Geology* 424, 106164.
- 712 Andrews, J.T. and Vogt, C. 2020b: Results of bulk sediment X-ray diffraction analysis and
 713 quantification of mineral phases based on the RockJock quantitative analysis. *Pangaea*.
 714 <https://doi.pangaea.de/10.1594/PANGAEA.923135>
- 715 Austin, W.E.N., Hibbert, F.D., 2012. Tracing time in the ocean: a brief review of
 716 chronological constraints (60-8 kyr) on North Atlantic marine event-based
 717 stratigraphies. *Quaternary Science Reviews* 36, 28-37.
- 718 Bai, Y., Chen, J.F., Sicre, M.-A., Jin, H., Ren, J., Li, H., Xue, B., Ji, Z., Zhuang, Y.,
 719 Klein, V., Zhao, M., 2019. Seasonal and spatial variability of sea ice and
 720 phytoplankton biomarker flux in the Chukchi Sea (Western Arctic). *Progress in*
 721 *Oceanography* 171, 22-37.
- 722 Bassis, J.N., Petersen, S.V. Cathles, L.M., 2017. Heinrich events triggered by ocean
 723 forcing and modulated by isostatic adjustment. *Nature* 542, 332-334.
- 724 Belt, S.T., 2018. Source-specific biomarkers as proxies for Arctic and Antarctic sea ice,
 725 *Organic Geochemistry* 125, 277–298, doi: 10.1016/j.orggeochem.2018.10.002.
- 726 Belt, S.T., Masse, G., Rowland, S.J., Poulin, M., Michel, C., LeBlanc, B., 2007. A novel
 727 chemical fossil of palaeo sea ice: IP₂₅. *Organic Geochemistry* 38, 16-27.

- 728 Belt, S.T., Müller, J., 2013. The Arctic sea ice biomarker IP₂₅: a review of current
 729 understanding, recommendations for future research and applications in palaeo sea
 730 ice reconstructions. *Quaternary Science Reviews* 79, 9-25.
 731
- 732 Belt, S.T., Cabedo-Sanz, P., Smik, L., Navarro-Rodriguez, A., Berben, S.M. P., Knies, J.,
 733 Husum, K., 2015. Identification of paleo Arctic winter sea ice limits and the
 734 marginal ice zone: optimised biomarker-based reconstructions of late Quaternary
 735 Arctic sea ice. *Earth and Planetary Science Letters* 431, 127-139.
- 736 Belt, S.T., Brown, T.A., Smik, L., Tatarek, A., Wiktor, J., Stowasser, G., Assmy, P.,
 737 Allen, C.A., Husum, K., 2017. Identification of C₂₅ highly branched isoprenoid
 738 (HBI) alkenes in diatoms of the genus *Rhizosolenia* in polar and non-polar marine
 739 phytoplankton. *Organic Geochemistry* 110, 65–72
- 740 Bendle, J., Rosell-Melé, A., 2004. Distributions of UK'₃₇ and UK₃₇ in the surface waters
 741 and sediments of the Nordic Seas: implications for paleoceanography. *Geochemistry,*
 742 *Geophysics, Geosystems*, Q11013. doi:10.1029/2004GC000741.
- 743 Bigg, G.R., 2016. Icebergs. Their Science and links to Global Change. Cambridge
 744 University Press.
- 745 Blaauw, M., 2012. Out of tune: the dangers of aligning proxy archives. *Quaternary Science*
 746 *Reviews* 36, 38-49.
- 747 Blaauw, M., Christen, J.A., Bacon Manual, 2016. -v2.2, p. 11 pp.
- 748 Blaauw, M., Christen, J.A., 2005. The problems of radiocarbon dating. *Science* 308,
 749 1552-1553.
- 750 Boers, N., Ghil, M., Rousseau, D.D., 2018. Ocean circulation, ice shelf, and sea ice
 751 interactions explain Dansgaard-Oeschger cycles. *Proceedings of the National*
 752 *Academy of Sciences of the United States of America* 115, E11005-E11014.
- 753 Boers, N., Goswami, B., Ghil, M., 2017. A complete representation of uncertainties in
 754 layer-counted paleoclimate archives. *Climate of the Past* 13, doi:10.5194/cp-13-
 755 1169-2017
- 756 Brakstad, A., Vage, K., Havik, L., Moore, G.W.K., 2019. Water Mass Transformation in the
 757 Greenland Sea during the Period 1986-2016. *Journal of Physical Oceanography* 49, 121-
 758 140.

- 759 Brendryen, J., Haflidason, H., Sejrup, H.P., 2011. Non-synchronous deposition of North
 760 Atlantic Ash Zone II in Greenland ice cores, and North Atlantic and Norwegian Sea
 761 sediments: an example of complex glacial-stage tephra transport. *Journal of*
 762 *Quaternary Science* 26, 739-745.
- 763 Brooks, C.K., Nielsen, T.F.D., 1982. The Phanerozoic development of the
 764 Kangerdlugssuaq area, East Greenland. *Meddelelser on Gronland, Geoscience* 9, 1-
 765 30.
- 766 Cabedo-Sanz, P., Belt, S.T., Jennings, A.E., Andrews, J.T., Geirsdóttir, Á., 2016.
 767 Variability in drift ice export from the Arctic Ocean to the North Icelandic Shelf over
 768 the last 8000 years: A multi-proxy evaluation. *Quaternary Science Reviews* 146, 99-
 769 115.
- 770 Chayes, F., 1971. Ratio correlation. University of Chicago Press, Chicago.
- 771 Clark, D.L., 1990a. Arctic Ocean ice cover; Geologic history and climatic significance,
 772 *The Geology of North America*. Geological Society of America, pp. 53-62.
- 773 Clark, D.L., 1990b. Stability of the Arctic Ocean ice-cover and Pleistocene warming
 774 events: Outlining the problem, in: Bleil, U., Thiede, J. (Eds.), *Geological History of*
 775 *the Polar Oceans: Arctic Versus Antarctic*. Kluwer Academic Publishers,
 776 Netherlands, pp. 273-287.
- 777 Clark, P. U., Hostetler, S. W., Pisias, N. G., Schmittner, A., and Meissner, K. J., 2007.
 778 Mechanisms for a ~7 kyr climate and sea-level oscillation during marine isotope
 779 stage 3. In Schmittner, A., Chiang, J., and Hemming, S. (eds.), *Ocean Circulation:*
 780 *Mechanisms and Impacts*. Geophysical Monograph 173. Washington, DC: AGU,
 781 pp. 209–246.
- 782 Conte, M. H., M.-A. Sicre, C. Rühlemann, J. C. Weber, S. Schulte, D. Schulz-Bull, T.
 783 Blanz, 2006. Global temperature calibration of the alkenone unsaturation index
 784 ($U^{K'_{37}}$) in surface waters and comparison with surface sediments, *Geochemistry,*
 785 *Geophysics, Geosystems* 7, Q02005, doi:10.1029/2005GC001054.
- 786 Darby, D.A., Andrews, J.T., Belt, S.T., Jennings, A.E., Cabedo-Sanz, P., 2017. Holocene
 787 cyclic records of ice-rafted debris and sea ice variations on the East Greenland and
 788 NW Iceland margins. *Antarctic, Arctic, and Alpine Research* 49, 649-672.

- 789 Darby, D.A., Zimmerman, P., 2008. Ice-rafted detritus events in the Arctic during the last
790 glacial interval, and the timing of the Innuitian and Laurentide ice sheet calving
791 events. *Polar Research* 27, 114-127.
- 792 Davies, S.M., Wastegard, S., Rasmussen, T.L., Svensson, A., Johnsen, S.J., Steffensen,
793 J.P., Andersen, K.K., 2008. Identification of the Fugloyarbanki tephra in the NGRIP
794 ice core: a key tie-point for marine and ice-core sequences during the last glacial
795 period. *Quaternary Science Reviews* 23, 409-414.
- 796 Divine, D.V., Dick, C., 2006. Historical variability of the sea ice edge position in the
797 Nordic Seas. *Journal of Geophysical Research* 111, 14pp.
798 doi:10.1029/2004JC002851.
- 799 Dokken, T.M., Nisancioglu, K.H., Li, C., Battisti, D.S., Kissel, C., 2013. Dansgaard-
800 Oeschger cycles: Interactions between ocean and sea ice intrinsic to the Nordic seas.
801 *Paleoceanography* 28, 491-502.
- 802 Domack, E.W., Jacobson, E.A., Shipp, S., Anderson, J.B., 1999. Late Pleistocene-
803 Holocene retreat of the West Antarctic Ice-Sheet system in the Ross Sea: Part 2 -
804 Sedimentologic and stratigraphic signature. *Geological Society of America Bulletin*
805 111, 1517-1536.
- 806 Dowdeswell, J.A., Elverhoi, A., Andrews, J.T., Hebbeln, D., 1999. Asynchronous de-
807 position of ice-rafted layers in the Nordic seas and North Atlantic Ocean. *Nature*
808 400, 348-351.
- 809 Eberl, D.D., 2003. User guide to RockJock: A program for determining quantitative
810 mineralogy from X-ray diffraction data. United States Geological Survey, Open File
811 Report 03-78, 40 pp, Washington, DC.
- 812 Elliot, M., Labeyrie, L., Dokken, T., Manthe, S., 2001. Coherent patterns of ice-rafted
813 debris deposited in the Nordic regions during the last glacial (10-60 ka). *Earth and*
814 *Planetary Science Letters* 194, 151-163.
- 815 Evans, J., Dowdeswell, J. A., Grobe, H., Niessen, F., Stein, R., Hubberten, H.-W. &
816 Whittington, R. J. 2002: Late Quaternary sedimentation in Kaiser Joseph Fjord and
817 the continental margin of East Greenland. In Dowdeswell, J. A. & O Cofaigh, C.
818 (eds.): *Glacier-Influenced Sedimentation on High-Latitude Continental Margins*,
819 *Special Publication* 203, 149-179. The Geological Society of London, London.

- 820 Evans, J., Dowdeswell, J. A., Grobe, H., Niessen, F., Stein, R., Hub-
 821 berten, H.-W. & Whittington, R. J. 2002: Late Quaternary sedi-
 822 mentation in Kejser Joseph Fjord and the continental margin of East
 823 Greenland. In Dowdeswell, J. A. & O Cofaigh, C. (eds.):
 824 Glacier-Influenced Sedimentation on High-Latitude Continental
 825 Margins, 149–179. The Geological Society of London, Special
 826 Publication 203, London.
- 827 Evans, J., O Cofaigh, C., Dowdeswell, J. A. & Wadhams, P. 2009:
 828 Marine geophysical evidence for former expansion and flow of the
 829 Greenland Ice Sheet across the north-east Greenland continental
 830 shelf. *Journal of Quaternary Science* 24, 279–293.
- 831 Evans, J., Dowdeswell, J. A., Grobe, H., Niessen, F., Stein, R., Hub-
 832 berten, H.-W. & Whittington, R. J. 2002: Late Quaternary sedi-
 833 mentation in Kejser Joseph Fjord and the continental margin of
 834 East Greenland. In Dowdeswell, J. A. & O Cofaigh, C. (eds.):
 835 Glacier-Influenced Sedimentation on High-Latitude Continental
 836 Margins, 149–179. The Geological Society of London, Special
 837 Publication 203, London.
- 838 Evans, J., O Cofaigh, C., Dowdeswell, J. A. & Wadhams, P. 2009:
 839 Marine geophysical evidence for former expansion and flow of the
 840 Greenland Ice Sheet across the north-east Greenland continental
 841 shelf. *Journal of Quaternary Science* 24, 279–293.
- 842 Evans, J., Dowdeswell, J. A., Grobe, H., Niessen, F., Stein, R., Hubberten, H.-W. &
 843 Whittington, R. J. 2002: Late Quaternary sedi- mentation in Kejser Joseph Fjord and
 844 the continental margin of East Greenland. In Dowdeswell, J. A. & O Cofaigh, C.
 845 (eds.): Glacier-Influenced Sedimentation on High-Latitude Continental Margins,
 846 149–179. The Geological Society of London, Special Publication 203, London.
- 847 Evans, J., O Cofaigh, C., Dowdeswell, J. A. & Wadhams, P. 2009: Marine geophysical
 848 evidence for former expansion and flow of the Greenland Ice Sheet across the north-
 849 east Greenland continental shelf. *Journal of Quaternary Science* 24, 279–293.

- 850 Funder, S., Goosse, H., Jepsen, H., Kaas, E., Kjaer, K.H., Korsgaard, N.J., Larsen, N.K.,
851 Linderson, H., Lysa, A., Moller, P., Olsen, J., Willerslev, E., 2011a. A 10,000-year
852 record of Arctic Ocean sea-ice variability-view from the beach. *Science*, 333, 747-
853 750.
- 854 Funder, S., Kjeldsen, K.K., Kjaer, K.H., O Cofaigh, C., 2011b. The Greenland Ice Sheet
855 during the past 300,000 years: A review. p. 699-713 In, Ehlers, J., Gibbard, P.L., and
856 Hughes, P.D., (Eds), *Quaternary Glaciations - Extent and Chronology: A Closer*
857 *Look*. Elsevier, Amsterdam
- 858 Greco, M., Jonkers, L., Kretschmer, K., Bijma, J., Kucera, M., 2019. Depth habitat of the
859 planktonic foraminifera *Neoglobobulimina pachyderma* in the northern high
860 latitudes explained by sea-ice and chlorophyll concentrations. *Biogeosciences* 16,
861 3425-3437.
- 862 Hammer, O, Harper, D.A.T., and Ryan, P.D., 2001. PAST: Paleontological statistics
863 software package for education and data analysis. *Palaeontologia Electronica*,
864 4.1.4:1-9. http://palaeo-electronica.org/2001_1/past/issue1_01.htm
- 865 Harden, B.E., Pickart, R.S., Valdimarsson, H., Vage, K., de Steur, L., Richards, C., Bahr,
866 F., Torres, D., Borve, E., Jonsson, S., Macrandar, A., Osterhus, S., Havik, L.,
867 Hattermann, T., 2016. Upstream sources of the Denmark Strait Overflow:
868 Observations from a high-resolution mooring array. *Deep-Sea Research Part I-*
869 *Oceanographic Research Papers* 112, 94-112
- 870 Hassani, H., 2007. Singular Spectrum Analysis: Methodology and comparison. *Journal of*
871 *Data Science* 5, 239-257.
- 872 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast
873 Atlantic Ocean during the past 130,000 years. *Quaternary Research* 29, 143-152.
- 874 Hemming, S.R., 2004. Heinrich Events: Massive late Pleistocene detritus layers of the
875 North Atlantic and their global climate imprint. *Reviews of Geophysics* 42,
876 RG1005/2004.
- 877 Henriksen, H., 2008. Geological history of Greenland. Geological Survey of Denmark
878 and Greenland, Copenhagen.
- 879 Hesse, R., 2016. Ice-proximal Labrador Sea Heinrich layers: a sedimentological
880 approach. *Canadian Journal of Earth Sciences* 53, 71-100.

- 881 Higgins, A.K., Gilotti, J.A., Smith, P.M., 2008. The Greenland Caledonides. Evolution of
882 the Northeast margin of Laurentia. Geological Society of America, Boulder, CO, p.
883 368.
- 884 Hoff, U., Rasmussen, T.L., Stein, R., Ezat, M.M., Fahl, K., 2016. Sea ice and millennial-
885 scale climate variability in the Nordic seas 90 kyr ago to present. *Nature*
886 *Communications* 7. 12247.
- 887 Howe, J.N.W., Piotrowski, A.M., Noble, T.L., Mulitza, S., Chiessi, C.M., Bayon, G.,
888 2016. North Atlantic Deep Water production during the Last Glacial Maximum.
889 *Nature Commun.* 7, 11765. doi: 10.1038/ncomms11765
- 890 Jennings, A.E., Andrews, J.T. et al., 2018. Baffin Bay paleoenvironments in the LGM
891 and HS1: Resolving the ice-shelf question. *Marine Geology.* 402, 5-16.
- 892 Jennings, A.E., Reilly, B., Andrews, J.T., Hogen, K., Walczak, M., Stoner, J., Mix, A.C.,
893 Jakobsson, M., 2019. Modern ice shelf facies and Early Holocene counterparts in
894 Petermann Fjord and Northern Nares Strait. (Abstract) International Association of
895 Sedimentologists IAS, Rome.
- 896 Jonsson, S., and Valdimarsson, H. 2004. A new path for the Denmark Strait overflow
897 water from the Iceland Sea to Denmark Strait. *Geophysical Research Letters* 31, 4pp.
898 doi:10.1029/2003GL019214, 012004.
- 899 Jonsson, S., Valdimarsson, H., 2005. The flow of Atlantic water to the North Icelandic
900 Shelf and its relation to the drift of cod larvae. *ICES Journal of Marine Science* 62,
901 1350-1359.
- 902 Jonsson, S., Briem, J., 2003. Flow of Atlantic water west of Iceland and onto the north
903 Atlantic shelf. *ICES Marine Science Symposia* 219, 326-328.
- 904 Keigwin, L.D., and Swift, S.A., 2017. Carbon isotope evidence for a northern source of
905 deep water in the glacial western North Atlantic. *Proceedings of the National*
906 *Academy of Sciences of the United States of America* 114, 2831-2835.
- 907 Knudsen, K.-L., Eiriksson, J., 2002. Application of tephrochronology to the timing and
908 correlation of palaeoceanographic events recorded in Holocene and Late Glacial
909 shelf sediments off North Iceland. *Marine Geology* 191, 165-188.

- 910 Konert, M., Vandenberghe, J., 1997. Comparison of laser grain size analysis with pipette
 911 and sieve analysis: a solution for the underestimation of the clay fraction.
 912 *Sedimentology* 44, 523-535.
- 913 Knudsen, K.-L., Jiang, D., Jansen, E., Eiriksson, J., Heinemeier, J., Seidenkrantz, M.-S.,
 914 2003. Environmental changes off North Iceland during the deglaciation and the
 915 Holocene: foraminifera, diatoms and stable isotopes. *Marine Micropaleontology* 953,
 916 1-33.
- 917 Knutz, P.C., H. Ebbesen, S. Christiansen, M.-A. Sicre, and A. Kuijpers, 2011. The triple
 918 stage deglacial retreat of the southern Greenland Ice Sheet driven steps by vigorous
 919 Irminger Current, and its significance for the Younger Dryas cooling,
 920 *Paleoceanography*, 26, PA3204, doi:10.1029/2010PA002053, 2011
- 921 Kristjansdottir, G.B., Stoner, J.S., Gronvold, K., Andrews, J.T., Jennings, A.E., 2007.
 922 Geochemistry of Holocene cryptotephra from the North Iceland Shelf (MD99-
 923 2269): Intercalibration with radiocarbon and paleomagnetic chronostratigraphies.
 924 *The Holocene* 17, 155-176.
- 925 Kristjansson, L., Saemundsson, K., Thorarinsson, S., Saemundsson, K., Thorarinsson, S.,
 926 Einarsson, P., Bjornsson, S., Simonarson, L., Fridleifsson, I., Jaksoobsson, S.P.,
 927 Bjornsson, H., 1979. Special Issue: Geology of Iceland. *Jökull* 29, 1-101.
- 928 Labeyrie, L., Jansen, E., Cortijo, E., 2003. Les rapports de campagnes a la mer
 929 MD114/IMAGES V. Institut Polaire Francais Paul-Emile Victor, Brest.
- 930 Labeyrie, L.D. and Cortijo, E. 2005: Physical properties of sediment core MD99-2274.
 931 *Pangaea*. <https://doi.org/10.1594/PANGAEA.253605>
- 932 Lakeman, T.R., Pienkowski, A.J., Nixon, F.C., Furze, M.F.A., Blasco, S., Andrews, J.T.,
 933 King, E.L., 2018. Collapse of a marine-based ice stream during the early Younger
 934 Dryas chronozone, western Canadian Arctic. *Geology* 46, 211-214.
- 935 Larsen, N.K., Levy, L.B., Carlson, A.E., Buizert, C., Olsen, J., Strunk, A., Bjork, A.A. and Skov,
 936 D.S. (2018) Instability of the Northeast Greenland Ice Stream over the last 45,000 years.
 937 *Nature Communications*, 9.doi: 10.1038/s41467-018-04312-7
- 938 Lind, E.W., Lilja, C., Wastegard, S., Pearce, N.J.G., 2016. Revisiting the Borrobol
 939 Tephra. *Boreas* 45, 693-643.

- 940 Liu, Y., Hallberg, R., Sergienko, O., Samuels, B.L., Harrison, M., & Oppenheimer, M.
 941 2018. Climate response to the meltwater runoff from Greenland ice sheet: evolving
 942 sensitivity to discharging locations. *Climate Dynamics*, 51, 1733–1751. doi
 943 10.1007/s00382-017-3980-7
- 944 Lohne, O.S., Mangerud, J., Birks, H.H., 2013. Precise C-14 ages of the Vedde and
 945 Saksunarvatn ashes and the Younger Dryas boundaries from western Norway and
 946 their comparison with the Greenland Ice Core (GICC05) chronology. *Journal of*
 947 *Quaternary Science* 28, 490-500.
- 948 Maffezzoli, N., Vallelonga, P., Edwards, R., Saiz-Lopez, A., Turetta, C., Kjaer, H.A.,
 949 Barbante, C., Vinther, B., Spolaor, A., 2019. A 120 000-year record of sea ice in the
 950 North Atlantic? *Climate of the Past* 15, 2031-2051.
- 951 Manley, W.F., Jennings, A.E., 1996. Radiocarbon Date List VIII: Eastern Canadian
 952 Arctic, Labrador, Northern Quebec, East Greenland Shelf, Iceland Shelf, and
 953 Antarctica. INSTAAR, University of Colorado, p. 163 pp.
- 954 Marcott, S.A., Clark, P.U., Padman, L., Klinkhammer, G.P., Springer, S.R. , Liu, Z., ,
 955 Otto-Bliesner, B.L., Carlson, A.E., Ungerer, A., Padman, J., He, F., Cheng, J. and
 956 Schmittner, A., 2011: Ice-shelf collapse from subsurface warming as a trigger for
 957 Heinrich events. *Proceedings of the National Academy of Sciences of the United*
 958 *States of America* 108, 13415-13419
- 959 Marshall, N.R., Piper, D.J.W., Saint-Ange, F., Campbell, D.C., 2014. Late Quaternary
 960 history of contourite drifts and variations in Labrador Current flow, Flemish Pass,
 961 offshore eastern Canada. *Geology Marine Letters* 34, 457-470. doi:10.1007/s00367-
 962 014-0377-z.
- 963 Mauritzen, C., 1996. Production of dense overflow waters feeding the North Atlantic
 964 across the Greenland-Scotland Ridge. 1. Evidence for a revised circulation scheme.
 965 *Deep-Sea Research I* 43, 769-806.
- 966 Massé, G., Rowland, S.J., Sicre, M.-A., Jacob, J., Jansen, E., Belt, S.T., 2008. Abrupt
 967 climate changes for Iceland during the last millennium: Evidence from high
 968 resolution sea ice reconstructions. *Earth and Planetary Science Letters* 269, 565–569.
- 969 Matthews, I.P., Birks, H.H., Bourne, A.J., Brooks, S.J., Lowe, J.J., Macleod, A., Pyne-
 970 O'Donnell, S.D.F., 2011. New age estimates and climatostratigraphic correlations for

- 971 the Borrobol and Penifiler Tephra: evidence from Abernethy Forest, Scotland.
 972 *Journal of Quaternary Science* 26, 247-252.
- 973 McCave, I.N., Andrews, J.T., 2019a. Distinguishing current effects in sediments
 974 delivered to the ocean by ice. I. Principles, methods and examples. *Quaternary*
 975 *Science Reviews* 212, 92-107.
- 976 McCave, I.N., Andrews, J.T., 2019b. Distinguishing current effects in sediments
 977 delivered to the ocean by ice. II. Glacial to Holocene changes in North Atlantic high
 978 latitude upper ocean flows. *Quaternary Science Reviews* 223, no. 105902, 21pp.
- 979 McCave, I.N., Hall, I.R., Bianchi, G.G., 2006. Laser vs settling velocity differences in silt
 980 grainsize measurements: estimation of palaeocurrent vigour. *Sedimentology* 53, 919-
 981 928.
- 982 McCave, I.N., Manighetti, B. and Robinson, S.G., 1995. Sortable silt and fine sediment
 983 size/composition slicing: parameters for palaeocurrent speed and palaeoceanography.
 984 *Paleoceanography* 10, 593-610.
- 985 McCave, I.N., Thornalley, D.J.R., Hall, I.R., 2017. Relation of sortable silt grain-size to
 986 deep-sea current speeds: Calibration of the 'Mud Current Meter'. *Deep-Sea Research*
 987 *Part I* 127, 1-12.
- 988 McCave, I.N., Syvitski, J.P.M., 1991. Principles and methods of geological particle size
 989 analysis, in: Syvitski, J.P.M. (Ed.), *Principles, methods and application of particle*
 990 *size analysis*. Cambridge University Press, pp. 3-21.
- 991 McKay, R., Golledge, N.R., Maas, S., Naish, T., Levy, R., Dunbar, G., Kuhn, G., 2016.
 992 Antarctic marine ice-sheet retreat in the Ross Sea during the early Holocene.
 993 *Geology* 44, 7-10.
- 994 Millo, C., Sarnthein, M., Erlenkeuser, H., Frederichs, T., 2005. Methane-driven Late
 995 Pleistocene delta C-13 minima and overflow reversals in the southwestern Greenland
 996 Sea. *Geology* 33, 873–876.
- 997 Moles, J.D., McGarvie, D., Stevenson, J.A., Sherlock, S.C., Abbott, P.M., Jenner, F.E.,
 998 Halton, A.M., 2019. Widespread tephra dispersal and ignimbrite emplacement from
 999 a subglacial volcano (Torfajökull, Iceland). *Geology* 47, 577-580.
- 1000 Moros, M., McManus, J., Rasmussen, T., Kuijpers, A., Dokken, T., Snowball, I., Nielsen,
 1001 T., Jansen, E., 2004. Quartz content and the quartz-to-plagioclase ratio determined

- 1002 by X-ray diffraction: a proxy for ice rafting in the northern North Atlantic? *Earth and*
 1003 *Planetary Science Letters* 218, 389-401.
- 1004 Newton, A.M.W., Huuse, M., and Brocklehurst, S.H. 2018. A persistent Norwegian
 1005 Atlantic Current through the Pleistocene glacials. *Geophysical Research Letters* 45,
 1006 5599–5608. <https://doi.org/10.1029/2018GL077819>
- 1007 Nam, S.-I., Stein, R., Grobe, H., Hubberten, H., 1995. Late Quaternary
 1008 glacial/interglacial changes in sediment composition at the East Greenland
 1009 continental margin and their paleoceanographic implications. *Marine Geology* 122,
 1010 243-262.
- 1011 Norðdahl, H., Ingolfsson, O., 2015. Collapse of the Icelandic ice sheet controlled by sea-
 1012 level rise? *Arktos*, 1 pp 1-18.
- 1013 O'Cofaigh, C., Taylor, J., Dowdeswell, J.A., Rosell-Mele, A., Kenyon, N.H., Evans, J.,
 1014 Mienert, J., 2002. Geological evidence for sediment reworking on high-latitude
 1015 continental margins and its implications for palaeoceanography: insights from the
 1016 Norwegian– Greenland Sea. In: Dowdeswell, J.A., Ó Cofaigh, C. (Eds.), *Glacier-*
 1017 *influenced sedimentation on high-latitude continental margins*. Geological Society
 1018 London Special Paper, 20. Geological Society, London, pp. 325–348.
- 1019 Paillard, D., Labeyrie, L., Yiou, P., 1996. Macintosh program performs time-series
 1020 analysis. *EOS* 77, 379.
- 1021 Patton, H., Hubbard, A., Bradwell, T., Schomacker, A., 2017. The configuration,
 1022 sensitivity and rapid retreat of the Late Weischelian Icelandic Ice Sheet. *Earth-*
 1023 *Science Reviews* 166, 223-245.
- 1024 Petersen, S.V., Schrag, D.P., Clark, P.U., 2013. A new mechanism for Dansgaard-
 1025 Oeschger cycles. *Paleoceanography* 28, 24-30.
- 1026 Phillips, R.L., Grantz, A., 2001. Regional variations in provenance and abundance of ice-
 1027 rafted clasts in Arctic Ocean sediments: implications for the configuration of late
 1028 Quaternary oceanic and atmospheric circulation in the Arctic. *Marine Geology* 172,
 1029 91-115.
- 1030 Pickart, R.S., Torres, D.J., Fratantoni, P.S., 2005. the East Greenland spill jet. *Journal of*
 1031 *Physical Oceanography* 35, 1037-1053.

- 1032 Prahl, F.G., Muelhausen, L.A., Zahnle, D.L., 1988, Further evaluation of long-chain
1033 alkenones as indicators of paleoceanographic conditions. *Geochimica et*
1034 *Cosmochmicia Acta* 52, 2303-2310.
- 1035 Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B.,
1036 Cvijanovic, I., Dahl-Jensen, D., Johnsen, S.J., Fischer, H., Gkinis, V., Guillevic, M.,
1037 Hoek, W.Z., Lowe, J.J., Pedro, J.B., Popp, T., Seierstad, I.K., Steffensen, J.P.,
1038 Svensson, A.M., Vallelonga, P., Vinther, B.M., Walker, M.J.C., Wheatley, J.J.,
1039 Winstrup, M., 2014. A stratigraphic framework for abrupt climatic changes during
1040 the Last Glacial period based on three synchronized Greenland ice-core records:
1041 refining and extending the INTIMATE event stratigraphy. *Quaternary Science*
1042 *Reviews* 106, 14-28.
- 1043 Rasmussen, T.L., Wastegard, S., Kuijpers, A., van Weering, T.C.E., Heinemeier, J.,
1044 Thomsen, E., 2003. Stratigraphy and distribution of tephra layers in marine sediment
1045 cores from the Faeroe Islands, North Atlantic. *Marine Geology* 199, 263-277.
- 1046 Reeh, N., 2004. Holocene climate and fjord glaciations in Northeast Greenland:
1047 implications for IRD deposition in the North Atlantic. *Sedimentary Geology* 165,
1048 333-342.
- 1049 Reeh, N., Mayer, C., Miller, H., Thomsen, H.H., Weidick, A., 1999. Present and past
1050 climate control on fjord glaciations in Greenland: Implications for IRD-deposition in
1051 the sea. *Geophysical Research Letters* 26, 1039-1042.
- 1052 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Grootes,
1053 P.M., Guilderson, T.P., Hafliðason, H., Hajdas, I., Hatt?, C., Heaton, T.J., Hoffmann, D.L.,
1054 Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer,
1055 R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M. and van der
1056 Plicht, J. (2013) IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0-
1057 50,000 Years cal BP. *Radiocarbon*, 55, 1869-1887.
- 1058 Robinson, S.G., Maslin, M.A., McCave, I.N., 1995. Magnetic susceptibility variations in
1059 Upper Pleistocene deep-sea sediments of the N.E. Atlantic: Implications for ice
1060 rafting and palaeocirculation at the Last Glacial Maximum. *Paleoceanography* 10,
1061 221-250.

- 1062 Roesch, A. and Schmidbauer, H.W.C., 2018: WaveletComp: Computational Wavelet
1063 Analysis. R package 1.1. <https://CRAN.R-project.org/package=Wavelet.Comp>.
- 1064 Ruddiman, W.F., McIntyre, A., 1981. The North Atlantic Ocean during the last
1065 deglaciation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 35, 145-214.
- 1066 Sarnthein, M., Winn, K., Jung, S., Duplessy, J. C., Labeyrie, L., Erlenkeuser, H., and
1067 Ganssen, G., 1994. Changes in East Atlantic deepwater circulation over the last
1068 30,000 years – An eight-time-slice record. *Paleoceanography* 9, 209–267.
- 1069 Sarnthein, M., Pflaumann, U., Weinelt, M., 2003. Past extent of sea ice in the northern
1070 North Atlantic inferred from foraminiferal paleotemperature estimates.
1071 *Paleoceanography* 18, doi:10.1029/2002PA000771.
- 1072 Schlitzer, R., 2011. Ocean Data View, <http://odv.awi.de>, 2011
- 1073 Sejrup, H.P., Sioholm, J., Furnes, H., Beyer, I., Eide, L., Jansen, E., Mangerud, J., 1989.
1074 Quaternary tephrochronology on the Iceland Plateau, north of Iceland. *Journal of*
1075 *Quaternary Science* 4, 109-114.
- 1076 Shaffer, G., S.M. Olsen, and C.J. Bjerrum, 2004. Ocean subsurface warming as a
1077 mechanism for coupling Dansgaard-Oeschger climate cycles and ice-rafting events.
1078 *Geophysical Research Letters* 31, L24202, doi:10.1029/2004GL020968.
- 1079 Sicre, M.-A., Bard, E., Ezat, U., Rostek, F., 2002. Alkenone distributions in the North
1080 Atlantic and Nordic sea surface waters. *Geochemistry, Geophysics, Geosystems*,
1081 Article 2001GC000159.
- 1082 Sicre, M.-A., Labeyrie, L., Ezat, U., Duprat, J., Turon, J-L, Schmidt, S., Michel, E.,
1083 Mazaud A., 2005. Southern Indian Ocean response to Northern Hemisphere Heinrich
1084 events. *Earth and Planetary Science Letters* 240, 724-731, doi: 10.1016.
- 1085 Sicre, M.-A., Jacob, J., Ezat, U., Rousse, S., Kissel, C., Yiou, P., Eiriksson, J., Knudsen,
1086 K.-L., Jansen, E., Turon, J.-L., 2008a. Decadal variability of sea surface temperatures
1087 off North Iceland over the last 2000 years. *Earth and Planetary Science Letters* 268,
1088 137-142.
- 1089 Sicre, M.-A., Yiou, P., Eiriksson, J., Ezat, U., Guimbaut, E., Dahhaoui, I., Knudsen, K.-
1090 L., Jansen, E., Turon, J.-L., 2008b. A 4500-year reconstruction of sea surface
1091 temperature variability at decadal time scales off North Iceland. *Quaternary Science*
1092 *Reviews* 27, 2041-2047.

- 1093 Simon, Q., Hillaire-Marcel, C., St-Onge, G., Andrews, J.T., 2014. North-eastern
1094 Laurentide, western Greenland and southern Inuitian ice stream dynamics during
1095 the last glacial cycle. *Journal of Quaternary Science* 29, 14-26.
- 1096 Skinner L.C. & I.N. McCave. Analysis and modelling of the behaviour of gravity and
1097 piston corers based on soil mechanical principles. *Marine Geology* 199, 181-204.
- 1098 Skinner, L. C., Muschitiello, F., & Scrivner, A. E., 2019. Marine reservoir age variability
1099 over the last deglaciation: Implications for marine carbon cycling and prospects for
1100 regional radiocarbon calibrations. *Paleoceanography and Paleoclimatology* 34,
1101 1807–1815. <https://doi.org/10.1029/2019PA003667>
- 1102 Stefansson, U., 1962. North Icelandic Waters. *Rit Fiskideildar III*. Bind, Vol 3, 269.
- 1103 Stein, R., 2008. Arctic Ocean Sediments. Processes, Proxies, and Paleoenvironment.
1104 Elsevier, New York.
- 1105 Stein, R., Nam, S.-I., Grobe, H., Hubberten, H., 1996a. Late Quaternary glacial history
1106 and short-term ice-rafted debris fluctuations along the East Greenland continental
1107 margin, in: Andrews, J.T., Austin, W.A., Bergsten, H., Jennings, A.E. (Eds.), *Late*
1108 *Quaternary paleoceanography of North Atlantic margins*. Geological Society,
1109 London, pp. 135-151.
- 1110 Stein, R., Nam, S.-Il., Grobe, H., Hubberten, Hans-Wolfgang, H., 1996b. Sedimentology
1111 and stable isotope ratios of cores from the East Greenland continental
1112 margin. *PANGAEA*, <https://doi.org/10.1594/PANGAEA.733965>,
- 1113 Stokes, C.R., Clark, C.D., Darby, D.A., Hodgson, D.A., 2005. Late Pleistocene ice export
1114 events into the Arctic Ocean from the M'Clure Strait Ice Stream, Canadian Arctic
1115 Archipelago. *Global and Planetary Change* 49, 139-162.
- 1116 Telford, R.J., Heegaard, E., Birks, H.J.B., 2003. All age-depth models are wrong: but
1117 how badly? *Quaternary Science Reviews* 23, 1-5.
- 1118 Trachsel, M., Telford, R.J., 2017 All age-depth models are wrong, but are getting better.
1119 *Holocene* 27, 860-869
- 1120 van Kreveld, S., Sarthein, M., Erlenkeuser, H., Grootes, P., Jung, S., Nadeau, M.J.,
1121 Pflaumann, U., Voelker, A., 2000. Potential links between surging ice sheets,
1122 circulation changes, and the Dansgaard-Oeschger cycles in the Irminger Sea, 60-18
1123 ka. *Paleoceanography* 15, 425-442.

- 1124 Vasskog, K., Langebroek, P.M., Andrews, J.T., Nilsen, J.E.O., Nesje, A., 2015. The
 1125 Greenland Ice Sheet during the last glacial cycle: Current ice loss and contribution to
 1126 sea-level rise from a palaeoclimatic perspective. *Earth-Science Reviews* 150, 45-67.
- 1127 Velay-Vitow, J., Peltier, W.R., and Stuhne, G. , 2019. Tides of the Glacial Ocean and
 1128 their role in Heinrich Event instability. *Geophysical Research Abstracts*. 21,
 1129 EGU2019-3733.
- 1130 Venkatesh, S., Murphy, D.L., Wright, G.F., 1994. On the deterioration of icebergs in the
 1131 marginal ice-zone. *Atmosphere-Ocean* 32, 469-484.
- 1132 Verplanck, E.P., Farmer, G.L., Andrews, J., Dunhill, G., Millo, C., 2009. Provenance of
 1133 Quaternary glacial and glacialmarine sediments along the southeast Greenland margin.
 1134 *Earth and Planetary Science Letters* 286, 52-62.
- 1135 Voelker, A.H.L., 1999. Zur Deutung der Dansgaard-Oeschger Ereignisse in ultra-
 1136 hochauflösenden Sedimentprofilen aus dem Europäischen Nordmeer, Dansgaard-
 1137 Oeschger events in ultra-high resolution sediment records from the Nordic Seas.
 1138 Universität Kiel, Kiel, p. 271.
- 1139 Voelker, A.H.L., Grootes, P.M., Nadeau, M-J., and Sarnthein, M., 2000. Radiocarbon
 1140 levels in the Iceland Sea from 25–53 kyr and their link to the earth’s magnetic field
 1141 intensity. *Radiocarbon* 42, p 437–452
- 1142 Voelker, A.H.L., Haflidason, H., 2015. Refining the Icelandic tephrochronology of the
 1143 last glacial period - The deep-sea core P52644 record from the southern Greenland
 1144 Sea. *Global and Planetary Change* 131, 35-62.
- 1145 Voelker, A.H.L., Sarnthein, M., Grootes, P.M., Erlenkeuser, H., Laj, C., Mazaud, A.,
 1146 Nadeau, M-J., Schleicher, M., 1998: Correlation of marine ¹⁴C ages from the Nordic
 1147 Seas with the GISP2 isotope record: implications for ¹⁴C calibration beyond 25 ka
 1148 BP. *Radiocarbon* 40, 517-534.
- 1149 Volkman, J.K., 1986. A review of sterol markers for marine and terrigenous organic
 1150 matter. *Organic Geochemistry* 9, 83–99.
- 1151 Vogt, C., 2017. Bulk mineral assemblage via Quantitative Phase Analysis with X-ray
 1152 diffraction of sediment core PS2644-5 KAL. PANGAEA.
 1153 <https://doi.org/10.1594/PANGAEA.875919>

- 1154 Walczuk, M.H. & 13 others, 2020. Phasing of millennial-scale climate variability in the
1155 Pacific and Atlantic oceans. *Science*, 370, 716–720.
- 1156 Watkins, S.J., Maher, B.A., 2003. Magnetic characterization of present-day deep-sea
1157 sediments and sources in the North Atlantic. *Earth and Planetary Science Letters*
1158 214, 379-394.
- 1159 White, L.F., Bailey, I., Foster, G.L., Allen, G., Kelley, S.P., Andrews, J.T., Hogan, K.,
1160 Dowdeswell, J.A., Storey, C.D., 2016. Tracking the provenance of Greenland-
1161 sourced, Holocene aged, individual sand-sized ice-rafted debris using the Pb-isotope
1162 compositions of feldspars and Ar-40/Ar-39 ages of hornblendes. *Earth and Planetary*
1163 *Science Letters* 433, 192-203.
- 1164 Xiao, X., Fahl, K., Stein, R., 2013. Biomarker distributions in surface sediments from the
1165 Kara and Laptev seas (Arctic Ocean): indicators for organic-carbon sources and sea-
1166 ice coverage. *Quaternary Science Reviews* 79, 40–52.
- 1167 Xiao, X., Zhao, M., Knudsen, K.L., Sha, L., Eiríksson, J., Gudmundsdóttir, E., Jiang, H.,
1168 Guo, Z., 2017. Deglacial and Holocene sea–ice variability north of Iceland and
1169 response to ocean circulation changes. *Earth and Planetary Science Letters* 472, 14–
1170 24.
- 1171 Zou, H., 2016. An X-Ray Diffraction Approach: Bulk Mineral Assemblages as
1172 Provenance Indicator of Sediments from the Arctic Ocean. University of Bremen,
1173 Bremen, pp. 116. <https://elib.suub.uni-bremen.de/edocs/00105354-1.pdf>.
1174
1175
1176

Methods

Magnetic susceptibility: Magnetic susceptibility was measured on-board the *Marion Dufresne* (Labeyrie and Cort, 2005) in 2-cm increments (hence ~150yr sampling on average). Measurements were taken on the 1.5 m core sections. In this area of Iceland, the marine deposits are strongly affected by erosion and transport of basalt, which results in very high values of magnetic susceptibility. The export of sediments from the erosion of bedrock with much lower magnetic susceptibilities, such as granites and other felsic-rich bedrock in NE Greenland and from more distant sources (Verplanck et al., 2009; White et al., 2016) will lower the magnetic susceptibility readings. It is important to note that although magnetic susceptibility is straightforward to measure, data interpretation is complex, being a product of sediment density, grain-size, and mineralogy (Robinson et al., 1995; Stoner and Andrews, 1999; Watkins and Maher, 2003).

Quantitative X-ray Diffraction (qXRD): The weight % (wt%) of the non-clay and clay mineral composition of the < 2 mm sediment fractions is based on the US Geological Survey method (Eberl, 2003), which has been used extensively in this region (e.g. Andrews et al., 2017; Andrews and Eberl, 2007; Andrews and Vogt, 2014). One gram of sediment (dry weight) is spiked with 0.111 g of zincite, prepared (Eberl, 2003), run in the X-ray diffractometer, and the resulting intensity data processed in the Excel macro-program Rockjock v6. We investigate the wt% and presence/absence of 34 minerals and reduced this number by combining individual mineral wt% into larger groups, such as k-feldspars, plagioclase, dolomite, and amorphous minerals. Importantly in the context of this paper we had earlier shown that qXRD can recognize the presence

1200 of tephra and volcanic glass, with some ability to distinguish between basaltic and
1201 rhyolitic glass (Andrews et al., 2013).

1202 To gain a better understanding of possible changes in the provenance of the
1203 mineral compositions we processed the mineral wt% data in a sediment unmixing
1204 program “SedUnMix” (Andrews and Eberl, 2012). Two models were considered, the first
1205 with qXRD results from #2274 with four appropriate bedrocks, namely: basalt, dolerite,
1206 gneiss, and granite; and secondly with the mineral compositions of glacial marine
1207 sediment samples from potential source areas, namely: NE Greenland, E. Greenland, and
1208 Iceland (Suppl. Table of bedrock and marine sediment sources). The program calculates a
1209 “degree of fit” and also derives error estimates on each source within a sample. Ideally,
1210 the sum of the sources should equal 100% but marked deviations from this suggest that
1211 one or more sources have not been included, and/or that the sources are not representative
1212 of the sediment samples.

1213 **Grain-size:** Sediment was wet-sieved at 2 mm and the grain-size volume
1214 percentages in 96 intervals between 0.01 and 2000 μm were obtained via a Malvern laser
1215 system. Comparisons between the Malvern and other grain-size systems have been
1216 documented and found comparable (McCave et al., 2006; McCave and Syvitski, 1991).
1217 However, the objections of McCave et al. (2006) to laser sizers on the grounds of grain
1218 shape (Konert and Vandenberghe, 1997) are not valid for equant grains such as those
1219 produced by glacial grinding, as pointed out by Piper (Marshall et al., 2014), and thus
1220 size data are believed valid in the setting of MD2274. Grain-size curves have provided
1221 vital information on sediment transport and deposition in this region, and methods have
1222 been developed to reconstruct variations in bottom current speed for sediments delivered

to the ocean from dominantly glacial sources (McCave and Andrews, 2019a, b) The calibration of sortable silt mean (mean of 10-63 μm), a sensitivity, by McCave et al., (2017) has been applied to changes in the grainsize record.

Biomarkers: Biomarkers were extracted from freeze-dried subsamples (~2-4 g). Prior to extraction, samples were spiked with an internal standard (9-octylheptadec-8-ene, 9-OHD, 10 μL ; 10 $\mu\text{g mL}^{-1}$) to permit quantification of the highly branched isoprenoid (HBI) biomarkers IP₂₅, HBI II and HBI III. 5 α -androstan-3 β -ol; (0.1 μg) was also added to permit quantification of brassicasterol in some cases. Samples were then saponified in a methanolic KOH solution (~5 mL H₂O:MeOH (1:9); 5% KOH) for 60 min (70 °C). Hexane (3 $\times$ 2 mL) was added to the saponified mixtures, with supernatant solutions, containing non-saponifiable lipids (NSLs), transferred by glass pipettes to glass vials, and solvent removed using a gentle stream of N₂. Dried NSLs were re-suspended in hexane (0.5 mL) and fractionated using column chromatography (SiO₂; 0.5 g). Non-polar lipids, including IP₂₅ and HBI II, were eluted with hexane (6 mL), while more polar lipid fractions containing alkenones were eluted with MeOH (6 mL). For a few horizons, additional NSLs were fractionated to yield non-polar (hexane; 6 mL) and polar fractions containing sterols (hexane:methyl acetate 4:1; 6 mL). Each non-polar fraction was further purified to remove saturated components using silver-ion chromatography (Belt et al., 2015), with saturated compounds eluted with hexane (2 mL) and unsaturated compounds, including IP₂₅ and other HBIs, collected in a subsequent acetone fraction (3 mL). Analysis of fractions containing IP₂₅ and other HBIs was carried out using gas chromatography–mass spectrometry (GC–MS) following the methods and operating conditions described previously (Belt et al., 2012). Mass spectrometric analysis was

carried out in total ion current (TIC) and selected ion monitoring (SIM) modes. The identification of IP₂₅ and HBI II was based on their characteristic GC retention indices (e.g. RI_{HP5MS} = 2081, 2082 and 2044 for IP₂₅, HBI II and HBI III, respectively) and mass spectra (Belt, 2018). Quantification of all HBIs was achieved by comparison of mass spectral responses of selected ions (e.g. IP₂₅, *m/z* 350; HBI II, *m/z* 348; HBI III, *m/z* 346) in SIM mode with those of the internal standard (9-OHD, *m/z* 350) and normalized according to their respective instrumental response factors, derived from solutions of known biomarker concentration, and sediment masses (Belt et al., 2012). Fractions containing sterols were derivatized with N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA; 100 µL; 70°C for 60 min) immediately prior to analysis by GC–MS. Sterols were identified by comparison with GC–MS responses compared to those of standards. Sterol quantification was achieved as per the approach described above for HBIs.

Polar fractions containing alkenones obtained from elution with MeOH (6 mL) were further purified with 2 mL of hexane:methyl acetate (95:5 v/v) and 2 mL of hexane:methyl acetate (90:10 v/v). Alkenones were analyzed using a Thermo Trace GC Ultra gas chromatograph equipped with a CPSil5 capillary column (50m length, 0.32 i.d. and 0.25 mm film thickness), an FID detector and a septum programmable injector (SPI). Helium was used as carrier gas. 5α-cholestane was added as an external standard prior to GC injection. SST estimates were determined using the following equation (Prahl et al., 1988).

$$U_{37}^{K'} = \frac{C_{37:2}}{C_{37:2} + C_{37:3}} = 0.034 T + 0.039$$

References for Methods

- Andrews, J.T., Dunhill, G., Vogt, C., Voelker, A.H.L., 2017. Denmark Strait during the Late Glacial Maximum and Marine Isotope Stage 3: Sediment sources and transport processes. *Marine Geology* 390, 181-198.
- Andrews, J.T., Eberl, D.D., 2007. Quantitative mineralogy of surface sediments on the Iceland shelf, and application to down-core studies of Holocene ice-rafted sediments. *Journal of Sedimentary Research* 77, 469-479.
- Andrews, J.T., Eberl, D.D., 2012. Determination of sediment provenance by unmixing the mineralogy of source-area sediments: The "SedUnMix" program. *Marine Geology* 291, 24-33.
- Andrews, J.T., Vogt, C., 2020. Variations in felsic- versus mafic-sources in the Western Nordic Seas during MIS 1 to MIS 4. *Marine Geology* 424, 106164.
- Belt, S.T., 2018. Source-specific biomarkers as proxies for Arctic and Antarctic sea ice, *Organic Geochemistry* 125, 277–298, doi: 10.1016/j.orggeochem.2018.10.002.
- Belt, S.T., Brown, T.A., Navarro-Rodriguez, A., Cabedo-Sanz, P., Tonkin, A., Ingle, R., 2012. A reproducible method for the extraction, identification and quantification of the Arctic sea ice proxy IP₂₅ from marine sediments. *Analytical Methods* 4, 705–713.
- Belt, S.T., Cabedo-Sanz, P., Smik, L., Navarro-Rodriguez, A., Berben, S.M. P., Knies, J., Husum, K., 2015. Identification of paleo Arctic winter sea ice limits and the marginal ice zone: optimised biomarker-based reconstructions of late Quaternary Arctic sea ice. *Earth and Planetary Science Letters* 431, 127-139.
- Eberl, D.D., 2003. User guide to RockJock: A program for determining quantitative mineralogy from X-ray diffraction data. United States Geological Survey, Open File Report 03-78, 40 pp, Washington, DC.
- Konert, M., Vandenberghe, J., 1997. Comparison of laser grain size analysis with pipette and sieve analysis: a solution for the underestimation of the clay fraction. *Sedimentology* 44, 523-535.
- Marshall, N.R., Piper, D.J.W., Saint-Ange, F., Campbell, D.C., 2014. Late Quaternary history of contourite drifts and variations in Labrador Current flow, Flemish Pass, offshore eastern Canada. *Geology Marine Letters* 34, 457-470. doi:10.1007/s00367-014-0377-z.

- 1301 McCave, I.N., Andrews, J.T., 2019a. Distinguishing current effects in sediments
 1302 delivered to the ocean by ice. I. Principles, methods and examples. *Quaternary*
 1303 *Science Reviews* 212, 92-107.
- 1304 McCave, I.N., Andrews, J.T., 2019b. Distinguishing current effects in sediments
 1305 delivered to the ocean by ice. II. Glacial to Holocene changes in North Atlantic high
 1306 latitude upper ocean flows. *Quaternary Science Reviews* 223, no. 105902, 21pp.
- 1307 McCave, I.N., Hall, I.R., Bianchi, G.G., 2006. Laser vs settling velocity differences in silt
 1308 grainsize measurements: estimation of palaeocurrent vigour. *Sedimentology* 53, 919-
 1309 928.
- 1310 McCave, I.N., Manighetti, B. and Robinson, S.G., 1995. Sortable silt and fine sediment
 1311 size/composition slicing: parameters for palaeocurrent speed and palaeoceanography.
 1312 *Paleoceanography* 10, 593-610.
- 1313 McCave, I.N., Thornalley, D.J.R., Hall, I.R., 2017. Relation of sortable silt grain-size to
 1314 deep-sea current speeds: Calibration of the 'Mud Current Meter'. *Deep-Sea Research*
 1315 *Part I* 127, 1-12.
- 1316 McCave, I.N., Syvitski, J.P.M, 1991. Principles and methods of geological particle size
 1317 analysis, in: Syvitski, J.P.M. (Ed.), *Principles, methods and application of particle*
 1318 *size analysis*. Cambridge University Press, pp. 3-21.
- 1319 Robinson, S.G., Maslin, M.A., McCave, I.N., 1995. Magnetic susceptibility variations in
 1320 Upper Pleistocene deep-sea sediments of the N.E. Atlantic: Implications for ice
 1321 rafting and palaeocirculation at the Last Glacial Maximum. *Paleoceanography* 10,
 1322 221-250.
- 1323 Stoner, J.S., Andrews, J.T., 1999. The North Atlantic as a Quaternary magnetic archive,
 1324 in: Maher, B., Thompson, R. (Eds.), *Quaternary Climates, Environments and*
 1325 *Magnetism*. Cambridge University Press, Cambridge, UK, pp. 49-80.
- 1326 Verplanck, E.P., Farmer, G.L., Andrews, J., Dunhill, G., Millo, C., 2009. Provenance of
 1327 Quaternary glacial and glacial marine sediments along the southeast Greenland margin.
 1328 *Earth and Planetary Science Letters* 286, 52-62.
- 1329 Watkins, S.J., Maher, B.A., 2003. Magnetic characterization of present-day deep-sea
 1330 sediments and sources in the North Atlantic. *Earth and Planetary Science Letters*
 1331 214, 379-394.

1332 White, L.F., Bailey, I., Foster, G.L., Allen, G., Kelley, S.P., Andrews, J.T., Hogan, K.,
1333 Dowdeswell, J.A., Storey, C.D., 2016. Tracking the provenance of Greenland-
1334 sourced, Holocene aged, individual sand-sized ice-rafted debris using the Pb-isotope
1335 compositions of feldspars and Ar-40/Ar-39 ages of hornblendes. Earth and Planetary
1336 Science Letters 433, 192-203.
1337
1338

Figures with Captions

Figure 1: A) location of MD99-2274 and some other cores noted in the paper (Table 1) (ODV, Schlitzer, 2011). The shaded areas represent the late glacial maximum (LGM) extent of the ice sheets north of Denmark Strait; the words “basalt” and “felsic” define the primary sediment mineral sources and the arrows show probable flow paths for icebergs. BB = Blosseville Basi; TMF = Scoresby Sund Trough Mouth Fan; B) Additional cores referenced in the paper (see also Table 1). Note that “Cald” on this figure references the southern outcrop of the Greenland Caledonides (Higgins et al., 2008). SS = Scoresby Sund; RIC = Renland Ice Cap. C) Surface and bottom currents and historical April sea-ice edge (1870-1920) (dashed white line; Divine and Dick, 2006). NIIC = North Iceland Irminger Current; EGC = East Greenland Current; EIC = East Iceland Current; Yellow lines: Bottom Currents DSOW = Denmark Strait Overflow Water; NIJ = North Iceland Jet., S = Separated East Greenland Current; OC = Iceland Sea Ocean Convection site (after Harden et al., 2016).

Figure 2: A) Downcore plot of magnetic susceptibility (SI^{-5}) and Bayesian ((Blaauw and Christen, 2016) depth age plots for MD99-2274 (see Table 2)---the red curve is for the initial available data blue curve is for the estimated ages with the addition of an estimated core top age and the presence of the Vedde and NAAZII tephras (see text). The Marine Isotope Stage (MIS) boundaries are indicated. Location of radiocarbon dates and tephras are noted. B) Plot of the departures from the median values of magnetic susceptibility ($2.03 \times 10^{-3} \text{ SI}$) and quartz wt% (5.3). Note that the quartz axis is reversed.

Figure 3: Variation in the Sortable Silt mean size (3-point 1-2-1 weighted smoothing with raw data dots) and IRD% >240 μm . Minima in $\overline{SS}$ are seen at the time of Hudson Strait H events -H6, -H4 and -H1 while -H4, -H2, early -H1 and the YD (-H0) are marked by elevated IRD %. Blue bars are regions where the data are unreliable indicators of flow speed according to the $\overline{SS}$ -SS% correlation criterion of McCave and Andrews, (2019a)

1371

1372

1373

1374 Figure 4: Plots of the variations in the weight% of minerals in MD99-2274, the
 1375 quartz/pyroxene ratio, and magnetic susceptibility. The green shaded areas represent
 1376 Holocene values, hence points above represent departures. Numbers 1 through 5 identify
 1377 IRD quartz peaks. The vertical blue shading areas represent times when the weight% of
 quartz exceeds Holocene limits.

Figure 5: Plots of the sediment source percentages and the degree of fit (DOF), that is the average absolute bias in the SedUnMix calculation of (observed mineral wt% - predicted mineral weight%) for each sample. The top panel shows the location of measurable quantities of gravel, and sites of tephra layers and the radiocarbon dates on near-surface planktonic foraminifera (Table 3). Numbers on the NE Greenland panel represent the peaks in that source and the yellow bars locate areas with minimal input from that area.

Figure 6: Biomarker data (A) IP_{25} and HBI II concentrations; (B) $\Sigma C_{37:3} + C_{37:2}$ alkenone and brassicasterol concentrations; C) SST $^{\circ}C$ estimates and the % $C_{37:4}$; and D) Weight % quartz and different coarse sediment fractions.

Figure 7: Schematic presentation of changes in sea ice and iceberg distribution. The first panel (upper left) shows core locations (see Table 1 and Fig. 1A and B) and the adjoining panel the inferred conditions during MIS 3 and 2 with pervasive sea ice and embedded icebergs. The remaining panels show the proposed evolution in the state of sea ice and iceberg supply (red triangles) during deglaciation into the Holocene (adapted from Cabedo-Sanz et al., 2016; Xiao et al., 2017). SS = Scoresby Sound, RIC = Renland Ice Cap.

Figure 8: Analysis of the quartz wt% records from PS2644 (Vogt, 2017) and MD99-2274 at a common 0.6 ky spacing. A) Original quartz data (black line) and the wavelet reconstructions for the two records; B) Cross-wavelet power spectrum of quartz wt% for PS2644 and MD99-2274. The cone of confidence indicated by the light grey areas; 0.05% probability area demarcated by white line. Arrows pointing to the right mean that the two records are in phase, arrows pointing down mean that x leads y, arrows pointing to the left indicate the records are anti-phase and pointing up indicates that #2274 leads PS2644. C) Cross-wavelet (Fig. 8B) average power. The 0.05 significance period is red and delimited by the dashed slanting line. The horizontal dashed line indicates the peak periodicity (~8.5 ky).