1 INSHORE OR OFFSHORE? THE NEOLITHIC FISHERMEN OF RA'S AL-HAMRA (RH-6, RH-5,
2 MUSCAT, SULTANATE OF OMAN)

3 Anaïs MARRAST¹, Philippe BÉAREZ¹, Emilie BADEL², Lapo Gianni MARCUCCI³.
4

5 ABSTRACT

6 Ra's al-Hamra 6 (RH-6) and Ra's al-Hamra 5 (RH-5) are two of the most important Neolithic
7 coastal sites in the Sultanate of Oman. Located in the capital Muscat, they have supplied an
8 important corpus of fish bones and revealed a high taxonomic richness in the different
9 assemblages. The results provided information on fishing practices that held an important role
10 in the subsistence strategies of the inhabitants during the two millennia of occupation (6th–4th
11 millennia BC). Pelagic species (jacks and kawakawa/tunas) were largely dominant throughout
12 both occupations, calling into question the interpretation of inshore or offshore fishing; these
13 fish could have been caught with a hook-and-line or by setting nets at a reasonable distance
14 from the coast. However, the inhabitants also had to supplement this subsistence by catching
15 demersal fish and small fish, caught from the cliff, in the estuary or the mangrove. At both sites,
16 the inhabitants clearly exploited the different nearby inshore environments: coastal open water,
17 reefs, estuary and mangrove. Moreover, the high fish richness (69 genera at RH-6; 35 genera at
18 RH-5) indicates that both sites could have been occupied on a year-round basis.
19

20 INTRODUCTION

21 Many shell middens have been identified in South Eastern Arabia for the Middle Holocene
22 (7th–4th millennia), and the important economic role of aquatic resources is well established
23 (e.g., Beech 2004; Biagi 1994, 2006). Fishing does appear to have been important to these
24 populations, described as fisher-hunter-gatherer communities (Charpentier 2008; Cleuziou and
25 Tosi 2007; Cleuziou 2005), yet surprisingly, fish remains from only a few sites in Oman have

26 been analysed and published (Cartwright 2004; Uerpmann and Uerpmann 2003; Wilkens
27 2005).

28 The ancient shell midden of Ra's al-Hamra 6 (RH-6) and the more recent one of Ra's al-
29 Hamra 5 (RH-5) are two of the most important Neolithic sites in the Gulf of Oman (Figure 1).
30 The total area of each of the two sites is estimated at approximately 3500 m² for RH-6 and 3000
31 m² for RH-5. Several successive campaigns have highlighted a deep stratigraphy of almost 2 m
32 with numerous finds, including animal remains, fishing gear, and burials (Biagi 1987, 1999;
33 Biagi and Maggi 1987; Biagi and Nisbet 1992; Biagi and Salvatori 1986; Coppa et al. 1985;
34 Marcucci et al. 2011, 2014; Salvatori 2007). The first fish bone analyses at Ra's al-Hamra were
35 carried out by Biagi and Travers (1985), followed by more recent studies by Uerpmann and
36 Uerpmann (2003) and Wilkens (2005); however, their results did not take into account the
37 chronology of the site.

38
39 *Figure 1. (A) Location, altitudinal and bathymetric mapping of the Qurum mangrove and*
40 *the Ra's al-Hamra's cape, with location of the archaeological sites; (B) Map of the excavated*
41 *sectors of RH-6 (Marcucci et al., 2014, fig. 1.A.); and (C) Map of the excavated sectors of*
42 *RH-5 (Marcucci et al., 2011, fig. 2.)*

43
44 The latest investigations on the Arabian Sea coastal fringe include new fish bone studies
45 from two Neolithic sites: Suwayh 1 (SWY-1) and Ruways 1 (RWY-1), which provide
46 information on specific subsistence strategies. At these sites, ancient fishermen specialized in
47 lagoon environment exploitation, although with very different patterns of fishing: primarily
48 sharks at SWY-1 (Marrast, Béarez, and Charpentier 2020), and sardines, mullets, and juvenile
49 fish at RWY-1 (Berger et al. 2020).

50 Based on recent excavations at RH-6 and RH-5 (Marcucci et al. 2011, 2014; Marcucci,
51 Badel, and Genchi 2021), this paper presents the results from new fish bone analyses (Marrast
52 2020) and sheds light on Neolithic fishing practices in the capital area of Muscat (southern coast
53 of the Gulf of Oman).

54

55 *Environmental setting*

56 The study area is located at the southeastern end of the sandy beaches in the Al Batinah
57 Region, where the Qurum mangrove swamp stands, and where the rocky shores of Muscat
58 begin. The settlements of Ra's al-Hamra are located on the eastern edge of the Qurum Natural
59 Reserve in the district of Qurum (Figure 1A). In this area, twelve settlements have been reported
60 and four have been excavated (Durante & Tosi 1977; Uerpmann and Uerpmann 2003). Today,
61 RH-6 is about 600 m away from the sea, and RH-5 is on top of the rocky headland overhanging
62 the Gulf of Oman. But, it is important to note that at the end of the fifth millennium BC, which
63 corresponds to the end of RH-6 and the beginning of RH-5 occupations, a higher sea level (+
64 1m) than today has been identified in the northern coast of Batinah (Vita-Finzi 1982; Berger et
65 al. 2005, 56). The sites are surrounded by a multifaceted natural setting which allowed Neolithic
66 communities quick access to the Qurum mangrove, the shallow sea, the estuary, the sandy
67 beach, and even the rocky shoreline along the cliffs. In these distinct environments, ancient
68 fishermen had access to various animal resources, such as seashells, crabs, and, of course, fish
69 (Biagi 1987; Biagi and Maggi 1987; Biagi and Travers 1985; Tosi 1975). Within the Gulf of
70 Oman, the Omani Coast is influenced by monsoonal winds; it is also indirectly affected by the
71 Arabian Sea upwellings, which bring nutrient rich cold waters to the surface (Al-Azri et al.
72 2014).

73

74 Both sites offer insight into the continuous occupation of this area from the middle of the
75 sixth millennium to the end of the fourth millennium BC. This occupation took place during the
76 Holocene Humid Period, which had a stronger summer monsoon than today, illustrating a
77 climatic optimum induced by a higher latitude position of the Intertropical Convergence Zone
78 (ITCZ) (Fleitmann et al. 2007; Lézine et al. 2017; McClure 1976). This situation was probably
79 favorable for the extension of mangrove and lagoon ecosystems, as confirmed by the numerous
80 *Terebralia palustris* shells and *Avicennia* sp. charcoal fragments radiocarbon dated from this
81 period (Berger et al. 2013; Tengberg 2005).

82

83 *Archaeological setting*

84 The occupation of RH-6 ranges from the sixth millennium to the middle of the fifth
85 millennium BC, but the graveyard is currently dated to the beginning of the fourth millennium
86 BC (Biagi 2006; Zazzo et al. 2016). Dating of the settlement makes RH-6 the oldest site in the
87 archaeological sequence of Qurum. Recent excavations identified several sectors (Marcucci et
88 al. 2014): Sector A (eastern edge of the midden); B (post-occupation burials of RH-6 that were
89 contemporaneous with the occupation of RH-5); C/TT-Z (eastern edge of the midden); and
90 Trench North (northern edge of the midden) (Figure 1B). Several post holes and hearths,
91 indicating the presence of huts and living structures, were uncovered. Sector C/TT-Z, a trench
92 with a maximum depth of about 1.75 m, was selected for bioarchaeological sampling. This
93 trench provided a lot of organic material: seeds and charcoal fragments, seashells, and remains
94 of terrestrial and marine mammals, sea turtles (Frazier et al. 2018), and fish. Each stratigraphic
95 unit (SU) was sieved using 3 mm and 1 mm mesh. Thirty-two samples of charcoal fragments,
96 shells, and human bones were selected for ¹⁴C dating (Zazzo et al. 2016). The same protocol
97 was then applied to the Trench North excavation. Six periods were defined in Sector C/TT-Z
98 and four in Trench North (Marcucci et al. 2014).

The occupation of RH-5 ranges from the end of the fifth to the end of the fourth millennium BC (Biagi 1994; Biagi and Nisbet 2006; Zazzo, Munoz, and Saliège 2014). During recent excavations, the study area was divided into six sectors (Figure 1.C). Sector B2, which is the subject of this study, dates from 3910-3640 cal BC (LTL12826A: 4943±45 BP, IntCal05) (Marcucci 2015) and covers an area of 25 m², under an archaeological deposit 10–20 cm deep; here, 200 post holes were detected related to six living structures and associated with six probable waste pits. Four phases were defined in this sector with multiple remains of non-permanent living structures in connection with hearths, storage, and dump pits, and the many intersections of these structures, which suggest intensive and prolonged occupation of the site. Each SU was sieved using a 1 mm mesh (Marcucci et al. 2011; Marcucci, Badel, and Genchi 2021).

Previous ichthyological studies

Several studies have been conducted on the fish bones recovered during the excavations of the 1980s and 1990s, while three have been carried out using the RH-6 material. Biagi and Travers (1985) undertook an exploratory study on material taken from the first excavation (1981). The *in situ* occurrence of small pelagic species, such as sardines (Clupeidae) and anchovies (Engraulidae), was noted, as was the presence of large pelagic species, such as tunas (*Thunnus* spp., Scombridae) and numerous coastal fishes from rocky or sandy-mud bottoms, and many marine catfishes (Ariidae). Another study by Uerpmann and Uerpmann (2003, table 9.7) (51 NISP) questioned some of the previous identifications (Table 1). They obtained only one marine catfish specimen but 22 identified specimens belonging to jacks and trevallies (Carangidae), and 23 identified specimens from tunas. A complementary study was subsequently carried out by Wilkens (2005) on Sector A (570 NISP) that highlighted the

presence of different pelagic fish, such as several scombrids and numerous benthic-demersal fish taxa. In all these studies, the main taxa identified were similar (i.e., Scombridae, Carangidae, Sparidae, Serranidae, Lethrinidae).

Table 1. Table synthesizing the main results from previous studies on the fish bone remains from RH-6 and RH-5 (for Uerpmann and Uerpmann 2003, NISP have been calculated from identified Actinopterygii and Chondrichthyes), and modern data of artisanal fishing from the governorate of Muscat for 2009-2018 (M.A.F. 2018)

The RH-5 material studied by Uerpmann and Uerpmann (2003, table 9.3) came from across the whole site, without precise context (Table 1). Based on an NISP of 12,891, the authors indicated that fishing activities were oriented mainly toward the exploitation of medium-sized tunas (NISP = 7596), and jacks/trevallies (NISP = 4065). The exploitation of demersal taxa, such as breams (Sparidae), emperors (Lethrinidae), or parrotfishes (Scaridae), was secondary. However, they noted the occurrence of a large grouper (*Epinephelus* sp.), estimated at a minimum of 150 cm, and a shark represented by a vertebra of 40 mm in diameter, which is very large. In contrast, they also indicated the presence of a high number of small fish in the 1 mm samples (NISP = 402 of 3920 remains counted).

MATERIAL AND METHODS

The material studied here came from Sector C/TT-Z and Trench North for RH-6, and from Sector B2 for RH-5 (Figure 1B-C). For the RH-6 material, each stratigraphic unit was sieved using 3 mm and 1 mm mesh. For the RH-5 material, each SU was sieved using 1 mm mesh. Cut marks were absent, and all burning marks were reported. However, due to the high fragmentation and the number of remains, we did not record the fragmentation pattern. To

facilitate the reader's understanding, the results are presented by period; more detailed results are available in Marrast (2020).

For the identification of the fish remains, we used the fish skeletons reference collection of the National Museum of Natural History of Paris, which contains for ca. 900 specimens representing ca. 500 species of Indo-Pacific fishes. Fish remains were identified at the most precise taxonomic level (species), but due to the high fragmentation of the remains, the identification was done mainly to genus or family level. Occasionally, we used higher taxa, such as the superfamily Sparoidea, which here includes all species of the families Sparidae, Lethrinidae, and Nemipteridae, as they have very similar vertebrae or teeth which were numerous in the small material and difficult to differentiate.

Grouping of fish skeletal elements is adapted from Wheeler and Jones (1989): cranial bones (neurocranium and branchiocranium elements); vertebrae; and others (fin elements, scales, and scutes). Fish systematic nomenclature follows Fricke, Eschmeyer, and Van der Laan (2021).

As for fish habitat and ecology, we followed the information from Randall (1995), and FishBase (Froese and Pauly 2021). To simplify our interpretations, the fishes have been classified into two main categories: pelagic fish, which include all species living in the water column (whether coastal or open sea, up to 200 m deep), and demersal fish, which are associated with coastal substrates. To discuss the seasonality of fishing activities and site occupation, we used information from the Ministry of Agriculture and Fisheries (M.A.F.) of the Sultanate of Oman, which provides data on monthly artisanal fisheries landings for the governorate of Muscat (M.A.F. 2013, 2015, 2016, 2018).

We mainly use the term NISP (Number of Identified Specimens), which corresponds to the number of fish remains identified to at least superorder level for Chondrichthyes (Batoidea for rays, Selachimorpha for sharks), and order level for Actinopterygii (e.g., Clupeiformes,

174 Siluriformes, Tetraodontiformes). We also use the term WISP (Weight of Identified Specimens)
175 for the weight of the identified bone remains.

176 We used the classical approach of comparative anatomy and chose to estimate the live weight
177 of the fish, rather than its size, because this seems to be a better indicator of the amount of flesh
178 the fish can provide. The live weights we provide are weight visual estimates, made by direct
179 comparison with specimens from the fish reference collection of known live weight and length
180 data, not reconstructions using osteometric models. The estimate of the MNI (Minimum
181 Number of Individuals) was performed on the main taxa (species or genus level) despite the
182 high number of vertebrae, which sometimes made it very difficult. MNI was estimated by
183 counting the most frequent anatomical element of a given species after considering laterality
184 and size (i.e., MNI of combination; Poplin, 1976). Several species of jacks and trevallies present
185 in the study area have characteristic swollen bones (hyperostosis) on the occipital crest or
186 cleithrum, which, due to their good preservation, were often found and used for the estimation
187 of the MNI.

188 In order to compare results from different chronological periods, chi square tests were
189 performed on the corresponding NISP to detect the dependency between two faunal
190 assemblages, and therefore the associated fishing patterns (Chenorkian, 1996). Tests were
191 performed on each pair of assemblages from two successive periods (H_0 : The two relative
192 proportions do not differ significantly). To respect the test conditions ($n > 5$) for each test, only
193 the most important fish families (between 5 or 7, depending on the assemblages) were tested.

194

195 RESULTS

196 *The RH-6 site*

197 Since Trench North is more representative of the beginning of the occupation and Sector
198 C/TT-Z is more representative of the end, we will present the results in this order. The

199 equivalence between the periods in the two sectors is adapted from Marcucci (2015: table
200 A2.13). For ease of presentation, we discuss the results for the main families and mostly at the
201 genus level, except for Thunnini. All the species identifications can be found in supplemental
202 files 1–3.

203

204

Trench North – 3 mm

205 For Trench North, located at the northern edge of the shell midden, vertebrae were the most
206 abundant (66.5% of the material anatomically identified), followed by cranial elements
207 (26.5%), and other post-cranial elements (7%). The dentary was the most represented cranial
208 bone (NISP = 528), followed by the premaxilla (NISP = 353) (Table 2). These jaw bones are
209 among the strongest bones in most teleosteans (bony fish) and consequently have a very good
210 conservation rate (Wheeler and Jones 1989, 90). Still, there is a major imbalance between the
211 remains of the skull and the spine because the vertebrae are thicker than the skull bones and
212 therefore better preserved.

213

214 *Table 2. Element distribution on the two sites according to the sector studied and the*
215 *sieving mesh size (Chondr. = Chondrichthyes)*

216

217 A total of 9116 fish bones (NISP) were identified. Most of this material corresponded to the
218 beginning of the occupation of RH-6, i.e., periods 1–4 of Sector C/TT-Z. In Trench North, four
219 periods were defined. However, as periods 3 and 4 were poor in fish (NISP = 9) these results
220 were not included in the discussion. Period 1 is taxonomically rich (32 families, 51 genera),
221 associated with a high NISP (4908). Period 2, with a lower NISP (4208), is less taxonomically
222 rich (23 families, 38 genera) (Figure 2A).

223

Figure 2. Taxonomic richness at family and genus levels for different sectors and time periods

Chi square tests were performed between the assemblages but did not reveal any significant differences (χ^2 : 6.6397, $p > 0.1$, df: 4) (Figure 3A; Table 3). Pelagic fish represent more than 91% of the assemblage NISP for the two periods, with trevallies (*Carangoides* spp.) as the main taxon (20%). Kawakawa (*Euthynnus affinis*) and jacks (*Caranx* spp.) were of similar importance throughout the site's occupation (15.5% and 12%), as was the golden trevally (*Gnathanodon speciosus*) (3.6%), while tunas seem to be the least important of the pelagic fishes; the two species accounting for just 5–6%. The contribution of demersal fish, mainly represented by different seabreams, emperors, and groupers (Serranidae), was very low (9%) (Table 3).

Figure 3. Illustration of the main taxa distribution (NISP).

(A) RH-6, Trench North, 3 mm; (B) RH-6, Sector C/TT-Z, 3 mm. Made with ACACIA (Nourissier, Nocus, and Dufraisse, 2010)

Table 3. Zooarchaeological fish remains from RH-6, Trench North, 3 mm, according to periods in NISP, % of NISP and WISP (g). Other pelagic = 12 families (Supplemental file 1); Other demersal = 19 families (Supplemental file 1).

The mean weight of the caught fish varied between periods. Indeed, while the tuna mean weight for period 1 is 7 kg, it falls to 4.5 kg, and has a greater distribution, in period 2 (Figure 4A). In contrast, jacks are larger in period 2 (2.5 kg versus 1 kg in period 1), while kawakawas, and trevallies weigh in at 3 to 3.5 kg for both periods (Figure 4B-D). However, the NISP of

249 kawakawa and jacks must be considered against the MNI (Table 4). Most of the kawakawa
 250 bones are vertebrae, which have high conservation rates because of their extremely robust
 251 constitution. On the other hand, many cranial elements of jacks were found, increasing their
 252 MNI. The difference in proportions between kawakawa and tunas is less when measured by
 253 MNI than by NISP.

254

255 *Figure 4. Illustration of the estimated weight of each main pelagic taxa from the richest*
 256 *RH-6 periods (P: Period), both sectors (TN: Trench North, C-TTZ: Sector C TT-Z). Min:*
 257 *Minimum, Max: Maximum, Q: Quartile. (A) Thunnus spp; (B) Euthynnus affinis; (C)*
 258 *Carangoides spp.; (D) Caranx spp.*

259

260 *Table 4. MNI estimates of the main pelagic taxa at RH-6, 3 mm, according to periods and*
 261 *comparison with the NISP*

262

263 Sector C/TT-Z – 3 mm

264 For Sector C/TT-Z, vertebrae were also the most abundant element (62% of the material
 265 anatomically identified), followed by cranial elements (30%), and other post-cranial elements
 266 (8%). Jaw bones, with 508 premaxillae and 496 dentaries, were the most frequently represented
 267 cranial bones (Table 2).

268 A total NISP of 8489 have been determined, but abundance of fish bones was very disparate
 269 from one period to another. Fish bones were much less abundant in the four oldest periods than
 270 in the two most recent ones (Figure 3B). Taxonomic richness has increased over time (Figure
 271 2B), in a trend that appears to be roughly independent of the NISP. Richness peaked in period
 272 6 (26 families, 52 genera), while its NISP (3464) was lower than in period 5 (4109).

273

274 In this sector, pelagic fish taxa were also dominant (Figure 3B, Table 5). Jacks were the main
275 taxa (18% of the total assemblage), followed by trevallies (11.5%), and kawakawas (10%). Chi
276 square tests were performed between all pairwise assemblages, with the exception of period 2
277 due to its small sample size but did not reveal any significant differences between periods 3 and
278 4 (Figure 3B) (χ^2 : 12.93, $p > 0.01$, df: 4). On the contrary, periods 4 and 5 exhibited a strong
279 statistical difference in faunal composition, supporting a genuine increase in the proportions of
280 the two main pelagic families (from 64 % to 82%) and a decrease in demersal fish (from 34%
281 to 16%) (χ^2 : 82.39, $p < 0.001$, df: 4). For the end of the occupation, between periods 5 and 6, a
282 chi square test indicated the opposite effect (χ^2 : 35.01, $p < 0.001$, df: 4), with an increase of
283 demersal fish (from 16% to 20.5%) and a decrease of pelagic fish, driven by the decrease of
284 scombrids (from 25% to 21%).

285
286 *Table 5. Zooarchaeological fish remains from RH-6, Sector C/TT-Z, 3 mm, according to*
287 *periods in NISP, % of NISP and WISP (g). Other pelagic = 5 families (Supplemental file 1);*
288 *Other demersal = 24 families (Supplemental file 1)*

289
290 In Sector C/TT-Z, an increase in MNI is associated with the end of occupation. Indeed,
291 periods 5 and 6 were represented by 386 and 294 main pelagic fishes, much more than in
292 previous periods (Table 4).

293
294 Mean fish weights were estimated for the two most recent periods (Figure 4), periods 5 and
295 6, because their sample sizes were large enough to allow comparison with the Trench North
296 periods. For kawakawa and tunas, specimens were bigger in period 6 with a mean weight
297 reaching 4 kg (versus 3.1 kg) and 5 kg (versus 3.5 kg), respectively (Figure 4A-B). Conversely,
298 for jacks and trevallies, there was a decrease in the mean weight at the end of RH-6 occupation,

with the mean weight of jacks decreasing from 2 kg to 700 g, and that of trevallies decreasing slightly from 3 kg to 2.6 kg (Figure 4C-D).

Sector C/TT-Z – 1 mm sampling

Here, the results are presented only for periods 2 to 6 of Sector C/TT-Z. Two samples were taken for each period (Table 6). A significant disparity was observed between the numbers of vertebrae (95%), cranial elements (4.5%), and other elements (0.5%) (Table 2).

Table 6. Zooarchaeological fish remains from RH-6, Sector C/TT-Z, 1 mm in NISP and % of NISP by period. Other pelagic = 9 families (Supplemental file 1); Other demersal = 13 families (Supplemental file 1)

A total of 9388 fish bones (NISP) were identified within 22 families and 16 genera (Figure 2C), reflecting the potential information added with fine mesh sieving. As sieving was not uniform, we chose not to perform chi square testing; however, a major change through time can still be observed. Sardines (Clupeidae) take on greater importance in the faunal assemblage until period 5 (Figure 5). Although the species could not be identified, it was probably mainly the Indian oil sardine (*Sardinella longiceps*), along with some goldstripe sardinella (*Sardinella gibbosa*), as is reported today among the traditional fishery in Oman (Strømme 1986). Different scads (Carangidae) were noteworthy, with bigeye scads (*Selar crumenophthalmus*) representing more than 90% of NISP. Their proportional abundance nevertheless declines from 29.6% in period 2 to 12% in period 6. The occurrence of silversides (Atherinidae) was noticeable, as was the presence of mullets and juvenile breams/emperors (Sparoidea).

Figure 5. Illustration of the main taxa distribution (NISP).

RH-6, Sector C/TT-Z, 1 mm (Periods 2, 3, 4, 5, 6) – RH-5, Sector B2, 1mm (Phases 2, 3, 4). Made with ACACIA (Nourissier, Nocus, and Dufraisse, 2010)

The RH-5 site

In Sector B2, all zooarchaeological material was recovered by sieving with a 1 mm mesh. There was a discrepancy in the anatomical distribution of vertebrae (95%), cranial elements (4.5%), and other elements (0.5%) (Table 2). A total of 8406 specimens were identified, weighing 92 g (Table 7). Assemblage abundance was variable over time and due to low specimen numbers in phase 1, these data are excluded from the interpretation of the results.

Table 7. Zooarchaeological fish remains from RH-5, Sector B2, 1 mm, according to periods in NISP, % of NISP and WISP (g). Other pelagic = 7 families (Supplemental file 1); Other demersal = 20 families (Supplemental file 1)

Since the beginning of occupation, richness was relatively high. For phase 1, despite a very low NISP (32), high richness was noted (7 families, 7 genera). It reached a peak in phase 3 (27 families, 33 genera), with the highest NISP (6640) counted. During phase 4 (15 families, 17 genera), there was a decrease in NISP (1492) (Figure 2D).

Chi square tests were performed between the assemblages and revealed significant proportional differences in faunal composition between phases 2 and 3 (χ^2 : 55.84, $p < 0.001$, df: 4). These changes are driven by the observed increases or decreases in several taxa. In phase 2, carangids, including bigeye scads, were the most abundant taxon, representing 44.5% of identified specimens (Table 7, Figure 5). They decreased to 36% in phase 3, replaced by sardines (55% in phase 3). For the phases 3 and 4, Chi square tests did not reveal significant

proportional difference in faunal composition (χ^2 : 6.68, $p < 0.001$, df: 5). Indeed, the sardines contributed a similar proportion to the assemblage in phase 4. Other pelagic (mulletts, barracudas) and demersal (Sparoidea) taxa declined in abundance over time.

The assemblages were composed mainly of small vertebrae, which complicated the MNI calculation. Estimating precise live weight from small bones is challenging, but most of the RH-5 fish (scads, sardines, mullets, Indian mackerel) belonged to the 150 g weight category. No fish larger than 5 kg were recorded from our study at RH-5.

DISCUSSION

Several trends clearly emerged concerning different aspects of the fishing economy across the Ra's al-Hamra sites, although the materials analyzed came from distinctly separate sectors.

Taphonomic remarks

The NISP of kawakawa and jacks must be considered alongside the MNI (Table 4). Most of the kawakawa bones are vertebrae, which have high conservation rates because of their extremely robust constitution. On the other hand, many cranial and hyperostotic elements of jacks and trevallies were found, which increases their MNI. The difference in proportions between kawakawa and jacks, therefore, no longer appears to be as great with the MNI as with the NISP.

As the cranial elements were highly represented in each SU and no traces of butchery were found, fish were probably complete and fresh when brought to the site. Apart from gutting, they were probably not processed before consumption. As for the burn marks on the bones, very few were noticed on both sites and no burning pattern could be discerned. The bones could be

completely burnt, indicating that they had probably simply been thrown into the fire after the fish had been consumed.

Fishing patterns

Our results indicate slight changes through time between the two sites. The RH-6 fishermen concentrated their efforts on catching medium-sized pelagic fish, comparable with previous RH-5 results. At both sites, pelagic fish accounted for more than 80 % over time (Tables 1, 3, 5), illustrating a subsistence strategy probably adapted to a marine environment rich in pelagic fish. However, when considering the proportions of the two main families of pelagic fish, the fishing pattern of the late RH-6 occupation (Period 6) and that of RH-5 (Table 1; Uerpmann and Uerpmann, 2003) appeared to be highly different (χ^2 : 2934.68, $p < 0.001$, df: 6). The inhabitants of RH-5 seem to have concentrated their efforts on catching different scombrids. This change may represent different cultural choices in fishing strategies.

The results from the 1 mm mesh fraction highlighted the importance of small fish, with a high number of remains counted at both sites (Tables 6, 7). It is particularly important to note here the very high proportion of vertebrae (95%) in the 1 mm fraction material at both sites. The preservation of vertebrae is often better than that of cranial bones, but this is even more true in the case of small fish. Based on the study of this material, fishing patterns appear to be very specific to each site (RH-6-P6 and RH-5-Ph3, χ^2 : 1400.76, $p < 0.001$, df: 6). Sardines constituted 45% of the fish remains at the beginning of RH-6 occupation, but by the end constituted 70% of the faunal assemblage. At RH-5, scads are abundant at the beginning of occupation (36% in phase 2, while sardines occupied 35%), but this pattern is reversed at the end of the occupation (sardines 58% and scads 32%). For RH-5, however, consideration of WISP provides a more nuanced understanding of the results. Indeed, if sardines, scads, and mackerel are dominant based on NISP, kawakawas, barracudas, and jacks/trevallies are the best

represented according to the WISP, especially jacks (*Caranx* spp.). Demersal fish were rare at RH-5, and it is interesting to note that most of the fish in Sector B2 were represented by specimens weighing less than 150 g.

Nowadays, artisanal fishing, by line, troll, or net, provides landings composed mainly of large pelagic species, such as longtail tuna (*Thunnus tonggol*), yellowfin tuna (*Thunnus albacares*), kawakawa (*Euthynnus affinis*), large jacks/trevallies (*Caranx* spp. and *Carangoides* spp.), and Talang queenfish (*Scomberoides commersonnianus*). An important part of the fishery, however, consists of small pelagic fish, such as sardines (Clupeidae), Indian mackerel (*Rastrelliger kanagurta*), and mullets (Mugilidae) (Table 1, M.A.F. 2018). However, while the species fished today were already present on the two archaeological sites, two pelagic species, the Talang queenfish and the Narrow-barred Spanish mackerel (*Scomberomorus commerson*), are absent or only very occasionally present in the mid-Holocene catch (see Supplemental file 1). Moreover, the proportions are very different today, with pelagic fish representing more than 80% of the catch at RH-6 and RH-5, compared to 70% in the modern catch.

Inshore or offshore?

Despite the occurrence of medium-sized pelagic fish in several sites throughout the Persian Gulf and Gulf of Oman for the Neolithic period, such as Delma (Lidour and Beech 2019), Akab (Lidour et al. 2020), Dosariyah (Uerpmann and Uerpmann 2018), and Khor Milkh (Uerpmann and Uerpmann 2003), the sites of Ra's al-Hamra appear to be distinctive given the high proportion of these fish in their faunal material.

The abundance of tuna and other pelagic fish raises questions about the existence of offshore fishing in the Neolithic sites of the Gulf of Oman. Since the first studies, interpretations have leaned towards offshore fishing, with the use of boats suggested for catching tunas/kawakawas

and jacks/trevallies (Biagi et al. 1984; Biagi and Nisbet 1992; Uerpmann and Uerpmann 2003: 198). The terminology of fishing areas was clarified by Pickard and Bonsall (2004) and adjusted by Béarez et al. (2016) who added water depth as an important parameter. We follow the latter to denote an inshore fishery as one that is conducted in the inshore and nearshore zone (down to 30 m depth), an offshore fishery as one that is conducted in deeper waters but within sight of land (< 5 km), and an open sea (deep-sea) fishery as one that is conducted out of sight of land.

Although jacks, trevallies, tunas and kawakawa are pelagic, these species can approach relatively close to the coast (Anderson 2013; Collette 2001). Tunas and other scombrids are pelagic fish known to form large, sometimes multi-species schools of thousands of individuals (Collette and Nauen 1983), a behavior which is also observed at a lesser scale in many carangids (scads and young to small adult trevallies and jacks). These open water schools can be observed by fishermen from the coast, especially when sea birds fly over them (Cartwright 2004; Lidour et al. 2020, fig. 7.A). Until recently, fishermen used small rafts or boats, such as *shasha*, to join a school of tunas, once sighted, and then deployed hook-and-lines or nets to catch them. These small boats are mainly made of date leaves and date palm fibers, which provide buoyancy (Al-Marshudi and Kotamaga 2006; Goddard 1985; Johnson 2016; Uerpmann and Uerpmann, 2003, 198-199). Similar, frail embarcations could have been made during the Middle Holocene with materials collected in the vicinity of the Ra's al-Hamra sites, such as Mazari palm (*Nannorrhops ritchieana*), common reed (*Phragmites australis*), or cattail (*Typha* spp.) (Biagi et al. 1984; Biagi and Nisbet 1999; M. Tengberg, pers. comm.). Stronger and sewn vessels are only attested for the third millennium (Cleuziou 2003; Vosmer 2019). As for the ropes of the lines and nets, they may also have been made from Mazari palm; indeed, ropes found at the Shahi Tump site (Iran, fourth millennium BC) were made from this species (Thomas et al. 2012). In addition, fishing cordage has traditionally been made from this species along the Omani Coast

(Richardson and Dorr 2003); it is favored by fishermen because its strength is increased when wet. Finally, on both RH-6 and RH-5 sites, different shell-hooks and sinkers have been recovered (Biagi, 1987; Marcucci et al., 2011, 2014).

Unfortunately, the difficulty in separating the two species of tuna, yellowfin and longtail, based on their osteology may explain the high rate of identifications at the genus level (*Thunnus* spp.) in our studies and those of Uerpmann and Uerpmann (2003), which limits the robustness of the interpretations. Normally, yellowfin tuna can reach larger sizes than the longtail tuna. However, at both archaeological sites, yellowfin tunas appear to be smaller than longtail tuna. Small schools of small tunas probably approached the coast on their own, while larger individuals remained offshore. In addition, kawakawa is the main scombrid found at both sites. It is known that among the schools of kawakawa targeted by fishermen, small tuna specimens may also have been present (Uerpmann and Uerpmann 2003, 199), so the mixed scombrids fishery may well have been inshore. Fishermen would only need a glittering hook, such as a mother-of-pearl hook, to catch voracious scombrids (Uerpmann and Uerpmann 2003, 198). The casting of nets from boats is suggested by the presence of large net sinkers, but this practice was probably difficult due to the vulnerability of their frail boats; also, this fishing technique may have been particularly dangerous (Lidour et al. 2020; Uerpmann and Uerpmann 2003, 199). Furthermore, it is possible to fish tuna from shore; indeed, this practice is conducted today for sport (Karachle, Dimarchopoulou, and Tsikliras 2020).

Large individuals of carangids can currently be found living solitary or in small groups along the coast (Grandcourt 2012; Grandcourt et al. 2004). In the Ra's al-Hamra material, three genera are important: *Carangoides*, *Caranx*, and *Gnathanodon*. The golden trevally (*G. speciosus*) is often associated with deep reefs and deep lagoons (Grandcourt et al. 2004). As they represent

the main family at RH-6, their presence could be connected with the numerous small line sinkers or the many complete or fragmented hooks found at RH-6 in different stages of manufacture (Biagi 1999). This could indicate a specialization in hook-and-line fishing. However, the large size of some sinkers (Durante and Tosi 1977) may suggest the use of big nets in open water, or fish traps (*gargour* in Arabic) (Uerpmann and Uerpmann 2003).

While offshore navigation is attested for the Neolithic of the Arabian Peninsula in the Persian Gulf, with the occupation of islands 10 km from the coast and the presence of bitumen (Beech 2004; Beech et al. 2016; Carter and Crawford 2010; Lidour et al. 2020), and in Oman with the occupation of Masirah Island (Charpentier et al. 2013), it is not clear whether the fish assemblage reflects offshore fishery. Indeed, previous studies have shown that the presence of scombrids is not necessarily a sign of this activity (Anderson 2013; Béarez, Gay, Lunniss 2012). As most of the species identified at RH-6 and RH-5 are found close to the coast (< 5 km), and there is an absence of large pelagic species such as Indo-Pacific sailfish (*Istiophorus platypterus*) or large tunas (> 10 kg), this might suggest inshore fishery. The presence of dolphins and whales at both sites could be an indication of offshore fishing, but their presence is too sporadic (NISP: 46 small dolphin, 34 big dolphin, and 31 small whale remains at RH-5) to indicate systematic fishing activity; however, scavenging of beached carcasses cannot be ruled out (Uerpmann and Uerpmann 2003).

Estuary and mangrove exploitation

In the underwater environment of Ra's al-Hamra, many demersal species could be fished, such as emperors, groupers, snappers, seabreams, and even parrotfish and porcupinefish, all of which have been found at both sites (Supplemental files 2–3). The main taxa from this category (seabreams, emperors, and groupers) appear to be able to be caught using the same fishing

techniques. These fish can be caught by net, trap, or hook-and-line, and are to be found all year-round, so they may have been used for daily subsistence.

Another environment exploited for fishing by the inhabitants of both sites was the mangrove. Indeed, using the 1 mm mesh size on both excavations allowed new information to come to light about this specific fishing ground, which is used as a nursery by many species, especially fish (Abu El-Regal and Ibrahim 2014; Laegdsgaard and Johnson 2001; Nagelkerken et al. 2002; Robertson and Duke 1987). The mangrove edges and shores can be considered excellent spots for positioning a fishing camp. These brackish-water ecosystems are normally characterized by their high biomass, with the presence of numerous mullets and sardines (Kulbicki 1997). Although Uerpmann and Uerpmann (2003, 198-199) have already noted the presence of many small fish at Ra's al-Hamra, they minimized their inclusion in the diet of ancient populations by suggesting that they could be tuna stomach contents. Our study, however, places greater importance on these fish as they correspond to an environment highlighted by previous environmental studies showing the most extensive development of mangrove at the end of the fifth millennium/beginning of the fourth millennium BC (Berger et al. 2013; Berger et al. 2005; Biagi and Nisbet 1992; Tengberg 2005). The many hooks found in situ do not coincide with this type of mangrove fishing, however, and the size of the fish suggests the use of small-mesh nets or beach seine. This accords with the small-sized sinkers (Biagi 1999, fig. 14: 1-7), possibly used for cast nets (Marrast 2020; Marrast, Béarez, and Charpentier 2020), or maybe the construction of a fish weir in the mangrove swamp.

Seasonality of fishing activities

Our results question the seasonality of site occupation. Today, the pelagic species observed in Ra's al-Hamra's faunal assemblage represent an important part of the fishery landings in the

Sultanate of Oman, especially in Muscat. Unfortunately, the data provided by the Ministry of Agriculture and Fisheries (M.A.F. 2013, 2015, 2016, 2018) are not precise enough, with many species (e.g., large jacks, emperors, seabreams) mixed into a single category, which does not allow for a detailed comparison of the data. However, it is clear that some species are caught all year round, while others are available in abundance on a seasonal basis (Figures 6–7).

Figure 6. Compiled data (in metric tons) for 4 years (2013, 2015, 2016, 2018) of artisanal fisheries in Muscat governorate for Thunnus tonggol, Thunnus albacares, Euthynnus affinis, and large jacks (Carangidae) (M.A.F. 2013, 2015, 2016, 2018). The thick line represents the mean. The filled area is the standard deviation of the mean

Figure 7. Ecological distribution of the main taxa found at both sites according to Froese and Pauly (2021), Randall (1995) and personal observations. Vector objects are from Vecteezy.com (Graphics RF)

Jacks and trevallies are regularly caught throughout the year, but this is not the case for tunas and kawakawas (Figure 6A-B, D). In the Muscat landings, kawakawas appear to be continuously caught, but between October and November catches peak (Figure 6C). For longtail tuna, the major peak is between May and June, but abundance is high until October (Figure 6A). Yellowfin tuna is mostly present between September and November (Figure 6B). The occurrence of scombrids in the summer months on the shores of the Gulf of Oman could be linked to the Southwest monsoon (June - September), which brings moist air masses. At the same time, an upwelling phenomenon occurs along the southern coast of Oman, which also affects the Gulf of Oman (Al-Azri et al. 2010; Elliot and Savidge 1990; Watanabe et al. 2017). Upwelling brings nutrient-rich waters to the surface, allowing an important trophic network to

develop, including schools of small pelagics, much sought after by scombrids and other pelagic species (Lee et al. 2000). As for sardines, even if they are available all year round, they are particularly important during the cooler season (Northeast monsoon), between November to March (Al-Jufaili and Al-Jahwari 2011). However, this could be partly due to the fact that weather (strong winds) and sea conditions (big waves) affect the sardine beach seine fishery during the summer. It is likely that coastal communities have been highly dependent on the summer upwelling phenomenon linked to the Southwest monsoon (Biagi and Salvatori 1986; Cavulli and Scaruffi 2013).

Since jacks were important at both sites, we can conclude that the fishery for coastal pelagic species could have been all year round, with tuna fishing most active in the autumn and winter. Though it should be noted that pelagic fishing is also associated with significant demersal fishing and a high exploitation of the mangrove and estuarine environment.

As previously mentioned, mangroves are used as a nursery by many species. In this region of the world, seabreams (*Acanthopagrus* spp. and *Rhabdosargus* spp.), scads (*Trachurus indicus*, *S. crumenophthalmus*), some jacks, trevallies, mojarras (Gerreidae), and goatfishes (Mullidae) spawn in the spring, suggesting that juveniles may be found in the mangrove in summer–autumn. In addition to these juveniles, which are characteristic of this environment, many adult fish are also found in the mangrove: jacks and trevallies, mojarras, and goatfishes all frequent the mangrove in autumn; sardines are spotted between winter and spring; and groupers, parrotfishes, and goatfishes in summer. In general, a peak in mangrove frequentation in this part of the world seems to be marked between winter and spring (Abu El-Regal 2013; Abu El-Regal and Ibrahim 2014; Laroche, Baran, and Rasoanandrasana 1997). Barracudas (Sphyraenidae) and small sharks can also be found throughout the year (Marrast, Béarez, and Charpentier 2020). All studies suggest that juveniles can be found in the mangrove at any time,

with a peak occurrence between spring and summer, suggesting that this was the best time for the exploitation of this marine resource in this ecosystem (Abu El-Regal 2013; Abu El-Regal and Ibrahim 2014; Lidour, Béarez, and Méry 2020; Robertson and Duke 1987).

Seasonality of site occupation

The exploitation of diverse resources during the 2,000 year occupation at Ra's al-Hamra implies that the communities had an excellent knowledge of the marine coastline, wildlife ecology, and climatic and environmental features (monsoon and upwelling). This fits well with fisher-hunter-gatherer patterns whose subsistence strategies are based on the acquisition of wild resources, implying a long familiarity with biotopes and wildlife ecology (Acheson 1981; Collet 1991). This multifaceted environment allows diversification in the resources consumed, enabling fluctuations in seasonal resources and any possible shortages to be offset (Biagi et al. 1984; Yesner et al. 1980).

Hunter-gatherers vary in their degree of mobility and may occupy different camps throughout the year. According to Yesner et al. (1980), this makes adaptation to resources possible, allowing the exploitation of several ecological niches (Figure 7). However, as hunters do not adapt to a single environment for their subsistence strategy, the long-term use of space can vary over the year. Fishing activities involve the making and handling of fishing gear, as well as fish preparation and storage activities, and require a high degree of social cooperation to manage the capture and arrival of large quantities of fish (Camelin 1995; Collet 1991). Such cooperation, therefore, fits better with a more sedentary lifestyle.

Until recently, traditional Bedouins on the coast of the Gulf of Oman had to move several times a year because none of the resources they exploited was permanent (Lancaster and Lancaster 1995; Miles 1910). Early ethnographic comparisons have established that shell

middens indicate winter camps and that Bedouin communities moved onto the piedmont in the primeval period and then into the mountain valleys in small, scattered groups (Cleuziou 2005; Cordes and Scholz 1980; Lancaster and Lancaster 1995). Within archaeological communities, some authors even talk of large-scale transhumance between coastal sites and inland pastures (al-Buhais 18, BHS-18) during the arid periods of the fourth millennium BC (Kutterer and Uerpmann 2017). Even if it is reasonable to consider the idea of small scale nomadism along the wadis between the coast and the foothills, due to the high temperatures recorded on the coast in summer, and because of the perishable character of the habitat structures that suggests structures were temporary (Cavulli and Scaruffi 2013), our data do not confirm such a scenario. This illustrates the complexity surrounding the seasonal occupation of Neolithic sites in the southeast Arabian Peninsula.

At RH-5, the community seems to be permanently settled (Munoz 2014; Salvatori 2007), observable in the high density of habitat structures (Marcucci et al. 2011) and no phases of abandonment (Béguier and Marcucci 2018; Marcucci et al. 2011; Marcucci, Badel, and Genchi 2021). Moreover, anthropological studies have established a certain genetic isolation (reduced tooth size) and increasing population density, suggesting the sedentary or semi-sedentary lifestyle of the RH-5 inhabitants (Macchiarelli 1989). This hypothesis is reinforced by geochemical analyses on human skeletons, which indicate a strong dependence on marine resources (Zazzo, Munoz, and Saliège 2014; Zazzo et al. 2016). This is consistent with the multi-niche ecological arguments and the slaughter pattern observed in the domestic animals present in the graves. Sometimes the remains indicated more than one and sometimes up to six animals; “*such slaughtering events require stable animal populations*” (Uerpmann 2003, 79).

CONCLUSION

623 At both sites, our results show that all the fish could have been caught in the vicinity of the
624 site. The residents of RH-6 and RH-5 intensively exploited pelagic fish, either by hook-and-
625 line or by setting nets at a reasonable distance from the coast. However, they also had to
626 supplement this subsistence by setting nets in the mangrove, which provided the opportunity to
627 catch many small fish.

628
629 For RH-6, our study is the first to provide results based on such a large corpus of remains,
630 highlighting a predominance of jacks over Thunnini during the entire period of occupation.
631 Fishing activities were mainly oriented toward the exploitation of medium-sized pelagic fish,
632 especially different Thunnini and jacks, which had approximately the same importance. While
633 these pelagic species dominate throughout the occupation, a gradual decrease in pelagic fish is
634 observed over time.

635 The results of our analysis show the fishing activities practiced during part of the first
636 occupation of the RH-5 site (3910-3640 cal BC), following the end of the RH-6 occupation.
637 The fishery during phases 1 and 2 mainly exploited small pelagic species, such as sardines or
638 bigeye scads. Their frequency is even greater during phases 3 and 4, where they constitute more
639 than 90% of the assemblage. This indicates fishing with small-meshed gillnets or beach seines,
640 as the hooks found at the site are too large to catch such small species.

641 However, a few medium-sized fish were also present, corresponding mainly to individuals
642 of pelagic species, such as tunas or various jacks. These taxa are easy to catch using hook-and-
643 lines thrown from a small boat or even from rocky points, such as those found at the foot of the
644 Ra's al-Hamra cliff. Our study also showed the importance of sieving with a small mesh, which
645 in this case revealed the diversity of the environments exploited and the diversity of fishing
646 techniques implemented at a fishing station in the Neolithic period. These results are extremely
647 promising for redefining the role of fish in the diet and daily life of Neolithic populations.

Indeed, although the importance of marine resources has always been considered by researchers, quantitative and detailed results are still scarce.

In contrast to many contemporary sites in the Persian Gulf (Beech 2004; Lidour, Béarez, and Méry 2020; Lidour and Beech 2019) and in the Arabian Sea, such as SWY-1 and RWY-1 (Berger et al. 2020; Marrast, Béarez, and Charpentier 2020) where lagoons have been mainly exploited, the Neolithic inhabitants of the capital area of Muscat appear to have specialized in the exploitation of coastal pelagic species. This kind of exploitation has already been highlighted at a few sites in the Persian Gulf, particularly at Akab, where carangids and scombrids represented 50% of the material (Lidour et al. 2020), and at Khor Milkh 1 (KM-1) (Uerpmann and Uerpmann 2003, 194), just south of Ra's al-Hamra. However, the specialization of this type of fishing (more than 80% at RH-6 and RH-5) in the Gulf of Oman is, for the moment, unique in this region. New ichthyoarchaeological studies in the region and their publication, especially from more recent sites (Bronze Age) or from areas that are still poorly known, will make it possible to complete the picture of ancient fisheries in this part of the world and, in particular, to better understand the reasons that led the coastal inhabitants to turn to pelagic resources. Similarly, the integration of analyses of other marine resources (crustaceans and crabs, sea turtles, cetaceans, etc.) would provide an accurate picture of the dietary habits during the Middle Holocene and improve our understanding of the role of the sea for the coastal populations of southeast Arabia.

SUPPLEMENTAL FILES

Supplemental file 1. List and NISP of taxa recovered at RH-6 and RH-5, arranged in taxonomic order. B: benthic; D: demersal; EP: epipelagic; P: pelagic. For fish behaviour, we followed the information from Randall (1995) and FishBase (Froese and Pauly 2021)

673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697

Supplemental file 2. Illustration of different bones from main taxa at both sites and possible fishing environment. Pelagic fishes: (A) Thunnus albacares, precaudal vertebra (ventral view); (B) Thunnus tonggol, caudal vertebra (lateral view); (C) Euthynnus affinis, precaudal vertebra (ventral view); (D) Caranx sexfasciatus, right cleithrum (lateral view); (E) Carangoides chrysophrys, supraoccipital hyperostosis (lateral view); (F) Gnathanodon speciosus, precaudal vertebra (ventral view). Demersal fishes: (G) Sufflamen fraenatum, left premaxilla (lateral view); (H): Scarus sp., left dentary (lateral view); (I) Epinephelus sp., right premaxilla (lateral view); (J) estuary view; (K) open sea view from the shore

Supplemental file 3. Illustration of different bones from main taxa at both sites and possible fishing environment. Small pelagic fishes: (A) Selar crumenophthalmus, precaudal vertebra (ventral view); (B) Clupeidae (cf. Sardinella longiceps), precaudal vertebra (ventral view); (C) Alepes djedaba, first vertebra (ventral view); (D) Rastrelliger kanagurta, caudal vertebra (lateral view); (E) Mugilidae, precaudal vertebra (ventral view). Juvenile demersal fishes: (F) Pomadasys sp., first vertebra (ventral view); (G) Sparoidea, teeth; (H-I) Qurum mangrove views

ACKNOWLEDGEMENTS

The authors (AM, PB) would like to thank the PEP's Mangrove and the ANR NeoArabia programs (dir. J.F. Berger) for funding field missions to the Sultanate of Oman. The authors are very grateful to the Minister of Heritage and Tourism of the Sultanate of Oman, H.E. Salim Mohammed Almahruqi, as well as Mr. Sultan Saif Al-Bakri, Director General for Archaeology, Mr. Khamis Al-Asmi, Director of the Department of Excavations and Archaeological Studies. Thanks are also due to K. Debue and B. David for assistance with the sorting of the 1 mm sieved

698 sediments, to A. Coppa for the funding of the date of sector B2 of RH 5, to A. Zazzo and M.
699 Tengberg for their advice, to H. David for the use of her map of the sites of Ra's al-Hamra, and
700 to J. Cucchi for copy editing. We would also like to thank the reviewers and the editor for their
701 contributions to our manuscript. Finally, the authors address their posthumous thanks to M.
702 Tosi who worked for a long time on the sites of Ra's al-Hamra and did much for the archaeology
703 of this area and the Sultanate.

704
705 **DECLARATION OF INTEREST STATEMENT**

706 The authors declare that they have no conflict of interest.
707

709 Abu El-Regal, M. 2013. Spawning seasons, spawning grounds and nursery grounds of some
710 Red Sea fishes. *The Global Journal of Fisheries and Aqua* 6(6): 105-125.

711 Abu El-Regal, M., and N. K. Ibrahim. 2014. Role of mangroves as a nursery ground for juvenile
712 reef fishes in the southern Egyptian Red Sea. *Egyptian Journal of Aquatic Research* 40:
713 71–78. doi:10.1016/j.ejar.2014.01.001.

714 Acheson, J. M. 1981. Anthropology of fishing. *Annual Review of Anthropology* 10: 275–316.
715 doi:10.1146/annurev.an.10.100181.001423.

716 Al-Azri, A. R., S. A. Piontkovski, K. A. Al-Hashmi, J. I. Goes, and H. R. do Gomes. 2010.
717 Chlorophyll a as a measure of seasonal coupling between phytoplankton and the
718 monsoon periods in the Gulf of Oman. *Aquatic Ecology* 44(2): 449–461.

719 Al-Azri, A. R., S. A. Piontkovski, K. A. Al-Hashmi, J. I. Goes, H. R. do Gomes, and P. M.
720 Glibert. 2014. Mesoscale and nutrient conditions associated with the massive 2008
721 *Cochlodinium polykrikoides* bloom in the Sea of Oman/Arabian Gulf. *Estuaries and*
722 *Coasts* 37 (2): 325–338. doi:10.1007/s12237-013-9693-1.

723 Al-Jufaili, S. M., and O. S. Al-Jahwari. 2011. The Omani coastal traditional sardine fishery
724 1994-2007: A review. *Agricultural and Marine Sciences* 16: 1–12.

725 Al-Marshudi, A. S., and H. Kotamaga. 2006. Socio-economic structure and performance of
726 traditional fishermen in the Sultanate of Oman. *Marine Resource Economics* 21 (2):
727 221–230. doi:10.1086/mre.21.2.42629506.

728 Anderson, A. 2013. The Antiquity of sustained offshore fishing. *Antiquity* 87 (337): 879–885.
729 doi:10.1017/S0003598X00049541.

730 Béarez, P., F. Fuentes-Mucherl, S. Rebolledo, D. Salazar, and L. Olguín. 2016. Billfish foraging
731 along the northern coast of Chile during the Middle Holocene (7400–5900cal BP).
732 *Journal of Anthropological Archaeology* 41: 185–195. doi:10.1016/j.jaa.2016.01.002.

733 Béarez, P., P. Gay, and R. Lunniss. 2012. Sea fishing at Salango (Manabí Province, Ecuador)
734 during the Middle Formative Machalilla phase. *Latin American Antiquity* 23 (2): 195-
735 214.

736 Beech, M. J. 2004. *In the land of the Ichthyophagi: Modelling fish exploitation in the Arabian*
737 *Gulf and Gulf of Oman from the 5th millennium BC to the Late Islamic period*. BAR
738 International Series 1217. Oxford: Archaeopress.

739 Beech, M. J., K. Strutt, L. Blue, A. K. Al-Kaabi, W. A. Omar, A. A. A. H. El-Faki, A. R.
740 Lingareddy, and J. Martin. 2016. Ubaid-related sites of the southern Gulf revisited: The
741 Abu Dhabi coastal heritage initiative. *Proceedings of the Seminar for Arabian Studies*
742 46: 9–23. doi:10.2307/45163414.

743 Béguier, I., and L. G. Marcucci. 2018. Première approche micromorphologique de la fin de
744 l’occupation néolithique d’un amas coquillier de la côte omanaise : données
745 préliminaires sur les modes d’occupation du site de RH-5 dans la seconde moitié du 4e
746 millénaire BCE [Initial micromorphological approach of the Neolithic occupation of a
747 shell midden on the Omani coastline: Preliminary data on the occupation patterns of the
748 RH-5 site in the second half of the 4th millennium]. *ArcheoSciences* 42 (2): 7–30.
749 doi:10.4000/archeosciences.5667.

750 Berger, J.-F., V. Charpentier, R. Crassard, C. Martin, G. Davtian, and J. A. López-Sáez. 2013.
751 The dynamics of mangrove ecosystems, changes in sea level and the strategies of
752 neolithic settlements along the coast of Oman (6000–3000 Cal. BC). *Journal of*
753 *Archaeological Science* 40 (7): 3087–3104. doi:10.1016/j.jas.2013.03.004.

754 Berger, J.-F., S. Cleuziou, G. Davtian, M. Cattani, F. Cavulli, V. Charpentier, M. Cremaschi, J.
755 Giraud, P. Marquis, C. Martin, et al. 2005. Evolution paléogéographique du Ja’alan
756 (Oman) à l’Holocène moyen : Impact sur l’évolution des paléomilieus littoraux et les

stratégies d'adaptation des communautés humaines [Paleogeographic evolution of Ja'alan (Oman) in the Middle Holocene: Impact on the evolution of coastal paleoenvironments and on the adaptative strategies of human communities]. *Paléorient* 31 (1): 46–63.

- Berger, J.-F., R. Guilbert-Berger, A. Marrast, O. Munoz, H. Guy, A. Barra, J. A. López-Sáez, S. Pérez-Díaz, M. Mashkour, K. Debue et al. 2020. First contribution of the excavation and chronostratigraphic study of the Ruways 1 Neolithic shell midden (Oman) in terms of neolithisation, palaeoeconomy, social-environmental interactions and site formation processes. *Arabian Archaeology and Epigraphy* 31 (1): 32–49. doi:10.1111/aae.12144.
- Biagi, P. 1987. The prehistoric fishermen settlements of RH5 and RH6 at Qurum, Sultanate of Oman. *Proceedings of the Seminar for Arabian Studies* 17: 15–19. doi:10.2307/41223038.
- Biagi, P. 1994. A radiocarbon chronology for the aceramic shell-middens of coastal Oman. *Arabian Archaeology and Epigraphy* 5: 17–31. doi:10.1111/j.1600-0471.1994.tb00053.x.
- Biagi, P. 1999. Excavations at the shell-midden of RH6 1986–1988 (Muscat, Sultanate of Oman). *Al-Rāfidān* 20: 57–84.
- Biagi, P. 2006. The shell-middens of the Arabian Sea and Gulf: Maritime connections in the seventh millennium BP? In *The city in the arab world in light of archaeological discoveries: Evolution and development*, ed. A. R. Al-Ansary, K. I. Al-Muaikel, and A. M. Alsharekh, 7–18. Al-Jouf: Abdul Rahman Al-sudairy Foundation.
- Biagi, P., and R. Maggi. 1987. Archaeological surveys along the Oman coast: Preliminary results of five years of research (1983–1987). *South Asian Archaeology* 1: 1990.
- Biagi, P., and R. Nisbet. 1992. Environmental history and plant exploitation at the aceramic sites of RH5 and RH6 near the mangrove swamp of Qurum (Muscat-Oman). *Bulletin de La Société Botanique de France. Actualités Botaniques* 139 (2–4): 571–578. doi:10.1080/01811789.1992.10827129.
- Biagi, P., and R. Nisbet. 1999. The shell-midden sites of RH-5 and RH-6 (Muscat, Sultanate of Oman) in their environmental setting. *Archaeologia Polona* 37: 31–47.
- Biagi, P., and R. Nisbet. 2006. The prehistoric fisher-gatherers of the western coast of the Arabian Sea: A case of seasonal sedentarization? *World Archaeology* 38 (2): 220–238. doi:10.1080/00438240600708188.
- Biagi, P., S. Salvatori. 1986. Gli scavi nell'insediamento preistorico e nella necropoli di Ra's al-Hamra 5 (Muscat, Oman), 1980–1985. *Rivista di Archeologia* 10: 5–14.
- Biagi, P., W. Torke, M. Tosi, and H.-P. Uerpmann. 1984. Qurum: A case study of coastal archaeology in northern Oman. *World Archaeology* 16 (1): 43–61. doi:10.1080/00438243.1984.9979915.
- Biagi, P., and R. A. Travers. 1985. Non-mammalian osteological remains and fishing implements at RH-5 and RH-6, Muscat. *East and West* 35 (4): 407–410.
- Camelin, S. 1995. Les pêcheurs de Shihr. Transmission du savoir et identité sociale. *Arabian Humanities*. 3. doi:10.4000/cy.80.
- Carter, R. A., and H. E. W. Crawford. 2010. *Maritime interactions in the Arabian Neolithic: The evidence from H3, As-Sabiyah, an Ubaid-related site in Kuwait*. Leiden: Brill.
- Cartwright, C. R. 2004. Reconstructing the use of coastal resources at Ra's Al-Hadd, Oman, in the Third Millennium BC. *Proceedings of the Seminar for Arabian Studies* 34: 45–51.
- Cavulli, F., and S. Scaruffi. 2013. Thoughts on nomadism in Middle Holocene Oman. *Arabian Archaeology and Epigraphy* 24: 15–27. doi:10.1111/aae.12023.
- Charpentier, V. 2008. Hunter-gatherers of the 'Empty Quarter of the Early Holocene' to the Last Neolithic societies: Chronology of the Late Prehistory of south-eastern Arabia (8000–3100 BC). *Proceedings of the Seminar for Arabian Studies* 38: 93–115.

- Charpentier, V., J.-F. Berger, R. Crassard, F. Borgi, G. Davtian, S. Méry, and C. S. Phillips. 2013. Conquering new territories: When the first black boats sailed to Masirah Island. *Proceedings of the Seminar for Arabian Studies* 43: 85–98.
- Chenorkian, R. 1996. *Pratique archéologique statistique et graphique*. Méthodes et techniques. Paris: Ed. Errance, Lattes: Association pour la diffusion de l'archéologie méridionale.
- Cleuziou, S. 2003. Early Bronze Age trade in the Gulf and the Arabian Sea: The Society behind the Boats. In *Archaeology of the United Arab Emirates*, edited by Daniel T. Potts, Hasan Al Naboodah, and Peter Hellyer, 133–150. Abu Dhabi, 2001: Trident Press Ltd.
- Cleuziou, S. 2005. Pourquoi si tard ? Nous avons pris un autre chemin. L'Arabie des chasseurs-cueilleurs de l'Holocène au début de l'Age du Bronze [Why so late? We have taken a different route. The Arabia of the hunter-gatherers from the Holocene to the Early Bronze Age]. In *Aux marges des grands foyers du Néolithique. Périphéries débitrices ou créatrices ?*, ed. J. Guilaine, 123–148. Collection des Hespérides. Paris: Editions Errance.
- Cleuziou, S., and M. Tosi. 2007. *In the shadow of the ancestors. The prehistoric foundations of the early Arabian civilization in Oman*. Muscat: Ministry of Heritage & Culture, Sultanat of Oman.
- Collet, S. 1991. Guerre et pêche : quelle place pour les sociétés de pêcheurs dans le modèle des chasseurs-cueilleurs ? [War and fishing: What place for fishing societies in the hunter-gatherer model?]. *Social Science Information* 30 (3): 483–522.
- Collette, B. B. 2001. Scombridae. Tunas (Also, albacore, bonitos, mackerels, seerfishes, and wahoo). *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific* 6: 3721–3756.
- Collette, B. B., and C. E. Nauen. 1983. *Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date*. Vol. 2. FAO Species Catalogue. Rome: Food & Agriculture Organization of the United Nations. <http://www.fao.org/docrep/009/ac478e/ac478e00.htm>.
- Coppa, A., R. Macchiarelli, S. Salvatori, and G. Santini. 1985. The prehistoric graveyard of Ra's al-Hamra (RH 5). *Journal of Oman Studies*, 8(1): 97–102.
- Cordes, R., and F. Scholz. 1980. *Bedouins, wealth, and change. A study of rural development in the United Arab Emirates and the Sultanate of Oman*. Tokyo: The United Nations University. <https://www.cabdirect.org/cabdirect/abstract/19811875308>.
- Durante, S., and M. Tosi. 1977. The aceramic shell middens of Ra's al-Hamra: A preliminary note. *The Journal of Oman Studies* 3 (2): 137–162.
- Elliott, A. J., and G. Savidge. 1990. Some features of the upwelling off Oman. *Journal of Marine Research* 48 (2): 319–333.
- Fleitmann, D., S. J. Burns, A. Mangini, M. Mudelsee, J. Kramers, I. Villa, U. Neff, A. A. Al-Subbary, A. Buettner, D. Hippler, et al. 2007. Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra). *Quaternary Science Reviews* 26 (1): 170–188. doi:10.1016/j.quascirev.2006.04.012.
- Frazier, J. G., V. Azzarà, O. Munoz, L. G. Marcucci, É. Badel, F. Genchi, M. Cattani, M. Tosi, and M. Delfino. 2018. Remains of leatherback turtles, *Dermochelys coriacea*, at Mid-Late Holocene archaeological sites in coastal Oman: Clues of Past Worlds. *PeerJ* 6: 1–34. doi:10.7717/peerj.6123.
- Fricke, R., W. N. Eschmeyer, and R. Van der Laan, eds. 2021. Eschmeyer's Catalog of Fishes: Genera, Species, References. Electronic version accessed 8 June 2021. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
- Froese, R., and D. Pauly, eds. 2021. FishBase. World Wide Web electronic publication. www.fishbase.org

- Goddard, D. 1985. Exeter maritime museum. *The Mariner's Mirror* 71 (2). Routledge: 201–214. doi:10.1080/00253359.1985.10656025.
- Grandcourt, E. M. 2012. Reef fish and fisheries in the Gulf. In *Coral Reefs of the Gulf: Adaptation to Climatic Extremes*, edited by Bernhard M. Riegl and Sam J. Purkis, 127–161. Coral Reefs of the World 3. Dordrecht: Springer Netherlands. doi:10.1007/978-94-007-3008-3_8.
- Grandcourt, E. M., T. Z. Al Abdessalaam, F. Francis, and A. T. Al Shamsi. 2004. Biology and stock assessment of the sparids, *Acanthopagrus bifasciatus* and *Argyrops spinifer* (Forsskal, 1775), in the southern Arabian Gulf. *Fisheries Research* 69 (1): 7–20.
- Johnson, D. V. 2016. Unusual date palm products: Prayer beads, walking sticks and fishing boats. *Emirates Journal of Food and Agriculture* 28 (1): 12–16. doi:10.9755/ejfa.2015-11-1021.
- Karachle, P. K., D. Dimarchopoulou, A. C. Tsikliras. 2020. Is shore-based recreational fishing in Greece an unregulated activity that increases catch uncertainty? *Regional Studies in Marine Science* 36: 101273
- Kulbicki, M. 1997. Bilan de 10 ans de recherche (1985-1995) par l'ORSTOM sur la diversité, la densité, la biomasse et la structure trophique des communautés des poissons lagonaires et récifaux en Nouvelle-Calédonie [Results of ten years of research (1985-1995) by ORSTOM on diversity, density, biomass, and community trophic structure of the lagoon and reef fishes in New Caledonia]. *Cybium* 21 (1): 47–79.
- Laegdsgaard, P., and C. Johnson. 2001. Why do juvenile fish utilise mangrove habitats? *Journal of Experimental Marine Biology and Ecology* 257 (2): 229–253. doi:10.1016/S0022-0981(00)00331-2.
- Lancaster, W., and F. Lancaster. 1995. Nomadic fishermen of Ja'alân, Oman. *Nomadic Peoples* 36/37: 227–243.
- Laroche, J., E. Baran, and N. B. Rasoanandrasana. 1997. Temporal patterns in a fish assemblage of a semiarid mangrove zone in Madagascar. *Journal of Fish Biology* 51 (1): 3–20. doi:10.1111/j.1095-8649.1997.tb02509.x.
- Lee, C. M., B. H. Jones, K. H. Brink, and A. S. Fischer. 2000. The upper-ocean response to monsoonal forcing in the Arabian Sea: Seasonal and spatial variability. *Deep-Sea Research Part II: Tropical Studies in Oceanography* 47 (7–8): 1177–1226. doi:10.1016/S0967-0645(99)00141-1.
- Lézine, A.-M., S. J. Ivory, P. Braconnot, and O. Marti. 2017. Timing of the southward retreat of the ITCZ at the end of the Holocene humid period in southern Arabia: Data-model comparison. *Quaternary Science Reviews* 164: 68–76. doi:10.1016/j.quascirev.2017.03.019.
- Lidour, K., P. Béarez, V. Charpentier, and S. Méry. 2020. The prehistoric fisheries of Akab Island (United Arab Emirates): New insights into coastal subsistence during Neolithic in eastern Arabia. *The Journal of Island and Coastal Archaeology* 15 (1): 80–103. doi:10.1080/15564894.2018.1531330.
- Lidour, K., P. Béarez, and S. Méry. 2020. Analysis of archaeological fish remains provides new insights into seasonal mobility in eastern Arabia during the Mid-Holocene. The subsistence fisheries of UAQ36, Umm al-Quwain lagoon (United Arab Emirates). *Journal of Archaeological Science: Reports* 30: 102223. doi:10.1016/j.jasrep.2020.102223.
- Lidour, K., and M. J. Beech. 2019. The numerous islands of Ichthyophagi': Neolithic fisheries of Delma Island, Abu Dhabi Emirate (UAE). *Proceedings of the Seminar for Arabian Studies* 49: 207–222.
- Macchiarelli, R. 1989. Prehistoric 'fish-eaters' along the eastern Arabian coasts: Dental variation, morphology, and oral health in the Ra's al-Hamra community (Qurum,

Sultanate of Oman, 5th–4th Millennium BC). *American Journal of Physical Anthropology* 78 (4): 575–594. doi:10.1002/ajpa.1330780411.

M.A.F. (Ministry of Agriculture and Fisheries). 2013. *Fisheries statistics book*. Mascate: Ministry of Agriculture and Fisheries - General directorate of Planning and Development Fisheries Statistic Department.

M.A.F. (Ministry of Agriculture and Fisheries). 2015. *Fisheries statistics book*. Mascate: Ministry of Agriculture and Fisheries - General directorate of Planning and Development Fisheries Statistic Department.

M.A.F. (Ministry of Agriculture and Fisheries). 2016. *Fisheries statistics book*. Mascate: Ministry of Agriculture and Fisheries - General directorate of Planning and Development Fisheries Statistic Department.

M.A.F. (Ministry of Agriculture and Fisheries). 2018. *Fisheries statistics book*. Mascate: Ministry of Agriculture and Fisheries - General directorate of Planning and Development Fisheries Statistic Department.

Marcucci, L. G. 2015. *I siti preistorici di RH-5 e RH-6 a Ra's al-Hamra (Muscat, Sultanato d'Oman): analisi comparativa della stratigrafia, delle strutture di abitato e della cultura materiale. Continuità e sviluppo di due villaggi di pescatori-raccoglitori [The prehistoric sites of RH-5 and RH-6 at Ra's al-Hamra (Muscat, Sultanate of Oman): Comparative analysis of the stratigraphy, the living structures and the material culture. Continuity and development of two villages of fisher-gatherers]*. PhD diss., Université Paris 1 Panthéon-Sorbonne, Università di Bologna.

Marcucci, L. G., É. Badel, and F. Genchi. 2021. *Prehistoric fisherfolk of Oman: The Neolithic village of Ras Al-Hamra RH-5*. The Archaeological heritage of Oman, Vol. 6. Oxford: Archaeopress.

Marcucci, L. G., É. Badel, F. Genchi, O. Munoz, A. Todero, and M. Tosi. 2014. New investigations at the prehistoric shell midden of Ra's al-Hamra 6 (Sultanate of Oman): Results of the 2012 and 2013 Excavation Seasons. *Proceeding for Arabian Studies* 44: 235–256.

Marcucci, L. G., F. Genchi, É. Badel, and M. Tosi. 2011. Recent investigations at the prehistoric site RH-5 (Ra's al-Hamra, Muscat, Sultanate of Oman). *Proceeding for Arabian Studies*. 41: 201–222.

Marrast, A. 2020. *Stratégies d'exploitation des environnements côtiers arides : les activités halieutiques du Néolithique à l'âge du Bronze dans le Sud Est de la péninsule arabique (Sultanat d'Oman) [Exploitation strategies in arid coastal environments: Fishing activities from the Neolithic to the Bronze Age in the south-eastern Arabian Peninsula (Sultanate of Oman)]*. PhD diss., Muséum national d'histoire naturelle.

Marrast, A., P. Béarez, and V. Charpentier. 2020. Sharks in the lagoon? Fishing exploitation at the neolithic site of Suwayh 1 (Ash Sharqiyah Region, Arabian Sea, Sultanate of Oman). *Arabian Archaeology and Epigraphy* 31 (1): 178–193. doi:10.1111/aae.12136.

McClure, H. A. 1976. Radiocarbon chronology of Late Quaternary lakes in the Arabian desert. *Nature* 263 (5580): 755–756. doi:10.1038/263755a0.

Miles, S. B. 1910. On the border of the great desert: A journey in Oman. *Geographical Journal* 36 (2): 159–178.

Munoz, O. 2014. *Pratiques funéraires et paramètres biologiques dans la péninsule d'Oman du Néolithique à la fin de l'âge du Bronze ancien (Ve-IIIe millénaires av. NE) [Funerary practices and biological parameters in the Oman Peninsula from the Neolithic to the end of the Early Bronze Age (5th-3rd mill. BCE)]*. PhD diss., Université Paris 1 Panthéon-Sorbonne. <http://www.theses.fr/2014PA010514>.

Nagelkerken, I., C. M. Roberts, G. Van der Velde, M. Dorenbosch, M. C. Van Riel, E. Cocheret De La Moriniere, and P. H. Nienhuis. 2002. How important are mangroves and seagrass

- 956 beds for coral-reef fish? The nursery hypothesis tested on an island scale. *Marine*
 957 *Ecology Progress Series* 244: 299–305.
- 958 Nourissier, A., N. Nocus, and A. Dufraisse. 2016. *ACACIA: Automatical Creation of*
 959 *Anthracological Diagram. The User Guide*. Available on
 960 <https://dendrac.mnhn.fr/spip.php?rubrique69>
- 961 Pickard, C., and C. Bonsall. 2004. Deep-sea fishing in the European Mesolithic: Fact or fantasy?
 962 *European Journal of Archaeology* 7 (3). Routledge: 273–290.
 963 doi:10.1177/1461957104056504.
- 964 Poplin, F. 1976. A propos du nombre de restes et du nombre d'individus dans les échantillons
 965 d'ossements [About the number of remains and the number of individuals in bone
 966 samples]. *Cahiers du Centre de recherches préhistoriques* 5: 61-75.
- 967 Randall, J. E. 1995. *Coastal fishes of Oman*. University of Hawaii Press.
- 968 Richardson N. and M. Dorr. 2003. *The craft heritage of Oman*. Dubai: Motivate Publishing.
- 969 Robertson, A. I., and N. C. Duke. 1987. Mangroves as nursery sites: Comparisons of the
 970 abundance and species composition of fish and crustaceans in mangroves and other
 971 nearshore habitats in tropical Australia. *Marine Biology* 96 (2): 193–205.
 972 doi:10.1007/BF00427019.
- 973 Salvatori, S. 2007. *The prehistoric graveyard of Ra's Al Hamrā 5, Muscat, Sultanate of Oman*.
 974 Mascate: Ministry of Heritage and Culture, Sultanate of Oman.
- 975 Strømme, T. 1986. The pelagic and demersal fish resources of Oman. Results of the R/V Dr.
 976 Fridtjof Nansen surveys in Oman 1983–84. Final report from surveys 1983–84. Bergen:
 977 Institute of Marine Research. [http://agris.fao.org/agris-](http://agris.fao.org/agris-search/search.do?recordID=XF2016072613)
 978 [search/search.do?recordID=XF2016072613](http://agris.fao.org/agris-search/search.do?recordID=XF2016072613).
- 979 Tengberg, M. 2005. Les forêts de la mer. Exploitation et évolution des mangroves en Arabie
 980 orientale du Néolithique à l'époque islamique [The forests of the sea. Exploitation and
 981 evolution of mangroves in Eastern Arabia from the Neolithic to the Islamic period].
 982 *Paléorient* 31 (1): 39–45.
- 983 Thomas R., M. Tengberg, C. Moulhérat, V. Marcon and R. Besenval. 2012. Analysis of a
 984 protohistoric net from Shahi Tump, Baluchistan (Pakistan). *Archaeological and*
 985 *Anthropological Sciences* 4: 15–23. doi: 10.1007/s12520-011-0078-8
- 986 Tosi, M. 1975. Notes on the distribution and exploitation of natural resources in ancient Oman.
 987 *Journal of Oman Studies* 1: 187–206.
- 988 Uerpmann, H.-P., and M. Uerpmann. 2003. *The capital area of northern Oman. Part III, Stone*
 989 *Age sites and their natural environment*. Wiesbaden: Dr. Ludwig Reicherd Verlag.
- 990 Uerpmann, M. 2003. The Dark Millennium: Remarks on the Final Stone Age in the Emirates
 991 and Oman. In *Archaeology of the United Arab Emirates*, ed. D. T. Potts, H. Al
 992 Naboodah, and P. Hellyer, 74–81. Abu Dhabi: Trident Press Limited.
- 993 Uerpmann M. and H.-P. Uerpmann, 2018. Chapter 20. Faunal remains and subsistence
 994 strategies. In *Dosariyah: Reinvestigating a Neolithic coastal community in eastern*
 995 *Arabia*. Ed. P. Drechsler: 384-422. British Foundation for the Study of Arabia
 996 Monographs 19. Oxford: Archaeopress.
- 997 Vita-Finzi, C. 1982. Recent coastal deformation near the Strait of Hormuz. *Proceedings of the*
 998 *Royal Society A: Mathematical, Physical and Engineering Sciences* 382 (1783): 441–
 999 457.
- 1000 von den Driesch, A., and H. Manhart, 2000. Fish bones from Al Markh, Bahrain. In
 1001 *Archaeozoology of the Near East IVB. Proceedings of the fourth international*
 1002 *symposium on the archaeozoology of southwestern Asia and adjacent areas*. Eds. M.
 1003 Mashkour, A. M., Choyke, H., Buitenhuis, and F. Poplin: 350-367. Groningen: ARC
 1004 Publicatie.

- Vosmer, T. 2019. Sewn boats in Oman and the Indian Ocean. *International Journal of Nautical Archaeology* 48 (2): 302–313. doi:10.1111/1095-9270.12371.
- Watanabe, T. K., T. Watanabe, A. Yamazaki, M. Pfeiffer, D. Garbe-Schönberg, and M. R. Claereboudt. 2017. Past summer upwelling events in the Gulf of Oman derived from a coral geochemical record. *Scientific Reports* 7(1). doi:10.1038/s41598-017-04865-5
- Wheeler, A., and A. K. G. Jones. 1989. *Fishes*. Cambridge, New York: Cambridge University Press.
- Wilkins, B. 2005. Fishing in the Arabian Sea: A short note on the prehistoric sites RH6 and Ra's al-Jinz 1 in Oman. *Paléorient* 31 (1): 126–130.
- Yesner, D. R., W. S. Ayres, D. L. Carlson, R. S. Davis, R. Dewar, M. R. González Morales, F. A. Hassan, B. Hayden, J. J. Lischka, P. D. Sheets, et al. 1980. Maritime hunter-gatherers: Ecology and prehistory [and Comments and Reply]. *Current Anthropology* 21 (6): 727–750. doi:10.1086/202568.
- Zazzo, A., O. Munoz, É. Badel, I. Béguier, F. Genchi, and L. G. Marcucci. 2016. A revised radiocarbon chronology of the aceramic shell midden of Ra's Al-Hamra 6 (Muscat, Sultanate of Oman): Implication for occupational sequence, marine reservoir age, and human mobility. *Radiocarbon* 58 (2): 383–395. doi:10.1017/RDC.2016.3.
- Zazzo, A., O. Munoz, and J.-F. Saliège. 2014. Diet and mobility in a Late Neolithic population of coastal Oman inferred from radiocarbon dating and stable isotope analysis. *American Journal of Physical Anthropology* 153 (3): 353–364. doi:10.1002/ajpa.22434.

Figure 1. (A) Location, altitudinal and bathymetric mapping of the Qurum mangrove and the Ra's al-Hamra's cape, with location of the archaeological sites; (B) Map of the excavated sectors of RH-6 (Marcucci et al., 2014, fig. 1.A.); and (C) Map of the excavated sectors of RH-5 (Marcucci et al., 2011, fig. 2.)

923x973mm (150 x 150 DPI)

Main taxa	RH-6 (Uerpmann and Uerpmann)		RH-6 (Wilkins 2005)		RH-5 (Uerpmann and Uerpmann 2003)			M.A.F. (2018)		
	NISP	WISP	NISP	WISP	NISP	NISP %	WISP	Catches (tons)	%	(without small pelagic)
Scombridae	23	9.1	311	-	7596	58.9	6527.4	49466	28.1	44.0
Carangidae	22	15.8	74	-	4065	31.5	5934.7	10572	6.0	9.4
Other pelagic fish	2	0.1	17	-	209	1.6	100.5	18203	10.3	16.2
Mugilidae	1	0.2	3	-	11	0.1	5.3	821	0.5	
Small pelagic fish	-	-	-	-	-	-	-	62897	35.7	
Sparidae	2	0.1	47	-	662	5.1	295.4	2533	1.4	2.3
Lethrinidae	-	-	19	-	150	1.2	117.4	5356	3.0	4.8
Serranidae	-	-	56	-	36	0.3	76.6	3359	1.9	3.0
Other demersal fish	1*	0.3	36	-	150	1.2	171.8	15243	8.6	13.5
Sharks/Rays	-	-	7	-	12	0.1	20.2	7777	4.4	6.9
Total NISP	51	25.6	570	0	12891	100	13249.3	176227	100	100

Table 1. Table synthesizing the main results from previous studies on the fish bone remains from RH-6 and RH-5 (for Uerpmann and Uerpmann 2003, NISP have been calculated from identified Actinopterygii and Chondrichthyes), and modern data of artisanal fishing from the governorate of Muscat for 2009-2018 (M.A.F. 2018). *Ariidae

Anatomical element		Total RH-6			Total RH-5
		Trench North (3 mm)	Sector C/TT-Z (3 mm)	Sector C/TT-Z (1 mm)	
Cranium	Dentary	528	496	6	23
	Premaxilla	353	508	10	31
	Maxilla	248	206	1	5
	Quadrate	179	263	4	12
	Opercle	187	209	6	7
	Anguloarticular	166	209	1	4
	Cleithrum	232	114	-	7
	Vomer	133	113	2	2
	Scapula	26	100	1	4
	Pharyngeal bone	49	63	-	1
	Tooth	41	50	406	215
	Hyomandibula	19	61	1	1
	Palatine	17	31	-	4
	Basioccipital/ Parasphenoid	66	50	13	50
	Otolith	-	1	21	2
	Other	248	289	-	44
Vertebral column	First vertebra	284	127	707	1417
	Precaudal vertebra	2069	2103	4065	4349
	Caudal vertebra	3211	3242	5257	2926
	Other	712	231	1	14
Others	Pterygiophore	493	704	17	44
	Scale	-	-	41	-
	Scute	188	1	-	4
Chondr.	Tooth	2	-	2	2
	Vertebra	4	3	-	-
	Other	-	-	1	-
Unidentified remains		113325	46275	472	8559
Total		122780	55449	11035	17727

Table 2. Element distribution on the two sites according to the sector studied and the sieving mesh size (Chondr. = Chondrichthyes).

Figure 2. Taxonomic richness at family and genus levels for different sectors and time periods

233x127mm (150 x 150 DPI)

Figure 3. Illustration of the main taxa distribution (NISP).

(A) RH-6, Trench North, 3 mm; (B) RH-6, Sector C/TT-Z, 3 mm. Made with ACACIA (Nourissier, Nocus, and Dufraisse, 2010)

214x79mm (300 x 300 DPI)

Main taxa		Total Period 1			Total Period 2			Total		
		NISP	%	WISP	NISP	%	WISP	NISP	%	WISP
Carangidae	<i>Carangoides</i> spp.	986	20.09	786.21	852	20.2	618.3	1838	20.2	1404.51
	<i>Caranx</i> spp.	602	12.27	413.07	489	11.6	677.97	1091	12	1091.04
	<i>Gnathanodon speciosus</i>	180	3.67	111.44	147	3.5	154.23	327	3.6	2,65.67
	Other Carangidae	1278	26.04	398.45	1079	25.6	409.43	2357	25.9	807.89
Scombridae	<i>Euthynnus affinis</i>	775	15.79	506.04	609	14.5	377.02	1384	15.2	883.06
	<i>Thunnus albacares</i>	21	0.43	14.07	73	1.7	47.05	94	1	61.13
	<i>Thunnus tonggol</i>	4	0.08	2.48	47	1.1	24.01	51	0.6	26.48
	<i>Thunnus</i> spp.	239	4.87	133.52	146	3.5	82.37	385	4.2	215.89
	Other Scombridae	280	5.7	90.31	319	7.6	106.41	599	6.6	196.72
Other pelagic		98	2	16.23	51	1.2	12.1	149	1.6	28.33
Sparidae	<i>Acanthopagrus</i> spp.	58	1.18	13.84	34	0.8	114.25	92	1	128.09
	<i>Rhabdosargus</i> spp.	20	0.41	5.74	8	0.2	8.28	28	0.3	14.01
	Other Sparidae	44	0.9	16.53	94	2.2	45.51	138	1.5	62.03
Lethrinidae	<i>Lethrinus</i> spp.	80	1.63	19.18	110	2.6	23.75	190	2.1	42.93
	Other Lethrinidae				1	0	0.84	1	0	0.84
Serranidae	<i>Cephalopholis</i> spp.	39	0.79	4.61	21	0.5	7.15	60	0.7	11.76
	<i>Epinephelus</i> spp.	23	0.47	13.75	26	0.6	300.96	49	0.5	314.71
	Serranidae	27	0.55	7.03	20	0.5	28.07	47	0.5	35.09
Other demersal		154	3.14	36.57	82	1.9	27.9	236	2.6	64.47
Total pelagic		4463		2471.81	3812		2508.9	8275		4980.71
Total demersal		445		117.24	396		556.69	841		673.94
Unidentified fish		80701		13144.07	32906		5685.25	113607		18829.32
TOTAL NISP		4908	100	2589.05	4208	100	3065.59	9116	100	5654.64

Table 3. Zooarchaeological fish remains from RH-6, Trench North, 3 mm, according to periods in NISP, % of NISP and WISP (g) - Other pelagic (12 families; see Appendix), Other demersal (19 families; see Appendix)

Figure 4. Illustration of the estimated weight of each main pelagic taxa from the richest RH-6 periods (P: Period), both sectors (TN: Trench North, C-TTZ: Sector C TT-Z). Min: Minimum, Max: Maximum, Q: Quartile. (A) *Thunnus* spp.; (B) *Euthynnus affinis*; (C) *Carangoides* spp.; (D) *Caranx* spp.

130x165mm (150 x 150 DPI)

Main taxa	Trench North				Sector C/TT-Z											
	P1		P2		P1		P2		P3		P4		P5		P6	
	MNI	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI	NISP
<i>Carangoides</i> spp.	180	986	168	852	39	92	1	2	9	19	14	32	109	527	68	308
<i>Caranx</i> spp.	110	602	133	489	23	67	2	3	14	31	14	18	128	876	99	542
<i>Gnathanodon speciosus</i>	35	180	35	147	9	13	-	-	-	-	-	-	22	62	18	50
<i>Euthynnus affinis</i>	72	775	58	609	14	89	1	1	6	24	6	24	53	388	58	342
<i>Thunnus albacares</i>	7	21	23	73	7	12	1	1	1	2	7	19	33	146	18	34
<i>Thunnus tonggol</i>	1	4	8	47	12	43	-	-	3	4	1	1	34	312	28	168
<i>Thunnus</i> spp.	34	239	23	146	2	15	-	-	5	10	5	12	7	111	5	92
Total	439	2807	448	2363	106	331	5	7	38	90	47	106	386	2422	294	1536

Table 4. MNI estimates of the main pelagic taxa at RH-6, 3 mm, according to periods and comparison with the NISP.

Main taxa		Period 1			Period 2			Period 3			Period 4			Period 5			Period 6			Total		
		NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP
Carangidae	<i>Carangoides</i> spp.	92	21	65.12	2	10	0.59	19	7.9	6.49	32	14.8	38.77	527	12.8	316.49	308	8.9	1060.29	980	11.5	1487.75
	<i>Caranx</i> spp.	67	15.3	69.84	3	15	1.29	31	12.8	5.67	18	8.3	9.85	876	21.3	487.14	542	15.6	239.08	1537	18.1	812.87
	<i>Gnathanodon speciosus</i>	13	3	13.93										62	1.5	49.16	50	1.4	41.32	125	1.5	104.41
	Other Carangidae	62	14.2	33.87	3	15	2.53	21	8.7	5.28	22	10.2	6.08	870	21.2	200.76	1086	31.4	682.25	2064	24.3	930.77
Scombridae	<i>Euthynnus affinis</i>	89	20.3	58.56	1	5	0.24	24	9.9	8.38	24	11.1	7.37	388	9.4	252.34	342	9.9	271.25	868	10.2	598.13
	<i>Thunnus albacares</i>	12	2.7	6.82	2	10	1.62	2	0.8	0.4	19	8.8	9.92	146	3.6	73.67	34	1	18.62	215	2.5	111.05
	<i>Thunnus tonggol</i>	43	9.8	20.83				4	1.7	2.07	1	0.5	0.73	312	7.6	134.48	168	4.8	103.21	528	6.2	261.33
	<i>Thunnus</i> spp.	15	3.4	9.58				10	4.1	3.61	12	5.6	3.89	111	2.7	341.48	92	2.7	49.39	240	2.8	407.94
	Other Scombridae	2	0.5	0.48				9	3.7	1.56	11	5.1	2.07	79	1.9	30.92	99	2.9	17.71	200	2.4	52.73
Other pelagic		4	0.9	1.96	1	5	0.03	23	9.5	2.04	3	1.4	1.56	89	2.2	11.13	30	0.9	10.72	150	1.8	27.45
Sparidae	<i>Acanthopagrus</i> spp.	6	1.4	6.47	1	5	0.26	30	12.4	5.13	8	3.7	1.29	127	3.1	25.99	136	3.9	13.46	308	3.6	52.6
	<i>Rhabdosargus</i> spp.	3	0.7	1.06	2	10	0.54	19	7.9	6.4	13	6	5.36	23	0.6	10.87	14	0.4	6.73	74	0.9	30.97
	Other Sparidae	2	0.5	0.75	3	15	0.31	8	3.3	1.76	8	3.7	0.53	41	1	14.23	53	1.5	10.01	115	1.4	27.6
Let.	<i>Lethrinus</i> spp.	16	3.7	5.36	1	5	0.05	20	8.3	2.32	16	7.4	3.38	218	5.3	41.79	208	6	27.23	479	5.6	80.13
Serranidae	<i>Cephalopholis</i> spp.	3	0.7	0.73				5	2.1	1.32	13	6	2.17	42	1	6.95	66	1.9	17.85	129	1.5	29.01
	<i>Epinephelus</i> spp.	4	0.9	1.88	1	5.0	0.21	7	2.9	0.9	1	0.5	0.12	62	1.5	11.59	61	1.8	16.23	136	1.6	30.94
	Other Serranidae	2	0.5	0.81				3	1.2	0.18	6	2.8	0.66	14	0.3	2.22	32	0.9	6.2	57	0.7	10.06
Other demersal		3	0.7	1.53				7	2.9	1.03	9	4.2	6.49	122	3	34.19	143	4.1	22.12	284	3.3	65.36
Total pelagic		399		280.97	12		6.31	143		35.5	142		80.24	3460		1897.57	2751		2493.83	6907		4794.42
Total demersal		39		18.6	8		1.37	99		19.05	74		19.99	649		147.84	713		119.83	1582		326.66
Unidentified fish					942		51.7	1336		96.17	1120		103.96	18337		2831.27	25225		3262.15	46960		6345.25
TOTAL NISP		438	100	299.569	20	100	7.65	242	100	54.549	216	100	100.22	4109	100	2045.4	3464	100	2613.67	8489	100	5121.09

Table 5. Zooarchaeological fish remains from RH-6, Sector C/TT-Z, 3 mm, according to periods in NISP, % of NISP and WISP (g) - Other pelagic (5 families; see Appendix), Other demersal (24 families; see Appendix)

Main taxa	Period 2		Period 3		Period 4		Period 5		Period 6		Total	
	NISP	%	NISP	%	NISP	%	NISP	%	NISP	%	NISP	%
Clupeidae	691	45.2	2486	64.7	539	72.8	825	83.4	1600	69.9	6141	65.4
Carangidae	452	29.6	822	21.4	85	11.5	90	9.1	274	12.0	1723	18.4
Atherinidae	212	13.9	155	4.0	34	4.6	9	0.9	237	10.4	647	6.9
Mugilidae	98	6.4	158	4.1	17	2.3	3	0.3	31	1.4	307	3.3
Other pelagic	1	0.1	6	0.2	44	5.9	22	2.2	35	1.5	108	1.2
Sparoidea	67	4.4	194	5.0	13	1.8	36	3.6	97	4.2	406	4.3
Other demersal	8	0.5	21	0.5	8	1.1	4	0.4	15	0.7	56	0.6
Total pelagic	1454		3627		719		949		2177		8926	
Total demersal	75		215		21		40		112		462	
Unidentified fish	224		417		399		139		468		1647	
Total NISP	1529	100	3842	100	740	100	989	100	2289	100	9388	100

Table 6. Zooarchaeological fish remains from RH-6, Sector C/TT-Z, 1 mm, according periods in NISP and % of NISP. Other pelagic (9 families; see Appendix), Other demersal (13 families; see Appendix)

Figure 5. Illustration of the main taxa distribution (NISP).
 RH-6, Sector C/TT-Z, 1 mm (Periods 2, 3, 4, 5, 6) – RH-5, Sector B2, 1mm (Phases 2, 3, 4). Made with
 ACACIA (Nourissier, Nocus, and Dufraisse, 2010)

68x84mm (300 x 300 DPI)

Taxa		Phase 1			Phase 2			Phase 3			Phase 4			Total		
		NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP	NISP	%	WISP
Clupeidae		3	9.4	0.00	85	35.1	0.18	3668	55.2	9.92	863	57.8	2	4619	54.9	12.10
Carangidae	<i>Carangoides</i> spp.				2	0.8	1.49	8	0.1	2.55	1	0.1	0.01	11	0.1	4.13
	<i>Caranx</i> spp.	1	3.1	7.47	4	1.7	0.43	31	0.5	11.61	0	0.0	0	36	0.4	19.52
	<i>Selar crumenophthalmus</i>	8	25.0	0.06	88	36.4	0.43	2175	32.8	10.66	484	32.4	1.99	2755	32.8	13.14
	<i>Trachurus indicus</i>				5	2.1	0.05	61	0.9	0.31	17	1.1	0.08	83	1.0	0.44
	Other Carangidae	5	15.6	0.45	9	3.7	0.95	97	1.5	7.1	14	0.9	1.56	125	1.5	10.06
Scombridae	<i>Auxis thazard</i>	2	6.3	0.1				12	0.2	1.03	3	0.2	0.07	17	0.2	1.20
	<i>Euthynnus affinis</i>				8	3.3	2.31	34	0.5	6.66	3	0.2	0.98	45	0.5	9.95
	<i>Rastrelliger kanagurta</i>	3	9.4	0.06	8	3.3	0.15	109	1.6	2.06	24	1.6	0.18	144	1.7	2.45
	Other Scombridae				3	1.2	0.63	10	0.2	3.1	1	0.1	0.26	14	0.2	3.99
Mugilidae		1	3.1	0.03	5	2.1	0.1	45	0.7	0.51	13	0.9	0.11	64	0.8	0.75
Atherinidae					1	0.4	0	13	0.2	0.01	4	0.3	0	18	0.2	0.01
Sphyraenidae					3	1.2	0.52	43	0.6	0.69	7	0.5	0.08	53	0.6	1.29
Other pelagic							0	13	0.2	0.31		0.0		13	0.2	0.31
Sparoidea		8	25.0	0.47	10	4.1	0.57	225	3.4	5.91	38	2.5	1.37	281	3.3	8.32
Other demersal		1	3.1	0.04	11	4.5	0.51	96	1.4	4.15	20	1.3	0.18	128	1.5	4.89
Total pelagic		23		8.17	221		7.24	6319		56.52	1434		7.32	7997		79.34
Total demersal		9		0.51	21		1.08	321		10.06	58		1.55	409		13.21
Unidentified fish		91		4.90	608		26.78	7251		192.53	1371		31.80	9321		256.00
Total NISP		32	100	8.68	242	100	8.32	6640	100	66.58	1492	100	8.87	8406	100	92.55

Table 7. Zooarchaeological fish remains from RH-5, Sector B2, 1 mm, according to periods in NISP, % of NISP and WISP (g) - Other pelagic (7 families; see Appendix), Other demersal (20 families; see Appendix)

Figure 6. Compiled data (in metric tons) for 4 years (2013, 2015, 2016, 2018) of artisanal fisheries in Muscat governorate for *Thunnus tonggol*, *Thunnus albacares*, *Euthynnus affinis*, and large jacks (Carangidae) (M.A.F. 2013, 2015, 2016, 2018). The thick line represents the mean. The filled area is the standard deviation of the mean

199x81mm (150 x 150 DPI)

Figure 7. Ecological distribution of the main taxa found at both sites according to Froese and Pauly (2021), Randall (1995) and personal observations. Vector objects are from Vecteezy.com (Graphics RF)

178x147mm (300 x 300 DPI)

Higher taxa	Genus/Species	RH-6 Trench North	RH-6 Sector C/TT-Z	RH-6 Sector C/TT-Z - 1mm	RH-5 Sector B2	Habitat - Behavior
Selachimorpha					2	P
Carcharhinidae		5		1		P
Batoidea			1	2		D
Dasyatidae		1	2			D
Albulidae	<i>Albula</i> sp.		2			D
Muraenidae		3				B
	<i>Gymnothorax</i> spp.	5	5			B
Ophichthidae		2			1	B
Clupeidae		5		6141	4619	EP
Engraulidae				49		P
Chirocentridae	<i>Chirocentrus nudus</i>	2	4	1		P
Pristigasteridae	<i>Ilisha compressa</i>			31	1	P
Silluriformes				1		B
Plotosidae				11		B
Ariidae		13	3		3	D
	<i>Netuma bilineata</i>		3		1	D
	<i>Netuma</i> spp.	10	5			D
	<i>Plicofolis</i> spp.	2			1	D
Holocentridae	<i>Sargocentron</i> sp.		4			D
Stromateidae	<i>Pampus argenteus</i>	1				EP
Pomatomidae	<i>Pomatomus saltatrix</i>	1				P
Scombridae		435	118		8	P
	<i>Auxis thazard</i>	53	14		17	P
	<i>Euthynnus affinis</i>	1384	868		45	P
	<i>Katsuwonus pelamis</i>	82	42			P
	<i>Rastrelliger kanagurta</i>	27	17	22	144	P
	<i>Sarda orientalis</i>	1	2		2	P
	<i>Scomber australasicus</i>		4			P
	<i>Scomberomorus commerson</i>	1	3			P
	<i>Thunnus albacares</i>	94	215			P
	<i>Thunnus tonggol</i>	51	528			P
	<i>Thunnus</i> spp.	385	240		4	P
Trichiuridae	<i>Trichiurus</i> sp.	1				P
Sphyrinaeidae	<i>Sphyræna acutipinnis</i>		3			P
	<i>Sphyræna jello</i>		10			P
	<i>Sphyræna putnamae</i>	3	4			P
	<i>Sphyræna qenie</i>	1				P
	<i>Sphyræna</i> spp.	81	97	2	53	P
Polynemidae	<i>Eleutheronema tetradactylum</i>				1	D
Paralichthyidae	<i>Pseudorhombus</i> sp.	1				P

Carangidae		2248	1797	27	82	P
	<i>Alectis indica</i>		7			P
	<i>Alepes djedaba</i>	27	2	8	3	P
	<i>Alepes vari</i>		1		1	P
	<i>Alepes</i> spp.	9	9		1	P
	<i>Carangoides bajad</i>		106			P
	<i>Carangoides chrysophrys</i>	14	79			P
	<i>Carangoides equula</i>	2				P
	<i>Carangoides ferdau</i>	10			1	P
	<i>Carangoides fulvoguttatus</i>	4	248			P
	<i>Carangoides gymnothetus</i>	10				P
	<i>Carangoides</i> spp.	1800	547		10	P
	<i>Caranx heberi</i>	1	180			P
	<i>Caranx ignobilis</i>		185			P
	<i>Caranx sexfasciatus</i>	107	386		8	P
	<i>Caranx</i> spp.	981	786	1	28	P
	<i>Decapterus macarellus</i>	2	1		1	P
	<i>Elagatis bipinnulata</i>	2	4		2	P
	<i>Gnathanodon speciosus</i>	327	125		1	P
	<i>Scomberoides commersonnianus</i>		7			P
	<i>Scomberoides tol</i>		7		1	P
	<i>Scomberoides</i> spp.	13	13	2		P
	<i>Selar crumenophthalmus</i>	50	188	1677	2755	P
	<i>Selaroides leptolepis</i>				31	P
	<i>Seriola dumerili</i>	1	14			P
	<i>Seriolina nigrofasciata</i>		2			P
	<i>Trachinotus</i> spp.	4		8	2	P
	<i>Trachurus indicus</i>	1	12		83	P
Rachycentridae	<i>Rachycentron canadum</i>	9				P
Atherinidae				647	18	P
Belonidae		12	10	2	3	P
	<i>Albennes hians</i>	10				P
	<i>Tylosorus choram</i>				3	EP
	<i>Tylosurus crocodilus</i>				3	EP
	<i>Tylosurus</i> spp.	8	5		1	EP
Mugilidae		9	17	307	64	P
Drepaneidae	<i>Drepane</i> sp.		1			D
Chaetodontidae		1		4	5	D
Ephippidae			1			D
Siganidae	<i>Siganus</i> sp.	1	1	2	1	D
Acanthuridae		1				D
	<i>Acanthurus</i> spp.	3	2			D
	<i>Naso</i> spp.		4			D
Tetraodontiformes			2			D
Diodontidae		1				D
	<i>Diodon hystrix</i>		1			D

Tertraodontidae			1			D
Balistidae		6	4		1	D
	<i>Abalistes stellatus</i>		1			D
	<i>Balistoides viridescens</i>	1	1			D
	<i>Canthidermis macrolepis</i>		1			D
	<i>Odonus niger</i>	1				D
	<i>Sufflamen fraenatum</i>	5	33			D
Terapontidae	<i>Terapon jarbua</i>	12	8	1	4	D
Ambassidae				15		D
Serranidae		47	56	7	8	D
	<i>Cephalopholis hemistiktos</i>	1	22			D
	<i>Cephalopholis sexmaculata</i>	2	23			D
	<i>Cephalopholis</i> spp.	57	84	1	5	D
	<i>Epinephelus coioides</i>		1			D
	<i>Epinephelus diacanthus</i>	1	2			D
	<i>Epinephelus radiatus</i>		3			D
	<i>Epinephelus stoliczkae</i>	4	6			D
	<i>Epinephelus</i> spp.	44	124		5	D
	<i>Plectropomus</i> sp.		1			D
Apogonidae	<i>Apogon</i> sp.	1				D
Sillaginidae	<i>Sillago</i> spp.	1		6	6	D
Lutjanidae	<i>Lutjanus coeruleolineatus</i>		6			D
	<i>Lutjanus ehrenbergii</i>		5			D
	<i>Lutjanus fulviflamma</i>		4			D
	<i>Lutjanus indicus</i>		2			D
	<i>Lutjanus</i> spp.	10	38	3	49	D
Caesionidae	<i>Caesio</i> sp.				1	D
Gerreidae	<i>Gerres</i> spp.	2		3	13	D
Haemulidae		10			4	D
	<i>Plectorhinchus gaterinus</i>		2			D
	<i>Plectorhinchus pictus</i>		3			D
	<i>Plectorhinchus schotaf</i>		2			D
	<i>Plectorhinchus</i> spp.	8	26		1	D
	<i>Pomadasys argenteus</i>		4			D
	<i>Pomadasys commersonnii</i>		1			D
	<i>Pomadasys</i> spp.	52	33		12	D
Sparoidea		23	43	265	179	D
Sparidae		80	94	130	37	D
	<i>Acanthopagrus bifasciatus</i>	1	6			D
	<i>Acanthopagrus</i> spp.	91	302		21	D
	<i>Argyrops spinifer</i>	20	14			D
	<i>Argyrops</i> spp.	35	2			D
	<i>Rhabdosargus haffara</i>		23			D
	<i>Rhabdosargus</i> spp.	28	51		1	D
	<i>Sparidentex hasta</i>	3	5			D
Lethrinidae	<i>Gnathodentex aureolineatus</i>	1				D

	<i>Lethrinus borbonicus</i>		4			D
	<i>Lethrinus lentjan</i>	2	26			D
	<i>Lethrinus mahsena</i>		1			D
	<i>Lethrinus microdon</i>	1	49			D
	<i>Lethrinus nebulosus</i>	3	29			D
	<i>Lethrinus</i> spp.	184	370	11	42	D
Nemipteridae			5		1	D
Sciaenidae		6	2			D
	<i>Argyrosomus</i> sp.		1			D
	<i>Umbrina</i> sp.	2				D
Mullidae			4			D
Pomacentridae					1	D
Labridae			3		4	D
	<i>Cheilinus</i> sp.	1				D
	<i>Choerodon robustus</i>		1			D
	<i>Coris</i> sp.		2			D
Scaridae		26	5		1	D
	<i>Chlorurus</i> sp.	4				D
	<i>Scarus ghobban</i>	3				D
	<i>Scarus</i> spp.	16	6			D
Platycephalidae		3				B
	Total NISP	9116	8489	9388	8406	

Supplemental file 1. List and NISP of taxa recovered at RH-6 and RH-5, arranged in taxonomic order. B: benthic; D: demersal; EP: epipelagic; P: pelagic. For fish behaviour, we followed the information from Randall (1995) and FishBase (Froese and Pauly 2021).

210x185mm (150 x 150 DPI)

204x184mm (150 x 150 DPI)