1
2 Type II photosensitized oxidation in senescent microalgal cells
3 at different latitudes: Does low under-ice irradiance in polar
4 regions enhance efficiency?
5

6 Rontani Jean-François^{1*}, Amiraux Rémi^{1,2,3}, Smik Lukas⁴, Wakeham Stuart G.⁵,
7 Paulmier Aurélien⁶, Vaultier Frédéric¹, Ha Sun-Yong⁷, Min Jun-oh^{7,8}, Belt
8 Simon T.⁴
9

10 ¹ Aix Marseille Université, Université de Toulon, CNRS/INSU/IRD, Mediterranean Institute of
11 Oceanography (MIO), UM 110, 13288 Marseille, France.

12 ² Takuvik Joint International Laboratory, Laval University (Canada) - CNRS, Département
13 de biologie, Université Laval, Québec G1V 0A6, Québec, Canada.

14 ³ UMR 6539 Laboratoire des Sciences de l'Environnement Marin (CNRS, UBO, IRD,
15 Ifremer) Institut Universitaire Européen de la Mer (IUEM) Plouzané, France

16 ⁴ Biogeochemistry Research Centre, School of Geography, Earth and Environmental
17 Sciences, University of Plymouth, Drake Circus, Plymouth, Devon PL4 8AA, UK.

18 ⁵ Skidaway Institute of Oceanography, Savannah, Georgia, USA.

19 ⁶ Laboratoire d'Etudes en Géophysique et Océanographie Spatiales (LEGOS),
20 IRD/CNRS/UPS/CNES, Université de Toulouse, 31401 Toulouse Cedex 9, France.

21 ⁷ Division of Ocean Sciences, Korea Polar Research Institute, 26 Songdomirae-ro, Incheon, 21990,
22 Republic of Korea.

23 ⁸ Department of Marine Science and Convergence Technology, Hanyang University, 55
24 Hanyangdaehak-ro, Ansan, 15588, Republic of Korea.

25
26
27
28 * Corresponding author. Tel.: +33-4-86-09-06-02; fax: +33-4-91-82-96-41. E-mail address:
29 jean-francois.rontani@mio.osupytheas.fr (J.-F. Rontani)

30
31
32 **Abstract**

33 Comparison of Type II photosensitized oxidation of lipids (the photodynamic effect) and
34 photodegradation of chlorophyll (sensitizer photobleaching) in samples of particulate matter
35 collected previously from locations representing a diverse range of latitudes reveals an
36 enhancement of the photooxidation of lipids at the expense of chlorophyll photodegradation in
37 the polar regions. The efficiency of the photodynamic effect appears to be particularly high in
38 sinking particles collected under sea ice and is attributed to the rapid settling of highly
39 aggregated sympagic algae to depths of low light transmission favouring the photodynamic
40 effect at the expense of photobleaching of the sensitizer. Paradoxically, the low efficiency of
41 Type II photosensitized oxidation of lipids observed in temperate and equatorial regions is
42 associated with high solar irradiances in these regions. Type II photosensitized oxidation of
43 lipids in senescent phytoplankton seems thus to be strongly dependent of the intensity of solar
44 irradiance.

45
46 **Keywords:** Photodynamic effect; Senescent phytoplankton; Latitude; Polar regions; Solar
47 irradiance.

1. Introduction

Reconstructions of sedimentary palaeoenvironments are essential to place the current global warming trends into the context of natural and long-term climate variability. Lipid biomarkers preserved in sediments are often used for this purpose since they are key indicators of organic matter (OM) sources (Volkman et al., 1998; Wakeham et al., 1997). Maximizing the reliability of these reconstructions requires careful consideration of the processes that affect the fate of OM – notably OM degradation and/or preservation – during its transport down the water column from the euphotic zone to the seafloor. Degradation may be biotic (i.e. induced by zooplankton and bacteria, Harvey et al., 1987; Grossart et al., 2007) or abiotic (i.e. induced by light or radicals, Rontani, 2012). Photooxidation (which has generally received little attention until now in the literature) that destroys most of the unsaturated components of biogenic OM initially present in the settling material (for a recent review see Rontani and Belt, 2020) can strongly alter the lipid signature of OM reaching the seafloor. It is thus essential to take into account the potential effects of abiotic degradation when making palaeoenvironmental reconstructions from sedimentary OM.

In healthy phytoplankton cells, light absorption by chlorophyll creates an excited singlet state (^1Chl), which leads to the classical fast reactions of photosynthesis (Foote, 1976). The energy of this excited state is then transferred to various substrates, where it promotes photosynthetic reactions, while a relatively small proportion of ^1Chl (<0.1%) undergoes intersystem crossing (ISC) to form the longer-lived triplet state (^3Chl ; Knox and Dodge, 1985) (Fig. 1). ^3Chl is not only potentially damaging in itself in Type I reactions (Knox and Dodge, 1985), but can also generate reactive singlet oxygen ($^1\text{O}_2$) by reaction with ground state oxygen ($^3\text{O}_2$) via Type II photoprocesses (Fig. 1) (Krieger-Liszkay, 2005). Despite the production of other Reactive Oxygen Species (ROS), it is generally considered that the photo-production of $^1\text{O}_2$ plays the major role in light-induced damage to plant cells (Triantaphylides et al., 2008).

In view of the susceptibility of plant cells to oxidative damage, there are many antioxidant protective mechanisms in chloroplasts. For example, carotenoids quench ^3Chl and $^1\text{O}_2$ by energy transfer mechanisms at very high rates (Fig. 1). Such antioxidants have a dual role: first, they limit $^1\text{O}_2$ formation, and second, they help remove any $^1\text{O}_2$ that does form (Foote, 1976; Tefler, 2002). Tocopherols and ascorbic acid are also efficient quenchers of $^1\text{O}_2$ (Halliwell, 1987; Havaux et al., 2005).

In senescent phototrophic organisms, the cessation of photosynthetic reactions results in an accelerated rate of formation of ^3Chl and ROS (mainly $^1\text{O}_2$) (Nelson, 1993; Triantaphylides et al., 2008). The rate of formation of these potentially damaging species then often exceeds the quenching capacity of the photoprotective system such that photodegradation of cell components can occur via the so-called photodynamic effect (Merzlyak and Hendry, 1994) (Fig. 1). Direct and irreversible reaction of ^3Chl with $^3\text{O}_2$ (photobleaching) gives photooxidation products (Harbour and Bolton, 1978) (Fig. 1), while $^1\text{O}_2$ reacts extremely rapidly with nearby biomolecules at near diffusion-controlled rates (photodynamic effect) (Knox and Dodge, 1985; Cadenas, 1989; Skovsen et al., 2005). The very high reactivity of $^1\text{O}_2$ with numerous cell components (unsaturated lipids, some amino acids, nucleic acids; Rontani, 2012, Devasagayam and Kamat, 2002) is a consequence of the loss of the spin restriction that normally hinders reaction of $^3\text{O}_2$ with these biomolecules (Zolla and Rinalducci, 2002). $^1\text{O}_2$ also reacts with the sensitizer (chlorophyll) inducing its photobleaching (Nelson, 1993; Rontani, 2012) (Fig. 1). It is important to note, however, that photobleaching of the sensitizer reduces $^1\text{O}_2$ production and is thus competitive with the photodynamic effect.

In this study, we investigated the photooxidation of chlorophyll (photobleaching) and some common algal unsaturated lipids (Δ^5 -sterols and monounsaturated fatty acids (MUFAs)) (photodynamic effect) in marine particulate matter from locations representing a diverse range of latitudes, from the Arctic to the Antarctic. In order to try to explain an increased, yet highly

variable, efficiency of the photodynamic effect observed in material from the polar regions, particular attention was given to samples from the Canadian Arctic representing sea ice, the water column under the ice, and regions of open water.

2. Experimental

2.1. Sample collection

Detailed descriptions (e.g. sampling dates, depths, volumes, etc) of the collection of samples of marine particulate matter from the Arctic, English Channel, Mediterranean Sea, Arabian Sea, Equatorial Pacific, Equatorial Atlantic, Peru upwelling and East Antarctica (Fig. 2) has been described previously (see references in Tables 1 and 2). Briefly, suspended particulate matter (SPM) samples were collected with Niskin bottles or an in situ multiple-unit large-volume filtration system (MULVFS), while sinking particles were collected with floating or fixed mooring sediment traps. Sub-samples for lipid analysis were then filtered onto GF/F filters and stored frozen (-80°C). Sympagic algae were obtained from the bottom-most layer of sea ice (0-10 cm) from Resolute Passage (Canadian Arctic) (Rontani et al., 2014). Samples of the sub-ice colonial diatom, *Melosira arctica*, which colonises the underside of sea ice and is widely distributed across the Arctic (Boetius et al., 2013), were collected from the Canadian icebreaker CCGS Amundsen under the ice (49 ± 4 cm) in Baffin Bay ($70^{\circ}28'32''\text{N}$, $64^{\circ}0'37''\text{W}$) in June 2016 (GreenEdge Campaign)) and on board of R/V Lance ($81^{\circ}14'22''\text{N}$, $21^{\circ}54'50''\text{E}$) in August 2017 as part of an oceanographic transect north of Svalbard from Rijpfjorden towards the Nansen Basin. Near-surface (<10 m) water column samples (ca. 1 – 4 l) were collected along a N-S transect terminating in the Amundsen Sea (Antarctica) and filtered (GF/F) on board the Korean Icebreaker RV Araon in January/February 2016.

2.2. Sample treatment

The whole material of the different samples was reduced with excess NaBH₄ in MeOH (25 ml; 30 min) to reduce labile hydroperoxides (resulting from Type II photooxidation) to their corresponding alcohols, which are more amenable to analysis using gas chromatography/electron ionization mass spectrometry (GC-EIMS), gas chromatography/electron ionization tandem mass spectrometry (GC-EIMS/MS) and gas chromatography/electron ionization quadrupole time of flight mass spectrometry (GC-QTOF). Water (25 ml) and KOH (2.8 g) were then added and the resulting mixture saponified by refluxing (2 h). After cooling, the mixture was acidified (HCl, 2 N) to pH 1 and extracted with dichloromethane (DCM; 3 x 20 ml). The combined DCM extracts were dried over anhydrous Na₂SO₄, filtered and concentrated via rotary evaporation at 40 °C to give total lipid extracts (TLEs). All the solvents (pesticide/glass distilled grade) and reagents (Puriss grade) were obtained from Rathburn and Sigma-Aldrich, respectively. TLEs were derivatized by dissolving them in 300 µL pyridine/bis-(trimethylsilyl)trifluoroacetamide (BSTFA; Supelco; 2:1, v/v) and silylated (50 °C, 1 h). After evaporation to dryness under a stream of N₂, the derivatized residue was dissolved in hexane/BSTFA (to avoid desilylation) and analysed by the aforementioned mass spectrometric methods.

2.3. Assignment and quantification of lipids and their degradation products

Lipids and their degradation products were identified by comparison of retention times and mass spectra with those of standards and quantified using GC-EIMS, GC-EIMS/MS and GC-QTOF based on calibrations with external standards. Operating conditions employed during these analyses were as per those described previously (Rontani et al, 2019 and references therein). Standards of phytol, palmitoleic acid, 24-methylcholesta-5,22-dien-3β-ol (brassicasterol) and 24-methylcholesta-5,(24/28)-dien-3β-ol (24-methylenecholesterol) were

obtained from Sigma-Aldrich. Standard oxidation products of these compounds were produced according to previously described procedures (Rontani and Marchand 2000; Marchand and Rontani 2001; Rontani and Aubert, 2005).

2.4. Data treatment

2.4.1 Chlorophyll photooxidation estimates

The molar ratio 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol (phytyldiol):phytol (Chlorophyll Phytyl side-chain Photodegradation Index, CPPI) has been previously proposed to estimate the extent of photodegradation of chlorophylls possessing a phytyl side-chain in natural marine samples through use of the empirical equation: chlorophyll photodegradation % = $(1 - [\text{CPPI} + 1]^{-18.5}) \times 100$ (Cuny et al., 1999).

2.4.1. Lipid photooxidation estimates

The extent of photooxidation (%) of sterols was estimated using the equation: Δ^5 -sterol photooxidation % = $(\Delta^4\text{-stera-}6\alpha/\beta\text{-diols \%} \times (1+0.3)/0.3)$ (Christodoulou et al., 2009).

Type II photosensitized oxidation of MUFAs was estimated after quantification of isomeric *trans* allylic hydroxyacids resulting from NaBH₄-reduction of the corresponding photochemically-produced hydroperoxides after subtraction of the amounts of these compounds arising from autooxidation (Marchand and Rontani, 2001).

2.4.3. Statistical analysis

Mann – Whitney – Wilcoxon tests were performed to identify any significant differences in: (i) the ratio of brassicasterol photo-oxidation percentage / chlorophyll photo-oxidation percentage with latitude (Table S1), and (ii) the ratios of 24-methylenecholesterol photooxidation

percentage / chlorophyll photooxidation percentage, palmitoleic acid photooxidation percentage / chlorophyll photooxidation percentage and brassicasterol photooxidation percentage / chlorophyll photooxidation percentage between different sample types (i.e. sea ice POM, SPM and sediment traps) (Table S2).

3. Results and discussion

Since photodynamic processes and sensitizer photobleaching are competitive processes (Fig. 1), we used the ratio of the % photooxidation of membrane lipids to that of chlorophyll to estimate the efficiency of Type II photosensitized oxidation processes (the photodynamic effect in Fig. 1) in senescent phytoplankton at different latitudes. In the case of brassicasterol, a phytoplanktonic Δ^5 -sterol widely distributed in the oceans (Volkman, 1986, 2003), values of this ratio obtained from previously published and unpublished data sets (Table 1) show a strong increase (albeit with high variability) in samples from the polar regions in comparison to samples collected from temperate and equatorial settings (Fig. 3). Such increases were found to be significant when the data from the temperate and equatorial settings were compared with those from the Arctic and Antarctic datasets (and the combined Arctic/Antarctic dataset), yet no significant difference was found between the high latitude datasets (Table S1). Low temperatures have previously been shown to reduce diffusion rate of $^1\text{O}_2$ through cell membranes (Ehrenberg et al., 1998), thus favouring the intra-cellular involvement of the photodynamic effect. More recently, Amiraux et al. (2016) confirmed these results in the case of the centric diatom *Chaetoceros neogracilis* (strain RCC2022) and observed an increase (3.0 ± 0.5 fold) of the ratio $k_{\text{camp}}/k_{\text{chl}}$ (where k_{camp} is the first-order photodegradation rate of campesterol and k_{chl} this of chlorophyll) when the temperature decreased from 17 to 7°C. The characteristic low temperatures of the Arctic and Antarctic could thus be the cause of the high values of the ratio % photooxidation brassicasterol / % photooxidation chlorophyll measured in

197 samples from these regions. However, low temperatures alone cannot explain the very high
198 variability of this ratio observed within these samples (Fig. 3).

199 In an effort to explain this strong variability, we examined, more closely, Arctic samples
200 collected from sea ice, under the ice (suspended and sinking particles) and from open water
201 conditions (sinking particles), which are all dominated by diatoms. The % photooxidation of
202 the common diatom lipids brassicasterol, 24-methylenecholesterol and palmitoleic acid to that
203 of chlorophyll are summarized in Table 2 and Fig. 4. Strong differences of photodynamic effect
204 efficiency were observed between these different kinds of particles. These differences were
205 found to be significant between surface samples (sea ice and SPM) and traps and between under
206 ice traps and open water traps (Table S2). These different samples having been collected at very
207 close temperatures (ranging from -1 to -2 °C in surface waters of Arctic), it is clear that the low
208 temperatures do not represent the main driver for the enhancement of photodynamic efficiency
209 in polar regions.

210 Riebesell et al. (1991) previously suggested that growing cells released by sea ice remain
211 largely unaggregated (i.e. mainly in suspension). The very low photooxidation state of lipids
212 and chlorophyll observed in SPM collected under the ice (Fig. 4) could thus result from the
213 healthy state of the algal cells present in these samples (Rontani et al., 2016), which are largely
214 unaffected by photooxidative damage. Indeed, in healthy cells, the greater part of the
215 photoexcited chlorophyll singlet state is used in the fast photochemical reactions of
216 photosynthesis.

217 The efficiency of the photodynamic effect appears to be considerably higher in material
218 collected in sediment traps in spring from under sea ice compared to those collected in summer
219 from open water (Fig. 4, Table S2), which is likely attributable to the respective contributions
220 of sympagic (i.e. living within ice) vs pelagic algae. Indeed, in contrast to healthy and largely
221 unaggregated sympagic algal cells (see above), less metabolically-active sea ice algae that are

released from the ice are generally concentrated into aggregates that become sinking particles (Riebesell et al., 1991). Moreover, sympagic diatoms exhibit a higher sensitivity towards light-induced stress than pelagic diatoms (Kvernvik et al., 2020). Such differences in sensitivity were attributed, in part, to the gradually changing and low amplitude irradiance typically experienced by sympagic algae (Hill et al., 2018), whereas pelagic cells in open water, in contrast, experience high amplitude changes in light intensity over much shorter timescales (MacIntyre et al., 2000). This interesting hypothesis is not supported, however, by: (i) the very low efficiency of the photodynamic effect observed in the bottom-most layer of sea ice (Table 2) and (ii) the lack of correlation between lipid photooxidation % and the concentration of the sea ice lipid biomarker IP₂₅ (a well-known sea ice proxy, for reviews see Belt and Müller, 2013; Belt, 2018) in under ice traps ($R^2 < 0.008$).

During in vitro experiments carried out on senescent cells of the centric diatom *Chaetoceros neogracilis* (strain RCC2022), Type II photosensitized oxidation of lipids was also observed to be strongly enhanced by low irradiance levels, whereas the opposite was true for the photodegradation of chlorophyll (Amiraux et al., 2016). The ratio k_{camp}/k_{chl} thus increased 4.2 ± 0.8 fold when irradiance decreased from 2038 to 165 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. In the Arctic, the mean PAR (Photosynthetically Active Radiation) irradiance in the surface mixed layer is considerably higher in open water than in ice-covered zones (365 ± 62 and $10.9 \pm 2.7 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively, Alou-Font et al., 2016). The intensity of solar irradiance may thus be at the origin of the contrasting efficiency of the photodynamic effect observed in sinking particles collected under sea ice compared to those from open water (Fig. 4). Further, the increase of this efficiency with depth in under ice samples (Fig.4, Table S2) could result from enhanced aggregation of senescent sympagic algae as they sink to deeper traps (Riebesell et al., 1991; Rontani et al., 2016) where light transmission is lower. In contrast, the relatively high irradiance observed at the ice-water interface (up to ca. 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ in Davis

Strait, Galindo et al., 2017) could explain the greatly diminished photodynamic effect in sympagic algae inhabiting the bottom-most layer of sea ice and in *Melosira arctica* (Table 2) (Fig. 6). Examination of near-surface (<10 m) SPM samples collected along a N-S transect terminating in the Amundsen Sea (Antarctica) in open water and seasonally ice covered zones further confirmed the increase of photodynamic efficiency in regions of sea ice cover (Fig. 5) ($W = 52$, $p\text{-value} = 0.003205$). In addition, the relatively higher photooxidation state of these suspended particles compared to those collected in the Arctic (Fig. 4, Table 2) may be attributed to the low contribution (typically 0.5 to 2%) of sympagic algae to primary production in January-February in the Antarctic (Lizotte, 2001).

Due to their strong aggregation capability (Macdonald et al., 1998; Ambrose et al., 2001, 2005) and the low remineralizing potential of their associated bacteria (Amiriaux et al., 2017, 2020), sympagic algae seem to contribute more significantly than open water phytoplankton to the export of carbon to Arctic sediments, providing an early season food source for benthic fauna (Macdonald et al., 1998; Mincks et al., 2005). Episodic massive falls of *M. arctica* can sink to the seafloor (Ambrose et al., 2005; Boetius et al., 2013; Lalande et al., 2019). The polyunsaturated fatty acid (PUFA) content in algal biomass that reached the seafloor is essential for growth and reproduction of benthic fauna (Brett and Müller-Navarra, 1997; McMahon et al., 2006). Due to the high photosensitized oxidation of MUFAs observed in samples sinking under sea ice cover (Fig. 4), and the well-known increasing photooxidation rates of fatty acids with their degree of unsaturation (Frankel, 1998; Rontani et al., 1998), the PUFA content of sympagic material reaching the sediment would thus be greatly reduced. *Melosira* strands that are weakly affected by abiotic degradation processes under the ice (ratio C_{20:5} acid vs palmitoleic acid ranging from 0.25 to 0.47), despite a significant photooxidation of chlorophyll (Table 2), and that sink very quickly to the seafloor (Syvertsen, 1991) in massive falls (Ambrose

et al., 2005; Boetius et al., 2013; Lalande et al., 2019) may be an important source of fresh material rich in OM (i.e., PUFAs) for the benthos.

Further, the expected reduction of sea ice cover resulting from increased global warming may result in a shift in the relative contributions of ice-associated vs pelagic algae (Carroll and Carroll, 2003) to the seafloor. The high flux of fast-sinking aggregates of strongly abiotically-altered sympagic algae and fast sinking well preserved *M. arctica* reaching Arctic sediments could thus be replaced, progressively, by a lower flux of weakly abiotically altered pelagic biomass. Such a shift should thus strongly impact the quality and quantity of food reaching benthic communities, as previously proposed by Sun et al. (2007), and could significantly alter the community structure and spatial distribution of the benthos.

4. Conclusions

During this work, Type II photosensitized oxidation of lipids (photodynamic effect) and photodegradation of chlorophyll (photobleaching) were compared in several samples of particulate matter from locations ranging from the Arctic to the Antarctic. The results obtained clearly showed an enhancement of the photooxidation of lipids at the expense of chlorophyll photodegradation in polar regions compared to temperate and equatorial regions. Careful examination of different samples of sympagic and epiphytic algae, suspended and sinking particles collected in Arctic allowed to show that the efficiency of photodynamic effect was particularly high in sinking particles collected under the ice (Fig. 6). This high efficiency was attributed to the rapid settling of highly aggregated sympagic algae to depths of low light transmission favouring photodynamic effect at the expense of photobleaching of the sensitizer. In contrast, in the case of samples exposed to relatively high solar irradiances - sympagic and epiphytic algae collected in the bottommost layer of ice or at the underside of ice, respectively, and particles collected from open waters - the efficiency of photodynamic effect appeared to be

relatively weak (Fig. 6). The low efficiency of Type II photosensitized oxidation of lipids observed in temperate and equatorial regions could, perhaps paradoxically, thus be attributed to the high solar irradiances received in these regions. The intensity of solar irradiance seems thus to be a key parameter during the Type II photosensitized oxidation of lipids in senescent phytoplankton.

Acknowledgements

The sediment trap samples off Peru were collected thanks to the AMOP (“Activity of research dedicated to the Minimum of Oxygen in the eastern Pacific”) project. We would like to thank the crews of the German R/V Meteor, Peruvian R/V Olaya and French R/V Atalante for the AMOP fixed mooring servicing. We would like to thank M. Soto (IRD, Peru), E. Carrasco (IMARPE, Peru) and J. Grelet (IMAGO/IRD, France), as well as O. De Gesincourt, L. Scouarnec, A. Royer and E. De Saint Léger (DT-INSU/CNRS, France) for general logistics and administrative support. We would like also to thank C. Panagiotopoulos (MIO, France) and F.O. Velazco (IMARPE, Peru) for the sediment trap sampling, N. Leblond (OOV, France) and M. Bretagnon (LEGOS) for the initial samples processing, as well as A. Oschlies for a complementary SFB754 funding. Equatorial Pacific and Arabian Sea samples were collected under the auspices of the U.S. Joint Global Ocean Flux Program through U.S. National Science Foundation grants OCE-90-22319 and OCE-9310364, respectively, to SGW. Sediment trap samples from the Beaufort Sea were obtained through the Long-Term Oceanic Observatories Program of the ArcticNet Network of Centres of Excellence (NCE) of Canada. Thanks are due to the scientific party and crew of cruise NBP1402 funded by NSF (ANT-1143836) for the sampling of Antarctic samples. Collection of Resolute Passage samples (sea ice, SPM and sinking particles) was supported through a University of Manitoba Start-up grant and Natural Sciences and Engineering Research Council of Canada (NSERC). We thank the Captain and

crew of the R.V ‘Tethys II’ (Institut National des Sciences de l’Univers, France) for assistance in the field during recovery and deployment of the trap mooring in Ligurian Sea. We acknowledge GBMF-MMI Grants 2293 and 2928 and NSF-OCE-0934095 for the financial support provided to Amazon plume collections, as well as the Brazilian government (Ministério da Marinha). Thanks are due to Philipp Assmy for collecting a sample of *Melosira arctica* during the Rjipfjorden transect which was funded by the former Centre for Ice, Climate and Ecosystems at the Norwegian Polar Institute. Thanks are also due to the scientific party and crew of cruise NBP1402 funded by NSF (ANT-1143836) for the sampling of East Antarctic samples. Samples from the Amundsen Sea transect were collected as part of the project ‘Variability of carbon pump and climate control in the Southern Ocean’ (PE20140). Further, we thank the FEDER 1166-39417 for the funding of the apparatus employed. We are also grateful to two anonymous reviewers for their useful and constructive comments.

References

- Alou-Font, E., Roy, S., Agusti, S., Gosselin, M., 2016. Cell viability, pigments and photosynthetic performance of Arctic phytoplankton in contrasting ice-covered and open-water conditions during the spring–summer transition. *Mar. Ecol. Prog. Ser.* 543, 89-106.
- Ambrose, W.G., Clough, L.M., Tilney, P.R., Beer, L., 2001. Role of echinoderms in benthic remineralization in the Chukchi Sea. *Mar. Biol.* 139, 937–949.
- Ambrose, W.G., Quillfeldt, Jr.C., Clough, L.M., Tilney, P.V.R., Tucker, T., 2005. The sub-ice algal community in the Chukchi Sea: large- and small-scale patterns of abundance based on images from a remotely operated vehicle. *Polar Biol.* 28, 784-795.
doi:10.1007/s00300-005-0002-8

344 Amiraux, R., Jeanthon, C., Vaultier, F., Rontani, J.-F., 2016. Paradoxical effects of temperature
 345 and solaire irradiance on the photodegradation state of killed phytoplankton. *J. Phycol.*
 346 52, 475-485.

347 Amiraux, R., Belt, S.T., Vaultier, F., Galindo, V., Gosselin, M., Bonin, P., Rontani, J.-F., 2017.
 348 Monitoring photo-oxidative and salinity-induced bacterial stress in the Canadian Arctic
 349 using specific lipid tracers. *Mar. Chem.* 194, 89–99.
 350 doi.org/10.1016/j.marchem.2017.05.006.

351 Amiraux R., Burot C., Bonin P., Massé, G., Guasco, S., Babin, M., Vaultier, F., Rontani, J.-F.,
 352 2020. Stress factors resulting from the Arctic vernal sea ice melt: impact on the viability
 353 of the bacterial communities associated to sympagic algae. *Elem. Sci. Anth.* (In press).

354 Belt, S.T., Müller, J., 2013. The Arctic sea ice biomarker IP₂₅: a review of current
 355 understanding, recommendations for future research and applications in palaeo sea ice
 356 reconstructions. *Quarter. Sci. Rev.* 79, 9–25. doi:10.1016/j.quascirev.2012.12.001

357 Belt, S.T., 2018. Source-specific biomarkers as proxies for Arctic and Antarctic sea ice. *Org.*
 358 *Geochem.* 125, 273-295.

359 Boetius, A., Albrecht, S., Bakker, K., Bienhold, C., Felden, J., Fernandez-Méndez, M.,
 360 Hendricks, S., Kathein, C., Lalande, C., Krumpen, T., Nicolaus, M., Peeken, I., Rabe, B.,
 361 Rogacheva, A., Rybakova, E., Somavilla, R., Wenzhöfer, F., RV Polarstern ARK27-3-
 362 shipboard Science Party., 2013. Export of algal biomass from the melting Arctic sea ice.
 363 *Science* 339, 1430–1432. doi:10.1126/science.1231346

364 Bretagnon, M., Paulmier, A., Garçon, V., Dewitte, B., Illig, S., Leblond, N., Coppola, L.,
 365 Campos, F., Velazco, F., Panagiotopoulos, C., Oschlies, A., Hernandez-Ayon, J.M.,
 366 Vergara, O., Montes, I., Martinez, P., Carrasco, E., Grelet, J., Depretz de Gesincourt, O.,
 367 Maes, C., Scouarnec, L. Modulation of the vertical particle transfer efficiency in the

368 Oxygen Minimum Zone off Peru., 2018. Biogeoscience 15, 5093-5111.
 369 doi.org/10.5194/bg-15-5093-2018

370 Brett, M.T., Müller-Navarra, D.C., 1997. The role of highly unsaturated fatty acids in aquatic
 371 food web processes. *Freshwater Biol.* 38, 483–499.

372 Cadenas, E., 1989. Biochemistry of oxygen toxicity. *Annu. Rev. Biochem.* 58, 79-110.

373 Carroll, M.L., Carroll, J., 2003. The Arctic seas. In: Black, K.D., Shimmield, G.B. (eds)
 374 Biogeochemistry of Marine Systems. Blackwell, Oxford, pp 127–156.

375 Christodoulou, S., Marty, J.-C., Miquel, J.-C., Volkman, J.K., Rontani, J.-F., 2009. Use of lipids
 376 and their degradation products as biomarkers for carbon cycling in the northwestern
 377 Mediterranean Sea. *Mar. Chem.* 113, 25–40. doi:10.1016/j.marchem.2008.11.003

378 Cuny, P., Romano, J.-C., Beker, B., Rontani, J.-F., 1999. Comparison of the photodegradation
 379 rates of chlorophyll chlorin ring and phytol side chain in phytodetritus: is the phytyldiol
 380 versus phytol ratio (CPPI) a new biogeochemical index? *J. Exp. Mar. Biol. Ecol.* 237,
 381 271–290. doi:10.1016/S0022-0981(99)00010-6

382 Devasagayam, T., Kamat, J., 2002. Biological significance of singlet oxygen. *Ind. J. Exp. Biol.*
 383 40, 680–692.

384 Ehrenberg, B., Anderson, J.L., Foote, C.S., 1998. Kinetics and yield of singlet oxygen
 385 photosensitized by hypericin in organic and biological media. *Photochem. Photobiol.* 68,
 386 135–140. doi:10.1111/j.1751-1097.1998.tb02479.x

387 Foote, C.S., 1976. Photosensitized oxidation and singlet oxygen: consequences in biological
 388 systems. In: Pryor, W.A. (Ed.), *Free Radicals in Biology*. Academic Press, New York,
 389 pp. 85-133.

390 Frankel, E.N., 1998. Lipid oxidation. The Oily Press, Dundee.

391 Galeron, M.-A., Radakovitch, O., Charriere, B., Vaultier, F., Volkman, J.K., Bianchi, T.S.,
 392 Ward, N.T., Medeiros, P., Sawakuchi, H., Tank, S., Kerhervé, P., Rontani J.-F., 2018.

Lipoxygenase-induced autoxidative degradation of terrestrial particulate organic matter in estuaries: a widespread process enhanced at high and low latitudes. *Org. Geochem.* 115, 78-92.

Galindo, V., Gosselin, M., Lavaud, J., Mundy, C.J., Else, B., Ehn, J., Babin, M., Rysgaard, S., 2017. Pigment composition and photoprotection of Arctic sea ice algae during spring. *Mar. Ecol. Progr. Ser.* 585: 49–69. doi.org/10.3354/meps12398

Grossart, H.-P., Tang, K.W., Kjørboe, T., Ploug, H., 2007. Comparison of cell-specific activity between free-living and attached bacteria using isolates and natural assemblages. *FEMS Microbiol. Let.* 266, 194–200. doi.org/10.1111/j.1574-6968.2006.00520.x

Halliwell, B., 1987. Oxidative damage, lipid peroxidation and antioxidant protection in chloroplasts. *Chem. Phys. Lipids* 44, 327-340.

Harbour, J. R., Bolton, J. R., 1978. The involvement of the hydroxyl radical in the destructive photooxidation of chlorophylls in vivo and in vitro. *Photochem. Photobiol.* 28, 231-234.

Harvey, H.R., Eglinton, G., O'hara, S.C.M., Corner, E.D.S., 1987. Biotransformation and assimilation of dietary lipids by *Calanus* feeding on a dinoflagellate. *Geochim. Cosmochim. Acta* 51, 3031–3040. doi.org/10.1016/0016-7037(87)90376-0

Havaux, M., Eymery, F., Porfirova, S., Rey, P., Dormann, P., 2005. Vitamin E protects against photoinhibition and photooxidative stress in *Arabidopsis thaliana*. *Plant Cell* 17, 3451-3469.

Hill, V.J., Light, B., Steele, M., Zimmerman, R.C., 2018. Light availability and phytoplankton growth beneath Arctic sea ice: integrating observations and modelling. *J. Geophys. Res. Oceans* 123, 3651–3667. doi.org/10.1029/2017JC013617

Korytowski, W., Bachowski, G.J., Girotti, A.W., 1992. Photoperoxidation of cholesterol in homogeneous solution, isolated membranes, and cells: comparison of the 5 α - and 6 β -

417 hydroperoxides as indicators of singlet oxygen intermediacy. *Photochem. Photobiol.* 56,
 418 1-8.

419 Knox, J.P., Dodge, A.D., 1985. Singlet oxygen and plants. *Phytochemistry* 24, 889-896.

420 Krieger-Liszkay, A., 2005. Singlet oxygen production in photosynthesis. *J. Exp. Bot.* 56, 337–
 421 346.

422 Kvernvik, A.C., Rokitta, S.D., Leu, E., Harms, L., Gabrielsen, T.M., Rost, B., Hoppe, C.J.M.,
 423 2018. Higher sensitivity towards light stress and ocean acidification in an Arctic sea-ice-
 424 associated diatom compared to a pelagic diatom. *New Phytol.* 226, 1708–1724. doi:
 425 10.1111/nph.16501

426 Lalande, C., Nöthig, E.-M., Fortier, L., 2019. Algal export in the Arctic Ocean in times of global
 427 warming. *Geophys. Res. Lett.* 46, 5959–5967. doi.org/10.1029/2019GL083167

428 Lizotte, M., 2001. The contributions of sea ice algae to Antarctic marine primary production.
 429 *Amer. Zool.* 1, 57-73.

430 MacDonald, R.W., Solomon, S.M., Cranston, R.E., Welch, H.E., Yunker, M.B., Gobeil, C.,
 431 1998. A sediment and organic carbon budget for the Canadian Mackenzie shelf. *Mar.*
 432 *Geol.* 144, 255-273.

433 MacIntyre, H.L., Kana, T.M., Geider, R.J., 2000. The effect of water motion on short-term rates
 434 of photosynthesis by marine phytoplankton. *Trends Plant Sci.* 5, 12-17.

435 MacMahon, K.W., Ambrose Jr, W.G., Johnson, B.J., Sun, M.Y., Lopez, G.R., Clough, L.M.,
 436 Carroll, M.L., 2006. Benthic community response to ice algae and phytoplankton in Ny
 437 Ålesund, Svalbard. *Mar. Ecol. Progr. Ser.* 310, 1-14.

438 Marchand, D., Rontani, J.-F., 2001. Characterisation of photo-oxidation and autoxidation
 439 products of phytoplanktonic monounsaturated fatty acids in marine particulate matter and
 440 recent sediments. *Org. Geochem.* 32, 287–304. doi.10.1016/S0146-6380(00)00175-3

441 Merzlyak, M.N., Hendry, G.A.F., 1994. Free radical metabolism, pigment degradation and lipid
 442 peroxidation in leaves during senescence. *Proc. Roy. Soc. Edinburgh* 102B, 459-471.
 443 Mincks, S.L., Smith, C.R., DeMaster, D.J., 2005. Persistence of labile organic matter and
 444 microbial biomass in Antarctic shelf sediments: Evidence of a sediment food bank. *Mar.*
 445 *Ecol. Progr. Ser.* 300, 3–19. doi.org/10.3354/meps300003.
 446 Nelson, J.R., 1993. Rates and possible mechanism of light-dependent degradation of pigments
 447 in detritus derived from phytoplankton. *J. Mar. Res.* 51, 155–179.
 448 doi.10.1357/0022240933223837
 449 Riebesell U., Schloss I., Smetacek V., 1991. Aggregation of algae released from melting sea
 450 ice -implications for seeding and sedimentation. *Pol. Biol.* 11:239–248
 451 Rontani, J.-F., Grossi, V., Faure, F., Aubert, C., 1994. “Bound” 3-methylidene-
 452 7,11,15-trimethylhexadecan-1,2-diol: a new isoprenoid marker for the photodegradation
 453 of chlorophyll-a in seawater. *Org. Geochem.* 21, 135–142. doi.org/10.1016/0146-
 454 6380(94)90150-3
 455 Rontani, J.-F., Cuny, P., Grossi, V., 1998. Identification of a pool of lipid photoproducts in
 456 senescent phytoplanktonic cells. *Org. Geochem.* 29, 1215–1225. doi.org/10.1016/S0146-
 457 6380(98)00073-4
 458 Rontani, J.-F., Marchand, D., 2000. Δ^5 -Stenol photoproducts of phytoplanktonic origin: a
 459 potential source of hydroperoxides in marine sediments? *Org. Geochem.* 31, 169-180.
 460 Rontani, J.-F., Aubert, C., 2005. Characterization of isomeric allylic diols resulting from
 461 chlorophyll phytyl side chain photo- and autoxidation by electron ionization gas
 462 chromatography/mass spectrometry. *Rapid Commun. Mass Spectrom.* 19, 637-646.
 463 Rontani, J.-F., Zabeti, N., Wakeham, S.G., 2011. Degradation of particulate organic matter in
 464 the equatorial Pacific Ocean: biotic or abiotic? *Limnol. Oceanogr.* 56, 333-349.

- Rontani, J.-F., 2012. Photo- and free radical-mediated oxidation of lipid components during the senescence of phototrophic organisms. In: T. Nagata Ed., *Senescence*. Intech, Rijeka, pp. 3-31.
- Rontani, J.-F., Charriere, B., Forest, A., Heussner, S., Vaultier, F., Petit, M., Delsaut, N., Fortier, L., Sempéré, R., 2012. Intense photooxidative degradation of planktonic and bacterial lipids in sinking particles collected with sediment traps across the Canadian Beaufort Shelf (Arctic Ocean). *Biogeosciences* 9, 4787–4802. doi.org/10.5194/bg-9-4787-2012
- Rontani, J.-F., Belt, S.T., Brown, T., Vaultier, F., Mundy, C.J., 2014. Identification of sequential photo- and autoxidation of diatom lipids in Arctic sea ice. *Org. Geochem.* 77, 59-71.
- Rontani, J.-F., Belt, S.T., Brown, T.A., Amiraux, R., Gosselin, M., Vaultier, F., Mundy, C.J., 2016. Monitoring abiotic degradation in sinking versus suspended Arctic sea ice algae during a spring ice melt using specific lipid oxidation tracers. *Org. Geochem.* 98, 82–97. doi.org/10.1016/j.orggeochem.2016.05.016
- Rontani, J.-F., Smik, L., Belt, S.T., Vaultier, F., Armbrrecht, L., Leventer, A., Armand L.K., 2019. Abiotic degradation of highly branched isoprenoid alkenes and other lipids in the water column off East Antarctica. *Mar. Chem.* 210, 34-47.
- Rontani, J.-F., Belt, S.T., 2020. Photo- and autoxidation of unsaturated algal lipids in the marine environment: An overview of processes, their potential tracers, and limitations. *Org. Geochem.* 139, 103941. doi.org/10.1016/j.orggeochem.2019.103941
- Rontani, J.-F., Smik, L., Vaultier, F., Widdicombe, C., Belt S.T., 2021. Seasonal monitoring of biotic and abiotic degradation of suspended particulate lipids in the western English Channel suggests a link between bacterial 10S-DOX activity and production of free fatty acids by wounded diatoms. *Mar. Chem.* (In Press)

490 Skovsen, E., Snyder, J.W., Lambert, J.D.C., Ogilby, P.R., 2005. Lifetime and diffusion of
 491 singlet oxygen in a cell. *J. Phys. Chem. B* 109, 8570–8573.

492 Sun, M.-Y., Carroll, M.L., Ambrose, W.G.Jr., Clough, L.M., Zou, L., Lopez, G.R., 2007. Rapid
 493 consumption of phytoplankton and ice algae by Arctic soft-sediment benthic
 494 communities: Evidence using natural and ¹³C-labeled food materials. *J. Mar. Res.* 65,
 495 571-588.

496 Syvertsen, E.E., 1991. Ice algae in the Barents Sea: types of assemblages, origin, fate and role
 497 in the ice edge phytoplankton bloom. *Polar Res.* 10, 277-287. doi:10.1111/j.1751-
 498 8369.1991.tb00653.x

499 Telfer, A., 2002. What is beta-carotene doing in the photosystem II reaction centre? *Philos.*
 500 *Trans. Roy. Soc. Lond. B Biol. Sci.* 357, 1431-1439.

501 Triantaphylides, C., Krischke, M., Hoeberichts, F.A., Ksas, B., Gresser, G., Havaux, M., Van
 502 Breusegem, F., Johannes Mueller, M., 2008. Singlet oxygen is the major reactive Oxygen
 503 species involved in photooxidative damage to plants. *Plant Physiol.* 148, 960-968.
 504 doi/10.1104/pp.108.125690

505 Volkman, J.K. 1986. A review of sterol markers for marine and terrigenous organic matter.
 506 *Org. Geochem.* 9, 83-99.

507 Volkman, J.K., Barrett, S.M., Blackburn, S.I., Mansour, M.P., Sikes, E.L., Gelin, F., 1998.
 508 Microalgal biomarkers: a review of recent research developments. *Org. Geochem.* 29,
 509 1163–1179.

510 Volkman, J.K. 2003. Sterols in microorganisms. *Appl. Microbiol. Biotechnol.* 60, 495-506.

511 Wakeham, S.G., Lee, C., Hedges, J.I., Hernes, P.J., Peterson, M.L., 1997. Molecular indicators
 512 of diagenetic status in marine organic matter. *Geochim. Cosmochim. Acta* 61, 5363–
 513 5369.

514 Wakeham, S.G., Peterson, M.L., Hedges J.I., Lee, C., 2002. Lipid biomarker fluxes in the
515 Arabian Sea: with a comparison to the Equatorial Pacific Ocean, Deep-Sea Res. 49, 2265-
516 2301.

517 Zolla, L., Rinalducci, S., 2002. Involvement of active oxygen species in degradation of light-
518 harvesting proteins under light stresses. Biochem. 41, 14391–14402.

519

Figure captions

Figure 1. Potential pathways for chlorophyll (Chl) excitation energy in senescent phytoplankton cells (simplified scheme limited to the formation of $^1\text{O}_2$ and photoprotective role of carotenoids (Car)).

Figure 2. Sampling locations.

Figure 3. Values of the ratio brassicasterol photooxidation %/chlorophyll photooxidation % in particulate matter samples collected at different latitudes.

Figure 4. Values of the ratios brassicasterol photooxidation %/chlorophyll photooxidation %, 24-methylenecholesterol photooxidation %/chlorophyll photooxidation % and palmitoleic acid photooxidation %/chlorophyll photooxidation % in particulate matter samples collected in Arctic ice-covered and open water zones.

Figure 5. Values of the ratio brassicasterol photooxidation %/chlorophyll photooxidation % in SPM samples collected in open and ice-covered zones along a N-S transect terminating in the Amundsen Sea (Antarctica).

Figure 6. Conceptual scheme summarizing the efficiency of photodynamic effect in sympagic, epiphytic and pelagic algae in polar regions. (To simplify the scheme only the Type II photosensitized oxidation of palmitoleic acid is shown) (nd = not detected).

Table captions

Table 1. Variation of the ratio brassicasterol photooxidation %/chlorophyll photooxidation % in particulate matter samples according to the latitude.

Table 2. Photooxidation of chlorophyll, brassicasterol, 24-methylenecholesterol and palmitoleic acid in different sea ice, *M. arctica* and particulate matter samples collected in the Arctic.

Supplementary material

Table S1. Results of Mann – Whitney – Wilcoxon analysis testing the effect of the latitude on the brassicasterol photooxidation percentage : chlorophyll a photooxidation percentage ratio. Bold values are indicative of significance.

Table S2. Results of Mann – Whitney – Wilcoxon analysis testing the effect of the sample type on the 24-methylenecholesterol photo-oxidation percentage / chlorophyll photo-oxidation percentage, palmitoleic acid photo-oxidation percentage / chlorophyll photo-oxidation percentage and brassicasterol photo-oxidation percentage / chlorophyll photo-oxidation percentage ratios. Bold values are indicative of significance.

Table 1

Variation of the ratio brassicasterol photooxidation %/chlorophyll photooxidation % in particulate matter samples according to the latitude.

Latitude	Location	Nature of particles	n	Brassicasterol hv %/Chlorophyll hv %	References
71-74°N	Canadian Arctic	Suspended and sinking	46	0.77 ± 1.10	Rontani et al., 2012; 2016
50°N	English Channel	Suspended	24	0.02 ± 0.04	Rontani et al., 2021
43°N	Rhône Prodelta	Suspended	21	0.21 ± 0.11	Galeron et al., 2018
43°N	Ligurian Sea	Sinking	12	0.21 ± 0.10	Christodoulou et al., 2009
17°N	Arabian Sea	Suspended and sinking	7	0.07 ± 0.04	Wakeham et al., 2002
0°	Equatorial Atlantic	Suspended	2	0.10 ± 0.12	Galeron et al., 2018
0°	Equatorial Pacific	Suspended and sinking	11	0.11 ± 0.07	Wakeham et al., 2002; Rontani et al., 2011
12°S	Peru Upwelling	Sinking	10	0.18 ± 0.04	Bretagnon et al., 2018
40-67°S	South Pacific	Suspended	5	0.09 ± 0.04	
67-75°S	Antarctica	Suspended	23	0.50 ± 0.45	Rontani et al., 2019

Table 2

Photooxidation of chlorophyll, brassicasterol, 24-methylenecholesterol and palmitoleic acid in different sea ice, *M. arctica* and particulate matter samples collected in the Arctic.

Samples	Location	n	Chlorophyll photooxidation %	Brassicasterol photooxidation %/ chlorophyll photooxidation %	24-Methylenecholesterol photooxidation %/ chlorophyll photooxidation %	Palmitoleic acid photooxidation %/ chlorophyll photooxidation %
Sea ice (0-3 cm)	Resolute Passage ^a and Davies Strait ^b	21	54.5 ± 38.4	0.02 ± 0.05	0.04 ± 0.10	0.05 ± 0.08
<i>Melosira arctica</i>	Baffin Bay and North of Svalbard	6	40.2 ± 12.8	-	-	0.03 ± 0.03
Under ice spm (<10 m)	Resolute Passage ^c	20	16.3 ± 13.9	0.05 ± 0.17	0.42 ± 0.96	1.00 ± 1.60
Under ice trap (5 m)	Resolute Passage ^c	11	55.7 ± 34.1	1.10 ± 0.62	1.22 ± 0.98	2.87 ± 2.78
Under ice trap (30 m)	Resolute Passage ^c	11	11.5 ± 8.6	2.00 ± 1.39	3.97 ± 2.22	8.39 ± 7.15
Open water trap (100 m)	Beaufort Sea ^d	12	99.8 ± 0.2	0.23 ± 0.07	0.48 ± 0.14	0.47 ± 0.19

^a Rontani et al., 2014

^b Amiraux et al., 2017

^c Rontani et al., 2016

^d Rontani et al., 2012

Table S1: Results of Mann – Whitney – Wilcoxon analysis testing the effect of the latitude on the brassicasterol photooxidation percentage : chlorophyll *a* photooxidation percentage ratio. Bold values are indicative of significance.

Source of variation	W	<i>p-value</i>
High latitudes ^a x low latitudes ^b	3861.5	1×10^{-7}
Arctic x Antarctic	583	0.71
Arctic x low latitudes	1107.5	7×10^{-4}
Antarctic x low latitudes	1513	2×10^{-9}

^a Sum of Arctic and Antarctic values

^b Sum of low latitude values (between 67°S and 50°N)

Table S2: Results of Mann – Whitney – Wilcoxon analysis testing the effect of the sample type on the 24-methylencholesterol photo-oxidation percentage / chlorophyll photo-oxidation percentage, palmitoleic acid photo-oxidation percentage / chlorophyll photo-oxidation percentage and brassicasterol photo-oxidation percentage / chlorophyll photo-oxidation percentage ratios. Bold values are indicative of significance.

Source of variation	24-Me-cholesterol		Brassicasterol		Palmitoleic acid	
	W	<i>p-value</i>	W	<i>p-value</i>	W	<i>p-value</i>
Sea ice x SPM	180	0.4372	212.5	0.5858	4	1.0×10^{-7}
Sea ice x all traps	1	3.0×10^{-10}	2	3.7×10^{-10}	3	7.2×10^{-10}
SPM x all traps	78	3.3×10^{-6}	27	2.0×10^{-8}	242	0.1034
Surface ^a x Traps ^b	79	5.6×10^{-12}	29	7.6×10^{-14}	245	1.4×10^{-6}
Traps 5 m x Traps 30 m	10	2.0×10^{-4}	29	0.0225	19	0.0028
Traps 5 m x Traps 100 m	16	2.3×10^{-3}	0	1.5×10^{-6}	0	1.5×10^{-6}
Traps 30 m x Traps 100 m	0	3.6×10^{-5}	0	7.4×10^{-7}	0	7.4×10^{-7}

^a Sum of sea ice and SPM

^b Sum of traps