Inversion breakpoints and the evolution of supergenes

Romain Villoutreix¹, Diego Ayala², Mathieu Joron¹, Zachariah Gompert³, Jeffrey L. Feder⁴,
Patrik Nosil¹.

Molecular Ecology 30 (12): 2738-2755. <https://doi.org/10.1111/mec.15907>

¹ CEFE, CNRS, Univ. Montpellier. 1919 Route de Mende. 34293 Montpellier. France

² UMR MIVEGEC, IRD, CNRS, Univ. Montpellier, 911 avenue Agropolis, BP 64501, 34394
Montpellier, France.

³ Department of Biology, Utah State University, Utah 84322, USA

⁴ Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556,
USA

Abstract

The coexistence of discrete morphs that differ in multiple traits is common within natural populations of many taxa. Such morphs are often associated with chromosomal inversions, presumably because the recombination suppressing effects of inversions helps maintain adaptive combinations for traits that are controlled by different genes (supergene hypothesis). However, inversions can also harbor adaptive mutations at their breakpoints, leading to rise in inversion frequency in addition to (or independent from) a role for recombination suppression (breakpoint-mutation hypothesis). In this review, we first

22 describe the myriad of ways that breakpoints can create mutations. We then critically
23 examine the evidence for both hypotheses in examples of putative supergenes. Despite
24 decades of focused study, we find that evidence that inversions spread due to
25 recombination suppression, as proposed by the supergene hypothesis, is often indirect.
26 Breakpoints may harbor mutations that are adaptive in many cases, but again the evidence
27 is indirect and characterization of inversion breakpoints at the sequence level is incomplete
28 in most systems. Direct tests of the role of suppressed recombination and breakpoint
29 mutation in inversion evolution are needed. Finally, we emphasize how the two hypotheses
30 can act in conjunction, with implications for understanding the dynamics of supergene
31 evolution. We conclude that breakpoint characterization could be essential for
32 understanding the origin of complex, discrete phenotypic forms in nature.

33
34 **Keywords:** chromosomal inversion, linkage, mutation, recombination, genome evolution.

36 ***Supergenes and complex phenotypes***

37 The origin and maintenance of discrete morphs within natural populations has long
38 fascinated evolutionary biologists (Darwin, 1862; Fisher, 1930; Ford, 1971). Because of the
39 discrete nature of the phenotypic differences observed, this variation was often assumed to
40 have a genetic origin, making its study of primary interest for early geneticists (see Chapter
41 6 of (Ford, 1971) for a good overview of genetic programs of the time). Moreover, the
42 diverse nature of the traits that usually differ between these morphs (color, morphology,
43 sexual-compatibility, etc.), led to the hypothesis that the genetic basis of this discrete
44 variation is a functional unit containing multiple loci, coined as a “Super-gene” (spelled
45 supergene hereafter; see glossary) by Darlington and Mather in 1949 (Fisher, 1930; Ford,
46 1965; Mather, 1955). Fisher anticipated that genetic linkage between multiple selected loci
47 could evolve under natural selection (Fisher, 1930) and Ford later suggested that linkage
48 could be increased via translocation or chromosomal inversions (Ford, 1971). Chromosomal
49 inversions can indeed strongly reduce recombination between the genomic segments they
50 contain, a property leading to their discovery in 1921 by Sturtevant (Sturtevant, 1921).

51 Examples of discrete morphs associated with chromosomal inversions (*i.e.*, candidate
52 supergenes; see glossary) are accumulating rapidly, leading to peaked interest in supergene
53 and inversion evolution. There are currently two core hypotheses for how and why
54 supergenes and inversions evolve. First and perhaps more accepted, inversions can be
55 selected for their capacity to impede recombination between sets of epistatic or locally
56 adapted genes (*i.e.*, supergene hypothesis; see glossary) (Kirkpatrick, 2010; Kirkpatrick &
57 Barton, 2006). This hypothesis predicts that loci underlying trait variation will generally be
58 distributed across the inversion. Second, inversions can create adaptive mutations at their

breakpoints (*i.e.* breakpoint-mutation hypothesis; see glossary; Fig. 1A) (Dobzhansky, 1947, 1970; Kirkpatrick, 2010), leading to their rise in frequency via selection on breakpoint variants. This hypothesis predicts that adaptive mutations will be localized at breakpoints, rather than more widely distributed throughout the inversion. Notably, a breakpoint mutation can, in principle, be pleiotropic and affect multiple traits. Therefore, even when multiple traits are associated with an inversion, a role for recombination suppression needs to be demonstrated rather than assumed. As we discuss in more detail below, these hypotheses are not mutually exclusive and might combine to drive supergene evolution (*i.e.* breakpoint-linkage hypothesis; see glossary; Fig. 1A). Indeed, theory implies that for new inversions to establish under the supergene hypotheses, they must capture all segregating variants at selected loci in a favorable combination (Kirkpatrick & Barton, 2006). Alternatively, an inversion with an adaptive breakpoint mutation may be able to initially rise in frequency and persist for a sufficient amount of time to allow for the sorting out of any initially captured maladapted alleles (e.g., via gene conversion and double-recombination events).

Distinguishing among these hypotheses (Fig. 1A) is important as it will help answer the following major questions concerning adaptation and genome rearrangement (Fig 1.B). Do divergent selected variants evolve first and subsequently become locked together via suppressed recombination within an inversion (Kirkpatrick & Barton, 2006) ? Do inversions establish first because of an adaptive breakpoint mutation such that they are only subsequently turned into supergenes by the emergence of new variants within them (Navarro & Barton, 2003) ? Or, in contrast, pre-existing divergently selected variants and adaptive breakpoint mutations tend to coincide when supergenes form via the emergence

of an inversion? The answers to these questions are important because they determine the roles for mutation, pre-standing genetic variation and recombination in adaptation.

In this review, we critically examine the evidence for the three hypotheses mentioned previously regarding inversion evolution. We begin with a discussion of the mechanisms by which breakpoints can create adaptive mutations, as this is often a less appreciated facet of inversions than is recombination suppression. We then empirically review well-known cases of candidate supergenes (Box 2 – Fig. 1). Our collective findings lead us to propose that characterization of breakpoints at the sequence level could be crucial for understanding the evolutionary dynamics of supergenes. Specifically, the most common molecular mechanism generating inversions also generates duplications at both end of the inverted segment (Ranz et al., 2007). This offers a unique opportunity to date inversions, independently from their content, and test if they emerged before or after the divergently selected loci they may contain. Thus, breakpoint characterization could be essential for understanding the origin and maintenance of complex, discrete phenotypic forms in nature.

Molecular mechanisms by which inversion breakpoints can create adaptive mutations

The idea that inversions might be selected because of an adaptive mutation at one of their breakpoints is not new (Dobzhansky, 1947). However, this hypothesis is yet to receive highly focused study in supergene evolution, despite known molecular mechanisms that are likely to create a breakpoint mutation whenever inversions form. Two main processes can generate chromosomal inversions, ectopic recombination and double strand staggered breaks. Ectopic recombination generates inversions via a recombination event between two homologous sequences oriented head to head in a sequence (Box 1 for further details; Box 1 – Fig. 1). Double strand staggered breaks can generate inversions via the complete

detachment of a DNA fragment and its subsequent reattachment in the opposite orientation (Box 1; Box 1 – Fig. 1). Most inversions are generated by this latter mechanism (Ranz et al., 2007).

Both the mechanisms mentioned above can generate ample phenotypic effects by changing the expression of a gene (or set of genes) at breakpoints (Fig. 2). Breakpoints can also create chimeric genes (see glossary), which might sometimes have a beneficial effect. Indeed, examples of the potential adaptive potential of chimeric genes are accumulating in a diverse set of taxa (Carvunis et al., 2012; Guillén & Ruiz, 2012; Jones et al., 2012; Stewart & Rogers, 2019). For example, in *Drosophila mojavensis* a chimeric gene at an inversion breakpoint is associated with the detoxification of toxic compounds in cacti (Guillén & Ruiz, 2012). Another example is found in the threespine stickleback *Gasterosteus aculeatus*, where a gene located at an inversion breakpoint on chromosome XI harbors different exon sequences depending on the inversion orientation, and is associated with differences in muscular development between lake and oceanic phenotypes (Jones et al., 2012). While speculative, an inversion breakpoint could create a new pleiotropic chimeric gene that affects multiple traits. Moreover, the most common inversion mechanism (double strand staggered breaks) often generate duplications at both breakpoints in the derived sequence (Box 1 – Fig. 1), allowing for the creation of chimeric gene without loss of the donor genes. This property will also facilitate neo-functionalization of any gene duplicated by this inversion generation mechanism.

As we will see below, direct evidence that breakpoint mutations drive supergene evolution is scarce, but the mechanisms by which they might do so are established. With these

considerations in place we turn to evaluating the role of suppressed recombination and breakpoint mutation in the evolution of candidate examples of supergenes.

The supergene hypothesis: what is the evidence for recombination suppression among multiple loci controlling selected trait variation?

A core assumption of the supergene hypothesis (*i.e.*, that suppressed recombination is advantageous) is that multiple, linked genetic variants control trait variation that differs between inversion forms. In contrast, if variation is controlled by a single locus with pleiotropic effects on several traits, suppressed recombination is not necessarily favored or required to explain the existence of discrete morphs that differ in these traits.

Our review of the literature revealed that we still know very little about the number, location and identity of selected loci within candidate supergenes. This is not surprising given that determining the number of adaptive loci within a region of suppressed recombination is difficult, because suppressed recombination frustrates attempts at fine-scale genetic mapping. The presence of two or more selected loci in a supergene was historically inferred from genetic crosses aimed at detecting rare recombinants (*i.e.*, intermediate phenotypic forms) (Cain & Sheppard, 1954; Mather, 1950). For example, Mather described a genetic model where multiple selected loci locked within an inversion control heterostyly in *Primula sinensis*, a classic example of a putative supergene (Mather, 1950). However, recent work using modern genomic approaches suggests that Mather's simple genetic model is erroneous and needs modification (Li et al., 2016). Specifically, a 178 kilo base-pair indel containing 5 predicted genes is associated with the different floral types (Li et al., 2016), with one floral type being hemizygous for this segment and the other completely lacking it (Table 1) (Li et al., 2016).

150 A revision of the classic genetic model also occurred for the iconic land snail *Cepaea*
151 *nemoralis* (Table 1), where cryptic shell color-pattern was long thought to be associated
152 with five loci locked within an inversion (Cain & Sheppard, 1954). Recent work showed an
153 absence of recombination in the putative supergene, casting doubt on the number and
154 linkage relationships of loci controlling shell color (Gonzalez, Aramendia, & Davison, 2019).
155 These recent developments urge for more rigorous tests on the true number of selected
156 variants within supergenes and their precise genomic location(s).

157 Indeed, to our knowledge, the existence of multiple selected loci within a supergene
158 associated with an inversion has been demonstrated in only two systems, both plants (Table
159 1; Box 2). In addition, indirect but fairly convincing evidence exists in two insect systems.
160 Details of these example are contained in Box 2. Clearly, further work determining the
161 number of loci within inversions and candidate supergenes that contribute to trait variation
162 is warranted. In this regard, emerging techniques that allow functional genetic manipulation
163 of structural variants (e.g., CRISPR-Cas9) hold promise for making such progress (Hopkins,
164 Tyukmaeva, Gompert, Feder, & Nosil; Kraft et al., 2015).

165 ***The breakpoint hypothesis: what is the evidence that functional loci reside in or near***
166 ***inversion breakpoints?***

167 A core assumption of the breakpoint hypothesis is that functional loci reside in or near
168 inversion breakpoints. We found indications for this hypothesis in three systems (*Papilio*
169 *polytes*, *Solenopsis invicta* and *Timema cristinae*), but expect this number to increase when
170 breakpoint characterization is completed in more systems (Table 1).

171 In the mimetic butterfly *Papilio polytes*, an inversion breakpoint disrupts a transcriptional
172 regulator and affects the expression in embryonic wings of three neighboring genes

potentially involved in wing color-pattern development (*UXT*, *U3T* and *prospero*; Table 1) (Nishikawa et al., 2015). In the fire ant *Solenopsis invicta*, a region of reduced recombination generated via three successive inversions (Yan et al., 2020) is associated with different social structures of colonies (Wang et al., 2013). A breakpoint of the second inversion is increasing the expression of a neighboring gene (*SI2.2.0_02248*), which is a strong candidate to explain differences in colony social structures (Huang, Dang, Chang, & Wang, 2018; Yan et al., 2020). In *Timema cristinae* a massive deletion occurring at one likely breakpoint of a putative inversion deleted multiple loci known to affect color in another species of the genus (Villoutreix et al., 2020).

Interestingly, with the exception of *Timema cristinae*, indications of putative selected loci within the inversion at locations other than the breakpoint exist for these systems. We discuss the implication of this observation in the following section.

The breakpoint-linkage hypothesis and implications for evolutionary dynamics

As mentioned previously, the supergene and breakpoint-mutation hypotheses are not mutually exclusive. The selective advantage of an inversion could come from its property of impeding recombination between at least one locus within the inversion and a favorable breakpoint mutation. We refer to this mixed hypothesis as the breakpoint-linkage hypothesis (see glossary), which predicts that loci controlling trait variation will be localized both at one (or both) breakpoint(s) and within the inversion.

In theory, this hypothesis could involve different processes and dynamics compared to the classic supergene hypothesis. A supergene composed of a selected locus within an inversion and an adaptive breakpoint mutation suggests two possible evolutionary scenarios. First, pre-existing genetic variants within the inversion could become associated with a new

adaptive breakpoint mutation. Although formal modeling is required, it is possible that such a coupling could help relax some of the restrictions of a model based strictly on suppressed recombination (Kirkpatrick & Barton, 2006). For example, such coupling might mean that the initial inversion needs not to capture essentially all favorable segregating alleles in the correct linkage phase with one another. Instead, such sorting could occur after the initial establishment of the inversion, with the initial establishment of the inversion being dependent on the sum of adaptive values of the breakpoint and other genetic variants.

Second, an inversion could establish first because of an adaptive breakpoint mutation (Guerrero, Rousset, & Kirkpatrick, 2012), and subsequently evolve into a supergene by the emergence of additional genetic variants within it. This latter scenarios could be initiated in allopatry, where an inversion with an adaptive breakpoint sweeps to fixation and is selected for its recombination suppressing properties only later upon secondary contact (or introgression) and gene flow. In this latter case, allopatry allows for the evolution and capture of alternative sets of favorable alleles in alternate arrangements, by passing difficulties of the supergene emergence when gene flow occurs and adaptive alleles are segregating within populations (Feder, Gejji, Powell, & Nosil, 2011).

In terms of empirical evidence, we found indications for the breakpoint-linkage hypothesis in two systems (Table 1). In the mimetic butterfly *Papilio polytes*, an inversion contains a locus shown to affect wing color-pattern (*doublesex*) and a breakpoint mutation affecting the expression of three other genes potentially also affecting wing color-pattern (*UXT*, *U3T* and *prospero*) (Kunte et al., 2014; Nishikawa et al., 2015). In the fire ant *Solenopsis invicta*, a region of reduced recombination generated via three successive inversions contains a strong candidate gene for odor generation/reception (*Gp-9*), a mechanisms at the root of

social structure differences in fire ants (Wang et al., 2013). A breakpoint of the second inversion increase the expression of another strong candidate gene for odor reception (*Sl2.2.0_02248*) (Huang et al., 2018; Yan et al., 2020).

While functional work is needed to better elucidate the role of these different loci in *P. polytes* and *S. invicta*, these two systems highlight the possibility for breakpoints to be one of several selected loci of a supergene. The vast majority of candidate supergene systems still lack a complete characterization of inversion breakpoints (usually one breakpoint is characterized but not the other; Table 1). For this reason we expect more systems to support the breakpoint-linkage hypothesis after further investigation.

Future directions

Future work could usefully focus on two directions. First, better information on the number, location and identify of loci controlling trait variation within inversion is required. This can be achieved using crosses to create recombinant forms, as for the two plant examples discussed in Box 2. The regions identified with this technique are likely still large, and the number of loci identified here is likely to be an under-estimation. Moreover, the identification of causal variants at breakpoints will be challenging as many variants will be linked due to the lack of recombination at breakpoints (Hoffmann & Rieseberg, 2008; Matzkin, Merritt, Zhu, & Eanes, 2005). Alternatively to controlled crosses, genome wide association mapping can be applied in species that exhibit more continuous variation in the same traits as those of close relatives with discrete morphs and inversions, but that lack inversion polymorphism. This will yield more precise estimates, but need to be complemented with functional manipulation in the species with the inversion. Ultimately, functional transformations such as those offered by CRISPR-Cas9 may be required to

precisely and firmly establish the variants functionally contributing to trait variation, possibly by ‘engineering’ recombinant alleles to validate the independent effects of a given variant on trait variation (Hopkins et al.; Kraft et al., 2015).

Second, once the selected loci within the inversion are identified, further work can be implemented to characterize supergene origin, more specifically to test if the genetic variants locked in the inversion predated the inversion. To do so, we need to date the inversion independently from the selected variants it contains. An estimation of inversion age by averaging the divergence time over the whole inverted sequence is likely to underestimate inversion age, possibly resulting in the erroneous conclusion that the variants appearance predated the inversion. Indeed, recombination still occurs between inverted segments, and will only be prevented for selected loci and breakpoints. To circumvent this caveat, one can take advantage of a useful feature of most inversions: duplication at breakpoints (Ranz et al., 2007). Dating the duplication event will allow one to accurately date the inversion independently from its genomic content. Such dating could also be done for genes within the inversion and near breakpoints, but this will be less precise, depending on the amount a recombination they experience.

Such dating was shown to be effective in *Heliconius numata* butterflies (Jay et al., 2018). It requires sequence-level characterization of breakpoints, a difficult endeavor because breakpoints often reside in highly repetitive regions of the genomes (Russell B. Corbett-Detig et al., 2019; da Silva et al., 2019), which are difficult to assemble using short sequencing read technologies (Fig. 3A). Nonetheless, creative approaches such as those applied by Corbett-Detig and colleagues in *Drosophila* (Russell B. Corbett-Detig & Hartl, 2012) hold promise (Fig. 3B), as do long reads technologies such as Nanopore and PacBio.

Concluding remarks

Rapidly emerging candidate supergenes systems urge a careful assessment of the state of research in this field. We outlined how inversions can facilitate supergene evolution either via suppressed recombination or by inducing adaptive mutation at breakpoints. To our knowledge, definitive evidence for the existence of multiple selected loci within candidate supergenes has only been gathered in two systems. Thus direct evidence even for the classic supergene hypothesis is still scarce, although many examples are consistent with this idea. Little is known about the origin of supergenes, specifically if genetic variants or inversion appeared first. As an alternative or in addition to the supergene hypothesis, inversions may spread due to adaptive breakpoint mutations. Observations in multiple candidate supergenes suggest a joint role for suppressed recombination and breakpoint mutation in supergene evolution. Thus, further tests of the breakpoint-linkage hypothesis are warranted. Distinguishing and confirming the role of breakpoints in supergene evolution will likely require characterization of inversion breakpoints at the sequence level, which has yet to be completed in most systems. We propose that further study of inversion breakpoints will facilitate understanding the origin and maintenance of complex, discrete phenotypic forms in nature, and the dynamics by which supergenes evolve. Although we focused here on morphs, similar processes could apply to the evolution of other discrete units of diversity such as ecotypes, species, and sexes.

284 ***Glossary:***

285 Breakpoint-linkage hypothesis: the selective advantage to an inversion stems from both a
286 favorable effect of mutation at one of its breakpoint and the recombination suppression
287 between this mutation and at least one other selected variant within the inversion.

288 Breakpoint-mutation hypothesis (also referred as 'position effect hypothesis' in the
289 literature): the selective advantage to an inversion stems from a favorable effect(s)
290 associated with a mutation at one or both of its breakpoints.

291 Candidate supergene: case where multiple co-varying and discrete phenotypes are
292 associated with an inversion; the existence of multiple loci associated with these traits must
293 be demonstrated to advance the example from a candidate to a bonafide supergene.

294 Chimeric gene: a new gene constituted of sequences not previously associated together, for
295 example of parts of two genes, or parts of genes and non-coding DNA. Here, we consider
296 cases where the different DNA segments are brought together by an inversion breakpoint
297 (see Fig. 2B).

298 Chromosomal inversion: a chromosomal rearrangement in which a segment of a
299 chromosome is reversed end to end. This causes the inverted segment to have an opposite
300 sequence order from the ancestral sequence and reduces recombination in inversion
301 heterozygotes. Recombination is especially reduced near chromosomal inversion
302 breakpoints.

303 Chromosomal inversion breakpoint (shortened to 'inversion breakpoint' or simply
304 'breakpoint' in the main text and hereafter): the position(s) on a chromosome where DNA
305 sequence orientation shifts after the occurrence of a chromosomal inversion (i.e., the end of

306 an inversion where the genome shifts from inverted back to collinear). An inversion
307 generally has two breakpoints.

308 (Chromosomal) breakpoint mutation: sequence modification(s) at breakpoints resulting
309 from a chromosomal inversion. Mutations may also occur near breakpoints after an
310 inversion has occurred, however, here, we refer here to changes associated with the
311 creation of the rearrangement itself.

312 Discrete morphs: individuals of the same species that exhibit discrete phenotypic
313 differences (e.g., in morphology, color, behavior, physiology, etc.). In this piece, we focus on
314 cases where morphs are differentiated by multiple co-varying phenotypic traits such that
315 multiple genes may be involved.

316 Supergene: a group of genes affecting different phenotypes so seldom separated by
317 crossing-over that in effect they operate as a single genetic entity (i.e., a non-recombining
318 locus). This end may be achieved by close genetic proximity on a chromosome (i.e., tight
319 physical linkage), by inclusion within an inversion, or by other molecular mechanisms
320 impeding crossing-over. This definition is modified from that of E. B. Ford (Ford, 1965).

321 Supergene hypothesis: supergenes evolve because suppressed recombination within them
322 maintains favorable allelic combinations at multiple loci that control different traits.

323 Switch gene: A gene controlling the development of alternative discrete morphs.

324

Acknowledgements

This manuscript greatly benefited from discussions with Pr. Russel Corbett-Detig, Pr. Angus Davison, Pr. Haruhiko Fujiwara, Dr. Clemens Küpper, Pr. Cheng-Ruei Lee, Pr. Thomas Mitchell-Olds and Dr. John Wang on their respective study systems. We are grateful to Pr. Cheng-Ruei Lee for providing us photographs of *Boechera stricta*. We are grateful to Dr. Venera Tyukmaeva for insights on CRISPR-Cas9 and functional genomics. R.V. and P.N. were supported by a grant from the European Research Council (EE-Dynamics 770826, <https://erc.europa.eu/>). D.A. was supported by an ANR grant (ANR-18-CE35-0002-01 – WILDING). M. J. was supported by an ANR grant (ANR-18-CE02-0019-02 supergène). Z. G. was supported by the NSF (DEB 1638768). J. F. was supported by grants from the USDA NIFA program and NSF. All authors contributed to writing the manuscript. The authors declare having no competing interests.

(A) Are inversions supergenes?

1) Supergene hypothesis

2) Breakpoint hypothesis

3) Breakpoint-linkage hypothesis

(B) Supergene origins.

1) variants first – inversion second

2) Inversion first – supergene second

3) Inversion and supergene

Figure 1 – Hypotheses and evolutionary scenarios: **A.** Hypotheses behind inversion evolution. **B.** Evolutionary scenarios for supergene evolution. Grey squares represents a chromosome; red dots are divergently selected loci; black bars are inversion breakpoints.

(A) Transcriptional changes.

(B) Chimeric gene creation.

Figure 2 – Possible mutations at inversion breakpoints: **A.** An inversion generated through ectopic recombination put two genes into a different transcription context, altering their level of expression. **B.** An inversion generated through the staggered break mechanism generates a new chimeric switch gene, while avoiding gene loss via duplication. Arrow represent genes, with orientation indicated via arrow direction; a, b, c, d represent single strand breaks, resulting here in two staggered breaks.

(A) Misassembly of inversion breakpoint duplications

(B) Characterizing breakpoints sequences.

Figure 3 – Characterizing breakpoint at the sequence level: **A.** Using short reads technologies can lead to misassembly of inversion breakpoints, for example due to duplications generated through the staggered breaks mechanisms (Ranz et al., 2007). **B.** A method to characterize breakpoint sequences using a single reference genome and short read data, as developed by (Russell B. Corbett-Detig, Cardeno, & Langley, 2012). Black arrows symbolize aligned reads on reference genome, while grey arrows symbolize their unaligned paired reads. Breakpoint sequence characterization is possible by using these read pairs to generate and *de novo* assembly of the region. Figure redrawn and modified from (Russell B. Corbett-Detig et al., 2012) with the permission of the authors.

Box 1 - Inversion breakpoints can create a diversity of types of genetic variation via mutation.

We here discuss how inversions are formed and the associated effects of induced mutations on genetic variation at breakpoints. Two known molecular mechanisms can generate inversions: (1) ectopic recombination and, (2) double strand staggered breaks (Box 1. - Fig1) (Ling & Cordaux, 2010; Ranz et al., 2007).

The first mechanism occurs from recombination between homologous sequences that are oriented head-to-head in the original sequence, subsequently resulting in inversion of the sequence located between these elements (Box 1 – Fig. 1A). This mechanism leaves little molecular traces and generates ‘cut-and-paste’ type breakpoints. The second mechanism involves double strand breaks occurring in two positions of the original sequences (Ranz et al., 2007). Such breaks are usually staggered, meaning that they will result in fragments with stretches of single stranded DNA at their extremities (Box 1. - Fig. 1B). The DNA repair mechanism that synthesizes the reverse complement of these single stranded stretches then join the DNA sequences back together in a non-homologous way, sometimes reinserting the DNA segment in the opposite orientation, creating an inversion. When the breaks are heavily staggered (*i.e.*, with long single strand stretches) this results in inversions with duplicated sequences at both breakpoints in the derived sequence (Box 1. - Fig. 1B) (Ranz et al., 2007). In contrast, when the breaks are blunt or lightly staggered, ‘cut-and-paste’ type breakpoints without duplications are created in the derived sequence.

Despite these clear mechanisms of inversion formation, we still know very little about genetic variation at breakpoints and the mutations that breakpoint generates (Table 1). This stems largely from the fact that inversion breakpoints often lie in very repetitive regions

(Russell B. Corbett-Detig et al., 2019; da Silva et al., 2019) which are difficult to sequence and assemble and thus challenging to study (Fig. 3A). Nonetheless, some progress has been made in *Drosophila spp.* (R. B. Corbett-Detig, 2016; Russell B. Corbett-Detig & Hartl, 2012; McBroome, Liang, & Corbett-Detig, 2020; Ranz et al., 2007), where double strand staggered breaks appear to be the most common mechanism generating inversions, very often resulting in duplications at breakpoints (Ranz et al., 2007).

We note, however, that locating breakpoints is not sufficient to identify their phenotypic effects. Obtaining the derived sequence is necessary to try to understand possible phenotypic effects. This is not a straightforward task, as disruption of gene sequence by breakpoints can sometimes leads to increase in their expression as seen in fire ants (Yan et al., 2020). Nonetheless, phenotypic effects are expected, for example because both the aforementioned mechanisms can change the expression context of set of genes (Fig. 2A). In addition, the double strand staggered breaks mechanism can have another potential effect, in which duplications of sequences leads to the creation of a new chimeric gene. This occurs via merging the initial part of a gene with another coding sequence such as another gene or pseudo-gene (Fig. 2B). This phenomenon can happen without the loss of the originally disrupted genes (via their duplication), mitigating costs.

Thus, there are several possibilities by which breakpoints can create genetic variation with likely phenotypic effects. The observation that similar genomic regions are often re-used repeatedly as breakpoints for different inversions, including in different taxa, raised the possibility for a recurrent and even somewhat predictable role for breakpoints in evolution (Pevzner & Tesler, 2003).

(A) Ectopic recombination. Cut and paste breakpoint.

(B) Staggered breaks. Duplication at both ends.

Box 1 - Figure 1 - Known molecular mechanisms generating inversions: Black and grey arrows represent genes, with orientation indicated by the direction of the arrow **A.** Ectopic recombination between similar elements (transposable elements or duplicated genes) in opposite orientation in a DNA segment leads to the inversion of the interspacing sequence. **B.** Four single strand breaks (a, b, c, and d) lead to two staggered breaks. Through end repair and reattachment in the opposite orientation an inversion is generated with sequence duplication at both breakpoints (black and grey genes).

Box 2 - Determining the number of loci within inversions that contribute to trait variation

Despite seventy years of research on supergene, evidence for the existence of multiple selected loci within an inversion has only been gathered in a handful of systems.

Evidence for multiple loci controlling trait variation within an inversion has been demonstrated in *Boechea stricta* (Box 2 – Fig. 1A), a perennial plant from the Rocky Mountains. Here, two subspecies adapted to different climates have entered in secondary contact (Lee et al., 2017). The subspecies differ in multiple traits and through crosses it was demonstrated that three different QTL are associated with these divergent traits (Box 2 – Fig. 2B). Interestingly, the inversion itself has an effect on these traits, independent of the genotype at the three QTL, but the reason for this is unknown. This supergene has also had its evolutionary origins characterized such that it is known that the inversion is relatively young and likely evolved after secondary contact (Lee et al., 2017). The existence of multiple selected loci within an inversion has also been demonstrated, again via crosses, in the *Mimulus guttatus* species complex, where two life strategies (perennial vs annual) are maintained through divergent selection (Lowry & Willis, 2010). Specifically, through crosses between the *Mimulus guttatus* annual type and a closely related perennial species with a collinear gene order, *Mimulus tilingii*, Coughlan and Willis (Coughlan & Willis, 2018) identified two QTL within the inversion associated with the traits experiencing divergent selection.

Less direct, but reasonably strong, evidence for multiple loci has also been reported in two insect systems (Table 1). Here, the approach was to compare the genetic basis of traits in closely related taxa that exhibit versus lack inversion polymorphism. In the polymorphic butterfly *Heliconius numata* a region of reduced recombination generated via multiple

successive inversions control variation in color morphs. The same genomic region is known to contain at least three recombining loci associated with color variation in other species of the genus (Box 1 – Fig. 1D) (Huber et al., 2015; Mathieu Joron et al., 2011; Van Belleghem et al., 2017). It is therefore likely that the inversion in *H. numata* harbor variants at these three loci that recombine in other *Heliconius* species. A similar example stems from *Timema*, a genus of stick insect that is endemic of the western USA and Mexico and that shows within-population variation for cryptic coloration (i.e., green versus melanistic morphs). In several species, morphs are highly discrete and controlled by a region of suppressed recombination on the linkage group eight (named ‘*Mel-Stripe*’) (Nosil et al., 2018; Villoutreix et al., 2020). However, a recent study in a species displaying more quantitative color variation (*Timema chumash*) revealed that multiple genetic variants, and likely multiple genes within *Mel-Stripe*, control color variation (Villoutreix et al., 2020).

(A) *Boechera stricta*

(B) Multiple QTL within *Bs1* inversion

(C) *H. numata* aposematic morphs

(D) *H. numata* colour supergene

454

455 **Box 2 - Figure 1 – Flagship examples of supergenes:** **A.** *Boechera stricta* subspecies types.
 456 The East subspecies has smaller leaves with more trichomes. Photo credit: Cheng-Ruei Lee.
 457 **B.** Multiple QTL were detected in the *Bs1* inversion. The red, blue dashed and grey dashed
 458 lines correspond to different composited traits obtained through discriminant function
 459 analysis. See (Lee et al., 2017) for more details. Black dashed lines correspond to inversion
 460 breakpoint positions. Breakpoint are of cut-and-paste type for the *Bs1* inversion. Figure
 461 redrawn and modified with the permission of the authors. **C.** *Heliconius numata* aposematic
 462 morphs, involved in Müllerian mimicry rings with other butterflies. **D.** Aposematic morphs in
 463 *H. numata* are associated with an inversion (named P) on linkage group LG15, encompassing
 464 at least three loci associated with aposematic color morph in *H. melpomene*: Yb, Sb and N.

Species	Multiple traits?	Selection on phenotypes?	Map to an inversion?	Multiple loci?	Selection on recomb. sup.?	Adaptive breakpoint mutation?	Breakpoints sequence?	Variants first?
Aves								
Ruff - <i>Philomachus pugnax</i>	Yes. (Kupper et al., 2016; Lamichhane et al., 2016)	Possibly. (Kupper et al., 2016; Lamichhane et al., 2016)	Yes. (Kupper et al., 2016; Lamichhane et al., 2016)	Likely. (Kupper et al., 2016; Lamichhane et al., 2016)	Not tested.	No. Counter-selected. (Kupper et al., 2016; Lamichhane et al., 2016)	Yes. (Kupper et al., 2016; Lamichhane et al., 2016)	Not known.
White-throated sparrow - <i>Zonotrichia albicollis</i>	Yes. (Tuttle et al., 2016)	Yes. (Tuttle et al., 2016)	Yes. (Thomas et al., 2008; Thorneycroft, 1966; Thorneycroft, 1975; Tuttle et al., 2016)	Likely. (Tuttle et al., 2016)	Not tested.	Not known.	No.	Not known.
Invertebrates								

Anopheles – <i>Anopheles gambiae</i>	Possibly. (Cheng, Tan, Hahn, & Besansky, 2018)	Possibly. (Fouet, Gray, Besansky, & Costantini, 2012; Gray, Rocca, Costantini, & Besansky, 2009; Rocca, Gray, Costantini, & Besansky, 2009)	Yes. (Cheng et al., 2018; Coluzzi, Sabatini, della Torre, Di Deco, & Petrarca, 2002; Coluzzi, Sabatini, Petrarca, & Di Deco, 1979; Fouet et al., 2012; Gray et al., 2009; Rocca et al., 2009)	Possibly. (Cheng et al., 2012)	Not tested.	No.	Partly characterized. (Cheng et al., 2018; Sharakhov et al., 2006)	Not known.
Grove snail - <i>Cepaea nemoralis</i>	Yes. (Cain & Sheppard, 1950, 1952)	Possibly. (Cain & Sheppard, 1950, 1954; Cameron & Cook, 2012)	Not known. (Gonzalez et al., 2019)	Not known.	Not known.	Not known.	No.	Not known.
Numata longwing - <i>Heliconius numata</i>	Yes. (Mathieu Joron et al., 2011)	Yes. (Benson, 1972; Chouteau, Llaurens, Piron-Prunier, & Joron, 2017; M. Joron,	Yes. (Mathieu Joron et al., 2011).	Very likely. (Huber et al., 2015; Mathieu Joron et al., 2011; Van Belleghem	Not tested.	No.	Partly characterized. (Jay et al., 2018)	Not known.

		Wynne, Lamas, & Mallet, 1999; Kapan, 2001; Mallet & Barton, 1989; Merrill et al., 2012)		et al., 2017)				
Common Mormon - <i>Papilio polytes</i>	Yes. (Kunte et al., 2014; Nishikawa et al., 2015)	Yes. (Clarke & Sheppard, 1972)	Yes. (Kunte et al., 2014; Nishikawa et al., 2015)	Likely. (Nishikawa et al., 2015)	Not tested.	Possibly. (Nishikawa et al., 2015)	Yes. (Nishikawa et al., 2015)	Not known.
Fire ant - <i>Solenopsis invicta</i>	Yes. (Wang et al., 2013).	Possibly. (Ross & Keller, 1995)	Yes. (Wang et al., 2013)	Possibly. (Wang et al., 2013)	Not tested.	Possibly. (Huang et al., 2018; Yan et al., 2020)	Yes. (Yan et al., 2020)	Not known.
Timema walking sticks - <i>Timema cristinae</i> .	Yes. (Comeault et al., 2015)	Yes. (Cristina P. Sandoval, 1994; C. P. Sandoval & Nosil, 2005)	Likely. (Villoutreix et al., 2020)	Very likely. (Villoutreix et al., 2020)	Not tested.	Likely. (Villoutreix et al., 2020)	Partly characterized. (Nosil et al., 2018; Villoutreix et al., 2020)	Not known.

<i>Mammals</i>								
Humans – <i>Homo sapiens</i>	Yes. (Puig, Casillas, Villatoro, & Caceres, 2015; Stefansson et al., 2005)	Likely. (Stefansson et al., 2005)	Yes (Puig et al., 2015; Stefansson et al., 2005)	Possibly (Puig et al., 2015)	Not tested.	Possibly. (de Jong et al., 2012)	Yes. (Stefansson et al., 2005)	Not known.
<i>Plants</i>								
Drummond's rockcress - <i>Boechea stricta</i>	Yes. (Lee et al., 2017)	Yes. (Anderson, Lee, & Mitchell-Olds, 2011; Anderson, Lee, Rushworth, Colautti, & Mitchell-Olds, 2013; Lee et al., 2017)	Yes. (Lee et al., 2017)	Yes. (Lee et al., 2017)	Yes, but debated. (Charlesworth & Barton, 2018; Lee et al., 2017)	No. (Lee et al., 2017).	Yes. (Lee et al., 2017).	Yes.

Common yellow monkeyflower - <i>Mimulus guttatus</i> species complex	Yes. (Coughlan & Willis, 2018; Lowry & Willis, 2010)	Yes. (Lowry & Willis, 2010)	Yes. (Coughlan & Willis, 2018; Lowry & Willis, 2010)	Yes. (Coughlan & Willis, 2018)	Not tested.	Not known.	No.	Not known.
Primrose - <i>Primula</i> sp.	Yes. (Darwin, 1862)	Likely. (Piper & Charlesworth, 2008)	No. (Li et al., 2016)	Possibly. (Li et al., 2016)	Not tested.	N.A. Indel. (Li et al., 2016)	Yes. (Li et al., 2016)	N. A.
Teleosts								
Atlantic Cod – <i>Gadus morhua</i> .	Yes. (Berg et al., 2016; Sinclair-Waters et al., 2018)	Likely. (Berg et al., 2016; Sinclair-Waters et al., 2018)	Yes. (Berg et al., 2016; Kirubakaran et al., 2016; Sinclair-Waters et al., 2018)	Possibly. (Berg et al., 2016; Kirubakaran et al., 2016; Sinclair-Waters et al., 2018)	Not tested.	Not known.	No.	Not known.
Threespine sticklebacks - <i>Gasterosteus aculeatus</i>	Yes. (Jones et al., 2012)	Likely. (Jones et al., 2012)	Yes. (Jones et al., 2012)	Possibly. (Albert et al., 2007;	Not tested.	Yes. (Jones et	Yes. (Jones et al., 2012)	Not known.

				Colosimo et al., 2004)		al., 2012)		
Steelhead/rainbow trout – <i>Oncorhynchus mykiss</i>	Yes. (Nichols, Edo, Wheeler, & Thorgaard, 2008; Pearse, Miller, Abadia-Cardoso, & Garza, 2014)	Not tested.	Possibly. [(Nichols et al., 2008; O'Malley, Sakamoto, Danzmann, & Ferguson, 2003; Pearse et al., 2014; Robison, Wheeler, Sundin, Sikka, & Thorgaard, 2001)	Likely. (Lemopoulos, Uusi-Heikkilä, Huusko, Vasemägi, & Vainikka, 2018)	Not tested.	Not known.	No.	Not known.

466

467

468 **Table 1 – Examples of candidate supergene. Species:** common and latin naming. **Multiple traits:** Does morphs different by multiple traits?
469 **Selection on phenotypes:** Has selection on these traits been shown? **Map to an inversion:** Are these traits associated with and inversion?
470 **Multiple loci:** Has the existence of multiple loci controlling the aforementioned traits been demonstrated in this system? **Selection on recomb.**
471 **sup.:** Has the selective advantage of an inversion been demonstrated to come from its property of impeding recombination between multiple
472 selected loci? **Adaptive breakpoint mutation:** Is their evidence for this inversion to be selected for or against because of one of its breakpoints.
473 **Breakpoint sequence:** Have the breakpoints been characterized at the sequence level. **Variants first?:** Do adaptive variants in the putative
474 supergene predate the origins of the inversion?

Bibliography

- Albert, A. Y. K., Sawaya, S., Vines, T. H., Knecht, A. K., Miller, C. T., Summers, B. R., . . . Schluter, D. (2007). The genetics of adaptative shape shift in stickleback: Pleiotropy and effect size. *Evolution*, 62(1), 76-85.
- Anderson, J. T., Lee, C. R., & Mitchell-Olds, T. (2011). Life-History Qtls and Natural Selection on Flowering Time in *Boechera Stricta*, a Perennial Relative of *Arabidopsis*. *Evolution*, 65(3), 771-787. doi:10.1111/j.1558-5646.2010.01175.x
- Anderson, J. T., Lee, C. R., Rushworth, C. A., Colautti, R. I., & Mitchell-Olds, T. (2013). Genetic trade-offs and conditional neutrality contribute to local adaptation. *Mol Ecol*, 22(3), 699-708. doi:10.1111/j.1365-294X.2012.05522.x
- Benson, W. W. (1972). Natural-Selection for Mullerian Mimicry in *Heliconius-Erato* in Costa-Rica. *Science*, 176(4037), 936-939. doi:DOI 10.1126/science.176.4037.936
- Berg, P. R., Star, B., Pampoulie, C., Sodeland, M., Barth, J. M. I., Knutsen, H., . . . Jentoft, S. (2016). Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. *Scientific Reports*, 6(23246). doi:10.1038/srep23246
- Cain, A. J., & Sheppard, P. M. (1950). Selection in the polymorphic land snail *Cepaea nemoralis*. *Heredity*, 4(3), 275-294. doi:10.1038/hdy.1950.22
- Cain, A. J., & Sheppard, P. M. (1952). The effects of natural selection on body colour in the land snail *Cepaea*

- nemoralis. *Heredity*, 6(2), 217-231.
doi:10.1038/hdy.1952.22
- Cain, A. J., & Sheppard, P. M. (1954). NATURAL SELECTION IN CEPAEA. *Genetics*, 39(1), 89-116.
- Cameron, R., & Cook, L. (2012). CORRELATED PHENOTYPIC RESPONSES TO HABITAT DIFFERENCE IN CEPAEA NEMORALIS (L.). *FOLIA MALACOLOGICA*, 20(4), 255-263.
- Carvunis, A.-R., Rolland, T., Wapinski, I., Calderwood, M. A., Yildirim, M. A., Simonis, N., . . . Vidal, M. (2012). Proto-genes and de novo gene birth. *Nature*, 487(7407), 370-374. doi:10.1038/nature11184
- Charlesworth, B., & Barton, N. H. (2018). The Spread of an Inversion with Migration and Selection. *Genetics*, 208(1), 377-382.
- Cheng, C. D., Tan, J. C., Hahn, M. W., & Besansky, N. J. (2018). Systems genetic analysis of inversion polymorphisms in the malaria mosquito *Anopheles gambiae*. *Proc Natl Acad Sci U S A*, 115(30), E7005-E7014. doi:10.1073/pnas.1806760115
- Cheng, C. D., White, B. J., Kamdem, C., Mockaitis, K., Costantini, C., Hahn, M. W., & Besansky, N. J. (2012). Ecological Genomics of *Anopheles gambiae* Along a Latitudinal Cline: A Population-Resequencing Approach. *Genetics*, 190(4), 1417-1432.
doi:10.1534/genetics.111.137794
- Chouteau, M., Llaurens, V., Piron-Prunier, F., & Joron, M. (2017). Polymorphism at a mimicry supergene maintained by opposing frequency-dependent selection pressures. *Proc Natl Acad Sci U S A*, 114(31), 8325-8329. doi:10.1073/pnas.1702482114
- Clarke, C. A., & Sheppard, F. R. S. (1972). The genetics of the mimetic butterfly *Papilio polytes* L. *Philosophical*

541 *Transactions of the Royal Society of London. B,*
542 *Biological Sciences*, 263(855), 431-458.
543 doi:10.1098/rstb.1972.0006

544 Colosimo, P. F., Peichel, C. L., Nereng, K., Blackman, B. K.,
545 Shapiro, M. D., Schluter, D., & Kingsley, D. M. (2004).
546 The Genetic Architecture of Parallel Armor Plate
547 Reduction in Threespine Sticklebacks. *Plos Biology*,
548 2(5), e109. doi:10.1371/journal.pbio.0020109

549 Coluzzi, M., Sabatini, A., della Torre, A., Di Deco, M. A., &
550 Petrarca, V. (2002). A polytene chromosome analysis of
551 the *Anopheles gambiae* species complex. *Science*,
552 298(5597), 1415-1418. doi:10.1126/science.1077769

553 Coluzzi, M., Sabatini, A., Petrarca, V., & Di Deco, M. A.
554 (1979). Chromosomal differentiation and adaptation to
555 human environments in the *Anopheles gambiae* complex.
556 *Royal Society of Tropical Medicine and Hygiene*, 73(5),
557 483-497.

558 Comeault, Aaron A., Flaxman, Samuel M., Riesch, R.,
559 Curran, E., Soria-Carrasco, V., Gompert, Z., . . . Nosil, P.
560 (2015). Selection on a Genetic Polymorphism
561 Counteracts Ecological Speciation in a Stick Insect.
562 *Current Biology*, 25(15), 1975-1981.
563 doi:10.1016/j.cub.2015.05.058

564 Corbett-Detig, R. B. (2016). Selection on Inversion
565 Breakpoints Favors Proximity to Pairing Sensitive Sites
566 in *Drosophila melanogaster*. *Genetics*, 204(1), 259-265.
567 doi:10.1534/genetics.116.190389

568 Corbett-Detig, R. B., Cardeno, C., & Langley, C. H. (2012).
569 Sequence-Based Detection and Breakpoint Assembly of
570 Polymorphic Inversions. *Genetics*, 192(1), 131-137.
571 doi:10.1534/genetics.112.141622

572 Corbett-Detig, R. B., & Hartl, D. L. (2012). Population
573 Genomics of Inversion Polymorphisms in *Drosophila*

- melanogaster. *PLoS Genet*, 8(12), e1003056.
doi:10.1371/journal.pgen.1003056
- Corbett-Detig, R. B., Said, I., Calzetta, M., Genetti, M.,
McBroome, J., Maurer, N. W., . . . Besansky, N. J.
(2019). Fine-Mapping Complex Inversion Breakpoints
and Investigating Somatic Pairing in the *Anopheles*
gambiae Species Complex Using Proximity-Ligation
Sequencing. *Genetics*, 213(4), 1495-1511.
doi:10.1534/genetics.119.302385
- Coughlan, J. M., & Willis, J. H. (2018). Dissecting the role of
a large chromosomal inversion in life history divergence
throughout the *Mimulus guttatus* species complex. *Mol*
Ecol, 28(6), 1343-1357. doi:doi:10.1111/mec.14804
- da Silva, V. H., Laine, V. N., Bosse, M., Spurgin, L. G.,
Derks, M. F. L., van Oers, K., . . . Groenen, M. A. M.
(2019). The Genomic Complexity of a Large Inversion in
Great Tits. *Genome Biology and Evolution*, 11(7), 1870-
1881. doi:10.1093/gbe/evz106
- Darwin, C. (1862). On the Two Forms, or Dimorphic
Condition, in the Species of *Primula*, and on their
remarkable Sexual Relations. *Journal of the Proceedings*
of the Linnean Society of London. Botany, 6(22), 77-96.
doi:10.1111/j.1095-8312.1862.tb01218.x
- de Jong, S., Chepelev, I., Janson, E., Strengman, E., van den
Berg, L. H., Veldink, J. H., & Ophoff, R. A. (2012).
Common inversion polymorphism at 17q21.31 affects
expression of multiple genes in tissue-specific manner.
BMC Genomics, 13(458). doi:10.1186/1471-2164-13-
458
- Dobzhansky, T. G. (1947). Adaptive Changes Induced by
Natural Selection in Wild Populations of *Drosophila*.
Evolution, 1(1/2), 1-16. doi:10.2307/2405399

- Dobzhansky, T. G. (1970). *Genetics of the Evolutionary Process*: Columbia University Press.
- Feder, J. L., Gejji, R., Powell, T. H. Q., & Nosil, P. (2011). ADAPTIVE CHROMOSOMAL DIVERGENCE DRIVEN BY MIXED GEOGRAPHIC MODE OF EVOLUTION. *Evolution*, 65(8), 2157-2170. doi:10.1111/j.1558-5646.2011.01321.x
- Fisher, R. A. (1930). *The Genetical Theory of Natural Selection*: Oxford.
- Ford, E. B. (1965). *Genetic Polymorphism*. London: Studies Faber & Faber.
- Ford, E. B. (1971). *Ecological Genomics* (3rd ed.). London: Chapman and Hall LTD.
- Fouet, C., Gray, E., Besansky, N. J., & Costantini, C. (2012). Adaptation to Aridity in the Malaria Mosquito *Anopheles gambiae*: Chromosomal Inversion Polymorphism and Body Size Influence Resistance to Desiccation. *PLoS One*, 7(4), e34841. doi:10.1371/journal.pone.0034841
- Gonzalez, D. R., Aramendia, A. C., & Davison, A. (2019). Recombination within the *Cepaea nemoralis* supergene is confounded by incomplete penetrance and epistasis. *Heredity*, 123(2), 153-161. doi:10.1038/s41437-019-0190-6
- Gray, E. M., Rocca, K. A. C., Costantini, C., & Besansky, N. J. (2009). Inversion 2La is associated with enhanced desiccation resistance in *Anopheles gambiae*. *Malaria Journal*, 8. doi:10.1186/1475-2875-8-215
- Guerrero, R. F., Rousset, F., & Kirkpatrick, M. (2012). Coalescent patterns for chromosomal inversions in divergent populations. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 367(1587), 430-438.

- Guillén, Y., & Ruiz, A. (2012). Gene alterations at *Drosophila* inversion breakpoints provide prima facie evidence for natural selection as an explanation for rapid chromosomal evolution. *BMC Genomics*, 13(1), 53. doi:10.1186/1471-2164-13-53
- Hoffmann, A. A., & Rieseberg, L. H. (2008). Revisiting the Impact of Inversions in Evolution: From Population Genetic Markers to Drivers of Adaptive Shifts and Speciation? *Annual Review of Ecology, Evolution, and Systematics*, 39(1), 21-42. doi:10.1146/annurev.ecolsys.39.110707.173532
- Hopkins, D. P., Tyukmaeva, V. I., Gompert, Z., Feder, J. L., & Nosil, P. Functional genomics offers new tests of speciation hypotheses. *in press*.
- Huang, Y.-C., Dang, V. D., Chang, N.-C., & Wang, J. (2018). Multiple large inversions and breakpoint rewiring of gene expression in the evolution of the fire ant social supergene. *Proceedings of the Royal Society B: Biological Sciences*, 285(1878), 20180221. doi:10.1098/rspb.2018.0221
- Huber, B., Whibley, A., Poul, Y. L., Navarro, N., Martin, A., Baxter, S., . . . Joron, M. (2015). Conservatism and novelty in the genetic architecture of adaptation in *Heliconius* butterflies. *Heredity*, 114(5), 515-524. doi:10.1038/hdy.2015.22
- Jay, P., Whibley, A., Frézal, L., Rodríguez de Cara, M. Á., Nowell, R. W., Mallet, J., . . . Joron, M. (2018). Supergene Evolution Triggered by the Introgression of a Chromosomal Inversion. *Current Biology*, 28(11), 1839-1845.e1833. doi:10.1016/j.cub.2018.04.072
- Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., . . . Kingsley, D. M. (2012). The genomic basis of adaptive evolution in threespine

672 sticklebacks. *Nature*, 484, 55-61.

673 doi:10.1038/nature10944

674 [https://www.nature.com/articles/nature10944#supplementary-](https://www.nature.com/articles/nature10944#supplementary-information)
675 [information](https://www.nature.com/articles/nature10944#supplementary-information)

676 Joron, M., Frezal, L., Jones, R. T., Chamberlain, N. L., Lee, S.
677 F., Haag, C. R., . . . ffrench-Constant, R. H. (2011).

678 Chromosomal rearrangements maintain a polymorphic
679 supergene controlling butterfly mimicry. *Nature*,
680 477(7363), 203-206. doi:10.1038/nature10341

681 Joron, M., Wynne, I. R., Lamas, G., & Mallet, J. (1999).

682 Variable selection and the coexistence of multiple
683 mimetic forms of the butterfly *Heliconius numata*.
684 *Evolutionary Ecology*, 13(7-8), 721-754.

685 doi:10.1023/A:1010875213123

686 Kapan, D. D. (2001). Three-butterfly system provides a field
687 test of mullerian mimicry. *Nature*, 409(6818), 338-340.

688 doi:10.1038/35053066

689 Kirkpatrick, M. (2010). How and Why Chromosome
690 Inversions Evolve. *Plos Biology*, 8(9), e1000501.

691 doi:10.1371/journal.pbio.1000501

692 Kirkpatrick, M., & Barton, N. (2006). Chromosome
693 Inversions, Local Adaptation and Speciation. *Genetics*,
694 173(1), 419-434. doi:10.1534/genetics.105.047985

695 Kirubakaran, T. G., Grove, H., Kent, M. P., Sandve, S. R.,
696 Baranski, M., Nome, T., . . . Andersen, O. (2016). Two
697 adjacent inversions maintain genomic differentiation
698 between migratory and stationary ecotypes of Atlantic
699 cod. *Mol Ecol*, 25(10), 2130-2143.

700 doi:10.1111/mec.13592

701 Kraft, K., Geuer, S., Will, Anja J., Chan, Wing L., Paliou, C.,
702 Borschiwer, M., . . . Andrey, G. (2015). Deletions,

703 Inversions, Duplications: Engineering of Structural

- 704 Variants using CRISPR/Cas in Mice. *Cell Reports*, 10(5),
 705 833-839. doi:<https://doi.org/10.1016/j.celrep.2015.01.016>
- 706 Kunte, K., Zhang, W., Tenger-Trolander, A., Palmer, D. H.,
 707 Martin, A., Reed, R. D., . . . Kronforst, M. R. (2014).
 708 doublesex is a mimicry supergene. *Nature*, 507(7491),
 709 229-232. doi:10.1038/nature13112
- 710 Kupper, C., Stocks, M., Risse, J. E., dos Remedios, N.,
 711 Farrell, L. L., Mcrae, S. B., . . . Burke, T. (2016). A
 712 supergene determines highly divergent male reproductive
 713 morphs in the ruff. *Nature Genetics*, 48(1), 79-83.
 714 doi:10.1038/ng.3443
- 715 Lamichhaney, S., Fan, G. Y., Widemo, F., Gunnarsson, U.,
 716 Thalmann, D. S., Hoepfner, M. P., . . . Andersson, L.
 717 (2016). Structural genomic changes underlie alternative
 718 reproductive strategies in the ruff (*Philomachus pugnax*).
 719 *Nature Genetics*, 48(1), 84-88. doi:10.1038/ng.3430
- 720 Lee, C. R., Wang, B. S., Mojica, J. P., Mandakova, T., Prasad,
 721 K. Y. S. K., Goicoechea, J. L., . . . Mitchell-Olds, T.
 722 (2017). Young inversion with multiple linked QTLs
 723 under selection in a hybrid zone. *Nature Ecology &*
 724 *Evolution*, 1(5). doi:10.1038/s41559-017-0119
- 725 Lemopoulos, A., Uusi-Heikkilä, S., Huusko, A., Vasemägi,
 726 A., & Vainikka, A. (2018). Comparison of Migratory and
 727 Resident Populations of Brown Trout Reveals Candidate
 728 Genes for Migration Tendency. *Genome Biology and*
 729 *Evolution*, 10(6), 1493-1503. doi:10.1093/gbe/evy102
- 730 Li, J., Cocker, J. M., Wright, J., Webster, M. A., McMullan,
 731 M., Dyer, S., . . . Gilmartin, P. M. (2016). Genetic
 732 architecture and evolution of the S locus supergene in
 733 *Primula vulgaris*. *Nature Plants*, 2(12), 16188.
 734 doi:10.1038/nplants.2016.188
- 735 Ling, A., & Cordaux, R. (2010). Insertion Sequence
 736 Inversions Mediated by Ectopic Recombination between

- Terminal Inverted Repeats. *PLoS One*, 5(12), e15654.
doi:10.1371/journal.pone.0015654
- Lowry, D. B., & Willis, J. H. (2010). A Widespread
Chromosomal Inversion Polymorphism Contributes to a
Major Life-History Transition, Local Adaptation, and
Reproductive Isolation. *Plos Biology*, 8(9), e1000500.
- Mallet, J., & Barton, N. H. (1989). Strong Natural-Selection
in a Warning-Color Hybrid Zone. *Evolution*, 43(2), 421-
431. doi:10.2307/2409217
- Mather, K. (1950). The Genetical Architecture of Heterostyly
in *Primula sinensis*. *Evolution*, 4(4), 340-352.
doi:10.2307/2405601
- Mather, K. (1955). Polymorphism as an Outcome of
Disruptive Selection. *Evolution*, 9(1), 52-61.
doi:10.2307/2405357
- Matzkin, L. M., Merritt, T. J. S., Zhu, C.-T., & Eanes, W. F.
(2005). The structure and population genetics of the
breakpoints associated with the cosmopolitan
chromosomal inversion In(3R)Payne in *Drosophila*
melanogaster. *Genetics*, 170(3), 1143-1152.
doi:10.1534/genetics.104.038810
- McBroome, J., Liang, D., & Corbett-Detig, R. (2020). Fine-
scale position effects shape the distribution of inversion
breakpoints in *Drosophila melanogaster*. *Genome*
Biology and Evolution. doi:10.1093/gbe/evaa103
- Merrill, R. M., Wallbank, R. W. R., Bull, V., Salazar, P. C.
A., Mallet, J., Stevens, M., & Jiggins, C. D. (2012).
Disruptive ecological selection on a mating cue.
Proceedings of the Royal Society B-Biological Sciences,
279(1749), 4907-4913. doi:10.1098/rspb.2012.1968
- Navarro, A., & Barton, N. H. (2003). Chromosomal
Speciation and Molecular Divergence--Accelerated

- Evolution in Rearranged Chromosomes. *Science*, 300(5617), 321-324. doi:10.1126/science.1080600
- Nichols, K. M., Edo, A. F., Wheeler, P. A., & Thorgaard, G. H. (2008). The genetic basis of smoltification-related traits in *Oncorhynchus mykiss*. *Genetics*, 179(3), 1559-1575. doi:10.1534/genetics.107.084251
- Nishikawa, H., Iijima, T., Kajitani, R., Yamaguchi, J., Ando, T., Suzuki, Y., . . . Fujiwara, H. (2015). A genetic mechanism for female-limited Batesian mimicry in *Papilio* butterfly. *Nature Genetics*, 47(4), 405-409. doi:10.1038/ng.3241
- Nosil, P., Villoutreix, R., de Carvalho, C. F., Farkas, T. E., Soria-Carrasco, V., Feder, J. L., . . . Gompert, Z. (2018). Natural selection and the predictability of evolution in *Timema* stick insects. *Science*, 359(6377), 765-770. doi:10.1126/science.aap9125
- O'Malley, K. G., Sakamoto, T., Danzmann, R. G., & Ferguson, M. M. (2003). Quantitative Trait Loci for Spawning Date and Body Weight in Rainbow Trout: Testing for Conserved Effects Across Ancestrally Duplicated Chromosomes. *Journal of Heredity*, 94(4), 273-284. doi:10.1093/jhered/esg067
- Pearse, D. E., Miller, M. R., Abadia-Cardoso, A., & Garza, J. C. (2014). Rapid parallel evolution of standing variation in a single, complex, genomic region is associated with life history in steelhead/rainbow trout. *Proceedings of the Royal Society B-Biological Sciences*, 281(1783). doi:10.1098/rspb.2014.0012
- Pevzner, P., & Tesler, G. (2003). Human and mouse genomic sequences reveal extensive breakpoint reuse in mammalian evolution. *Proceedings of the National Academy of Sciences*, 100(13), 7672. doi:10.1073/pnas.1330369100

- Piper, J., & Charlesworth, B. (2008). The evolution of distyly in *Primula vulgaris*. *Biological Journal of the Linnean Society*, 29(2), 123-137. doi:10.1111/j.1095-8312.1986.tb01827.x
- Puig, M., Casillas, S., Villatoro, S., & Caceres, M. (2015). Human inversions and their functional consequences. *Briefings in Functional Genomics*, 14(5), 369-379. doi:10.1093/bfpg/elv020
- Ranz, J. M., Maurin, D., Chan, Y. S., von Grotthuss, M., Hillier, L. W., Roote, J., . . . Bergman, C. M. (2007). Principles of Genome Evolution in the *Drosophila melanogaster* Species Group. *Plos Biology*, 5(6), e152. doi:10.1371/journal.pbio.0050152
- Robison, B. D., Wheeler, P. A., Sundin, K., Sikka, P., & Thorgaard, G. H. (2001). Composite Interval Mapping Reveals a Major Locus Influencing Embryonic Development Rate in Rainbow Trout (*Oncorhynchus mykiss*). *Journal of Heredity*, 92(1), 16-22. doi:10.1093/jhered/92.1.16
- Rocca, K. A. C., Gray, E. M., Costantini, C., & Besansky, N. J. (2009). 2La chromosomal inversion enhances thermal tolerance of *Anopheles gambiae* larvae. *Malaria Journal*, 8. doi:10.1186/1475-2875-8-147
- Ross, K. G., & Keller, L. (1995). Ecology and Evolution of Social-Organization - Insights from Fire Ants and Other Highly Eusocial Insects. *Annual Review of Ecology and Systematics*, 26, 631-656. doi:DOI 10.1146/annurev.es.26.110195.003215
- Sandoval, C. P. (1994). Differential visual predation on morphs of *Timema cristinae* (Phasmatodeae:Timemidae) and its consequences for host range. *Biological Journal of the Linnean Society*, 52(4), 341-356. doi:doi:10.1111/j.1095-8312.1994.tb00996.x

- Sandoval, C. P., & Nosil, P. (2005). Counteracting selective regimes and host preference evolution in ecotypes of two species of walking-sticks. *Evolution*, 59(11), 2405-2413.
- Sharakhov, I. V., White, B. J., Sharakhova, M. V., Kayondo, J., Lobo, N. F., Santolamazza, F., . . . Besansky, N. J. (2006). Breakpoint structure reveals the unique origin of an interspecific chromosomal inversion 2La in the *Anopheles gambiae* complex. *Proceedings of the National Academy of Sciences*, 103(16), 6258-6262. doi:10.1073/pnas.0509683103
- Sinclair-Waters, M., Bradbury, I. R., Morris, C. J., Lien, S., Kent, M. P., & Bentzen, P. (2018). Ancient chromosomal rearrangement associated with local adaptation of a postglacially colonized population of Atlantic Cod in the northwest Atlantic. *Mol Ecol*, 27(2), 339-351. doi:10.1111/mec.14442
- Stefansson, H., Helgason, A., Thorleifsson, G., Steinthorsdottir, V., Masson, G., Barnard, J., . . . Stefansson, K. (2005). A common inversion under selection in Europeans. *Nature Genetics*, 37(2), 129-137. doi:10.1038/ng1508
- Stewart, N. B., & Rogers, R. L. (2019). Chromosomal rearrangements as a source of new gene formation in *Drosophila yakuba*. *PLoS Genet*, 15(9), e1008314. doi:10.1371/journal.pgen.1008314
- Sturtevant, A. H. (1921). A Case of Rearrangement of Genes in *Drosophila*. *Proc Natl Acad Sci U S A*, 7(8), 235-237. doi:10.1073/pnas.7.8.235
- Thomas, J. W., Cáceres, M., Lowman, J. J., Morehouse, C. B., Short, M. E., Baldwin, E. L., . . . Martin, C. L. (2008). The Chromosomal Polymorphism Linked to Variation in Social Behavior in the White-Throated Sparrow (*Zonotrichia albicollis*) Is a Complex Rearrangement and

868 Suppressor of Recombination. *Genetics*, 179(3), 1455-
869 1468. doi:10.1534/genetics.108.088229

870 Thornycroft, H. B. (1966). Chromosomal Polymorphism in
871 the White-Throated Sparrow, *Zonotrichia albicollis*
872 (Gmelin). *Science*, 154(3756), 1571-1572.

873 Thornycroft, H. B. (1975). A Cytogenetic Study of the
874 White-Throated Sparrow, *Zonotrichia albicollis*
875 (Gmelin). *Evolution*, 29(4), 611-621.
876 doi:10.2307/2407072

877 Tuttle, E. M., Bergland, A. O., Korody, M. L., Brewer, M. S.,
878 Newhouse, D. J., Minx, P., . . . Balakrishnan, C. N.
879 (2016). Divergence and Functional Degradation of a Sex
880 Chromosome-like Supergene. *Current Biology*, 26(3),
881 344-350.

882 Van Belleghem, S. M., Rastas, P., Papanicolaou, A., Martin,
883 S. H., Arias, C. F., Supple, M. A., . . . Papa, R. (2017).
884 Complex modular architecture around a simple toolkit of
885 wing pattern genes. *Nature Ecology & Evolution*, 1,
886 0052. doi:10.1038/s41559-016-0052

887 [http://www.nature.com/articles/s41559-016-](http://www.nature.com/articles/s41559-016-0052#supplementary-information)
888 [0052#supplementary-information](http://www.nature.com/articles/s41559-016-0052#supplementary-information)

889 Villoutreix, R., de Carvalho, C. F., Soria-Carrasco, V.,
890 Lindtke, D., De-la-Mora, M., Muschick, M., . . . Nosil, P.
891 (2020). Large-scale mutation in the evolution of a gene
892 complex for cryptic coloration. *Science*, 369(6502), 460-
893 466. doi:10.1126/science.aaz4351

894 Wang, J., Wurm, Y., Nipitwattanaphon, M., Riba-Grognuz,
895 O., Huang, Y. C., Shoemaker, D., & Keller, L. (2013). A
896 Y-like social chromosome causes alternative colony
897 organization in fire ants. *Nature*, 493(7434), 664-668.
898 doi:10.1038/nature11832

899 Yan, Z., Martin, S. H., Gotzek, D., Arsenault, S. V., Duchon,
900 P., Helleu, Q., . . . Keller, L. (2020). Evolution of a

901 supergene that regulates a trans-species social
902 polymorphism. *Nature Ecology & Evolution*, 4, 240-249.
903