Negative mass hypothesis in cosmology and the nature of dark energy

Jean-Pierre Petit, Gilles d'Agostini

► To cite this version:

Jean-Pierre Petit, Gilles d'Agostini. Negative mass hypothesis in cosmology and the nature of dark energy. *Astrophysics and Space Science*, 2021, 354, pp.611 - 615. 10.1007/s10509-014-2106-5 . hal-03426503

HAL Id: hal-03426503

<https://hal.science/hal-03426503>

Submitted on 12 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Negative mass hypothesis in cosmology and the nature of dark energy

Petit J.P.^{*} and D'Agostini G.[†]
(Dated:)

The observed acceleration of the universe arises a puzzling question. What is the nature of a dark energy that would cause this phenomenon? We recall the arguments against the existence of negative matter on General Relativity grounds. They vanish if the universe is considered as a manifold M_4 associated to two coupled metrics, solutions of a coupled field equation system. We build a non-steady solution where the positive species accelerates while the negative one decelerates. So that dark energy is replaced by (dominant) negative matter action.

Keywords: Dark energy; Bimetric model; Negative mass.

INTRODUCTION

As pointed out by Riess et al. [24], Perlmutter et al. [17], reviewed by Riess [22], Filippenko and Riess [9], Leibundgut [16], observations of type Ia supernovae at redshift $z < 1$ rise a serious problem, showing that the universe accelerates, this phenomenon being attributed to some puzzling “dark energy”. Expanding this investigation on SNe Ia comforts such conclusion (Knop et al. [15]). This has been extended to larger redshifts samples (Tonry et al. [28]; Barris et al. [2]). See also Riess [23]. Nobody anticipated such startling phenomenon. The magnitude of the observed acceleration was not anticipated by theory and continues to defy a post facto explanation. What this newly discovered component of the universe could be made of?

CAN NEGATIVE MASS PLAY A ROLE IN PHYSICS?

The observed acceleration of the universe suggests the effect of a negative pressure, associated to a negative mass content. Is it possible that the Universe would contain negative energy particles? The dynamic group theory (Souriau [27]) shows that such concept comes from the antichron components of the Poincaré group. This last contains four connex components. The first four form the orthochron subgroup, or restricted Poincaré group. The last four form the antichron subset, whose elements reverse time. The coadjoint action of this group on its momentum shows that time inversion goes with energy, and mass inversion. Quantum theory of fields analyzes the inversion of space and time too. In dynamical groups theory groups are built with real components. In QFT the T operator become complex, so that it can be either linear and unitary, either antilinear and antiunitary. Weinberg [29, pp. 75–76] writes, we quote:

“At this point we have not *decided* whether P and T are linear and unitary or antilinear and antiunitary. The *decision* is an easy one. Setting $\rho = 0$ in Eq. (2.6.4) gives $PiHP^{-1} = -iH$, where $H \equiv P^0$ is the energy operator. If P were antiunitary and antilinear then it would anticommute with is, so $PHP^{-1} = -H$. But for any state Ψ of energy $E > 0$, there would be another state $P^{-1}\Psi$ of energy $-E$. *There are no states of negative energy* (energy less than that of the vacuum), so we are *forced to choose* the other alternative: P is *linear* and *unitary* and commutes than anticommutes with H. On the other hand, setting $\rho = 0$ in Eq. (2.6.6) yields $TiHT^{-1} = -iH$. If we supposed that T is linear and unitary we could simply cancel the is, and find $THT^{-1} = -H$, with the again *disastrous conclusion* that for any state Ψ of energy E there is another state $T^{-1}\Psi$ of energy $-E$. *To avoid this, we are forced to conclude* that T is antilinear and antiunitary”. We see than the quantum theory of fields is built from *a priori choices about fundamental operators*, which are done in order *to avoid the disastrous fact to include negative energy particles in the theory*”.

^{*} jppetit1937@yahoo.fr; BP 55 84122 Pertuis France

[†] dagostini.gilles@laposte.net

This absence is nothing but a pure hypothesis. A stronger critic comes from General Relativity theory and was pointed out in Bondi [3]. Newton's law emerges naturally from Einstein's equation, through Newtonian approximation. Considering the action of a particle (2) on a particle (1), Bondi defines three kinds of masses:

$$m_i^{(1)} \frac{d^2 r_1}{dt^2} = \frac{G(r_2 - r_1) m_p^{(1)} m_a^{(2)}}{|r_2 - r_1|} \quad (1)$$

m_i is the inertial mass, m_p the passive gravitational mass and m_a the active gravitational mass. Immediately the equivalence principle gives $m_i = m_p$ so that (1) becomes:

$$\frac{d^2 r_1}{dt^2} = \frac{G(r_2 - r_1) m_a^{(2)}}{|r_2 - r_1|} \quad (2)$$

which shows that positive mass attracts both positive and negative matter. Negative mass repels the two. So that two negative masses repel each other and cannot form clumps by gravitational instability. On another hand, if two masses with same absolute values and opposite signs encounter, the positive one escapes, while the negative one runs after it. The two experience a constant acceleration, but energy and momentum are conserved, because of the two masses are negative. Such preposterous features banned negative mass from cosmology during more than half a century.

REGIONS THAT BEHAVE AS IF THEY WERE FILLED BY NEGATIVE MATTER

Several authors suggested that the voids are a fundamental element of the large scale structure (LSS), with bubble-like structure (De Lapparent et al. 5; Kirshner et al. 14; Rood 25; Geller and Huchra 10; Petit 18, 19; Piran 21; El-Ad et al. 7, 8). El-Ad et al. [6] suggested that initial tiny perturbations, in the primordial density field, "acting as if the had a negative mass", could extend and play a role in the LSS. He made the hypothesis, without justification, that negative mass could own a negative active mass (from Bondi's definition), but positive inertial and passive masses, so that two negative masses would mutually attract. Negative material could form clusters. Positive and negative masses would repel each other, which was studied in references of Petit [18, 19]. Piran developed a Void Finder Algorithm which identifies voids in reshift surveys and measure their size and underdensity. In a more recent paper Izumi et al. [13] consider gravitational lensing shear by the magnifying lens models such as negative mass compact objects causing gravitational repulsion on light, like a concave lens. They suggest, as on reference of Knop et al. [15] that negative masses might attract each other, and form negative massive clumps, so that they could reside in cosmological voids (and there, they mention Piran's work). The demagnifying effect was previously predicted in Knop et al. [15], due to *negative lensing effect*. In addition they have shown that the images by the lens models are radially distorted, due to the repulsive effect (instead tangentially, when due to positive masses).

Classically, the observed distortion of the images of galaxies (weak lensing effect) is used to determine the three-dimensional distribution of some hypothetical CDM. Conversely, when decoding the data, if one assumes that the observed distortion is radial, instead tangential, it could provide three-dimensional negative matter distribution. If this last is found as a regular set of clumps, located at the center of the observed big voids that would be in favor of the presence of such negative mass clumps in our universe.

TOWARDS A BIMETRIC DESCRIPTION OF THE UNIVERSE

General relativity is based on a M_4 manifold, associated to a single Riemannian metric, considered as solution of Einstein's field equation. On such grounds, negative matter cannot cohabit with positive matter. What about a different geometrical framework? A first bimetric description was proposed in Petit [18, 19] and Petit et al. [20], then in Henry-Couannier [11] and Hossenfelder [12]. Let us consider the following coupled field equations system:

$$R_{\mu\nu}^{(+)} - \frac{1}{2} R^{(+)} g_{\mu\nu}^{(+)} = \chi \left(T_{\mu\nu}^{(+)} + \varphi T_{\mu\nu}^{(-)} \right) \quad (3a)$$

$$R_{\mu\nu}^{(-)} - \frac{1}{2}R^{(-)}g_{\mu\nu}^{(-)} = -\chi \left(\phi T_{\mu\nu}^{(+)} + T_{\mu\nu}^{(-)} \right) \quad (3b)$$

$g_{\mu\nu}^{(+)}$ metric refers to positive energy (and mass if they have one) particles. $g_{\mu\nu}^{(-)}$ refers to *negative* energy (and mass if they have one) particles. $R_{\mu\nu}^{(+)}$ and $R_{\mu\nu}^{(-)}$ are Ricci tensors, built from metrics $g_{\mu\nu}^{(+)}$ and $g_{\mu\nu}^{(-)}$. For sake of brevity we will sometimes use the notation: $f \in \{+, -\}$. In mixed form, we will write the tensors:

$$T^{(f)}_{\nu}{}^{\mu} = \begin{pmatrix} \rho^{(f)}c^2 & 0 & 0 & 0 \\ 0 & -p^{(f)} & 0 & 0 \\ 0 & 0 & -p^{(f)} & 0 \\ 0 & 0 & 0 & -p^{(f)} \end{pmatrix} \quad \text{with} \begin{cases} \rho^{(f)} > 0 \text{ and } p^{(f)} > 0 & \text{for } f = "+" \\ \rho^{(f)} < 0 \text{ and } p^{(f)} < 0 & \text{for } f = "-" \end{cases} \quad (4)$$

φ and ϕ are functions that will be defined further (to ensure energy conservation). Each metric produces its own geodesics family. The two are distinct, so that opposite energy particles cannot encounter, on pure geometric grounds. They interact only through gravitation. We assume that positive light, composed by positive energy photons, cruises along null geodesics of metric $g_{\mu\nu}^{(+)}$, while negative light, composed by negative energy photons, cruises along null geodesics of metric $g_{\mu\nu}^{(-)}$. We assume that structures made of positive masses can emit and capture positive energy photons. Similarly, structures made of negative masses can emit and capture negative energy photons. So that on geometric grounds, negative structures would be invisible to us, through our eyes or astronomic instruments.

NON-STEADY SOLUTION FOR DUST UNIVERSE

We assume this bimetric universe is homogeneous and isotropic, so that the Riemannian metrics become:

$$\left(ds^{(f)} \right)^2 = c^2 dt^2 - \left(a^{(f)} \right)^2 \frac{du^2 + u^2 (d\theta^2 + \sin^2 \theta d\varphi^2)}{\left(1 + k^{(f)} \frac{u^2}{4} \right)^2} \quad (5)$$

Introducing these metrics in the system (3a) and (3b), with $p^{(+)} \cong 0$ and $p^{(-)} \cong 0$ we get classical systems:

$$\frac{3}{c^2 (a^{(+)})^2} \left(\frac{da^{(+)}}{dt^2} \right)^2 + \frac{3k^{(+)}}{c^2 (a^{(+)})^2} = -\chi c^2 \left[\rho^{(+)} + \varphi \rho^{(-)} \right] \quad (6a)$$

$$\frac{2}{c^2 (a^{(+)})^2} \frac{d^2 a^{(+)}}{dt^2} + \frac{1}{c^2 (a^{(+)})^2} \left(\frac{da^{(+)}}{dt} \right)^2 + \frac{k^{(+)}}{c^2 (a^{(+)})^2} = 0 \quad (7a)$$

$$\frac{3}{c^2 (a^{(-)})^2} \left(\frac{da^{(-)}}{dt^2} \right)^2 + \frac{3k^{(-)}}{c^2 (a^{(-)})^2} = +\chi c^2 \left[\phi \rho^{(+)} + \rho^{(-)} \right] \quad (6b)$$

$$\frac{2}{c^2 (a^{(-)})^2} \frac{d^2 a^{(-)}}{dt^2} + \frac{1}{c^2 (a^{(-)})^2} \left(\frac{da^{(-)}}{dt} \right)^2 + \frac{k^{(-)}}{c^2 (a^{(-)})^2} = 0 \quad (7b)$$

Applying classical mathematical methods (Adler et al. 1) the compatibility conditions of sets (6a) and (7a) and (6b) and (7b) gives:

$$3 \frac{da^{(+)}}{a^{(+)}} + \frac{d \left[\rho^{(+)} + \varphi \rho^{(-)} \right]}{\left[\rho^{(+)} + \varphi \rho^{(-)} \right]} = 0 \quad (8a)$$

$$3 \frac{da^{(-)}}{a^{(-)}} + \frac{d[\phi \rho^{(+)} + \rho^{(-)}]}{[\phi \rho^{(+)} + \rho^{(-)}]} = 0 \quad (8b)$$

The conservation of the energy (and mass):

$$E = \rho^{(+)} c^2 \left(a^{(+)}\right)^3 + \rho^{(-)} c^2 \left(a^{(-)}\right)^3 \quad (9)$$

is ensured if:

$$\varphi = \left(\frac{a^{(-)}}{a^{(+)}}\right)^3, \quad \phi = \left(\frac{a^{(+)}}{a^{(-)}}\right)^3, \quad \phi = \varphi^{-1} \quad (10)$$

So that coupled field equation system becomes:

$$R_{\mu\nu}^{(+)} - \frac{1}{2} R^{(+)} g_{\mu\nu}^{(+)} = \chi \left[T_{\mu\nu}^{(+)} + \left(\frac{a^{(-)}}{a^{(+)}}\right)^3 T_{\mu\nu}^{(-)} \right] \quad (11a)$$

$$R_{\mu\nu}^{(-)} - \frac{1}{2} R^{(-)} g_{\mu\nu}^{(-)} = -\chi \left[\left(\frac{a^{(+)}}{a^{(-)}}\right)^3 T_{\mu\nu}^{(+)} + T_{\mu\nu}^{(-)} \right] \quad (11b)$$

Let:

$$\chi = -\frac{8\pi G}{c^4} \quad (12)$$

$$\left(a^{(+)}\right)^2 \frac{d^2 a^{(+)}}{dt^2} = -\frac{4\pi G}{c^4} E \quad (13a)$$

$$\left(a^{(-)}\right)^2 \frac{d^2 a^{(-)}}{dt^2} = +\frac{4\pi G}{c^4} E \quad (13b)$$

THE NATURE OF “DARK ENERGY”

Assume $E < 0$. Then $a''^{(+)} > 0$ and $a''^{(-)} < 0$. If we suppose that our visible part of the universe corresponds to positive mass, then it accelerates, while the negative species decelerates. Equation (13b) identifies to Friedman equation, while (13a) identifies to Bonnor's [1989] solution:

$$\begin{aligned} a^{(+)}(u) &= \alpha^2 \cosh^2 u \\ t^{(+)}(u) &= \alpha^2 \left(1 + \frac{\sinh 2u}{2} + u \right) \end{aligned} \quad (14)$$

From dynamic group theory (Souriau 27) all particles, including photons, may have negative mass. If L_o is the element of the orthochron restricted Lorentz group, let us consider the following extension of the Poincaré group.

$$\begin{pmatrix} \lambda\mu & 0 & \phi \\ 0 & \lambda L_o & C \\ 0 & 0 & 1 \end{pmatrix} \quad \text{with } \lambda = \pm 1 \text{ and } \mu = \pm 1 \quad (15)$$

which acts on a five-dimensional Kaluza space. The dimension of the group is 11. Then the momentum gets an extra scalar q , identified by J.M. Souriau as the electric charge (Souriau 26). The coadjoint action of the group gets an additional relation:

$$q' = \lambda\mu q \quad (16)$$

Matter-antimatter duality holds in the negative mass world. The negative matter, which acts as “dark energy” is composed by negative energy (and negative mass, if they have one) particles: electrons, protons, neutrons, neutrinos, quarks, an so on, plus their antiparticles, plus photons.

NEWTONIAN APPROXIMATION

To develop approximation we need to expand the metrics into series around Lorentzian metrics. The Universe is not homogeneous. When we develop classical Newtonian approximation we assume that the zeroth order term is Lorentzian, which means that this metric is solution of the field equation with a null second member, describing an empty portion of space. It is not necessary that this zeroth order solution corresponds to a description of the whole universe, only a restricted portion of space, small enough to encompass the interaction of particles, planets or stars, considered as mass-points. Similarly, whatever is the time-dependent model, it is possible to consider that the system is quasi steady. In our bimetric description, we consider a finite portion of quasi-steady empty space-time (null curvature, Lorentzian metrics).

$$g^{(+)} = \eta^{(+)} + \varepsilon\eta^{(+)}, \quad g^{(-)} = \eta^{(-)} + \varepsilon\eta^{(-)} \quad (17)$$

The two scale factor $a^{(+)}$ and $a^{(-)}$ are considered as constants. Then, field equations are expanded into a series. Neglecting the second order terms, we find:

$$\varepsilon\gamma_{00|\beta|\beta}^{(+)} = -\chi c^2 \left[\delta\rho^{(+)} + \left(\frac{a^{(-)}}{a^{(+)}} \right)^3 \delta\rho^{(-)} \right] \quad (18a)$$

$$\varepsilon\gamma_{00|\beta|\beta}^{(-)} = \chi c^2 \left[\left(\frac{a^{(+)}}{a^{(-)}} \right)^3 \delta\rho^{(+)} + \delta\rho^{(-)} \right] \quad (18b)$$

Defining the potentials:

$$\varphi^{(+)} = \varepsilon\gamma_{00}^{(+)} \frac{c^2}{2}, \quad \varphi^{(-)} = \varepsilon\gamma_{00}^{(-)} \frac{c^2}{2} \quad (19)$$

we get:

$$\frac{dx^{(+)\alpha}}{dt^2} = -\frac{\partial\varphi^{(+)}}{\partial x_{\alpha}^{(+)}}, \quad \frac{dx^{(-)\alpha}}{dt^2} = -\frac{\partial\varphi^{(-)}}{\partial x_{\alpha}^{(-)}} \quad (20)$$

$$\sum_{\alpha=1}^3 \frac{\partial^2 \varphi^{(+)}}{\partial x^{(+)\alpha} \partial x_{\alpha}^{(+)}} = 4\pi G \left[\delta\rho^{(+)} + \left(\frac{a^{(-)}}{a^{(+)}} \right)^3 \delta\rho^{(-)} \right] \quad (21a)$$

$$\sum_{\alpha=1}^3 \frac{\partial^2 \varphi^{(-)}}{\partial x^{(-)\alpha} \partial x_{\alpha}^{(-)}} = -4\pi G \left[\left(\frac{a^{(+)}}{a^{(-)}} \right)^3 \delta \rho^{(+)} + \delta \rho^{(-)} \right] \quad (21b)$$

which are Poisson-like equations, associated to our bimetric model. Building the dynamics from this set of equations would perhaps require a more refined solution, involving different speeds of light $c^{(+)}$ and $c^{(-)}$, and possibly time scale factors $T^{(+)}$ and $T^{(-)}$. Anyway, it is possible to derive those laws in the simplified case $a^{(+)} = a^{(-)}$. Then:

$$\Delta \varphi^{(+)} = -\Delta \varphi^{(-)} = 4\pi G \left(\delta \rho^{(+)} + \delta \rho^{(-)} \right) \quad (22)$$

$$\frac{d^2 x^{\alpha}}{dt^2} = -\frac{\partial \varphi^{(+)}}{\partial x_{\alpha}}, \quad \frac{d^2 x^{(-)\alpha}}{dt^2} = -\frac{\partial \varphi^{(-)}}{\partial x_{\alpha}} = \frac{\partial \varphi^{(+)}}{\partial x_{\alpha}} \quad (23)$$

The subsequent interaction laws differ from GR's: positive masses attract together through Newton's law. Negative masses attract together through Newton's law. Opposite signs masses repel each other through an anti-Newton's law.

Referring to the previous works of Bondi [3] and Bonnor [4] the puzzling problem of runaway phenomenon is eliminated. By the way this model fits classical verification associated to GR. In effect, as the two species repel each other, where positive mass is present, negative mass is almost absent and the coupled field equations system reduces to:

$$R_{\mu\nu}^{(+)} - \frac{1}{2} R^{(+)} g_{\mu\nu}^{(+)} \simeq \chi T_{\mu\nu}^{(+)} \quad (24a)$$

$$R_{\mu\nu}^{(-)} - \frac{1}{2} R^{(-)} g_{\mu\nu}^{(-)} \simeq -\chi \left(\frac{a^{(+)}}{a^{(-)}} \right)^3 T_{\mu\nu}^{(+)} \quad (24b)$$

On another hand, considering portions of space containing a sphere filled by constant density matter (positive or negative), surrounded by void makes “internal” and “external” Schwarzschild solutions features clearer, as firstly presented in Petit [19], introducing negative gravitational lensing effect, evoked in Izumi et al. [13].

-
- [1] Adler, R., Bazin, R., Schiifer, M.: Introduction to General Relativity. McGraw-Hill, New York (1967)
 - [2] Barris, B., et al.: *Astrophys. J.* **602**, 571 (2004)
 - [3] Bondi, H.: Negative mass in general relativity. *Rev. Mod. Phys.* **29** (3), 423 (1957)
 - [4] Bonnor, W.B.: Negative mass and general relativity. *Gen. Relativ. Gravit.* **21** (11) (1989)
 - [5] De Lapparent, V., Geller, M.H., Huchra, J.P.: *Astrophys. J. Lett.* **302**, L1 (1986)
 - [6] El-Ad, H., Piran, T.: *Astrophys. J.* **491**, 421 (1997)
 - [7] El-Ad, H., Piran, T., da Costa, L.N.: *Astrophys. J. Lett.* **462**, L13 (1996)
 - [8] El-Ad, H., Piran, T., da Costa L.N.: *Mon. Not. R. Astro. Soc.*, **287**, 790 (1997)
 - [9] Filippenko, A.V., Riess, A.G.: In: *AIP Conf. Proc.*, vol. 540, pp. 227–246 (2001), Particle
 - [10] Geller, M.J., Huchra, J.P.: *Science* **246**, 897 (1989)
 - [11] Henry-Couannier, F.: *Int. J. Mod. Phys. A.* **20**, 2341 (2005)
 - [12] Hossenfelder, S.: A Bimetric theory with exchange symmetry. *Phys. Rev. D* **78**, 044015 (2008)
 - [13] Izumi, K., Hagiwara, C., Nakajima, K., Kitamura, T., Asada, H.: Gravitational lensing shear by an exotic lens with negative convergence or negative mass. *Phys. Rev. D, Part. Fields* **88**, 024049 (2013)
 - [14] Kirshner, R.P., Oemler, A. Jr., Schlechter, P.L., Schectman, S.A.: *Astrophys. J.* **314**, 496, (1987)
 - [15] Knop, R., et al.: *Astrophys. J.* **598**, 102, (2003)

- [16] Leibundgut, B.: *Annu. Rev. Astron. Astrophys.*, **39**, 67, (2001)
- [17] Perlmutter, S., et al.: *Astrophys. J.* **517**, 565 (1999)
- [18] Petit, J.P.: The missing mass problem. *Il Nuovo Cimento B* **109**, 697–710 (1994)
- [19] Petit, J.P.: Twin universe cosmology. *Astrophys. Space Sci.* **226**, 273–307 (1995)
- [20] Petit, J.P., Midy, P., Landhseat, F.: Twin matter against dark matter. In: *Conf. on Astr. and Cosm., “Where Is the Matter?”*, Marseille (2001).
- [21] Piran, T.: On gravitational repulsion. *Gen. Relativ. Gravit.*, **29** (11), 1363 (1997)
- [22] Riess, A.G.: *Publ. Astron. Soc. Pac.* **112**, 1284 (2000)
- [23] Riess, A.G.: *Astrophys. J.* **607**, 665—687 (2004)
- [24] Riess, A.G. et al.: *Astron. J.* **116**, 1009 (1998)
- [25] Rood, H.J.: *Annu. Rev. Astron. Astrophys.* **26**, 245 (1988)
- [26] Souriau, J.M.: *Géométrie et Relativité*. Hermann, Paris (1964) (French edition only)
- [27] Souriau, J.M.: *Structure des Systèmes Dynamiques*, Dunod, France (1970) (*Structure of Dynamical Systems*, Birkhäuser, Boston, 1997)
- [28] Tonry, J.T., et al.: *Astrophys. J.* **594**, 1 (2003)
- [29] Weinberg, S.: *The Quantum Theory of Fields*. Cambridge University Press, Cambridge (2005)