Estimating extreme sea levels combining systematic observed skew surges and historic record sea levels

Laurie Saint Crieg, Yasser Hamdi, Eric Gaume, Taha Ouarda

► To cite this version:

Laurie Saint Crieg, Yasser Hamdi, Eric Gaume, Taha Ouarda. Estimating extreme sea levels combining systematic observed skew surges and historic record sea levels. European Geosciences Union General Assembly, vEGU21, Apr 2021, VIRTUAL, Austria. 2021. hal-03431752

HAL Id: hal-03431752

<https://hal.science/hal-03431752>

Submitted on 16 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Estimating extreme sea levels combining systematic observed skew surges and historic record sea levels

Laurie Saint Cricq¹, Yasser Hamdi¹, Eric Gaume² and Taha B.M.J. Ouarda³
¹IRSN, BEHRIG, France, ²Université Gustave Eiffel, GERS, France, ³INRS, ETE, Canada
 Contact: laurie.saintcricq@irsrn.fr

INTRODUCTION

The estimation of the sea levels corresponding to a high return period is crucial for coastal planning. High tide levels can be seen as the combination of the predicted astronomical high tide and the skew surge (i.e. the difference between the predicted astronomical high tide and the observed maximum sea level during a tidal cycle). This presentation is a contribution to the integration of historical information (i.e. events observed before the beginning of systematic tide gauge recordings) to better estimate extreme sea levels. It is largely recommended by the literature to separately model astronomical high tides and skew surges. Historic records consist of record sea level values. Their corresponding skew surges may be estimated but the exhaustiveness of historic skew surge series, which is an essential criterion for an unbiased statistical inference¹, cannot be guaranteed. Indeed, extreme skew surges can easily remain unnoticed if they occur with low or moderate high tides and do not generate extreme sea levels². This study proposes to combine, in a single Bayesian inference procedure, information of two different nature: measured systematic skew surges and observed historic extreme sea levels to estimate the statistical distribution of skew surges. The proposed approach is compared to two previously published methods aiming at valuating historical reconstructed skew surges^{3,4}.

PROPOSED MODEL

- The global likelihood expression of systematic skew surges and historic sea levels is:

$$L(X_{sys}, Z_{hist} | \theta, \lambda) = L(X_{sys} | \theta, \lambda) L(Z_{hist} | \theta, \lambda)$$

Extreme skew surges above the threshold u are chosen to be modelled by a General Pareto (GP) distribution:

$$\forall x > u, F_{\theta}(x) = \begin{cases} 1 - \left[1 + \xi \frac{(x-u)}{\sigma}\right]^{-\frac{1}{\xi}}, & \xi \neq 0, \\ 1 - \exp\left(-\frac{x-u}{\sigma}\right), & \xi = 0. \end{cases}$$

where $\theta = (\sigma, \xi)$ are the scale ($\sigma > 0$) and shape ($\xi \in \mathbb{R}$) parameters.

- The likelihood formulation of systematic skew surges ($>u$) is:

$$L(X_{sys} | \theta, \lambda) = P_{\lambda}(N = n) \prod_{i=1}^n f_{\theta}(x_{sys,i})$$

with f_{θ} the GP density function.

The number of systematic skew surges exceeding u during the systematic period w_S follows a Poisson process of intensity λw_S .

- The likelihood formulation of historic sea levels accounts for:

- [A] the $N - h$ ordinary historical sea levels that did not exceed η_H ...
- [B] the h extreme historic sea levels that exceeded the threshold η_H ... during the historical period

$$L(Z_{hist} | \theta, \lambda) = \underbrace{G_{\theta, \lambda}(\eta_H)^{N-h}}_{[A]} \underbrace{\left[1 - G_{\theta, \lambda}(\eta_H)\right]^h \prod_{i=1}^h \frac{g_{\theta, \lambda}(Z_{hist,i})}{1 - G_{\theta, \lambda}(\eta_H)}}_{[B]}$$

where $g_{\theta, \lambda}$ and $G_{\theta, \lambda}$ are respectively the density and the cumulative distribution functions of sea levels which can be numerically computed given the statistical distributions of the skew surges and of the maximum astronomical tidal levels.

METHOD

- Data and Monte Carlo simulations

Time series of hourly sea level observations are recorded by tide gauges of the SHOM (Service Hydrographique et Oceanographique de la Marine) and are processed to retrieve series of astronomical high tides and skew surges. Record historic sea levels are contained in the database « Tempêtes et submersions marines » developed by IRSN. For a given site, a statistical distribution is adjusted to the observed series of skew surges, with parameters (λ, σ, ξ) . 1000 Random samples of skew surges and maximum sea levels (after a convolution with the empirical series of observed high tides), with the same characteristics than the observed systematic and historical series are then generated through Monte Carlo simulations, to test the proposed statistical estimation method.

- Properties of the estimated 100-year quantile and parameters with maximum likelihood (ML)

The estimated 100-year quantiles and model parameters corresponding to the ML ($\hat{Q}_{100}^{ML}$, $\hat{\lambda}^{ML}$, $\hat{\sigma}^{ML}$, $\hat{\xi}^{ML}$) obtained for the 1000 random samples are compared to their real values (Q_{100} , λ , σ , ξ): see Fig.1.

- Reliability of the posterior credibility intervals

In the Bayesian framework, the result is the posterior density of parameters conditionally to the data which allow to compute credibility intervals for the estimated quantiles. The reliability of the posterior credibility intervals is evaluated with a diagnosis method named the rank histogram⁵ (Fig. 2). It consists in estimating the non exceedance probability of the real quantile $P(\hat{Q}_{100} < Q_{100})$, according to the computed credibility intervals for each random sample. $P(\hat{Q}_{100} < Q_{100})$ should be uniformly distributed if the credibility intervals are reliable.

CASE STUDY AND RESULTS

Case study: La Rochelle (tide gauge located on the French Atlantic Coast)

- Systematic period (1941 – 2017): $w_S = 32.58$ years, $\mu = 0.62$ m, $n = 34$
- Historical period (1866 – 1940): $w_H = 80$ years, $\eta_H = 7.15$ m, $h = 4$ (7.15m - 7.15m - 7.30m - 7.40m)

Fig 1. Accuracy of the 100-year quantile and the parameters estimated based on the maximum likelihood

- The proposed model provides unbiased estimates with the ML (compared to the previously published methods) and reduces the uncertainties especially of the 100-year quantile Q_{100} and the shape parameter ξ of the skew surge distribution.

Fig 2. Rank histograms

- $P(\hat{Q}_{100} < Q_{100})$ is uniformly distributed. The posterior credibility intervals for the 100-year quantile seem accurate with the proposed model but not with the previously published methods.

Fig 3. Adjusted statistical models for the observed series at La Rochelle

- The credibility intervals obtained with the proposed approach well fit the data.
- Integrating historic sea levels with the proposed model allows to decrease the width of the credibility intervals.

CONCLUSION

- The proposed approach combines, in a single Bayesian inference, skew surges for the recent period and record sea levels for the historical period.
- It has been successfully tested and applied to four different sites located on the French Channel and Atlantic coast (La Rochelle illustrated here). Unlike some other previously published methods, it provides unbiased quantile estimates and accurate credibility intervals.
- For more details, an article will come soon...

References

- E. Gaume, "Flood frequency analysis: The Bayesian choice", Wiley Interdis-ciplinary Reviews: Water 5 (2018), 10.1002/wat2.1290
- S. Outten et al., "Re-assessing extreme sea level events through interplay of tides and storm surges" (2020), 10.5194/egusphere- egu2020- 8000
- Y. Hamdi et al., "Use of historical information in extreme-surge frequency estimation: the case of marine flooding on the La Rochelle site in France" (2015), 10.5194/nhess-15-1515-2015
- R. Frau et al., "The use of historical information for regional frequency analysis of extreme skew surge" (2018), 10.5194/nhess-18-949-2018
- J. Bellier, "Prévisions hydrologiques probabilistes dans un cadre multivarié: quels outils pour assurer abilité et coherence spatio-temporelle?" (2018), PhD Thesis