The update of the line positions and intensities in the line list of carbon dioxide for the HITRAN2020 spectroscopic database

E.V. Karlovets, I.E. Gordon, L.S. Rothman, R. Hashemi, R.J. Hargreaves, G.C. Toon, A. Campargue, V.I. Perevalov, P. Čermák, M. Birk, et al.

► To cite this version:

E.V. Karlovets, I.E. Gordon, L.S. Rothman, R. Hashemi, R.J. Hargreaves, et al.. The update of the line positions and intensities in the line list of carbon dioxide for the HITRAN2020 spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 2021, 276, pp.107896. 10.1016/j.jqsrt.2021.107896 . hal-03433127

HAL Id: hal-03433127

<https://hal.science/hal-03433127>

Submitted on 17 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal of Quantitative Spectroscopy and Radiative Transfer

The update of the line positions and intensities in the line list of carbon dioxide for the HITRAN2020 spectroscopic database

--Manuscript Draft--

Manuscript Number:	
Article Type:	VSI:HITRAN2020
Keywords:	Carbon dioxide; CO2 line lists; HITRAN database; HITRAN2020; spectroscopic line parameters
Corresponding Author:	Ekaterina Karlovets Harvard-Smithsonian Center for Astrophysics Cambridge, MA United States
First Author:	E.V. Karlovets
Order of Authors:	E.V. Karlovets I.E. Gordon L.S. Rothman R. Hashemi R.J. Hargreaves G.C. Toon A. Campargue V.I. Perevalov P. Čermák M. Birk G. Wagner J.T. Hodges J. Tennyson S.N. Yurchenko
Abstract:	<p>This paper describes the updates of the line positions and intensities for the carbon dioxide transitions in the 2020 edition of the HITRAN spectroscopic database. The new line list for all 12 naturally abundant isotopologues of carbon dioxide replaces the previous one from the HITRAN2016 edition. This update is primarily motivated by several issues related to deficient HITRAN2016 line positions and intensities that have been identified from laboratory and atmospheric spectra. Critical validation tests for the spectroscopic data were carried out to find problems caused by inaccuracies in CO₂ line parameters. New sources of data were selected for the bands that were deemed problematic in the HITRAN2016 edition. Extra care was taken to retain the consistency in the data sources within the bands. The comparisons with the existing theoretical and semi-empirical databases (including ExoMol, NASA Ames, and CDSD-296) and with available experimental works were carried out. The HITRAN2020 database has been extended by including additional CO₂ bands above 8000 cm⁻¹, and magnetic dipole lines of CO₂ were introduced in HITRAN for the first time by including the v₂+v₃ band in the 3.3-μm region. Although the main topic of this article are line positions and intensities, for consistency a recent algorithm for the line-shape parameters proposed in Hashemi et al. JQSRT (2020) was reapplied (after minor revisions) to the line list.</p>
Suggested Reviewers:	Yan Tan tanyan@ustc.edu.cn Livio Gianfrani livio.gianfrani@unicampania.it Keyoon Sung

Highlights

- The new HITRAN2020 line list for all 12 naturally abundant isotopologues of carbon dioxide replaces the previous one from the HITRAN2016 edition;
- Evaluation of the HITRAN CO₂ line lists by comparison to laboratory and atmospheric spectra below 8310 cm⁻¹ was carried out;
- The updates made for the bands having mixed CDSD and UCL-IAO line intensities in the HITRAN2016 line list below 8000 cm⁻¹ are presented;
- The CO₂ bands in the (1.4-2.1) μm region were improved;
- New CO₂ bands above 8000 cm⁻¹ were added to the HITRAN2020 edition from the new high-temperature UCL-4000 ¹²C¹⁶O₂ line list (ExoMol database) and the CRDS spectra for the ¹⁶O¹²C¹⁸O isotopologue;
- The magnetic dipole 01111-00001 band of the ¹²C¹⁶O₂ isotopologue in the 3.3 μm region was introduced into HITRAN for the first time;

The update of the line positions and intensities in the line list of carbon dioxide for the HITRAN2020 spectroscopic database

E.V.Karlovets^{1,2*}, I.E.Gordon^{1*}, L.S.Rothman¹, R. Hashemi¹, R.J.Hargreaves¹, G.C. Toon³, A.Campargue⁴, V.I. Perevalov⁵, P. Čermák⁶, M.Birk⁷, G.Wagner⁷, J.T.Hodges⁸, J. Tennyson⁹, and S.N.Yurchenko⁹

¹ Harvard-Smithsonian Center for Astrophysics, Atomic and Molecular Physics Division, 60 Garden St, Cambridge, MA, USA

² Tomsk State University, Laboratory of Quantum Mechanics of Molecules and Radiative Processes, 36, Lenin Avenue, 634050, Tomsk, Russia

³ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

⁴ University of Grenoble Alpes, CNRS UMR 5588, LIPhy, Grenoble F-38000, France

⁵ V.E. Zuev Institute of Atmospheric Optics, 1, Academician Zuev square, 634055 Tomsk, Russia

⁶ Department of Experimental Physics, Faculty of Mathematics, Physics and Informatics, Comenius University, Mlynská dolina, 842 48 Bratislava, Slovakia

⁷ Remote Sensing Technology Institute, German Aerospace Center (DLR), D-82234 Wessling, Germany

⁸ National Institute of Standards and Technology, Gaithersburg, MD 20899, USA

⁹ Department of Physics and Astronomy, University College London, Gower Street, WC1E 6BT London, UK

Number of pages: 44

20 May 2021

Number of tables: 2

Number of figures: 17

Key words: Carbon dioxide; CO₂ line lists; HITRAN; spectroscopic line parameters.

* Corresponding author:

E-mails: ekarlovets@gmail.com (E.V.Karlovets), igordon@cfa.harvard.edu (I.E.Gordon)

Abstract

This paper describes the updates of the line positions and intensities for the carbon dioxide transitions in the 2020 edition of the HITRAN spectroscopic database. The new line list for all 12 naturally abundant isotopologues of carbon dioxide replaces the previous one from the HITRAN2016 edition. This update is primarily motivated by several issues related to deficient HITRAN2016 line positions and intensities that have been identified from laboratory and atmospheric spectra. Critical validation tests for the spectroscopic data were carried out to find problems caused by inaccuracies in CO₂ line parameters. New sources of data were selected for the bands that were deemed problematic in the HITRAN2016 edition. Extra care was taken to retain the consistency in the data sources within the bands. The comparisons with the existing theoretical and semi-empirical databases (including ExoMol, NASA Ames, and CDSD-296) and with available experimental works were carried out. The HITRAN2020 database has been extended by including additional CO₂ bands above 8000 cm⁻¹, and magnetic dipole lines of CO₂ were introduced in HITRAN for the first time by including the $\nu_2+\nu_3$ band in the 3.3- μ m region. Although the main topic of this article are line positions and intensities, for consistency a recent algorithm for the line-shape parameters proposed in Hashemi et al. JQSRT (2020) was reapplied (after minor revisions) to the line list.

1. Introduction

A plethora of critical applications drives a constant demand for ever higher-quality spectroscopic parameters of carbon dioxide (CO₂) transitions. Atmospheric CO₂ is the key gaseous contributor to the greenhouse effect in the terrestrial atmosphere. Its continuous increase in the atmosphere over the last two centuries from anthropogenic sources is considered the principal driver of climate change. The Martian and Venusian (and those of many rocky exoplanets) atmospheres consist mostly of carbon dioxide with mole fractions of over 95%. Monitoring CO₂ is important in combustion. Currently, there are several satellite-based observatories, including GOSAT[1–3], GOSAT-2 [4], OCO-2 [5,6], OCO-3 [7], MIPAS [8], and ACE [9] as well as ground-based networks TCCON [10] and NDACC [11], monitoring atmospheric CO₂. Accurate interpretation of the data from these missions is necessary to make informed decisions

71 regarding controlling of the anthropogenic contribution to the greenhouse effect and ultimately
72 the climate change. All aforementioned CO₂ remote sensing activities depend on the reliability
73 of the reference CO₂ spectroscopy.

74 The HITRAN molecular spectroscopic database is the widely recognized standard
75 providing the spectroscopic data for atmospheric applications. Among many other molecules, it
76 includes the necessary calculated and experimental parameters for all naturally abundant
77 isotopologues of carbon dioxide. The HITRAN2016 [12] CO₂ line list for 12 stable isotopologues
78 contains 559874 transitions. Most of the issues related to spectral completeness (at least below
79 8000 cm⁻¹) and insufficient accuracy of line parameters were resolved in HITRAN2016 [12].
80 However, there is still a lot of room for improvement to meet the ever-increasing demands of
81 the scientific community. Since the release of the HITRAN2016 database[12], many new
82 experimental and theoretical data have become available, and they can be used as sources for
83 improving line positions, line intensities, and line-shape parameters of CO₂. The HITRAN-related
84 updates, including the CO₂-air and CO₂-CO₂ line-shape parameters together with the speed
85 dependence of the broadening parameters, their temperature-dependent exponents, the full
86 and first-order line mixing as well as their temperature dependencies have been published in
87 Hashemi et al. [13].

88 In the present work, we will mainly focus on the issues associated with deficiencies in line
89 positions and intensities in HITRAN2016. Most of these problematic cases have been identified
90 in new laboratory and atmospheric spectra, mainly from the Kitt Peak National Observatory,
91 MkIV balloon, cavity ring down spectroscopy (CRDS) measurements, and the Total Carbon
92 Column Observing Network (TCCON) [10,14–16]. A general review of the CO₂ bands affected by
93 these issues was performed and alternative sources of data for each CO₂ problematic band were
94 considered. Apart from new experimental data, the additional resources for improvements were
95 provided by two recent comprehensive CO₂ line lists, namely the 2019 version of the semi-
96 empirical CDSD-296 [17] list and the ExoMol variational nuclear motion line list [18], hereafter
97 referred to as UCL-4000. Using these spectroscopic data, we improved and extended the CO₂ line
98 lists for all 12 stable isotopologues of carbon dioxide.

The updates of the line positions, intensities, and line-shape parameters were incorporated into the carbon dioxide line list of the 2020 edition of the HITRAN spectroscopic database [16]. The remainder of this paper is organized as follows. In Section 2, we present an overview of CO₂ line lists in the HITRAN2016, ExoMol, Ames, and CDSD-296 spectroscopic databases. In Section 3, we describe evaluations of the HITRAN CO₂ line list based on laboratory and atmospheric measurements between 670 and 8310 cm⁻¹. For wavenumbers below 8000 cm⁻¹, we discuss line intensity updates to the HITRAN2016 line list for bands provided by the Carbon Dioxide Spectroscopic Databank (CDSD) based on global fits of semi-empirical models to measurements and those obtained by the *ab initio* calculations of University College London (UCL). In Section 4, we discuss new measurements with sub-percent relative uncertainty [19–22] that were used to improve some CO₂ bands in the 1.4 to 2.1-μm region. In Section 5, we present new CO₂ bands above 8000 cm⁻¹ that have been included in the HITRAN2020 edition from the latest high-temperature line list of the ExoMol [18] database (with appropriate cut-off) for the main CO₂ isotopologue and the CRDS spectra [14] for the ¹⁶O¹²C¹⁸O isotopologue. The line parameters of the ν₂+ν₃ magnetic dipole band of the ¹²C¹⁶O₂ isotopologue, which have been introduced into HITRAN for the first time, will be presented in Section 6. The revision which updates the air- and self-broadened line-shape parameters of CO₂, described in Ref. [13], will be discussed in Section 7. The presentation of the final version of the HITRAN2020 CO₂ line list will be given in Section 8, followed by closing discussion and conclusions in Section 9.

2. Review of the current carbon dioxide line lists

The present section describes the status of the most currently used spectroscopic databases of carbon dioxide: HITRAN2016 [12], NASA Ames [23], and the recent ExoMol [18] and CDSD-296 [17] line lists. New laboratory and atmospheric spectra helped to expose several problems in the HITRAN2016 CO₂ line list occurring in various spectral regions (for more details, see Section 3). The CDSD-296, ExoMol, and NASA Ames line lists were used to update, extend or improve the line positions and intensities, including a number of problematic bands, in the 2020 version of the HITRAN spectroscopic database [24].

The CO₂ line list for the 12 stable isotopologues in the 2016 version of HITRAN [12] contained 559874 CO₂ transitions and covered the wavenumber range of 0.757-14075.298 cm⁻¹. The HITRAN2016 data for the ¹⁴C¹⁶O₂ radioactive isotopologue in HITRAN2016 were provided as static files rather than through the relational-database structure featured in HITRAN*online* [25]. The HITRAN2016 carbon dioxide line list, below 8000 cm⁻¹, comprised CDSD line positions [26] and *ab initio* line intensities from UCL [27–29] except for the bands that were identified as "sensitive"[27], where CDSD (circa 2015) intensity values were preferred. Most of the line parameters for wavenumbers in the region above 8000 cm⁻¹ were taken from the 2015 version of the CDSD-296 database [26]. An extended and improved version of the CDSD for atmospheric applications (CDSD-296) was published in 2019 [17]. This database contains a computed line list based on global modelling of the line positions and intensities using the method of effective operators [30,31]. The global model of the CO₂ absorption spectrum is a polyad model with a block diagonal structure, each polyad being characterized by the polyad number $P = 2V_1 + V_2 + 3V_3$ (where V_i are the vibrational quantum numbers). For each CO₂ isotopologue, the effective Hamiltonians and effective dipole moment parameters were fitted to measured line positions and intensities, respectively. More than 530000 spectral lines for the 12 stable isotopologues of CO₂ covering the 345-14075 cm⁻¹ spectral range were included in this database. The reference temperature is 296K, and the intensity cut-off is 10⁻³⁰ cm/molecule. A comparison between the line positions of common lines from the HITRAN2016 and CDSD-296 [17] databases versus wavenumber in the 0-14075 cm⁻¹ spectral region is presented in **Fig. 1**. This plot shows the good agreement between the line positions from HITRAN2016 (mostly based on the previous CDSD version [26]) and those from the new version of the CDSD-296 [17]. As a result, most of the line positions for the 12 CO₂ isotopologues between 345 and 14075 cm⁻¹ in HITRAN2020 have been updated using the CDSD-296 [17] line positions. It should be noted that the slightly updated version of the CDSD-296 database (uploaded at ftp.iao.ru as pub/CDSD-296/cdsd-296_version_1.rar) was used. This version differs from the CDSD-296 published in Tashkun et al. [17] in the following ways: (i) the $\Delta P = 6$ region of the ¹⁶O¹²C¹⁸O isotopologue was recalculated using the new set of effective dipole moment parameters; (ii) a total of 226 lines with vibrational quantum number $\Delta l_2 = 4$ of the ¹²C¹⁶O₂, ¹³C¹⁶O₂, and ¹⁶O¹²C¹⁸O isotopologues were added. The

lower-state energies and the uncertainty codes were also transferred from the CDSD-296 line list to the HITRAN2020 line list.

The recent high-temperature line list from the ExoMol database (named UCL-4000) for the main isotopologue of CO₂ (¹²C¹⁶O₂) was published in Yurchenko et al. [15]. This line list contains almost 2.5×10^9 transitions involving about 3.5×10^6 states. The wavenumber range of the CO₂ list is from 0 to 20000 cm⁻¹ with the lower-state energies up to 16000 cm⁻¹ and $J \leq 202$. These results were generated using UCL's *ab initio* dipole moment surface (DMS) [32] and the semi-empirical potential energy surface (PES) Ames-2 [22] with the variational program TROVE [33]. The UCL-4000 line list was converted to HITRAN format and compared with the HITRAN2016 line list. As a result, more than 3600 spectral lines above 8000 cm⁻¹ were added to the HITRAN2020 spectroscopic database [24] (see Section 5).

Fig. 1. Differences between the line positions from the HITRAN2016 [12] and CDSD-296 [17] databases.

The NASA Ames line lists for 13 CO₂ isotopologues were published by Huang et al. [23]. They were computed at 296 K using the empirically refined Ames-2 potential energy surface and pure *ab initio* DMS-2 dipole moment surface with $J \leq 150$ and the lower-state energy up to 24000 cm⁻¹. Critical validation tests of the available spectroscopic data were carried out using the existing theoretical and semi-empirical databases, including ExoMol, NASA Ames, and CDSD-296 [17,18,23] and with published experimental works [14,34–42] .

Accordingly, the present work allowed revealing some problems with the HITRAN2016 CO₂ line list and thus allowed improving and extending the line lists for all 12 stable isotopologues in the final version of the HITRAN2020 CO₂ line list.

3. Evaluation of the HITRAN CO₂ line lists by laboratory and atmospheric spectra

In this section, we present the general review of problematic bands identified in new laboratory and atmospheric spectra, mainly those obtained from the Kitt Peak National Observatory, MkIV balloon, CRDS, and TCCON [10,14–16]. Most of the issues have been associated with deficient line positions and intensities for which critical validation tests were performed and alternative sources of data for each problematic band were suggested.

Several CO₂ line lists, including HITRAN 2008 [43], HITRAN2012 [44], and HITRAN2016 [12], were evaluated by fitting laboratory (mainly Kitt Peak) and atmospheric solar absorption spectra (MkIV and TCCON) between 670 and 8310 cm⁻¹ [15,16]. The studied region was divided into several windows, most encompassing at least one complete CO₂ absorption band or a sub-branch. The GFIT spectral fitting algorithm [45] was used in all cases assuming a Voigt line profile and no line mixing. The line lists were evaluated in terms of the RMS fitting residuals and the window-to-window consistency of the retrieved gas amounts. There was no analysis of the separate isotopologues. The results showed progressive overall RMS fit improvements through use of each successive HITRAN version. This analysis showed that the HITRAN2020 CO₂ line list is substantially better than all previous HITRAN versions. In particular, in HITRAN2016, there were a few regions where it is worse than a previous version in terms of RMS fitting residuals (around 1800 cm⁻¹, 4800 cm⁻¹, and 6400 cm⁻¹). As an example, **Fig. 2** shows the absolute RMS residuals of the fits for 41 windows averaged over the 137 Kitt Peak and 12 JPL laboratory spectra in the 670-

214 8310 cm^{-1} spectral region. The pressure range is from 0.013 kPa-93 kPa (0.1 Torr to 700 Torr)
215 except for 2 Kitt Peak spectra (1.71 kPa and 1.89 kPa) covering 600-1400 cm^{-1} . However, in this
216 plot, the HITRAN2020 line_list gives worse fits in the 3730 cm^{-1} window in Kitt Peak lab spectra
217 than most previous line_lists. This is due to the SBHWs being too small in this region (the Kitt Peak
218 lab spectra include some with 200 Torr of pure CO_2). This SBHW issue has no effect on fits to
219 atmospheric spectra. The large discrepancies were also identified in the 4800 cm^{-1} regions, where
220 the HITRAN2016 line list was worse than previous editions in terms of RMS fitting residuals. The
221 same situation in this region was observed in the MkIV balloon spectra (low pressure). The Kitt
222 Peak laboratory spectrum revealed rotationally-dependent errors at the 10-15% level in the *ab*
223 *initio* intensities for the 40002-01101 band of the $^{12}\text{C}^{16}\text{O}_2$ isotopologue. Fits to a Kitt Peak
224 laboratory spectrum in the 4800 cm^{-1} region and zooms into four large residuals are presented
225 in **Fig.3**. Also, it was shown that the scatter factor of the 40002-01101 band is slightly elevated
226 (1.4 to 1.7). In addition to that, for low J , around $J=1$ to 3, there is a visible spike in the scatter
227 factor, which suggests a J -localized resonance. Indeed, by analyzing predictions within the
228 effective operator model [30], it was found that the 40002 and 21113 vibrational levels of the $P=$
229 8 polyads are interacting through a Coriolis interaction. This interaction causes systematic
230 problems with the *ab initio* intensities [27] for the transitions involving either of these two
231 vibrational states.

Fig. 2. The upper panel shows the absolute value of the mean *RMS* fit for 5 different CO₂ line lists: HIT2008: HITRAN2008 [43]; HIT2012: HITRAN2012 [44], HIT2016: HITRAN2016[12], ATM2018: 523230 CO₂ lines [15], HIT2020: HITRAN2020 [24]. The lower panel shows the differences from HITRAN2012 [44].

Fig.3. The fits to a Kitt Peak laboratory spectrum in the 4800 cm⁻¹ region using the HITRAN2016 linelist [12]. The positive residuals arise from the 40002-01101 band of the ¹²C¹⁶O₂ isotopologue.

For the HITRAN2016 CO₂ line list, there are 24 bands involving the 40002 and 21113 vibrational levels affected by Coriolis interaction. To quantify the impact of this perturbation on intensities, validation tests for the 40002 - 01101, 21113 - 01101, and 21113 - 11102 bands of the ¹²C¹⁶O₂ isotopologue were carried out using the published experimental values [34,35] and data from the various carbon dioxide spectroscopic databases [12,17,23] (**Fig. 4**). This comparison showed that the CDSD-296 line intensities [17] are preferable for the 40002 - 01101 and 21113 - 11102 bands whereas, in the case of the 21113-01101 band, the *ab initio* intensities from AMES [23] are the best choice. However, because of the lack of measurements we could not make similar comparisons for the numerous remaining bands affected by these Coriolis interactions. We hope that new experimental data will become available for further validation tests of bands involving the 40002 and 21113 vibrational levels.

Fig.4. Ratio of the experimental line intensities of the 40002 - 01101, 21113 - 01101, and 21113 - 11102 bands (centers: 4808 cm^{-1} , 4809 cm^{-1} , and 3544 cm^{-1} , respectively) from Refs [34,35] to those from the various CO_2 spectroscopic databases [12,17,23] .

In the 1800-2000 cm^{-1} spectral region, CO_2 retrievals of mixing ratio based on the HITRAN2016 database produces CO_2 retrieved amounts that are 5% larger than those based on previous line lists. The VMR Scale Factors (VSF) obtained by averaging each retrieved single-spectrum VSF value over all 19 MkIV balloon spectra that were fitted for each window are plotted versus wavenumber in **Fig. 5**. The MkIV instrument simultaneously records the region 600 cm^{-1}

to 5650 cm^{-1} , so derived VSFs should have good window-to-window consistency. The upper circle indicates anomalously high HITRAN2016 values in the 1800-2000 cm^{-1} spectral region corroborating observation in laboratory spectra.

Fig.5. The VMR Scale Factors (VSF) obtained by averaging each retrieved single-spectrum VSF value over all 19 MkIV balloon spectra that were fitted for each window. The measured VSF values are plotted *versus* wavenumber. See the designations of the databases in the caption of Fig. 2.

It was found that the most significant deviations in this region correspond to the 11102 - 00001 band near 1933 cm^{-1} . The 11101 - 00001 and 11102 - 00001 bands borrow intensities from the strong 00011 - 00001 band *via* Coriolis interaction. It was shown in Ref. [17] that the *ab initio* AMES line intensities [23] for the 11101 - 00001 band deviate considerably from the observations. A comparison of the 11102 - 00001 line intensities with the published measurements [36,37] and the different CO_2 databases [12,17,23] for the $^{12}\text{C}^{16}\text{O}_2$, $^{13}\text{C}^{16}\text{O}_2$, and $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ isotopologues is presented in **Fig. 6**. Replacement of line intensities of the CO_2 bands considered in the HITRAN2016 database with CDSD-296 [17] rectifies the above-mentioned discrepancy in the 1800-2000 cm^{-1} region. For consistency, the line intensities of the 11101 -

00001 band for other CO₂ isotopologues were also replaced by the intensities from the CDSD database in the HITRAN2020 line list. The 20003-01101 and 12202-01101 hot bands of ¹²C¹⁶O₂ located near 1900 cm⁻¹ were also increased by 5% to match the ATM values [15]. The intensity origin for these bands in HITRAN2016: 20003-01101 – UCL *ab initio* [27] and 12202-01101 – UCL *ab initio* and CDSD2015 [26,27]. Also, the comparison of the ATM and CDSD2019 line intensities showed a deviation up to 30% for the 20003- 01101 band of the ¹²C¹⁶O₂ isotopologue.

Fig. 6. The comparison of line intensities between the published measurements from Refs [36,37] and the different CO₂ databases [12,17,23] for the 11102-00001 bands of the ¹²C¹⁶O₂, ¹³C¹⁶O₂, and ¹⁶O¹²C¹⁸O isotopologues with centers: 1933 cm⁻¹, 1897 cm⁻¹, and 1903 cm⁻¹, respectively.

The TCCON observations considered here cover the 3950 to 9500 cm^{-1} spectral region. These measurements are made with a Fourier transform spectrometer with an InGaAs detector and optical path difference (OPD) of 45 cm, corresponding to a resolution of 0.02 cm^{-1} . Use of the HITRAN2016 line lists reduces the CO_2 retrieved from the 6220 and 6338 cm^{-1} windows by 0.5% and 1.5% respectively, by comparison to results obtained with the other line lists, thus introducing a new inconsistency of 1.5%. TCCON CO_2 windows are circled in **Fig.7**. This issue is associated with the 30012 - 00001 and 30013 - 00001 line parameters and is discussed in detail in Section 4.

Fig.7. Retrieved VMR Scale Factors (VSF) obtained by averaging single-spectrum based on fitting each spectral window from the TCCON spectra. Results are plotted versus wavenumber. See the designations of the databases in the caption of Fig. 2.

The comparison of the fits to 136 Kitt Peak lab spectra with those modeled using the HITRAN2016 CO_2 line list shows large residuals in the 3500 and 4800 cm^{-1} spectral windows due to inaccuracies of line positions of the 10012-00001, 20012-00001, and 20013-00001 bands of the $^{13}\text{C}^{16}\text{O}_2$ isotopologue. The comparison showed that the HITRAN2012 [44] line positions for the 20012-00001 and 20013-00001 bands led to better residuals than those of the HITRAN2016 and CDSD-296 [17] databases. As a result, the line positions of the 20012 - 00001 and 20013 - 00001 bands of the $^{13}\text{C}^{16}\text{O}_2$ isotopologue were replaced with the corresponding HITRAN2012 line positions into the HITRAN2020 CO_2 line list. In the case of the 10012-00001 band, we subtracted

0.0009 cm⁻¹ from the CDSD [17] line positions used for this band in HITRAN2020 to match the ATM values [15].

Also, inaccuracies in the positions of some ¹⁶O¹²C¹⁸O lines of the 01121-01101 hot band (P13e, P14e, P14f, Q12f, Q13e, Q13f, R11e, R12e, and R12f) were evidenced in the HITRAN2016 line list between 4600 and 4625 cm⁻¹. This issue was identified by comparison with fits to Kitt Peak spectra and was solved by replacing the line positions with positions from the CDSD-296 database [17] in the HITRAN2020 CO₂ line list.

As mentioned above, below 8000 cm⁻¹, most line intensities in the HITRAN2016 database came from the UCL-IAO line lists [27–29] which are based on *ab initio* calculations. Because of excessive uncertainty in the *ab initio* line positions, the HITRAN 2016 line list is, in general, a combination of the CDSD- based line positions [26] and the UCL variational line intensities [27–29]. In the case of the so-called "sensitive bands" (as defined by Zak et al. [27–29]) *ab initio* intensities were replaced with those from CDSD when available. This substitution resulted in some cases where transitions in a given band would have intensities from both *ab initio* calculations as well as those from CDSD. Although this approach was justified in many cases, a number of inconsistencies in the rotational distribution of intensities in HITRAN2016 were revealed in CRDS spectra of ¹⁸O and ¹³C enriched and natural CO₂ near 1.74 μm [39,40,46]. In these works, the anomalies involve the perpendicular bands of the Δ*P*= 9 series of transitions. As shown in **Fig 8**, the mixing of CDSD and UCL *ab initio* intensities for the 41104-00001 bands of ¹²C¹⁶O₂ and ¹³C¹⁶O₂ isotopologues leads to an apparent inconsistency in the rotational dependence with strong intensity variation between successive *J* values or even missing transitions. A similar situation was found in the case of the 41104-00001 band of ¹⁶O¹²C¹⁸O isotopologue (See **Fig. 9**). A comparison of the measurements from Refs [39,40,46] to various carbon dioxide spectroscopic databases is presented in **Fig. 10**. As can be seen from this figure, the AMES intensity values for this band are all close to the experimental ones. As a result, for HITRAN2020, the AMES intensities of the 41104-00001 bands of the ¹²C¹⁶O₂, ¹³C¹⁶O₂, and ¹⁶O¹²C¹⁸O isotopologues were preferred to the mixed intensity values in the HITRAN2016 line list.

Similar validation tests using the values available in the literature and from the different CO₂ databases were carried out for other bands affected by the problem due to the mixing of

intensities of CDSD and *ab initio* origin in the HITRAN line list. These bands were systematically searched (see Fig. 5 of Ref.[39]), and when problems were identified, alternative sources of data for each such bands were suggested. For the $^{12}\text{C}^{16}\text{O}_2$, $^{13}\text{C}^{16}\text{O}_2$, $^{16}\text{O}^{12}\text{C}^{18}\text{O}$, and $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ isotopologues, we present in Table 1 the set of bands that were checked and affected by this issue along with the alternative intensity sources for these bands used in HITRAN2020. The substitution of the suggested intensity sources for the bands presented in Table 1 was performed for the corresponding bands of all the CO_2 isotopologues in HITRAN2020 where experimental data exists. In the absence of experimental reference data, the replacement of the intensity source was performed only for the corresponding band of the specific isotopologue. In the case of the 31104e-01101e and 11112e-11101e hot bands of the $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ and $^{13}\text{C}^{16}\text{O}_2$ isotopologues, respectively (see **Fig.11**), we cannot offer an alternative source for replacing the mixed line intensities, since there is no experimental data to verify them. Nevertheless, we kept the HITRAN2016 line intensities for the 31114e-01101e band of $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ isotopologue in HITRAN2020. In the future we will consider using UCL *ab initio* [28] line intensities for this band. In the case of the 11112e-11101e hot band of the $^{13}\text{C}^{16}\text{O}_2$ isotopologue, the UCL and CDSD-296 databases [17,27] give similar line intensities although there is an inconsistency in the rotational dependence with substantial intensity variation between successive J values of the P branch. Validation of this band is a challenge for future experiments.

Fig. 8. Comparison of the CRDS measurements (purple squares) of the 41104 - 00001 band of $^{13}\text{C}^{16}\text{O}_2$ [46] and $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ [40] to various databases: HITRAN2016 [44], CDSD-296 (CDSD2015) [26], CDSD-296 (CDSD2019) [17], UCL [27,28], and AMES [23] line lists.

Fig.9. Overview comparison of the Q-branch of the 41104 - 00001 band of $^{12}\text{C}^{16}\text{O}_2$ [39] to the HITRAN2016[12], CDSD-296 (CDSD2015) [26], CDSD-296 (CDSD2019) [17], UCL [27] and AMES [23] line lists. The lower panel highlights the Q branch where the HITRAN2016 list uses UCL and CDSD2015 as intensity sources for J below 50 and above 52, respectively, leading to the observed intensity jump between $Q(50)$ and $Q(52)$. The CRDS measurements [39] validate the AMES intensity values which were taken as the unique source for HITRAN2020 of this band.

Table 1. The bands for the $^{12}\text{C}^{16}\text{O}_2$, $^{13}\text{C}^{16}\text{O}_2$, $^{16}\text{O}^{12}\text{C}^{18}\text{O}$, and $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ isotopologues where intensities were from mixed sources in HITRAN2016 and the new sources of intensities chosen for HITRAN2020.

Band	Band center (cm ⁻¹)	Intensity origin		Experimental works
		HITRAN2016	HITRAN2020	
¹² C ¹⁶ O ₂ isotopologue				
21103-02201	1846.33	[26,27]	[26,27]	-
13302-02201	1907.85	[26,27]	[27]	-
20002-01101	2003.76	[26,27]	[17]	[41]
21102-10002	2054.72	[26,27]	[27,28]	-
22201-11101	2120.50	[26,27]	[17]	[42]
40002-21103	2295.65	[26,27]	[27]	-
30004-01101	3125.10	[26,27]	[17]	[34]
31104-10002	3131.52	[26,27]	[27]	-
22203-01101	3156.20	[26,27]	[27]	-
31103-10002	3306.48	[26,27]	[27]	-
22202-01101	3342.10	[26,27]	[27]	-
23302-02201	3344.01	[26,27]	[27]	-
31102-10001	3365.31	[26,27]	[17]	[34]
30002-01101	3398.45	[26,27]	[27]	-
21101-00001	3501.45	[26,27]	[27]	-
40002-11102	3544.64	[26,27]	[27]	-
01121-10002	4031.07	[26,27]	[27]	-
31104-00001	4416.15	[26,27]	[17]	[35]
41103-10001	4613.12	[26,27]	[27]	-
25501-02201	4696.61	[26,27]	[27]	-
10021-01101	5346.31	[26,27]	[17]	[47]
50006-01101	5572.65	[26,27]	[27]	-
50005-01101	5768.12	[26,27]	[27]	[39]
42204-01101	5805.64	[26,27]	[17]	[39]
41104-00001	5830.79	[26,27]	[23]	[39]
50004-01101	5920.94	[26,27]	[27]	-
50003-01101	6057.89	[26,27]	[27]	[48]

$^{13}\text{C}^{16}\text{O}_2$ isotopologue				
04421-01101	1946.12	[26,29]	[27,28]	-
11112-11101	2110.13	[26,27]	[27]	-
$^{16}\text{O}^{12}\text{C}^{18}\text{O}$ isotopologue				
31104-30004	616.12	[26,28]	[17,28]	-
21101-10002	2195.21	[27,28]	[28]	-
21101-00001	3454.63	[27,28]	[28]	-
31104-01101	3677.44	[26,28]	[28]	-
$^{16}\text{O}^{13}\text{C}^{18}\text{O}$ isotopologue				
12202-01101	1856.75	[26,28]	[28]	-

416

417

418

419

420

421

422

423

424

425

426

427

428

429

Fig.10. Ratios of the measured line intensities of the 41104 - 00001 band of the $^{12}\text{C}^{16}\text{O}_2$, $^{13}\text{C}^{16}\text{O}_2$ and $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ [39,40,46] isotopologues to the HITRAN2016 [12], CDSD-296 (CDSD2019) [17], UCL [27,28] and AMES [23] line lists.

Fig. 11. The 31104e-01101e and 11112e-11101e hot bands of ¹⁶O¹²C¹⁸O (628) and ¹³C¹⁶O₂ (636) isotopologues, respectively, having the mixed line intensities in the HITRAN2016 database [12].

4. New experimental data with sub-percent uncertainty

Accurate line parameters of the spectral lines are generally required for the most demanding atmospheric applications. In particular, the accurate values of the line intensities with uncertainties as low as 0.3%-0.5% [49] are necessary for forward models used in retrievals of CO₂ concentrations from some remote sensing missions. A number of very accurate measurements have become available after the release of HITRAN2016 in the NIR region: the most recent CRDS measurements from NIST [19,20] and FTS measurements from DLR reported in Birk et al. [22] (with the corresponding measurements and line parameter database provided on Zenodo [21]). The results obtained in these works were used to improve the HITRAN line intensities of the ¹²C¹⁶O₂ isotopologue.

The accurate CRDS line intensity measurements for the 3001*i*–00001 (*i* = 2-4) bands reported by Long et al. [19] were used to refine the calculated HITRAN2016 CO₂ band intensities near 1.6 μm. This region is actively used for spectroscopic measurements of atmospheric CO₂ concentrations. For instance, the 30013–00001 band is targeted by the OCO-2 and OCO-3 missions [5,7], while the 30012–00001 band is used for the LIDAR missions [50]. It was shown in Ref. [19] that the band-integrated CRDS intensity measurements and the *ab initio* calculations of Zak et al. [27] agree at the 0.06% level for the 30013–00001 and 30014–00001 bands, but there is a systematic discrepancy of about 1.1% for the 30012–00001 band. We note that the HITRAN2016 line intensities are from the *ab initio* UCL results [27], and they have relative uncertainties < 2 % for the 30012 - 00001 and 30013 - 00001 bands and < 5 % for the 30014 - 00001 band, respectively. In contrast, the current CO₂ spectroscopic databases [17,23] show a significant spread in their line intensities at about or above 1%. Following these results [19], the HITRAN2016 line intensities for the 3001*i*–00001 (*i*=2-4) bands were scaled to the recommended band-dependent scaling constant factors, while preserving the original *J*-dependence of the *ab initio* calculations. The comparison of the HITRAN2016 and HITRAN2020 line intensities of the 30012-00001 band of ¹²C¹⁶O₂ isotopologue to the experimental values reported in Long et al. [19] is plotted in **Fig.12**.

In Fleurbaey et al. [20], it was shown that a constant scaling of 1.0069±0.0002 of HITRAN2016 values in the 20013–00001 band of CO₂ near 2.06 μm is consistent with the

experiment. Therefore, we have performed this scaling in HITRAN2020. In Refs [19,20], the reported relative uncertainty in intensity is better than 0.1%, thus the uncertainty code 8 (< 1%) for the line intensities was given for the corresponding bands in the HITRAN2020 line list.

Bruker IFS 125HR Fourier-transform spectrometer measurements have been conducted at the German Aerospace Center (DLR) to measure pure carbon dioxide transmittance spectra in the 6000–7000 cm⁻¹ spectral region, including the 30011 - 00001, 30012 - 00001, 30013 - 00001, 30014 - 00001, and 00031 - 00001 bands [22]. In this work, line intensity uncertainties of 0.15% were reported. The scaling factor 1.0061 was used for the HITRAN2016 line intensities of the 30011 - 00001 band of the ¹²C¹⁶O₂ isotopologue to match with DLR measurements [22]. The 30013 - 00001 and 30014 - 00001 HITRAN2016 line intensities and NIST measurements were found to be in good agreement with the DLR measurements. However, for the 30012 - 00001 band, the differences outside of the stated uncertainties with both NIST measurements and HITRAN2016 were reported. The difference for the 30012-00001 band was 1.6%, whereas for Long et al. [19] the difference was about 1%. It should be noted that these differences are outside the stated uncertainty budget. In addition, the DLR data showed systematic differences of relative line intensities to HITRAN2016 within the 3001*i*-00001 (*i*=2,3) bands on the order of 0.3%.

In the HITRAN2016 edition, the line intensities of the 00031–00001 band of the ¹²C¹⁶O₂ isotopologue near 1.4 μm used CDSD-296 [26] as an intensity source because this band was identified as "sensitive" in the *ab initio* calculations [27]. The comparison of the CDSD line intensities with the DLR measurements [20] showed rotationally-dependent deviations up to 4%. To improve the 00031–00001 line intensities in the HITRAN2020 line list, we scaled the UCL line intensities [27] by the factor of 1.1217 to match the line intensities reported by Birk et al. [22] (See **Fig. 12**). It should be noted that two lines (P20 and R18) were absent in the UCL line list [27], so that the intensities of these corresponding lines were updated from Birk et al. [22].

Fig. 12. Upper panel: Comparison of the HITRAN2016 and HITRAN2020 line intensities of the 30012-00001 band of $^{12}\text{C}^{16}\text{O}_2$ isotopologue to the experimental values measured by Long et al. [19]. Lower panel: Comparison of the measured line intensities reported in Birk et al. [22] and the UCL intensities scaled by the factor of 1.1217 for the 00031-00001 band of the $^{12}\text{C}^{16}\text{O}_2$ isotopologue.

The weak 10032 - 10002 and 01131 - 01101 hot bands of the $^{12}\text{C}^{16}\text{O}_2$ isotopologue located near 6900 cm^{-1} were also compared to the DLR measurements [22]. It was found that the HITRAN2016 line intensities of the 10032 - 10002 band should be scaled by a factor of 1.1346 while the line intensities of the 01131 - 01101 band having the intensity origin from CDSD-296K [26] should be scaled by a factor of 1.0022.

5. New bands above 8000 cm^{-1}

The recent high-temperature ExoMol UCL-4000 line list containing almost 2.5×10^9 transitions at $T=4000\text{K}$ for the $^{12}\text{C}^{16}\text{O}_2$ isotopologue was published in Ref. [18]. To compare this line list with HITRAN2016, we converted UCL-4000 to HITRAN format at 296 K and applied an intensity cut-off of $10^{-30}\text{ cm/molecule}$ using the ExoMol_to_HITRAN.py program downloaded from <http://exomol.com/software/>[51]. This code transfers the position, intensity, Einstein A, lower state energy, and upper/lower level statistical weights into the correct location for a HITRAN default format ".par" file. It works by looking at the "trans" file to get the upper and lower state ID. We edited this program to extract the m_1, m_2, l_2, m_3 HITRAN quantum numbers from the ExoMol files. In the case where the states were not assigned, the vibrational assignments were replaced with "-2-2-2-20". When it was possible, we also performed additional assignments using the CDSD-296 [17] and Ames [23] databases. **Fig. 13** gives an overview comparison of the HITRAN 2016 line list to the UCL-4000 line list for the $^{12}\text{C}^{16}\text{O}_2$ isotopologue from 0 to 20000 cm^{-1} . It should be noted that the $^{12}\text{C}^{16}\text{O}_2$ line list in HITRAN2016 was limited to the spectral range 158.302 to 14075.298 cm^{-1} . In **Fig.13**, only the new bands from the UCL-4000 line list, which are missing in the HITRAN spectroscopic database, are presented. They are highlighted in blue. Most of these bands are located above 8000 cm^{-1} . Over 3600 spectral lines with line intensities from 1.8×10^{-28} to $9.9 \times 10^{-30}\text{ cm/molecule}$ from the UCL-4000 line list were included in the HITRAN2020 spectroscopic database. The line positions in the UCL-4000 dataset [18] are calculated from the energy levels derived from the HITRAN2016 database. Due to diversity of data in HITRAN some inconsistencies could occur when deriving upper state energy levels, which could result in the deviations. It is important to point out that the UCL-4000 line list has an issue regarding the matching of line positions. As an example, this case is shown in **Fig.**

14, where the comparison of the line positions for the 50015 - 00001 and 60014 - 10002 bands of the $^{12}\text{C}^{16}\text{O}_2$ isotopologue is presented. We traced the source of the discrepancies in UCL-4000 to the procedure used to substitute the variational (upper) state energies with the empirical values. The substituted energies were taken from HITRAN2016, either as the lower state energy term values E'' or as $E' = E'' + \nu$ (where ν is the transition wavenumber). The examples of these upper state energies, including those shown in **Fig. 14**, were affected by inconsistency of the original line positions from different sources. Most of such inconsistency cases identified in the present analysis have been improved in HITRAN2020 and will be propagated to UCL-4000 after the release of HITRAN 2020. In case of the small number of states (~ 20) that are still affected by small inconsistencies in HITRAN 2020's upper states, all for the transitions well above the 8000 cm^{-1} , the CDSD synthetic data [17] will be used.

Fig. 13. Overview comparison of the HITRAN2016 [12] and ExoMol UCL-4000 [18] carbon dioxide line lists in the 0-20000 cm^{-1} spectral region.

Fig. 14. Difference between the HITRAN2020 and UCL-4000 ExoMol line positions for the 50015 - 00001 and 60014 - 10002 bands of the main CO₂ isotopologue.

The weak 30022 - 00001 and 30023 - 00001 bands of the ¹⁶O¹²C¹⁸O isotopologue with band centres as 8374.23 cm⁻¹ and 8497.45 cm⁻¹, respectively, were missing in the HITRAN2016 [12] and CDSD-296 [17] spectroscopic databases. These bands were assigned in the analysis of CRDS spectra of natural CO₂ near 1.18 μm [14] and included in the new CO₂ line list. We also

calculated the energy levels (thereby obtaining line positions) for the 30022 - 00001 and 30023 - 00001 bands of $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ up to $J_{\text{max}}=34$ using the spectroscopic constants of the lower and upper state obtained in Refs [14,52] and equation (1).

$$F_v(J) = G_v + B_v J(J+1) - D_v J^2(J+1)^2 + H_v J^3(J+1)^3 \quad (1)$$

where G_v is the vibrational term value, B_v is the rotational constant, D_v and H_v are the centrifugal distortion constants, J is the angular momentum quantum number.

Also, it was shown in Ref. [14] that the R-branch intensities of the very weak 00041 - 01101 hot band of the $^{12}\text{C}^{16}\text{O}_2$ isotopologue, which are missing in CDSD-296 [17], are in good agreement with the NASA Ames intensities [23] while HITRAN2016 values were largely overestimated. For the HITRAN2020 database, the line intensities of the 30022-00001, 30023-00001, and 00041-01101 bands were updated using the line intensities from NASA Ames [23]. **Fig.15** shows the good agreement between the CRDS line intensities from Ref. [14] and the calculated values from the AMES line list [23]. The uncertainty codes for line positions and intensities were updated to 4 for these bands in the HITRAN2020 line list, corresponding to ($\geq 0.0001 \text{ cm}^{-1}$ and $< 0.001 \text{ cm}^{-1}$) and ($\geq 10 \%$ and $< 20 \%$), respectively.

Fig.15. Comparison CRDS [14] and AMES [23] line intensities of the 30022 - 00001 and 30023 - 00001 bands of the $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ isotopologue.

6. Addition of magnetic dipole band of $^{12}\text{C}^{16}\text{O}_2$ at 3.3 μm

The HITRAN2020 database has been extended by including the 01111 - 00001 ($\nu_2+\nu_3$) magnetic dipole band of the principal isotopologue of carbon dioxide. These line parameters were introduced into HITRAN for the first time; all previous editions of HITRAN provided only CO_2 electric dipole transitions. Interestingly, the first observation of the $\nu_2+\nu_3$ band has been reported at 3.3 μm in the atmosphere of Mars [53] by the ExoMars Trace Gas Orbiter ACS instrument [54]. This band is forbidden as an electric dipole absorption. However, it is allowed through electric quadrupole and the magnetic dipole mechanisms, which are typically much weaker than those

allowed through the electric dipole mechanism. The maximum line intensities in the 01111-00001 band are on the order of 6×10^{-28} cm/molecule. The detailed studies of this band providing the selection rules for the vibration-rotation transitions are presented in Refs [53,55]. The vibrational transition magnetic dipole moment of the 01111-00001 band was fitted to the line intensities measured with a Bruker IFS 125 HR FTS and a 30 m base multipass gas cell of the V.E.Zuev Institute of Atmospheric Optics SB RAS [56]. The line positions and intensities of this band were computed using the vibrational transition magnetic dipole moment and the set of the effective Hamiltonian parameters reported by Majcherova et al. [57]. We included the calculated line parameters up to $J=64$ of the 01111-00001 band of $^{12}\text{C}^{16}\text{O}_2$ in the HITRAN2020 line list. The calculated line intensities were consistent with an intensity cut-off 10^{-30} cm/molecule at 296K. The calculated line intensities agree well with the values measured independently by Optical-Feedback-Cavity Enhanced Absorption Spectroscopy (OFCEAS) in Fleurbaey et al. [58] for five R-branch lines of this band (R26-R32 and R36). An overview of the 01111-00001 magnetic dipole band of the main CO_2 isotopologue with band center 3004.012 cm^{-1} is displayed in **Fig. 16**. The corresponding uncertainty codes for the line positions and intensities were used for the 01111-00001 band of $^{12}\text{C}^{16}\text{O}_2$: code 4 (≥ 0.0001 and < 0.001) and code 4 ($\geq 10\%$ and $< 20\%$), respectively. It should be noted, that to distinguish the magnetic dipole transitions in the traditional HITRAN ".par" format, the letter "m" was introduced into the quantum field dedicated to upper state rotational ("local") quanta (see HITRAN2004 paper [59]).

We should also mention that in their OFCEAS study of the $\nu_2+\nu_3$ band, Fleurbaey et al. [58] reported together with the detection of the magnetic dipole lines, the measurement of weaker electric quadrupole of the same $\nu_2+\nu_3$ band. This detection was made possible by accurate *ab initio* predictions of the E2 line intensities $^{12}\text{C}^{16}\text{O}_2$ in the 0-10000 cm^{-1} range [60]. Overall, the *ab initio* intensities of the very weak quadrupole lines are validated by this OFCEAS values (maximum line intensities on the order of 10^{-28} cm/molecule). There is also evidence for these quadrupole lines in the ACS Mars spectrum [60]. The weak electric quadrupole lines are not included in the present version of the HITRAN database and will be considered for forthcoming editions.

Fig. 16. The 01111-00001 magnetic dipole band of the main CO₂ isotopologue [56,58]. The band center of this band is 3004.012 cm⁻¹.

7. The line-shape parameters for the HITRAN2020 CO₂ line list

A major update of the line-shape parameters of CO₂ broadened by air and CO₂ is described in Ref. [13]; this involved both enhancing the HITRAN database and improving completeness. A systematic extrapolation method was introduced for producing the air- and self- half width (broadening) Voigt profile parameters for unmeasured transitions [61], based on the measurements reported in Ref. [62]. The temperature-dependent exponents of the air- and self-broadening parameters were generated based on existing measurements and a semi-empirical calculation method using the measurements given in Refs [63,64]. Air- and self-pressure shifts for every line of CO₂ were calculated using the semi-empirical approach proposed by Hartmann [65] and fits using carefully selected experimental data.

Besides providing the standard 160-character ".par" parameter, the HITRAN2020 database will include additional parameters for the speed-dependent Voigt (SDV) profile [66,67]

in a separate set. The list of the SDV parameters was provided in Table 1 of Ref. [13]. The air- and self-speed dependence of the broadening parameters with their temperature dependence will be added for all the lines of CO₂, and they can be downloaded using customized output format on www.hitran.org. One can also retrieve these parameters with the HITRAN Application Programming Interface (HAPI) [68] using the SDVoigt parameters group.

The CO₂ line-mixing package, developed by Lamouroux et al. [69] and available at HITRAN*Online*, was updated using the new spectroscopic parameters obtained in the present study and the parameters found in Ref. [13]. The full and the first-order line-mixing parameters have been calculated with the latter being and the first-order line-mixing parameters will be provided for every line of CO₂ in HITRAN. The new data were verified using different sets of laboratory spectra to compare with the absorption coefficient calculated by the line-mixing package program. The uncertainties better than 0.5% were achieved for the regions examined compared to HITRAN2016 (we refer to figures 16-19 in Ref. [13]), which should be beneficial for CO₂ retrieval missions, including OCO-2, ACE, and GOSAT.

8. Overview summary of the HITRAN2020 CO₂ line list

The HITRAN2020 line list for the twelve stable isotopic species of carbon dioxide contains 545084 transitions. It covers the spectral range from 0.757 cm⁻¹ to 19908.186 cm⁻¹ with $J \leq 128$ and with the lower-state energies up to 6533.030 cm⁻¹. Most of the line positions were replaced with line positions from CDSD-296 [17]. For the problematic line intensities identified in new laboratory and atmospheric spectra, critical validation tests were performed to improve their accuracy of the CO₂ lists. The line-shape parameters of CO₂ broadened by air and CO₂ were updated. A summary of the CO₂ line list in the HITRAN2020 edition compared to the HITRAN2016 is given in **Table 2**. The HITRAN2020 carbon dioxide line list generated in HITRAN format is available on the HITRAN*Online* website www.hitran.org. and Supplementary material of this paper. An overview of the line lists in HITRAN2020 and HITRAN2016 is plotted in **Fig.17**.

Fig. 17. An overview of the HITRAN2016 and HITRAN2020 line lists for the 12 stable CO₂ isotopologues: ¹²C¹⁶O₂(626), ¹³C¹⁶O₂(636), ¹⁶O¹²C¹⁸O(628), ¹⁶O¹²C¹⁷O(627), ¹⁶O¹³C¹⁸O(638), ¹⁶O¹³C¹⁷O(637), ¹²C¹⁸O₂ (828), ¹⁸O¹²C¹⁷O(827), ¹²C¹⁷O₂(727), ¹³C¹⁸O₂(838), ¹⁸O¹³C¹⁷O(837), and ¹³C¹⁷O₂(737) (The numbers in parentheses are the AFGL shorthand code for the isotopologues).

673 **Table 2.** Comparison of HITRAN2020 and HITRAN2016 line lists for the 12 stable CO₂ isotopologues.

674

Formula	AFGL code	Abundance	HITRAN2016			HITRAN2020		
			Number of lines	Spectral region (cm ⁻¹)	Q(296K)	Number of lines	Spectral region (cm ⁻¹)	Q(296K)
¹² C ¹⁶ O ₂	626	0.984204	173024	158.302-14075.298	286.094	174446	158.302-19908.186	286.094
¹³ C ¹⁶ O ₂	636	0.011057	70577	332.649-13734.963	576.644	69870	332.649-13734.963	576.644
¹⁶ O ¹² C ¹⁸ O	628	0.003947	127850	1.473-12677.181	607.713	122140	1.473-12677.182	607.713
¹⁶ O ¹² C ¹⁷ O	627	7.339890×10 ⁻⁴	77941	0.757-12726.562	3542.610	73942	0.757-12726.562	3542.610
¹⁶ O ¹³ C ¹⁸ O	638	4.434460×10 ⁻⁵	43782	2.945-9212.609	1225.270	41058	2.945-9212.608	1225.270
¹⁶ O ¹³ C ¹⁷ O	637	8.246230×10 ⁻⁶	25175	9.086-8061.741	7140.024	23607	9.086-8061.739	7140.024
¹² C ¹⁸ O ₂	828	3.957340×10 ⁻⁶	10522	482.813-8162.743	323.424	10498	482.814-8162.752	323.424
¹⁷ O ¹² C ¹⁸ O	728	1.471800×10 ⁻⁶	15878	491.181-8193.172	3766.044	14623	498.617-8193.172	3766.044
¹² C ¹⁷ O ₂	727	1.368470×10 ⁻⁷	6518	535.384-6932.693	10971.91	6493	535.384-6932.693	10971.91
¹³ C ¹⁸ O ₂	838	4.446000×10 ⁻⁸	2916	539.626-6686.983	652.242	2926	539.620-6686.983	652.242
¹⁷ O ¹³ C ¹⁸ O	738	1.653540×10 ⁻⁸	4190	549.473-4914.496	7593.900	3980	549.473-4914.496	7593.900
¹³ C ¹⁷ O ₂	737	1.537500×10 ⁻⁹	1501	575.853-3614.084	22129.96	1501	575.853-3614.084	22129.96
Total			559874			545084		

675

676 **Note:** AFGL code is the shorthand notation for the isotopologue, abundance is the terrestrial value assumed by HITRAN, and Q(296
677 K) is the partition sum at the reference temperature of 296 K.

678

9. Conclusion

This work presents an improved and extended version of the HITRAN2020 spectroscopic database for the carbon dioxide molecule. The database includes updates of the line positions and intensities described in this work, and line shape parameters (described in Hashemi et al. [13]) for 12 stable CO₂ isotopologues. Critical validation tests for the spectroscopic data, including the comparisons with the most advanced theoretical and semi-empirical databases [17,18,23,27] and accurate experimental measurements, were carried out. Evaluation of the HITRAN CO₂ line lists by comparison to laboratory and atmospheric spectra below 8310 cm⁻¹ and the description of the updates made for the bands having mixed CDS and UCL-IAO line intensities in the HITRAN2016 line list below 8000 cm⁻¹ are presented. Recent experimental measurements with sub-percent uncertainty [19–22] were used to improve the CO₂ bands in the (1.4–2.1) μm region. Several new CO₂ bands (more than 3600 transitions) above 8000 cm⁻¹ were added to the HITRAN2020 edition from the new high-temperature UCL-4000 ¹²C¹⁶O₂ line list from the ExoMol [18] database. Also, the updated CO₂ line list has been extended by including new experimentally observed bands [14] corresponding to the ¹⁶O¹²C¹⁸O isotopologue. The magnetic dipole 01111-00001 band of the ¹²C¹⁶O₂ isotopologue in the 3.3 μm region was introduced into HITRAN for the first time. Updated self- and air-broadened line shape parameters of CO₂, as described in Ref. [13] and involved in the enhancement of the HITRAN databases, were also revised. These updates of the CO₂ line parameters in HITRAN2020 are expected to have an important impact on the capabilities of current and future remote-sensing missions. The updated line lists for 12 CO₂ isotopologues are available as Supplementary material of this paper or can be downloaded from the HITRAN website (www.hitran.org).

Acknowledgments

This work was supported by NASA grant funding from AURA NNX17AI78G. The National Institute of Standards and Technology (NIST) received support from the NASA Science Team for the OCO Missions (NRA) NNH17ZDA001N-OCO2 and the NIST Greenhouse Gas and Climate Science Measurements Program. This work was also supported by the Ministry of Science and Higher Education of the Russian Federation. The work performed at UCL was supported by the STFC Projects No. ST/M001334/1 and ST/R000476/1. Part of this work was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

References

- [1] Butz A, Guerlet S, Hasekamp O, Schepers D, Galli A, Aben I, et al. Toward accurate CO₂ and CH₄ observations from GOSAT. *Geophysical Research Letters* 2011;38:2–7. <https://doi.org/10.1029/2011GL047888>.
- [2] Alexe M, Bergamaschi P, Segers A, Detmers R, Butz A, Hasekamp O, et al. Inverse modelling of CH₄ emissions for 2010–2011 using different satellite retrieval products from GOSAT and SCIAMACHY. *Atmospheric Chemistry and Physics* 2015;15:113–33. <https://doi.org/10.5194/acp-15-113-2015>.
- [3] Yokota T, Yoshida Y, Eguchi N, Ota Y, Tanaka T, Watanabe H, et al. Global Concentrations of CO₂ and CH₄ Retrieved from GOSAT : First Preliminary Results. *Scientific Online Letters on the Atmosphere* 2009;5:160–3. <https://doi.org/10.2151/sola.2009-041>.
- [4] Nakajima, M., Suto, H., Yotsumoto, K., Abe, M., Kuze, A., Shiomi, K. et al. Overview of the GOSAT-2 Mission, *Proceedings of the 29th ISTS Nagoya-Aichi, Japan* 2013:60.
- [5] Fabiano Oyafuso, Vivienne H. Payne, Brian J. Drouin, V. Malathy Devi, D. Chris Benner. High accuracy absorption coefficients for the Orbiting Carbon Observatory-2 (OCO-2) mission: Validation of updated carbon dioxide cross-sections using atmospheric spectra. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2017;203:213–23. <https://doi.org/10.1016/j.jqsrt.2017.06.012>.
- [6] Eldering, A., Wennberg, P. O., Crisp, D., Schimel, D. S., Gunson, M. R., Chatterjee, A. et al. The Orbiting Carbon Observatory-2 early science investigations of regional carbon dioxide fluxes. *Science* 2017;358:188. <https://doi.org/10.1126/science.aam5745>.
- [7] Eldering A, Taylor TE, O'Dell CW, Pavlick R. The OCO-3 mission: Measurement objectives and expected performance based on 1 year of simulated data. *Atmospheric Measurement Techniques* 2019;12:2341–70. <https://doi.org/10.5194/amt-12-2341-2019>.
- [8] Fischer H, Birk M, Blom C, Carli B, Carlotti M, von Clarmann T, et al. MIPAS: An instrument for atmospheric and climate research. *Atmospheric Chemistry and Physics* 2008;8:2151–88. <https://doi.org/10.5194/acp-8-2151-2008>.
- [9] Bernath PF, McElroy CT, Abrams MC, Boone CD, Butler M, Camy-Peyret C, et al. Atmospheric chemistry experiment (ACE): Mission overview. *Geophysical Research Letters* 2005;32:1–5. <https://doi.org/10.1029/2005GL022386>.
- [10] Wunch D, Toon GC, Blavier JFL, Washenfelder RA, Notholt J, Connor BJ, et al. The total carbon column observing network. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 2011;369:2087–112. <https://doi.org/10.1098/rsta.2010.0240>.
- [11] Hase F. Improved instrumental line shape monitoring for the ground-based, high-resolution FTIR spectrometers of the Network for the Detection of Atmospheric Composition Change. *Atmospheric Measurement Techniques* 2012;5:603–10. <https://doi.org/10.5194/amt-5-603-2012>.
- [12] Gordon IE, Rothman LS, Hill C, Kochanov RV, Tan Y, Bernath PF, et al. The HITRAN2016 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2017;203:3–69. <https://doi.org/10.1016/j.jqsrt.2017.06.038>.

- [13] Hashemi R, Gordon IE, Tran H, Kochanov RV, Karlovets EV, Tan Y, Lamouroux J, et al. Revising the line-shape parameters for air- and self- broadened CO₂ lines toward a sub-percent accuracy level. *JQSRT* 2020;256:107283. <https://doi.org/https://doi.org/10.1016/j.jqsrt.2020.107283>.
- [14] Karlovets EV, Kassi S, Campargue A. Journal of Quantitative Spectroscopy & Radiative Transfer High sensitivity CRDS of CO₂ in the 1.18 μ m transparency window . Validation tests of current spectroscopic databases 2020;247:1–8. <https://doi.org/10.1016/j.jqsrt.2020.106942>.
- [15] Toon GC. CO₂ Spectroscopy Evaluation: 670 to 8310 cm⁻¹, Reports and Presentations for the HITRAN meeting, Jun 2020, Jet Propulsion Laboratory, California Institute of Technology (2020).
- [16] Toon GC. CO₂ Spectroscopy Evaluation: 670 to 7000 cm⁻¹ , Reports and Presentations for the ACE STM, Oct 2018, Jet Propulsion Laboratory, California Institute of Technology (2018).
- [17] Tashkun SA, Perevalov VI, Gamache RR, Lamouroux J. CDSD-296, high-resolution carbon dioxide spectroscopic databank: An update. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2019;228:124–31. <https://doi.org/10.1016/j.jqsrt.2019.03.001>.
- [18] Yurchenko SN, Mellor TM, Freedman RS, Tennyson J. ExoMol line lists-XXXIX. Rovibrational molecular line list for CO₂. *Monthly Notices of the Royal Astronomical Society* 2020;496:5282–91. <https://doi.org/10.1093/mnras/staa1874>.
- [19] Long DA, Reed ZD, Fleisher AJ, Mendonca J, Roche S, Hodges JT. High-Accuracy Near-Infrared Carbon Dioxide Intensity Measurements to Support Remote Sensing. *Geophysical Research Letters* 2020;47:e2019GL086344. <https://doi.org/10.1029/2019GL086344>.
- [20] Fleurbaey H, Yi H, Adkins EM, Fleisher AJ, Hodges JT. Cavity ring-down spectroscopy of CO₂ near λ = 2.06 μ m: Accurate transition intensities for the Orbiting Carbon Observatory-2 (OCO-2) “strong band.” *Journal of Quantitative Spectroscopy and Radiative Transfer* 2020;252:107104. <https://doi.org/10.1016/j.jqsrt.2020.107104>.
- [21] Birk M, Röske C, Wagner C. Measurement and line parameter database CO₂ 6000-7000 cm⁻¹, Zenodo 2021.
- [22] Birk M, Röske C, Wagner C. High accuracy CO₂ Fourier transform measurements in the range 6000–7000 cm⁻¹. *Journal of Quantitative Spectroscopy and Radiative Transfer* Submitted to HITRAN2020 Special Issue 2021.
- [23] Huang X, Richard S, Freedman B, Lee TJ. Ames-2016 line lists for 13 isotopologues of CO₂: Updates, consistency, and remaining issues. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2017;203:224–41. <https://doi.org/10.1016/j.jinf.2020.02.020>.
- [24] Gordon IE, Rothman LS, Hargreaves, Hashemi R, Karlovets EV, Skinner FM, Conway EK, Hill C, et al. The HITRAN2020 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer* Submitted to HITRAN2020 Special Issue 2021.
- [25] Hill C, Gordon IE, Kochanov RV, Barrett L, Wilzewski JS, Rothman LS. HITRANonline: An online interface and the flexible representation of spectroscopic data in the HITRAN database. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2016;177:4–14. <https://doi.org/10.1016/j.jqsrt.2015.12.012>.

- [26] Tashkun SA, Perevalov VI, Gamache RR, Lamouroux J. CDSD-296, high resolution carbon dioxide spectroscopic databank: Version for atmospheric applications. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2015;152:45–73. <https://doi.org/10.1016/j.jqsrt.2014.10.017>.
- [27] Zak E, Tennyson J, Polyansky OL, Lodi L, Zobov NF, Tashkun SA, et al. A room temperature CO₂ line list with ab initio computed intensities. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2016;177:31–42. <https://doi.org/10.1016/j.jqsrt.2015.12.022>.
- [28] Zak EJ, Tennyson J, Polyansky OL, Lodi L, Zobov NF, Tashkun SA, et al. Room temperature linelists for CO₂ asymmetric isotopologues with ab initio computed intensities. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2017;203:265–81. <https://doi.org/10.1016/j.jqsrt.2017.01.037>.
- [29] Zak EJ, Tennyson J, Polyansky OL, Lodi L, Zobov NF, Tashkun SA, et al. Room temperature linelists for CO₂ symmetric isotopologues with ab initio computed intensities. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2017;189:267–80. <https://doi.org/10.1016/j.jqsrt.2015.12.022>.
- [30] Teffo JL, Sulakshina ON, Perevalov VI. Effective Hamiltonian for rovibrational energies and line intensities of carbon dioxide. *Journal of Molecular Spectroscopy* 1992;156:48–64. [https://doi.org/10.1016/0022-2852\(92\)90092-3](https://doi.org/10.1016/0022-2852(92)90092-3).
- [31] Teffo JL, Lyulin OM, Perevalov VI, Lobodenko EI. Application of the effective operator approach to the calculation of ¹²C¹⁶O₂ line intensities. *Journal of Molecular Spectroscopy* 1998;187:28–41. <https://doi.org/10.1006/jmsp.1997.7455>.
- [32] Polyansky OL, Bielska K, Ghysels M, Lodi L, Zobov NF, Hodges JT, et al. High-Accuracy CO₂ Line Intensities Determined from Theory and Experiment. *Physical Review Letters* 2015;114:1–5. <https://doi.org/10.1103/PhysRevLett.114.243001>.
- [33] Yurchenko SN, Thiel W, Jensen P. Theoretical ROVibrational Energies (TROVE): A robust numerical approach to the calculation of rovibrational energies for polyatomic molecules. *Journal of Molecular Spectroscopy* 2007;245:126–40. <https://doi.org/10.1016/j.jms.2007.07.009>.
- [34] Chris Benner, V. Malathy Devi, Curtis P. Rinsland and Penelope S. Ferry-Leeper. Absolute intensities of CO₂ lines in the 3140–3410-cm⁻¹ spectral region. *Applied Optics* 1988;27:1588–97. <https://doi.org/10.1364/AO.27.001588>.
- [35] Toth RA, Brown LR, Miller CE, Devi VM, Benner DC. Spectroscopic database of CO₂ line parameters: 4300–7000 cm⁻¹. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2008;109:906–21. <https://doi.org/10.1016/j.jqsrt.2007.12.004>.
- [36] Tomoaki Tanaka, Masashi Fukabori, Takafumi Sugita, Hideaki Nakajima et al. Spectral line parameters for CO₂ bands in the 4.8- to 5.3-μm region. *Journal of Molecular Spectroscopy* 2006;239:1–10. <https://doi.org/10.1016/j.jms.2006.05.013>.
- [37] Rinsland CP, Benner DC, Devi VM. Measurements of absolute line intensities in carbon dioxide bands near 5.2 μm. *Applied Optics* 1985;24:1644. <https://doi.org/10.1364/ao.24.001644>.
- [38] Karlovets EV, Kassi S, Tashkun SA, Perevalov VI, Campargue A. High sensitivity Cavity Ring Down spectroscopy of carbon dioxide in the 1.19-1.26μm region. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2014;144:137-153. <https://doi.org/10.1016/j.jqsrt.2014.04.001>.

- [39] Čermák P, Karlovets EV, Mondelain D, Kassi S, Perevalov VI, Campargue A. High sensitivity CRDS of CO₂ in the 1.74 μm transparency window. A validation test for the spectroscopic databases. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2018;207:95–103. <https://doi.org/10.1016/j.jqsrt.2017.12.018>.
- [40] Karlovets EV, Čermák P, Mondelain D, Kassi S, Campargue A, Tashkun SA, et al. Analysis and theoretical modeling of the ¹⁸O enriched carbon dioxide spectrum by CRDS near 1.74 μm. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2018;217:73–85. <https://doi.org/10.1016/j.jqsrt.2018.05.017>.
- [41] Rinsland CP, Benner DC. Absolute intensities of spectral lines in carbon dioxide bands near 2050 cm⁻¹. *Applied Optics* 1984;23:4523–8. <https://doi.org/https://doi.org/10.1364/AO.23.004523>.
- [42] Rinsland CP, Benner DC and Devi VM. Absolute line intensities in CO₂ bands near 4.8 μm. *Applied Optics* 1986;25:1204–14. <https://doi.org/https://doi.org/10.1364/AO.25.001204>.
- [43] Rothman LS, Gordon IE, Barbe A, Benner DC, Bernath PF, Birk M, et al. The HITRAN 2008 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2009;110:533–72. <https://doi.org/10.1016/j.jqsrt.2009.02.013>.
- [44] Rothman LS, Gordon IE, Babikov Y, Barbe A, Chris Benner D, Bernath PF, et al. The HITRAN2012 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2013;130:4–50. <https://doi.org/10.1016/j.jqsrt.2013.07.002>.
- [45] Wunch D, Toon GC, Blavier JFL, Washenfelter RA, Notholt J, Connor BJ, et al. GFIT. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 2011;369:2087–112. <https://doi.org/10.1098/rsta.2010.0240>.
- [46] Karlovets EV, Sidorenko AD, Mondelain D, Kassi S, Campargue A, Sidorenko AD, et al. The ¹³CO₂ absorption spectrum by CRDS near 1.74 μm. *Journal of Molecular Spectroscopy* 2018;354:54–9. <https://doi.org/10.1016/j.jms.2018.10.003>.
- [47] Lawrence P.Giver, Linda R.Brown, Charles Chackerian, Richard S. Freedman. The rovibrational intensities of five absorption bands of ¹²C¹⁶O₂ between 5218 and 5349 cm⁻¹. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2003;78:417–36. [https://doi.org/10.1016/S0022-4073\(02\)00277-7](https://doi.org/10.1016/S0022-4073(02)00277-7).
- [48] Perevalov BV, Kassi S, Romanini D, Perevalov VI, Tashkun SA, Campargue A. CW-cavity ringdown spectroscopy of carbon dioxide isotopologues near 1.5 μm. *Journal of Molecular Spectroscopy* 2006;238:241–55. <https://doi.org/10.1016/j.jms.2006.05.009>.
- [49] Miller CE, Crisp D, DeCola PL, Olsen SC, Randerson JT, Michalak AM, et al. Precision requirements for space-based XCO₂ data. *Journal of Geophysical Research Atmospheres* 2007;112:1–19. <https://doi.org/10.1029/2006JD007659>.
- [50] Campbell JF, Lin B, Dobler J, Pal S, Davis K, Erxleben W, McGregor D, O'Dell C, Bell E, et al. Field Evaluation of Column CO₂ Retrievals From Intensity-Modulated Continuous-Wave Differential Absorption Lidar Measurements During the ACT-America Campaign. *Earth and Space Science* 2020;7:e2019EA000847. <https://doi.org/10.1029/2019EA000847>.
- [51] Tennyson J, Yurchenko SN, Al-Refaie AF, Clark VHJ, Chubb KL, Conway EK, Dewan A, Gorman MN, Hill C, et al. The 2020 release of the ExoMol database: molecular line lists for exoplanet and other hot atmospheres. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2020; 255, 107228. <https://doi.org/10.1016/j.jqsrt.2020.107228>.

- [52] Rothman LS, Hawkins RL, Wattson RB, Gamache RR. Energy levels, intensities, and linewidths of atmospheric carbon dioxide bands. *Journal of Quantitative Spectroscopy and Radiative Transfer* 1992;48:537–66. [https://doi.org/10.1016/0022-4073\(92\)90119-O](https://doi.org/10.1016/0022-4073(92)90119-O).
- [53] Trokhimovskiy A, Perevalov V, Korablev O, Fedorova A, Olsen KS, Bertaux JL, et al. First observation of the magnetic dipole CO₂ main isotopologue absorption band at 3.3 μm in the atmosphere of Mars by the ExoMars Trace Gas Orbiter ACS instrument. *Astronomy & Astrophysics* 2020;639:1–7. <https://doi.org/10.1051/0004-6361/202038134>.
- [54] Korablev O, Montmessin F, Trokhimovskiy A, Fedorova AA, Shakun AV, Grigoriev AV, et al. The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter. *Space Science Reviews* 2018;214:7. <https://doi.org/10.1007/s11214-017-0437-6>.
- [55] Perevalov VI, Trokhimovskiy AY, Lukashevskaya AA, Korablev OI, et al. Magnetic dipole and electric quadrupole absorption in carbon dioxide. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2021;259:107408. <https://doi.org/10.1016/j.jqsrt.2020.107408>.
- [56] Borkov YG, Solodov AM, Solodov AA, Perevalov VI. Line intensities of the 01111–00001 magnetic dipole absorption band of ¹²C¹⁶O₂: Laboratory measurements. *Journal of Molecular Spectroscopy* 2021;376:111418. <https://doi.org/https://doi.org/10.1016/j.jms.2021.111418>.
- [57] Majcherova Z, MacKo P, Romanini D, Perevalov VI, Tashkun SA, Teffo JL, et al. High-sensitivity CW-cavity ringdown spectroscopy of ¹²CO₂ near 1.5 μm. *Journal of Molecular Spectroscopy* 2005;230:1–21. <https://doi.org/10.1016/j.jms.2004.09.011>.
- [58] Fleurbaey H, Grilli R, Mondelain D, Kassi S, Yachmenev A, Yurchenko SN, Campargue A. Electric-quadrupole and magnetic-dipole contributions to the ν₂+ν₃ band of carbon dioxide near 3.3 μm. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2021;266:107558. <https://doi.org/10.1016/j.jqsrt.2021.107558>.
- [59] Rothman LS, Jacquemart D, Barbe A, Benner DC, Birk M, Brown LR, et al. The HITRAN 2004 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2005;96:139–204. <https://doi.org/10.1016/j.jqsrt.2004.10.008>.
- [60] Yachmenev A, Küpper J, Campargue A, Yurchenko SN, and Tennyson. Quadrupole transitions in carbon dioxide. *J Chem Physics*, 2021 (in press).
- [61] Voigt W. Über das Gesetz Intensitätsverteilung innerhalb der Linien eines Gasspektrums. München; Berlin: Sitzber. Bayr Akad., 1912, 603 p.
- [62] Hashemi R, Rozario H, Ibrahim A, Predoi-Cross A. Line shape study of the carbon dioxide laser band. *Can J Phys* 2013;91(11):924–36. <https://doi.org/10.1139/cjp-2013-0051>.
- [63] Ma H, Sun M, Zha S, Liu Q, Cao Z, Huang Y et al. Temperature dependence of line parameters of CO₂ near 2.004 μm studied by tunable diode laser spectroscopy. *Chin Phys B* 2018;27(2):023301. <https://doi.org/10.1088/1674-1056/27/2/023301>.
- [64] Predoi-Cross A, Liu W, Murphy R, Povey C, Gamache R, Laraia A et al. Measurement and computations for temperature dependences of self-broadened carbon dioxide transitions in the 30012-00001 and 30013-00001 bands. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2010;111(9):1065–79. <https://doi.org/10.1016/j.jqsrt.2010.01.003>.

- [65] Hartmann J-M. A simple empirical model for the collisional spectral shift of air-broadened CO₂ lines. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2009;110(18):2019–26. <https://doi.org/10.1016/j.jqsrt.2009.05.016>.
- [66] Pickett HM. Effects of velocity averaging on the shapes of absorption lines. *J Chem Phys* 1980;73(12):6090–4. <https://doi.org/10.1063/1.440145>.
- [67] Pine AS. Line shape asymmetries in Ar-broadened HF ($\nu = 10$) in the Dicke arrowing regime. *J Chem Phys* 1994;101(5):3444–52. <https://doi.org/10.1063/1.467529>.
- [68] Kochanov RV, Gordon IE, Rothman LS, Wcisło P, Hill C, Wilzewski JS. HITRAN Application Programming Interface (HAPI): A comprehensive approach to working with spectroscopic data. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2016;177:15–30. <https://doi.org/10.1016/j.jqsrt.2016.03.005>.
- [69] Lamouroux J, Régalia L, Thomas X, Auwera JV, Gamache RR, Rothman LS, Gordon IE, Hartmann J-M. CO₂ Line-mixing database and software update and its tests in the 2.1 μm and 4.3 μm regions. *Journal of Quantitative Spectroscopy and Radiative Transfer* 2015;151:88–96. <https://doi.org/10.1016/j.jqsrt.2014.09.017>.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 11.

Fig. 12.

Fig. 13.

Fig. 14.

Fig. 15.

Fig. 16.

Fig. 17.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author Statement

E.V.Karlovets: Formal analysis, Methodology, Software, Writing - original draft., I.E.Gordon: Supervision, Conceptualization, Methodology, Writing - review & editing, L.S.Rothman: Writing - review & editing, R. Hashemi: Data curation, Writing - review & editing, R.J.Hargreaves: Software, G.C. Toon: Data curation, Software, A.Campargue: Data curation, Writing - review & editing, V.I. Perevalov: Data curation, Writing - review & editing, P. Čermák: Software, M.Birk: Data curation, G.Wagner: Data curation, J.T.Hodges: Data curation, Writing - review & editing, J. Tennyson: Writing - review & editing, S.N.Yurchenko: Data curation, Writing - review & editing.

Click here to access/download
Supplementary Material
Supplementary material.txt