1 **Sensitivity to cadmium of the endangered freshwater pearl mussel *Margaritifera***
2 ***margaritifera* from the Dronne river (France): experimental exposure**

3
4 Magalie Baudrimont, Patrice Gonzalez, Nathalie Mesmer-Dudons, Alexia Legeay

5
6 University of Bordeaux, UMR CNRS EPOC 5805, F-33120 Arcachon, France

7
8 **Corresponding author:**

9 Magalie Baudrimont

10 University of Bordeaux, UMR CNRS EPOC 5805, Aquatic Ecotoxicology team, Place du Dr Peyneau, F-
11 33120 Arcachon, France

12 Phone + 33(0)5 56 22 39 27

13 magalie.baudrimont@u-bordeaux.fr
14

15 **Abstract**

16 *Margaritifera margaritifera* is a critically endangered species in Europe. Among the causes
17 explaining its decline, metal pollution had never been deeply studied. Thus, an ecotoxicological
18 investigation was developed on this species which comes from the Dronne river (South-West of
19 France). Cadmium (Cd) exposure of mussels at 2 and 5 µg/L for 7 days was conducted to test their
20 vulnerability to this metal, but also the potential endocrine disruption power of Cd. Morphometric
21 analyses, gonad histological observations, metal bioaccumulation, metallothionein (MTs) production,
22 measures of malondialdehyde (MDA) and finally quantitative relative expression analysis of genes
23 involved in various metabolic functions were performed.

24 The main results showed Cd accumulation increasing in a dose dependent manner, especially
25 in the gills. The same trend was observed for gene expression relative to oxidative stress. Histological
26 analysis of the gonads highlighted a predominance of hermaphrodite individuals, but after 7d of
27 exposure to Cd, the percentage of female was largely increased compared to controls, from 17 to
28 33%. These results demonstrate the endocrine disruption effect of Cd on freshwater pearl mussels.

29
30
31 The pearl mussel *Margaritifera margaritifera* is sensitive to cadmium since the metallothioneins are
32 poorly induced, gene expression reveals oxidative stress and gonads tend to be feminized.

33
34 **Key words**

35 *Margaritifera margaritifera*, cadmium, metallothionein, gene expression, endocrine disruption,
36 gonad histology

1. INTRODUCTION

Indigenous freshwater mollusks are threatened by extinction in most parts of the world (Vaughn and Taylor, 1999; Strayer et al., 2004), and because little is known about their physiology and behavior, conservation efforts are difficult (Rotchell and Ostrander, 2003; Geist, 2010). Among them, the freshwater pearl mussel *Margaritifera margaritifera* was in the past the most common mussel type in Europe (Bauer, 1988). But populations have declined by more than 90% and only a small number of recruiting (functional) populations remain (Cosgrove and Hastie, 2001; Ziuganov, 2005; Hastie et al, 2008).

Several factors can be advanced to explain their disappearance during the XIXth and XXth centuries: pearl fishing, general deterioration of water quality, reduced host fish density, reduction of the habitats (Gumpinger et al., 2015), filling of sediments (Hauer, 2015), pollution by pesticides, phosphates and nitrogen responsible for eutrophication (Scheder et al., 2015), siltation and impoundments (Addy et al., 2012). Indeed, this species has particular ecological requirements (see appendix 1 in Varandas et al., 2013). The quality of the water must be excellent and the substrate healthy in particular for the survival of the juvenile mussels which are sensitive to the changes of their surrounding environment (Geist and Kuehn, 2005). For adults, the top of the valves (the oldest part of the mussel pearl-bearing) is the most damaged part and can prove to be fragile with the eventual chemical or physical attacks of water (Bensettiti & Gaudillat, 2004; Motte, 2005). Moreover, this mussel has a very particular life cycle since it has a very long lifespan (150 years on average in Europe) and requires the presence of a fish host. This life cycle is divided in four stages of development: glochidia (larval stage), parasitic stage, juvenile and adult stage. The sexes are normally separated, but females can become hermaphrodite when isolated and under stress (Bauer, 1987). First of all, the male releases sperm (June/July), which are trained by the current and then recovered by the female through its filtration system, after producing eggs. Fertilization produces a cell that evolves into a larva called glochidia. These larvae are incubated for four to six weeks in a part of the mussel mantle called marsupium. Then the glochidia are released into the current and, to continue their development, must be fixed in the week on the branchial system of a salmonid with a preference for brown trout (*Salmo trutta f. fario*) in central European populations (Wächtler et al., 2001). This favorable parasitic part of the cycle constitutes both a phase of larval development, but also a phase of dissemination of the species with the displacement of the host fish (Treasurer et al, 2006). Once fixed, glochidia parasitize their host for a period of several months. When they leave the gills of the host, the young mussels measure about half a millimeter and have become true bivalves. They then bury themselves in the sediment for a period of four to five years in order to continue

1 their growth and they progressively rise to position at the sediment-water interface. The fish host
2 can also be affected by water quality and a lack in ecological continuity, which participate in the
3 disappearance of the freshwater pearl mussel. Among pollutants introduced in aquatic systems by
4 anthropogenic activities, metals could weaken populations of *M margaritifera* and jeopardize its
5 survival.

6 Currently, this species is only little represented in the rivers of France. Approximately
7 100,000 individuals are listed. The department of the Dordogne (South-West of France) shelters two
8 of these populations, on the Bandiat and the Dronne rivers. The population of the Dronne is
9 estimated at 15,000 individuals. Until now, the catchment area of the Dronne was relatively free of
10 pollution. However, some observations made by agents of the AFB (Agence Française de la
11 Biodiversité) or PNRPL (Parc Naturel Régional Périgord-Limousin) question a possible deterioration of
12 the river quality.

13 In this catchment area, there are several potential sources of impacts (agriculture, release of
14 suspended matter, artificial ponds which generate eutrophication,...). Among them, a source of
15 specific pollution was listed in the commune of Saint Saud Lacoussière: an illegal waste dump which
16 had received all kinds of waste for many years, including for instance batteries and drugs. This
17 discharge is located on a slope on which streams the resurgence and rainwater. The site was
18 destabilized by work and trace metals or chemicals can be brought out by scrubbing in the Dronne
19 river downstream. However at this place the river shelters a population of approximately 1,000 to
20 1,500 individuals of freshwater pearl mussels. An exceptional authorization (Agreement N°60/2008)
21 delivered by the French ministry of the environment allowed us to collect several individuals in order
22 to conduct an experiment and to evaluate if this discharge may have an impact on mussels. Recently,
23 a study using some of these collected mussels demonstrated the presence of metals (As, Cd, Co, Cr,
24 Ni and Zn) accumulated in tissues of *M. margaritifera* located downstream from this dump and
25 impacting their gene expressions (Bertucci et al, 2017).

26 In fact, filter-feeders such as bivalves are known to accumulate metals and other pollutants
27 in their tissues (Couillard et al, 1993, 1995; Baudrimont et al., 1999; Giguère et al, 2003; Marie et al.,
28 2006a,b; Paul-Pont et al, 2010; Arini et al., 2014). After accumulation, metals can generate toxic
29 effects, like in particular the production of oxidative stress which can poison the animal until death
30 (Cossu et al, 2000; Gonzalez et al., 2006; Farcy et al, 2011; Gillis, 2012; Gillis et al, 2014). Among
31 these metals, cadmium (Cd) is considered as a priority substance in Europe (European Water
32 Framework Directive 2008/105/EC) as it is a known toxic metal that accumulates and exerts its toxic
33 action primarily in the kidneys (nephrotoxicity) and liver (hepatotoxic) in aquatic organisms, in
34 conditions of chronic exposure. Cd is well known to provoke cellular or physiological dysfunctioning
35 in varying degrees according to the species (Viarengo, 1989; Perceval et al. 2004). It is responsible for

a disturbance of the calcium metabolism in the freshwater pearl mussel (Frank and Gerstmann, 2007), because of its ability to compete with Ca^{2+} for membrane carriers of this trace element, due to an atomic radius very close to the latter (0.97 and 0.99 Å). Cd is one of the main factor correlated with genes transcription levels, with effects on translation, apoptosis, immune response, response to stimulus, and transport pathways in *M. margaritifera* (Bertucci et al., 2017). Cd has also been described as an endocrine disruptor in fish (Pierron et al, 2008; Pierron et al, 2009), and as described in *Margaritifera margaritifera*, a gene of estrogen receptor was present (Bertucci et al., 2017) which can ask the question of this mode of action in the pearl mussel too. Among the possible lines of defense, the synthesis of small proteins such as the metallothioneins (MTs) can be induced under the influence of metals (Baudrimont et al., 2003; Amiard et al., 2006). These cytosolic proteins are able to sequester the metal ions and thus to neutralize their potential toxic impacts on other cellular targets. They are in the same way able to act directly against the generation of oxidative stress by protecting the cell from the presence of free radicals (Yang et al., 2009). This defense mechanism can be measured as well at the protein as at the gene level in tissues of aquatic organisms (Marie et al., 2006a,b; Paul-Pont et al., 2010). Among the effects of oxidative stress, the oxidative degradation of fatty acids (lipid peroxidation) at the level of the cellular membranes leads to the formation of an end product, malondialdehyde (MDA) which accumulates in the cell, and that we can easily quantify in tissues, as a lipidic marker of peroxidation (Legeay et al., 2005). Metals exert other toxic effects, like the disturbance of the mitochondrial metabolism, the apoptosis (programmed cellular death), or the attacks of the DNA. All these effects can be followed by quantitative real time RT-PCR of specific gene mRNA implied in all these functions (Gonzalez et al., 2006; Paul-Pont et al., 2010).

The objectives of this work were thus to study the potential impacts of cadmium on several individuals of the freshwater pearl mussel *Margaritifera margaritifera* collected around the dumping site during the reproductive period. Owing to the fact that Cd can be a potential endocrine disruptor, in parallel we exposed freshwater pearl mussels to estradiol as a positive control to compare its effect with Cd. The synthesis of detoxification proteins such as MTs was quantified, in parallel with MDA concentrations and relative gene expressions by quantitative PCR. Some histological preparations were also carried out in order to observe the gonads of the freshwater pearl mussels during Cd exposure in controlled conditions.

2. MATERIALS AND METHODS

2.1. Sampling of the freshwater pearl mussels in the Dronne river

Adult *Margaritifera margaritifera* sampling was carried out during July 2009, upstream and downstream from the illegal waste dump of Saint Saud Lacoussière (44°33.019; 44°51.008), following

1 a request for authorization of sampling which was given by the DIREN (French Ministry of the
2 Environment, Agreement N°39/2016) in 2008.

3 4 2.2. Experimental study of Cd exposure during the reproductive period

5
6 Thirty individuals of *M. margaritifera* were used to conduct the exposure of mussels to Cd under
7 controlled conditions. Pearl mussels were 89 ± 7 mm long (ranging from 70 to 99 mm) and their
8 mean age was estimated at 39 years old according to Bertucci et al., 2017. Five experimental
9 conditions were performed, with 6 individuals per experimental unit chosen in order to have the
10 same average size in each of them: « Upstream control, UC », corresponding to individuals collected
11 upstream from the wild waste dump; “Downstream control, DC”, corresponding to individuals
12 collected downstream from the wild waste dump ; Cd exposure in a direct way with 2 or 5 µg/L with
13 individuals coming from upstream of the wild waste dump; and estradiol exposure with 100 µg/L
14 with individuals coming from upstream of the wild waste dump, in order to get a positive control of
15 endocrine disruption, and notably of reproductive hormones (Flynn and Spellman, 2009). The
16 contamination exposure lasted 7 days, after an acclimatization period of 14 days. Note that just
17 before the beginning of the experiment, one individual died in the DC experimental unit,
18 corresponding to only 3% of mortality for the entire experiment.

19 An experimental glass unit (volume = 40 L) consisted of 7 cm of ultrapure sand SILAQ (0,7-1,4 mm)
20 which was rinsed beforehand and deposited on the bottom of the unit previously protected with a
21 food standard plastic bag (Plastiluz). Each unit was filled using water of the Dronne river taken
22 upstream from the dump and was placed at 19°C, near to the temperature measured in the river at
23 this period, using a system of thermic regulation. The photoperiod was controlled with 16h/8h
24 light/dark and oxygenation was ensured thanks to a system of ventilation by air pumps. Water was
25 permanently filtered in order to avoid the intoxication of the freshwater pearl mussels by their own
26 waste and the mussels were fed for the period of acclimatization using phytoplanktonic green algae
27 *Scenedesmus subspicatus*. Cd was added in the experimental units every day using a stock solution
28 of CdCl₂. The concentration was measured in water samples collected every morning and adjusted
29 daily by the compensated additions method, in order to maintain it constant for all the duration of
30 the experiment. Temperature, pH and oxygen concentrations were daily measured in the
31 experimental units and remained constant ($T^{\circ}\text{C} = 18.76 \pm 0.13$; $\text{pH} = 7.2 \pm 0.05$; $\text{O}_2 = 9.11 \pm 0.03$
32 mg/L).

33 34 2.3. Cd bioaccumulation in tissues of *Margaritifera margaritifera*

Cd bioaccumulation in 3 tissues of mussels (gills, visceral mass and rest of the body) was analyzed by atomic absorption spectrophotometry (AAS) with graphite furnace (M6 solar AA spectrometer, Thermoptec). After dissection, weighing and drying at 50°C during 48h00, samples were analyzed. Tissues were first digested for 3h at 100°C after addition of nitric acid 63% (3mL for 100 mg of dry tissue) in polypropylene tubes. Samples were diluted 6 fold with pure water (15 mL) and stored at 4°C until analysis. For each mineralization procedure, 6 certified samples (50 mg of Tort-2 - Lobster Hepatopancreas Certified Reference Material for Trace Metals - and 50 mg of Dolt-3 - Dogfish Liver Certified Reference Material for Trace Metals -, NRC-CNRC) and 3 blanks were analyzed in order to test the validity of the method. The values obtained were systematically within the expected range. Results are expressed in µg of metal per gram of dry tissue. The detection limit is 0.1 µg/L for Cd.

2.4. Histological analyses

In order to determine precisely the sex of individuals, histological analyses were carried out on all the individuals used for the experiment (6 individuals per condition). Samples of gonads not exceeding 5 mm thickness were delicately cut using a scalpel, in a standardized manner. The fixing of tissues was carried out immediately after sampling by immersion of samples in liquid of Bouin (picric acid 0.9% + formol 7.4% + acetic acid 5%) for one minimal period of 48 H at 4°C. After several rinsings, the samples were dehydrated by successive alcohol baths of increasing degree. Complete dehydration was obtained by immersion of the samples during one day in a pure butanol bath renewed every 2 h. The samples were then placed in three successive paraffin baths (56°C) at a rate of 15 min per bath. At the end of the last paraffin treatment, samples were left on the straw mattress at ambient temperature. After solidification, blocks of paraffin were cut out in ribbon of 10 µM thickness using a microtome (Reichert). Before coloring, the paraffin was eliminated by immersion of the blades in two baths of xylene for 5 min. The cuts were then rehydrated. Gonads were colored according to the Hematoxylin/Eosin method described by Gabe (1968). Determination of the stage of evolution: Stage 0: sexual rest; Stage 1: proliferation of the gonies; Stage F: female sexual maturity; Stage MF: male and female sexual maturity.

2.5. Metallothioneins (MTs) quantification

Aliquots of tissues of *M. margaritifera* stored beforehand at -80°C under nitrogen atmosphere were analyzed by the method of saturation by inorganic mercury, coupled with Hg determination by cold vapor AAS (Baudrimont et al., 2003). After defrosting and drying on absorbing paper, the tissues aliquots were weighed then immersed in 500 µL to 1 mL of Tris 25mM (pH 7,2) buffer. Under nitrogen atmosphere, they were homogenized using an ultra-Turrax mechanical crusher (IKA T10). The homogenates were centrifuged to 20,000 g at 4°C during 1h, the supernatants were then

1 saturated with mercury (50 mg/L Hg (II) - TCA 10%). The mercury not fixed to MTs was recovered by
2 addition of porcine hemoglobin in excess. After centrifugation of 20 min to 20,000 g, the supernatant
3 containing only MTs fixing mercury was recovered, weighed, and 100 µL were analyzed by cold vapor
4 AAS under O₂ flow (LECO AMA 254, Altec, Prague, Czech Republic) in order to determine the
5 quantities of mercury contained in the samples. The detection limit of the device is 1 ng mercury. By
6 calculation, the number of nmoles of sites having fixed Hg per gram of fresh weight of sample was
7 deduced as an estimation of the MT concentration.

8

9 2.6. Malonedialdehyde (MDA) determination

10 In experiments, 200 µL of supernatant was mixed with 650 µL of solution of NMPI (N-methyl-2-
11 phenylindole, according to the LPO-586 method), the whole was homogenized using a vortex, then
12 150 µL of chlorhydric acid (HCl) 12N was added. The mixture was incubated 1h at 45°C then
13 centrifuged at 15,000 g at 4°C during 10 min. The supernatant was recovered and its absorbance
14 (corresponding to the presence of carbocyanine formed by the presence of a molecule of MDA and
15 two molecules of NMPI) was measured by spectrophotometry at 586 nm, thus giving the quantity of
16 MDA present in µmol per liter of supernatant. To be able to compare various samples, the
17 concentration of MDA was reported to the quantity of proteins (µmol per gram of protein; Legeay et
18 al., 2005).

19

20 2.7. Gene analyses

21

22 *2.7.1. Total RNAs extraction*

23 Total RNAs were extracted from 20 to 40 mg of gills (preserved beforehand in RNA-later and glycerol
24 at -80°C) by using the agilent kit "Absolutely RNA Miniprep Kit" following the manufacturer's
25 instructions.

26

27 *2.7.2. Reverse transcription of RNA into cDNA*

28 Reverse transcription was carried out with the "Stratascript first strand synthesis system" kit
29 (stratagene) using total RNA (3 to 5 µg). 1 µL of oligo-dT (1 µM), 1 µL of hexanucleotides (1 µM), 0.8
30 µL of dNTP (10 mM) and 2 µL of 10X activity buffer was added to the RNA. This mix spent 5 minutes
31 at 65°C in the thermocycler in order to linearize RNAs. Then, 1 µL of reverse transcriptase (1 U) and
32 0.5 µL of RNase-block (0.5 U) are added then heated at 42°C during 1h with the thermocycler to
33 allow reverse-transcription.

2.7.3. Real time quantitative PCR (qPCR)

The qPCR was carried out using the thermocycler LightCycler (Roche) and LightCyclerRun software 5.32. In this device, the follow-up in real time of the quantity of products of amplification is made possible by the addition of an intercalating agent: SybrGreen in the reactional medium. The reaction takes place in capillaries of glass of 20 µL prepared with: (I) 17 µL of a mix of reagent consisted 12.8 µL water, 3.2 µL of MgCl₂ (25 mM) and 1 µL of reactional mixture (Taq with 5 U/mL, dNTP, 10X buffer and SybrGreen); (II) 2 µL of a mix primers with 3 µM specific from each gene (Table 2) and (III) 1 µL of cDNA resulting from the RT. The protocol used is as follows: activation of the enzyme (95°C, 10 min), then 50 cycles of PCR (denaturation 95°C, 5s; fixing of the primers 60°C, 5s and elongation 72°C, 20s). Reaction specificity was determined for each reaction from the dissociation curve of the PCR product. This dissociation curve was obtained by following the SyberGreen fluorescence level during gradual heating of the PCR products from 60 to 95°C. Relative quantification of each gene expression level was normalized according to the β-actin gene using the 2^{-ΔCT} method as described by Livak and Schmittgen (2001).

2.9. Statistical treatment of data

The data processing was carried out using Statistica software (Statistica 12). After checking of the homogeneity of the variances, the factors “conditions” and “tissues” were tested by a two-way factor ANOVA or a one-way ANOVA for histological analyses. In the event of significance for one or the other of the factors taken into account, the mean comparison test (LSD Fisher) was carried out. In the event of nonconformity of the homogeneity of the variances, the non-parametric test of Kruskal Wallis was used in order to detect the significant differences between the sites or the tissues or conditions and this, for each parameter. The different letters mentioned on the graphs correspond to significant differences with the threshold α = 0.05.

3. RESULTS AND DISCUSSION

3.1. Cadmium bioaccumulation

When exposed during seven days to Cd, freshwater mussels showed significant accumulations in all the studied organs. In gills, Cd concentrations reached values of 12.4 and 17.2 µg/g (dw) under the conditions 2 and 5 µg/L respectively, compared to 1.1 µg/g for the upstream control individuals (Figure 1A). These accumulations corresponded to a direct route which involved exposure to the metal transferred from the dissolved phase of the water through the gill epithelia during filtration for respiratory and nutritional purposes. Cd bioaccumulation was also effective in the visceral mass and

the rest of the tissues, but less pronounced, attesting a blood transfer of Cd from the gills to the internal organs. Such concentrations were dependent on the exposure dose with highest values measured for the condition of 5 µg/L compared to 2 µg/L, especially in the gills and the rest of the body. In contrast, concentrations measured in the conditions "Downstream Control" and "Estradiol" showed no accumulation of this metal. The freshwater pearl mussel *M. margaritifera* therefore accumulated relatively large quantities of Cd, since in seven days, it multiplied its concentration in the gills by a factor of about 15. According to Frank and Gerstmann (2007) the freshwater pearl mussel indeed accumulated trace metals including Cd easily, with concentrations in Germany of about 50 µg/g (dw) in some contaminated sites in the visceral mass, compared to measures conducted in Finland, showing concentrations of this metal around 7 µg/g (dw) in these tissues. By comparison, here we measured values in control individuals acclimated for 14 days in the laboratory at around 2.5 µg/g (dw), which appears to be relatively low and consistent with measurements made *in situ* in the Dronne river upstream from the dump (Bertucci et al, 2017).

3.2. Metallothionein response

In freshwater pearl mussels *M. margaritifera*, concentrations of MTs measured in organs exposed to Cd 2 or 5 µg/L, showed a weak response to this metal in comparison to the condition "Upstream Control", which seemed very surprising given the bioaccumulation results of this metal for the species (Figure 1B). This means that unlike most species, the freshwater pearl mussel seemed to react very little in terms of detoxification by these proteins in the presence of Cd after a 7 days exposure period. This result could suggest a particular sensitivity of the freshwater pearl mussel to this metal, given the very strong bioaccumulation levels measured in this species in the field (Frank and Gerstmann, 2007).

However, a strong induction of these proteins was measured in the visceral mass of *M. margaritifera* downstream from the dump on the Dronne river. The induction of these proteins was at a factor of 3 compared to individuals from upstream, which was very high. Given the absence of Cd accumulation in these individuals downstream relative to those upstream, we were not able to explain this very strong induction which was probably related to the presence of other metals as observed by Bertucci et al (2017) *in situ*, or organic contaminants.

3.3. Malondialdehyde (MDA) quantification

MDA is a marker of oxidative degradation of membrane lipids (peroxidation) following the generation of oxidative stress. The accumulation of this molecule demonstrates a severe breach of physiological functioning at the cell level, which can lead to death. In our experiment under laboratory conditions, we did not observe any accumulation of this marker, whatever the conditions

studied (data not shown). This result is positive in the sense that it shows no severe toxic attack of Cd after 7 days of laboratory exposure on membrane lipids.

3.4. Gene expression

Five major genes were analyzed (Table 1). Whatever the organ, the expression of the main gene involved in mitochondrial metabolism (*cox1*) was higher in individuals from downstream compared to upstream individuals (Table 2). For the *mt* gene, the highest overexpression was noted in the kidney with a very significant induction factor (1085), followed by the gills where this gene was found 36 times more expressed than in individuals from upstream, and the visceral mass (15 fold higher). The *mt* gene expressions are consistent with MT synthesis we observed before, notably in the visceral mass. Some differences are on the contrary observed in the gills where MT proteins are not significantly produced compared to upstream despite a significant gene induction. These results can be explained by the delays between gene expression and MT production, which was already observed in mollusks (Paul-Pont et al, 2010). In gills, a higher expression of the *sodMn* gene (factor 6) was also demonstrated. The results in these two groups of individuals suggested different levels of basic expression. This suggests that animals from downstream were probably adapted to contamination, but the nature of which remains unclear. This contamination impacted the mitochondria and detoxification systems. The fact that the gills and kidney were the two most affected organs suggests contamination by both the direct and trophic routes. Indeed, gills constitute the first biological barrier crossed by contaminants present in the water. The kidneys in turn are the organs involved in the elimination and thus the detoxification of these compounds. The fact that *sodMn* was overexpressed in the gills of downstream individuals also strengthens this hypothesis. This overexpression reflects a greater generation of ROS in the mitochondria of this organ.

The expression levels observed in conditions of Cd contamination were compared with those obtained from control individuals from upstream. A dose-dependent response was determined in the three organs studied. For example, in the gills Cd caused overexpression of *cox1* and *mt* genes, but also a repression of cytoplasmic *sod*. The same trends were found in the kidney. However, once again, the factors of inductions raised for this organ showed the highest values; 22 compared to 18 for the *mt* gene and 181 compared to 19.5 for the *cox1* gene in the kidney and gills respectively after exposure to 5 µg/L. In the visceral mass, the same dose dependent overexpression of mitochondrial and detoxification genes was found. However, an induction of *sodMn* for a factor of 6 was also visible. These results were consistent with previous exposure to Cd studies in model organisms such as the zebrafish *Danio rerio* (Gonzalez et al, 2006; Orioux et al, 2011), the amphibian *Xenopus laevis* (Mouchet et al, 2006) or bivalves *Corbicula fluminea*, *Crassostrea gigas* or even *Cerastoderma edule*

(Baudrimont et al, 2003; Legeay et al, 2005; Marie et al, 2006a, b; Paul Pont et al, 2010). Thus, the production of defense proteins such as metallothioneins allows the sequestering of large quantities of metal ions at intracellular level in order to reduce the toxicity of these elements with respect to other cellular targets. These proteins are inducible by many other factors, such as cytotoxic agents, reproductive factors, the presence of parasites, bacterial infections, other xenobiotics etc ... (Kägi, 1991; Baudrimont et al, 1997; 2006), which leads to then being considered more as representative of a comprehensive response to stress (Amiard et al., 2006). Cd, once in the cell, is able to inhibit the transfer of electrons along the mitochondrial respiratory chain, but also to induce the production of reactive oxygen species (Wang et al 2004; Gonzalez et al, 2006). The oxidative stress thus generated is at the origin of DNA breaks and induction of apoptotic mechanisms leading to cell death (Bagchi et al 2000; Chan and Cheng 2003; Mouchet et al, 2006). The differential expression observed in *M. margaritifera* can account for these phenomena.

Unlike the Cd, the response to estradiol was fundamentally different depending on the organ. In the gills, a single increase of the 12S gene (factor 4), indicating a greater number of mitochondria and therefore an increasing energy demand, was visible. Instead, repression of this gene and of the *sod* gene was observed in the kidneys, associated with an overexpression of *coxI*. This suggests an inhibitory effect of estradiol on this organ. The increase of *coxI* surely compensates for this decrease in activity and thus maintains a sufficient level of ATP in the cells. These results are consistent with studies of estradiol and its metabolites which showed that these compounds can inhibit mitochondrial respiration (Hagen et al, 2004). In the kidney, estradiol caused in hamsters, through deletion and mutation mechanisms, a decline in the number of copies of mitochondrial DNA leading to a decrease in energy capacity of these organelles and an increase in the sensitivity of the kidney (Bhat, 2002). Thus, this type of damage would be involved in the onset and progression of cancers in this organ. In the liver, only *coxI* and *sodMn* genes were found induced after exposure to estradiol, suggesting a greater demand for energy and the generation of oxidative stress in mitochondria. In this context, the production of ROS has recently been observed in the metabolism of estradiol, especially in the liver (Spencer et al, 2012). This production can generate ROS damage to DNA.

3.5. Histological analyses

The histological analysis of gonads was performed on individuals from this experiment of exposure to estradiol and Cd to evaluate the potential endocrine disruptive effect of Cd, compared to the positive control condition represented by "Estradiol" condition (Table 3).

1 The results obtained allowed us to demonstrate for control individuals, whether from
2 upstream or downstream, a predominance of hermaphrodite individuals (0% of female individuals).
3 It is largely accepted that freshwater pearl mussels have separate sexes, however, in adverse
4 conditions (poor water quality, for example, or low population density), the female individuals can
5 develop hermaphroditism to maintain their reproduction (Bauer, 1987). Concerning the mussels of
6 the Dronne river, it seems that we are in this situation, which is positive in the sense that these
7 individuals are able to reproduce (mature gonads oocytes and sperm - Figure 2) contrary to what we
8 would have expected following the direct observation of the gonads (data not shown) showing a
9 predominant presence of female individuals. However, these results suggest a reaction or adaptation
10 of populations to their current location in the river, which can be judged in a rather unfavorable
11 manner. We can also add from this point of view that the estradiol and Cd (2 and 5µg/L) conditions
12 of exposure are less adverse conditions than downstream condition, which is also confirmed by gene
13 expressions (Table 2). Nevertheless, we must consider carefully these first results since only 11
14 individuals were analyzed (6 in the condition "Upstream Control" and only 5 in the condition
15 "Downstream Control").

16 For the results obtained from exposure to estradiol, the female sex hormone stimulating
17 oocyte development, we saw that after only 7 days' exposure, the percentage of strictly female
18 individuals was largely increased compared to control conditions (33% against 0%). We have
19 therefore a marked effect of this hormone on the oocyte development of *M. margaritifera*, as has
20 already been demonstrated by Flynn and Spellman (2009) in the freshwater bivalve *Elliptio*
21 *complanata*. Following exposure to Cd, we observed the same phenomenon, with a range effect of
22 this metal, since the percentage of female individuals increased from 17 to 33% for 2 and 5 µg/L
23 respectively. From this analysis we can observe that the Cd mimics the effect of estradiol on the
24 development of female sexual characteristics of the freshwater pearl mussel, which attests to an
25 endocrine disruptor effect in this bivalve. This is important given the effects already observed
26 previously in terms of cell and molecular impact on the tissues of the freshwater pearl mussel, and
27 facing significantly higher accumulations of Cd downstream relative to upstream from the waste
28 dump of St Saud Lacoussière on the Dronne river in April 2009. Measured Cd accumulation still
29 remained low compared to levels found in Germany or Finland, but may become a problem,
30 especially concerning endocrine disruption, because of other metals accumulated in parallel in the
31 field, or other potential contaminants, including organics that might amplify this phenomenon as
32 previously observed in freshwater mussels exposed to municipal effluents (Blaise et al, 2003; Gagné
33 et al, 2004).

34 35 **4. CONCLUSION**

1 This study features a novel approach in the potential toxicological impacts of trace metals,
2 and especially Cd, on the freshwater pearl mussel *Margaritifera margaritifera* from the Dronne river
3 in France. We measured metallothionein concentrations and malondialdehyde production for the
4 first time in this species and we quantified the relative expression of 5 genes implied in mitochondrial
5 metabolism, oxidative stress and detoxification. The results showed, after Cd exposure of the
6 freshwater pearl mussels in the laboratory, significant ability of mussels to accumulate this metal
7 after only 7 days of exposure, but at the same time, the relative slow response of MTs (only at the
8 gene level but not at the protein level) suggesting a certain sensitivity of these mussels to metals. We
9 observed also that the majority of the mussels from the Dronne river seemed to be hermaphrodite,
10 probably in relation with the decrease of the water quality of the river or with the low density of the
11 population. But after exposure to Cd or estradiol, we observed an increase of the amount of female
12 individuals, confirming for this species the endocrine disruptor role of Cd. These results thus bring
13 new knowledge on the sensitivity of this species to metal contamination, which can help to improve
14 reintroduction strategies on the European scale, taking into account the presence of this type of
15 contamination in the field.

16 17 **Acknowledgements**

18 We want to acknowledge the PNRPL (Parc Naturel Régional Périgord Limousin) for their cooperative
19 assistance in the field and their interest for this study. We also thank Eric Goursolle, Julie Chevalier,
20 Bénédicte Pécassou and Romain Papin-Vincent for their participation in realization of the different
21 analyses.

22 23 **References**

- 24 Addy S, Cooksley SL and Sime I. 2012. Impacts of flow regulation on freshwater pearl mussel
25 (*Margaritifera margaritifera*) habitat in a Scottish montane river. *Science of the Total*
26 *Environment*, 432: 318-328.
- 27 Amiard JC, Amiard-Triquet C, Barka S, Pellerin J and Rainbow P.S. 2006. Metallothioneins in aquatic
28 invertebrates: Their role in metal detoxification and their use as biomarkers. *Aquatic*
29 *Toxicology*, 76: 160–202.
- 30 Arini A, Daffe G, Gonzalez P, Feurtet-Mazel A and Baudrimont M. 2014. Detoxification and genetic
31 recovery capacities of *Corbicula fluminea* after an industrial metal contamination (Cd and Zn):
32 a one-year depuration experiment. *Environmental Pollution*, 192: 74-82.
- 33 Bagchi D, Joshi SS, Bagchi M, Balmoori J, Benner EJ, Kuszynski CA and Stohs SJ. 2000. Cadmium- and
34 chromium-induced oxidative stress, DNA damage, and apoptotic cell death in cultured human
35 chronic myelogenous leukemic K562 cells, promyelocytic leukemic HL-60 cells, and normal

- human peripheral blood mononuclear cells. *Journal of Biochemistry and Molecular Toxicology*, 14: 33-41.
- Baudrimont M, Andrès S, Durrieu G and Boudou A. 2003. The key role of metallothioneins in the bivalve *Corbicula fluminea* during the depuration phase, after an *in situ* exposure to Cd and Zn. *Aquatic Toxicology*, 63: 89-102.
- Baudrimont M, Andrès S, Métivaud J, Lapaquellerie Y, Ribeyre F, Maillet N, Latouche C and Boudou A. 1999. Field transplantation of the freshwater bivalve *Corbicula fluminea* along a polymetallic contamination gradient (river Lot, France) - Part II: Metallothionein response to metal exposure. *Environmental Toxicology and Chemistry*, 18: 2472-2477.
- Baudrimont M, de Montaudouin X and Palvadeau A. 2006. Impact of digenean parasite infection on metallothionein synthesis by the cockle (*Cerastoderma edule*): multivariate field monitoring. *Marine Pollution Bulletin*, 52: 494-502.
- Baudrimont M, Lemaire-Gony S, Métivaud J, Ribeyre F and Boudou A. 1997. Seasonal variations of metallothionein concentrations in the Asiatic clam (*Corbicula fluminea*). *Comparative Biochemistry and Physiology*, 118C (3): 361-367.
- Bauer G. 1987. Reproductive strategy of the freshwater pearl mussel *Margaritifera margaritifera*. *Journal of Animal Ecology*, 56: 691-704.
- Bauer G. 1988. Threats to the freshwater pearl mussel *Margaritifera margaritifera* L. in central Europe. *Biological Conservation*, 45: 239-53.
- Bensettiti F and Gaudillat V. 2004. Cahiers d'habitats Natura 2000. Connaissance et gestion des habitats et des espèces d'intérêt communautaire. Tome 7. Espèces animales. La Documentation française, pp 318-321.
- Bertucci A, Pierron F, Thébaud J, Klopp C, Bellec J, Gonzalez P and Baudrimont M. 2017. Transcriptomic responses of the endangered freshwater mussel *Margaritifera margaritifera* to trace metal contamination in the Dronne River, France. *Environmental Science and Pollution Research*, DOI 10.1007/s22356-017-0294-6.
- Bhat HK. 2002. Depletion of mitochondrial DNA and enzyme in estrogen-induced hamster kidney tumors: a rodent model of hormonal carcinogenesis. *Journal of Biochemistry and Molecular Toxicology*, 16: 1-9.
- Blaise C, Gagné F, Salazar M, Salazar S, Trottier S and Hansen PD. 2003. Experimentally-induced feminisation of freshwater mussels after long term exposure to a municipal effluent. *Fresenius Environmental Bulletin*, 12:865- 870.
- Chan PK and Cheng SH. 2003 Cadmium-induced ectopic apoptosis in zebrafish embryos. *Archives of Toxicology*, 77: 69-79.

- 1 Cosgrove PJ and Hastie LC. 2001. Conservation of threatened freshwater pearl mussel populations:
2 river management, mussel translocation and conflict resolution. *Biological Conservation*, 99:
3 183–90.
- 4 Cossu C, Doyotte A, Babut M, Exinger A and Vasseur P. 2000. Antioxidant biomarkers in freshwater
5 bivalves, *Unio tumidus*, in response to different contamination profiles of aquatic sediments.
6 *Ecotoxicology and Environmental Safety*, 45, 106-121.
- 7 Couillard Y, Campbell PGC and Tessier A. 1993. Response of metallothionein concentrations in a
8 freshwater bivalve *Anodonta grandis* along an environmental cadmium gradient. *Limnology
9 and Oceanography*. 38:299-313.
- 10 Couillard Y, Campbell PGC, Pellerin-Massicotte J and Auclair JC. 1995. Field transplantation of a
11 freshwater bivalve, *Pyganodon grandis*, across a metal contamination gradient. II.
12 Metallothionein response to Cd and Zn exposure, evidence for cytotoxicity, and links to effects
13 at higher levels of biological organization. *Canadian Journal of Fisheries and Aquatic Sciences*,
14 52:703-715.
- 15 Farcy E, Gagné F, Martel L, Fortier M, Trépanier S, Brousseau P and Fournier M. 2011. Short-term
16 physiological effects of a xenobiotic mixture on the freshwater mussel *Elliptio complanata*
17 exposed to municipal effluents. *Environmental Research*, 111:1096-1106.
- 18 Flynn K and Spellman T. 2009. Environmental levels of atrazine decrease spatial aggregation in the
19 freshwater mussel, *Elliptio complanata*. *Ecotoxicology and Environmental Safety*, 72: 1228–
20 1233.
- 21 Frank H and Gerstmann S. 2007. Declining Populations of Freshwater Pearl Mussels (*Margaritifera
22 margaritifera*) are burdened with heavy metals and DDT/DDE, *Ambio*, 36(7): 571-4.
- 23 Gabe M.1968. Techniques histologiques. Edition Masson et Cie.
- 24 Gagné F, Blaise C and Hellou J. 2004. Endocrine disruption and health effects of caged mussels,
25 *Elliptio complanata* placed downstream from a primary-treated municipal effluent plume for 1
26 year. *Comparative Biochemistry and Physiology C Toxicology and Pharmacology*, 138 (1), 33-
27 44.
- 28 Geist J. 2010. Strategies for the conservation of endangered freshwater pearl mussels (*Margaritifera
29 margaritifera*; L.): a synthesis of conservation genetics and ecology. *Hydrobiologia*, 644: 69–88.
- 30 Geist J and Kuehn R. 2005. Genetic diversity and differentiation of central European freshwater pearl
31 mussel (*Margaritifera margaritifera* L.) populations: implications for conservation and
32 management. *Molecular Ecology*, 14(2): 425-439.
- 33 Giguère A, Couillard Y, Campbell PGC, Perceval O, Hare L, Pinel-Alloul B. 2003. Steady-state
34 distribution of metals among metallothionein and other cytosolic ligands and links to
35 cytotoxicity in bivalves living along a polymetallic gradient. *Aquatic Toxicology*, 64, 185-200.

- 1 Gillis PL. 2012. Impacts of urban runoff and municipal wastewater effluents on the health of the
2 freshwater mussel, *Lasmigona costata*. *Science of the Total Environment*, 431:348-356.
- 3 Gillis PL, Higgins SK, Jorge MB. 2014. Evidence of oxidative stress in wild freshwater mussels
4 (*Lasmigona costata*) exposed to urban-derived contaminants. *Ecotoxicology and*
5 *Environmental Safety*, 102, 62-69.
- 6 Gonzalez P, Baudrimont M, Boudou A and Bourdineaud JP. 2006. Comparative effects of direct
7 cadmium contamination on gene expression in gills, liver, skeletal muscles and brain of the
8 zebra fish (*Danio rerio*). *Biometals*, 19: 225-235.
- 9 Gumpinger C, Hauer C and Scheder C. 2015. The current status and future challenges for the
10 preservation and conservation of freshwater pearl mussel habitats. *Limnologica*, 50: 1-3.
- 11 Hagen T, D'Amico G, Quintero M, Palacios-Callender M, Hollis V, Lam F and Moncada S. 2004.
12 Inhibition of mitochondrial respiration by the anticancer agent 2-methoxyestradiol.
13 *Biochemistry and Biophysics Research Communication*.322: 923-929.
- 14 Hastie LC, Young MR, Boon PJ, Cosgrove PJ and Henniger B. 2008. Sizes, densities and age structures
15 of Scottish *Margaritifera margaritifera* (L.) populations. *Aquatic Conservation: Marine and*
16 *Freshwater Ecosystems*, 10: 229-247.
- 17 Hauer C. 2015. Review of hydro-morphological management criteria on a river basin scale for
18 preservation and restoration of freshwater pearl mussel habitats. *Limnologica*, 50: 40-53.
- 19 Kägi JHR. 1991. Overview of metallothionein. In: Abelson, J.N., Simon, M.I. (Eds.), *Methods in*
20 *Enzymology*. Academic Press, San Diego, CA, pp. 613-626.
- 21 Legeay A, Achard-Joris M, Baudrimont M, Massabuau JC and Bourdineaud JP. 2005. Impact of
22 cadmium contamination and oxygen levels on biochemical responses in the Asiatic clam
23 *Corbicula fluminea*. *Aquatic Toxicology*, 74: 242-253.
- 24 Livak KJ and Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
25 quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25(4): 402-408.
- 26 Marie V, Baudrimont M and Boudou A. 2006a. Cadmium and zinc bioaccumulation and
27 metallothionein response in two freshwater bivalves (*Corbicula fluminea* and *Dreissena*
28 *polymorpha*) transplanted along a polymetallic gradient. *Chemosphere*, 65: 609-617.
- 29 Marie V, Gonzalez P, Baudrimont M, Bourdineaud JP and Boudou A. 2006b. Metallothionein response
30 to cadmium and zinc exposures compared in two freshwater bivalves, *Dreissena polymorpha*
31 and *Corbicula fluminea*. *Biometals*, 19: 399-407.
- 32 Mouchet F, Baudrimont M, Gonzalez P, Cuenot Y, Bourdineaud JP, Boudou A and Gauthier L. 2006.
33 Genotoxic and stress inductive potential of cadmium in *Xenopus laevis* larvae. *Aquatic*
34 *Toxicology*, 78: 157-166.
- 35 Motte G. 2005. Biologie et statut de la moule perlière en Wallonie. Forêt wallonne No 74, pp 10-16.

- Orieux N, Cambier S, Gonzalez P, Morin B, Adam C, Garnier-Laplace J and Bourdineaud JP. 2011. Genotoxic damages in zebrafish submitted to a polymetallic gradient displayed by the Lot River (France). *Ecotoxicology and Environmental Safety*.74: 974-983.
- Paul-Pont I, Gonzalez P, Baudrimont M, Nili H and de Montaudouin X. 2010. Short-term metallothionein inductions in the edible cockle *Cerastoderma edule* after cadmium or mercury exposure: discrepancy between mRNA and protein responses. *Aquatic toxicology*, 97:260-267
- Perceval O, Couillard Y, Pinel-Alloul B, Giguère A and Campbell PGC. 2004. Metal-induced stress in bivalves living along a gradient of Cd contamination: relating sub-cellular metal distribution to population-level responses. *Aquatic Toxicology*, 69 : 327–345.
- Pierron F, Baudrimont M, Dufour S, Elie P, Bossy A, Baloché S, Mesmer-Dudons N, Gonzalez P, Bourdineaud J-P and Massabuau J-C. 2008. How cadmium could compromise the completion of the European eel's reproductive migration. *Environmental Science and Technology*, 42: 4607–4612.
- Pierron F, Baudrimont M, Dufour S, Elie P, Bossy A, Lucia M and Massabuau JC. 2009. Ovarian gene expression and effects of cadmium pre-exposure during artificial sexual maturation of the European eel (*Anguilla anguilla*). *BioMetals*, 22: 985-994.
- Rotchell JM and Ostrander GK. 2003. Molecular markers of endocrine disruption in aquatic organisms. *Journal of Toxicology and Environmental Health*, 6: 453–495.
- Scheder C, Lerchegger B, Flodl P, Csar D, Gumpinger C and Hauer C. 2015. River bed stability versus clogged interstitial: Depth-dependent accumulation of substances in freshwater pearl mussel (*Margaritifera margaritifera* L.) habitats in Austrian streams as a function of hydromorphological parameters. *Limnologica*, 50: 29-39.
- Spencer WA, Vadhanam MV, Jeyabalan J and Gupta RC. 2012. Oxidative DNA damage following microsome/Cu(II)-mediated activation of the estrogens, 17 β -estradiol, equilenin, and equilin: role of reactive oxygen species. *Chemistry Research and Toxicology*, 25: 305-314.
- Strayer DL, Downing JA, Haag WR, King TL, Layzer JB and Newton TJ. 2004. Changing perspectives on pearly mussels, North America's most imperiled animals. *Bioscience*, 54: 429–39.
- Treasurer JW, Hastie LC, Hunter D, Duncan F, Treasurer CM, 2006. Effects of *Margaritifera margaritifera* glochidial infection on performance of tank-reared Atlantic salmon (*Salmo salar*). *Aquaculture journal*, 256: 74-79.
- Varandas S, Lopes-Lima M, Teixeira A, Hinzmann M, Reis J, Cortes R, Machado J, Sousa R. 2013. Ecology of southern European pearl mussels (*Margaritifera margaritifera*): first record of two new populations on the rivers Terva and Beça (Portugal). *Aquatic Conserv: Mar. Freshw. Ecosyst*. 23: 374–389.

- 1 Vaughn CC and Taylor CM. 1999. Impoundments and the decline of freshwaternussels: a case study
2 of an extinction gradient. *Conservative Biology*, 13: 912–20.
- 3 Viarengo A. 1989. Heavy metals in marine invertebrates: mechanisms of regulation and toxicity at
4 the cellular level. *Aquatic Sciences*, 1: 295-317.
- 5 Wächtler K, Dreher-Mansur MC and Richter T. 2001. Larval types and early postlarval biology in
6 naiads (Unionoida). In: Bauer G, Wächter K (eds) Ecology and evolution of the freshwater
7 mussels Unionoidea. Ecological Studies, 145. Springer Verlag Heidelberg, pp. 93-125.
- 8 Wang Y, Fang J, Leonard SS and Rao KMK. 2004. Cadmium inhibits the electron transfer chain and
9 induces reactive oxygen species. *Free Radical Biology and Medicine*, 36: 1434-1443.
- 10 Yang HY, Wang YM and Peng SQ. 2009. Basal expression of metallothionein suppresses butenolide-
11 induced oxidative stress in liver homogenates in vitro. *Toxicol*, 53: 246-253.
- 12 Ziuganov VV. 2005. A long-lived parasite extending the host life span: the pearl mussel *Margaritifera*
13 *margaritifera* elongates host life by turns out the program of accelerated senescence in salmon
14 *Salmo salar*. *Dokl Biological Science*, 403:291-4.
- 15

Table 1: Nucleotidic sequences of the primer-pairs used for the qPCR of *Margaritifera margaritifera*.
a: sens primer, b: anti-sens primer

Function	Gene	Nucleotidic primer sequence (5'-3')
Reference gene	<i>β-actin</i>	TATCCACGCTCCGTGAGG ^a GTTGGGATTCAGGCTGTGTT ^b
Oxydative stress	<i>sod</i>	ATGTGCATGAATTTGGAGATAAACT ^a TTGGCCCTGTGAGGGT ^b
	<i>sodMn</i>	TGGGAGGTGCTCAGCC ^a GTAGTGTTTATACGCAGTCTGCC ^b
Mitochondrial metabolism	<i>12S</i>	CTTGCTCAGAGAAGACCCAACA ^a GGAGTGACGGGCGATT ^b
Detoxification	<i>mt</i>	AGATTTTGAAGCACCGAAGATGT ^a CTGTGCAGCCCTCACCA ^b

Table 2: Differential expression factors relative to control individuals (> 1 induced, < 1 repressed) in the organs of *Margaritifera margaritifera* (n = 6) after exposure (i) to the water of the Dronne river downstream an illegal waste dump, (ii) to Estradiol at 100 µg/L and (iii) to cadmium at 2 and 5 µg/L during 7 days.

	Downstream	Estradiol	Cd 2 µg/L	Cd 5 µg/L
Gills				
<i>sod</i>	0.51	0.15	0.34	0.12
<i>sodmt</i>	6.00	1.34	0.62	0.55
<i>12S</i>	1.76	3.85	0.56	7.63
<i>mt</i>	36.32	0.99	3.65	18.12
<i>coxI</i>	3.31	1.36	4.41	19.51
Digestive gland				
<i>sod</i>	0.47	0.72	0.71	0.64
<i>sodmt</i>	1.94	4.33	6.05	1.77
<i>12S</i>	16.88	2.26	0.96	0.64
<i>mt</i>	15.19	1.60	4.77	9.38
<i>coxI</i>	10.51	6.52	3.41	7.44
Kidney				
<i>sod</i>	0.51	0.04	0.81	0.48
<i>sodmt</i>	1.08	0.80	0.33	0.73
<i>12S</i>	13.67	0.17	0.82	1.37
<i>mt</i>	1085.04	0.57	6.81	22.48
<i>coxI</i>	3.31	2.61	35.25	181.20

- 1 Table 3: Sex determination of individuals of *Margaritifera margaritifera* according to histological
- 2 analyses. The percentage of females is significantly dependent of the condition (ANOVA, $p = 0.0277$).

Condition	Hermaphrodites	Females	Undetermined	Percentage of females	<i>n</i>
Upstream	6	0	0	0	6
Downstream	4	0	1	0	5
Estradiol	4	2	0	33	6
Cd 2 µg/L	5	1	0	17	6
Cd 5 µg/L	4	2	0	33	6

3

Figure 1: Cadmium bioaccumulation (A) and metallothionein concentrations (B) in tissues (G: gills, VM: visceral mass and R: rest of the tissues) of *Margaritifera margaritifera* for control individuals upstream (UC) and downstream (DC) the wild waste dump in the Dronne river, the exposure conditions with estradiol (100 µg/L) and Cd (2 and 5 µg/L) (mean ± SE, n = 6). The letters on the graph indicate significant differences between conditions.

- 1 Figure 2: Histological observations of the gonads of freshwater pearl mussels *Margaritifera*
2 *margaritifera* in function of the different exposure conditions : Upstream control, Estradiol, Cd 5µg/L.
3 H : hermaphrodite individual, F : female, O : oocytes, SPZ : spermatozooids.

- 4
- 5
- 6
- 7