Environmentally relevant mixture of S-metolachlor and its two metabolites affects thyroid metabolism in zebrafish embryos

Eliška Rozmánková, Marek Pípal, Lucie Bláhová, Naveen Chandran Njattuvetty, Bénédicte Morin, Patrice Gonzalez, Luděk Bláha

► To cite this version:

Eliška Rozmánková, Marek Pípal, Lucie Bláhová, Naveen Chandran Njattuvetty, Bénédicte Morin, et al.. Environmentally relevant mixture of S-metolachlor and its two metabolites affects thyroid metabolism in zebrafish embryos. *Aquatic Toxicology*, 2020, 221, pp.105444. 10.1016/j.aquatox.2020.105444 . hal-03434081

HAL Id: hal-03434081

<https://hal.science/hal-03434081>

Submitted on 25 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Environmentally relevant mixture of S-metolachlor and its two metabolites affects thyroid metabolism in zebrafish embryos

AUTHORS

Rozmankova Eliska^{1,2}, Pipal Marek¹, Blahova Lucie¹, Njattuvetty Chandran Naveen¹, Morin Benedicte², Gonzalez Patrice², Blaha Ludek¹

1 Masaryk University, Faculty of Science, RECETOX, Kamenice 753/5, 625 00 Brno, Czech Republic

2 University of Bordeaux, EPOC, UMR 5805, 33400 Talence, France

HIGHLIGHTS (3-5, max 85 characters per highlight)

- Herbicide toxicity analysis on embryo-larval stages of zebrafish
- Reduced spontaneous movements in embryos exposed to S-metolachlor alone and in mixture
- S-metolachlor caused no effect on molecular level, unlike its metabolites
- Low concentration of pesticide mixture overexpressed thyroid genes *dio2*, *thra*, *thrb*
- Larvae locomotion, heart rate not affected; minor developmental malformations found

KEYWORDS (6 max)

- Zebrafish embryo, embryotoxicity, pesticide mixture, environmental concentration, sublethal effect, metabolite

Abstract <400 words

Herbicides and their metabolites are found in water bodies where they may harm non-target organisms. Their effects at environmentally relevant concentrations are often unclear, especially concerning mixture of pesticides. This study thus investigated the impacts of one of the most used herbicides: S-metolachlor and its two metabolites, metolachlor oxanilic acid (MOA) and metolachlor ethanesulfonic acid (MESA) on the development of zebrafish embryos (*Danio rerio*). Embryos were exposed to the pure substances and their environmentally relevant mixture until 120 hpf (hours post-fertilization). The focus was set on sublethal endpoints such as malformations, hatching success, length of fish, spontaneous movements, heart rate and locomotion of zebrafish larvae. Moreover, gene expression levels of eight genes linked to the thyroid system disruption, oxidative stress defense, mitochondrial metabolism, regulation of the cell cycle and retinoic acid signaling pathway were analyzed.

35 Exposure to the S-metolachlor (1 µg/L) and to the pesticide mixture (1 µg/L of each
36 substance) significantly reduced spontaneous tail movements of 21 hpf embryos. Few rare
37 developmental malformations were observed, but only in larvae exposed to more than
38 100 µg/L of pure substances (craniofacial deformation, non-inflated gas bladder, yolk sac
39 malabsorption) and to 30 µg/L of each substance in the pesticide mixture (spine deformation).
40 However, no effect on hatching success, length of larvae, heart rate or larvae locomotion were
41 found. Strong responses were detected at the molecular level, especially on the *p53* gene
42 regulating the cell cycle, caused by the pesticide mixture (1 µg/L of each substance), MESA
43 (30 µg/L) and MOA (100 µg/L), as well as on the *cyp26a1* – gene encoding for cytochrome
44 P450 after exposure to the pesticide mixture (1 µg/L of each substance). Genes implicated in
45 the thyroid system disruption (*dio2*, *thra*, *thrb*) were all overexpressed following exposure to
46 the environmentally relevant concentrations of the pesticide mixture (1 µg/L of each
47 substance) and to the metabolite MESA (1 µg/L). In conclusion, the zebrafish thyroid system
48 disruption was revealed not only by the overexpressed genes, but also by some of the
49 developmental malformations considered as markers of this type of disruption (mainly gas
50 bladder and yolk sac abnormalities) and spontaneous tail movements linked to the T4
51 hormone influencing its neurodevelopment. Thus, the thyroid system disruption represents a
52 likely hypothesis behind the effects caused by the low environmental concentrations of
53 S-metolachlor, its two metabolites and their mixture.

Introduction

S-metolachlor is one of the most commonly used pesticides in the world (Atwood and Paisley-Jones, 2017). This selective pre-emergent herbicide from the chloroacetanilides family was first registered in 1997 (Heydens et al., 2010). It is an S-enantiomer of 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(1-methoxypropan-2-yl)acetamide. It replaced the racemic metolachlor mixture due to its higher herbicidal activity (Müller et al., 2001; Thomas Poiger, 2002), thus reducing the quantity of pesticide needed by 40% (Blaser and Spindler, 1997). S-metolachlor inhibits the biosynthesis of very-long-chain fatty acids (Götz and Böger, 2004) and interferes with gibberellin synthesis enzymes (Rose et al., 2016), thereby inhibiting the growth of target plants such as grass weeds in corn fields (Heydens et al., 2010). However, S-metolachlor can easily contaminate surface water by agricultural runoff (Zemolin et al., 2014) and it is thus often detected in surface or coastal waters (Accinelli et al., 2002; De Liguoro et al., 2014; Fauvelle et al., 2018; Glinski et al., 2018; Kapsi et al., 2019; Meffe and de Bustamante, 2014; Ryberg et al., 2014). In soil, S-metolachlor is microbially degraded to two principal metabolites (Zemolin et al., 2014): metolachlor oxanilic acid (MOA) and metolachlor ethanesulfonic acid (MESA), which are mobile and persistent in the environment (Sidoli et al., 2016). As a result, these degradation products are often detected in higher concentrations than the parent compound itself in water samples (Elliott and VanderMeulen, 2017; Hladik et al., 2005). However, there is a general lack of information about the toxicity of individual pesticide metabolites and their mixtures occurring in the environment. Only recently, some studies highlighted the importance of aquatic toxicity of environmentally relevant pesticide mixtures (Gustavsson et al., 2017; Tian et al., 2018). Furthermore, the importance of evaluating mixture toxicity is emphasized by the European Chemicals Agency (2014).

The contamination of aquatic environments by herbicides may adversely affect non-target organisms such as fish. Fish are among the most sensitive organisms in aquatic ecosystems and are used as standard testing species in the hazard assessment of chemicals and their mixtures (Hayes et al., 2019). To the best of our knowledge, studies on the effects of S-metolachlor or its metabolites on fish are scarce. Most use high concentrations (mg/L) to obtain LC_{50} or EC_{50} values, and these are far from being environmentally relevant (Dobšíková et al., 2011; Munn et al., 2006; Quintaneiro et al., 2017). Unlike these studies, Jin et al. (2011) observed that lower concentrations ($\mu\text{g/L}$) of metolachlor (mixture of both R- and S-enantiomers) altered the thyroid system in Japanese medaka. Thyroid system disruption was

also caused by low concentrations of acetochlor ($\mu\text{g/L}$), another member of the chloroacetanilide family (Yang et al., 2015). In other non-target aquatic organisms, S-metolachlor and its metabolites MOA and MESA induced developmental abnormalities in Pacific oyster larvae and sperm at low environmental concentrations (Mai et al., 2014). Moreover, the deleterious effects of environmental concentrations of S-metolachlor and MOA on crayfish were also reported (Velisek et al., 2018a, 2018b). Although several studies pointed out the toxicity of high concentrations of these substances on non-target aquatic organisms (Gutiérrez et al., 2019; Munn et al., 2006; Quintaneiro et al., 2018, 2017), comprehensive information on sublethal effects caused by environmental concentrations is still missing.

Early life stage zebrafish (*Danio rerio*) are widely used as sensitive and reliable alternative toxicity models with multiple advantages including high-throughput screening, embryo transparency, quick breeding and development, a sequenced genome (Love et al., 2004) and are in compliance with the 3Rs principle (Russell et al., 1959). Zebrafish development is highly influenced by its thyroid metabolism (Jarque and Piña, 2014). Moreover, the early development of the zebrafish thyroid system is comparable to its development in humans (Alt et al., 2006), complete with a similar hypothalamus–pituitary–thyroid (HPT) axis (Porazzi et al., 2009). This similarity makes the zebrafish a suitable model for evaluating thyroid gland disruption (Raldúa et al., 2012). In a recent review, Spaan et al. (2018) write that the complexity of thyroid disruption assessment calls for the evaluation of multiple endpoints. Morphological effects, transcription level of several genes interfering with the HPT axis, hatching, heart rate, behavior and swim bladder inflation were cited as the most common.

Although the toxicity of high concentrations of S-metolachlor on non-target aquatic organisms has been previously studied, the effects posed by low environmentally relevant concentrations of this compound and its metabolites have yet to be examined. The present study investigated the sublethal toxicity of environmentally relevant concentrations of S-metolachlor, its two metabolites MOA and MESA and their mixture on embryo-larval stages of *Danio rerio*. We evaluated apical endpoints of zebrafish embryos, conducted a neurobehavioral study measuring swimming activity to assess potential neurotoxicity and assessed biochemical responses by measuring the transcription changes of selected genes. This integrative approach allowed for detailed examination of sublethal toxicity, including potential thyroid metabolism disruption.

2. Materials and methods

2.1 Chemicals

S-metolachlor (S-M, CAS 87392-12-9, Pestanal, purity $\geq$ 98.0%), metolachlor oxanilic acid (MOA, CAS 152019-73-3, Pestanal, purity $\geq$ 98.0%), metolachlor ethanesulfonic acid (MESA, CAS 947601-85-6, Pestanal, purity $\geq$ 95.0%) were purchased from Sigma-Aldrich. Their stock solutions (50 mg/L) were prepared in Milli-Q water and were stored at 4 °C. The hormone triiodothyronine (T3, dissolved in methanol, CAS 6893-02-3, purity $\geq$ 95%, purchased from Sigma-Aldrich, used immediately) was used as positive control in the gene modulation experiments. Ethanol absolute (CAS 64-17-5, purity $\geq$ 99.8%, purchased from VWR Chemicals) was used as positive control in the spontaneous movement analysis and for the PCR analysis. ISO medium (ISO, 1996) ($\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (294 mg/L), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (123.3 mg/L), NaHCO_3 (63 mg/L), KCl (5.5 mg/L) in Milli-Q water) was used to prepare the final dilutions for the tests. RNAlater® (Ambion) was purchased from Sigma-Aldrich.

2.2 Test organism

Zebrafish (*Danio rerio*) embryos were collected from an unspecified wild type zebrafish strain maintained at RECETOX, Masaryk University (Czech Republic). Adult fish are held in aquariums with tap water at 26 ± 1 °C, at 14:10h light:dark photoperiod and fed 3 times a day with commercially available feeds (of which at least once with live brine shrimp *Artemia salina*). About 10-14 hours before the experimentation a box for collecting the embryos is placed in the aquarium and the spawning is induced in the morning by turning the light on. The eggs are collected 30 minutes after, rinsed and transferred to the ISO medium. Only fertilized and normally developing eggs are picked for the tests using a stereomicroscope (OLYMPUS, Japan) and exposed at 3-4 hpf (hours post fertilization). The embryos were then maintained at 26 °C, at 14:10h light:dark photoperiod till 120 hpf without feeding or renewal of test solutions.

2.3 FET (Fish embryo toxicity) test

The experiments were conducted in accordance with the fish early-life stage toxicity guideline (OECD, 2013). Selected eggs were exposed to the chosen herbicide S-metolachlor and its two metabolites MOA and MESA. The experiments were done with individual compounds and then with the mixture of the three. Chosen concentrations are based on environmental

151 concentrations detected in the rivers in the Czech Republic and in the Arcachon Bay in France
152 during 2010-2014 (Table 1).
153

Table 1

Maximum concentrations of metolachlor and its metabolites MOA and MESA detected in Czech rivers (CHMU (Czech Hydrometeorological Institute), 2018) and Arcachon Bay in France (source) during 2010-2014. Number of samples: N=703 values for every substance in Arcachon Bay; N=13 175, 7 503 and 7 502 for metolachlor, MOA and MESA respectively in surface water)

	Metolachlor	MOA	MESA	Location
Maximum concentration (ng/L)	5800	4200	4200	Surface water (CZ)
	1696	1609	1059	Arcachon Bay (FR)
% of values > 100 ng/L	2.5%	4.1%	19.1%	Surface water (CZ)
	36.0%	6.1%	36.6%	Arcachon Bay (FR)

The lowest used test concentration (1 µg/L) reflects the actual environmental concentration, and three higher (non-lethal) concentrations were also tested (30, 100 and 300 µg/L). Corresponding concentrations in nmol/L are showed in Supplementary Table S2. The mixture exposure included two concentration levels: 1 µg/L of each substance and 30 µg/L of each substance. ISO medium was used as negative control. The tests were conducted in glass crystallization dishes with transparent lids containing 20 ml of media and 20 embryos in three replicates. Every test was repeated independently three times with eggs from different spawning. Mortality, hatching success and malformations were controlled daily using a stereomicroscope. Eventual dead embryos/larvae were immediately removed from dishes to prevent contamination. Length of larvae (without the caudal fin) was measured on the last, fifth, day of the experiment on photos of larvae mounted in mixture of cellulose and anesthetics MS-222. Fifteen fish per replicate in 3 independent tests were measured. Photos were analyzed using the software QuickPhoto Micro 2.3.

“Morphological score” endpoint was assessed in order to evaluate global state of individual fish using slightly modified approach as previously published (Panzica-Kelly et al., 2010). Every fish embryo is classified using 0-3 point scores (description is in Supplementary Figure S1).

2.4 Light: dark locomotor test

Embryos for the locomotor activity analysis were handled as described above (2.3). The evening before measurements (to allow for acclimation), larvae were transferred to transparent 96-well microplates (Gama, Czech Republic), one larva per well. The distribution was random, 32 fish per exposure condition were used. Only morphologically normal, i.e.

non-malformed embryos were used. The experimentations were performed during the 5 dpf between 9 am and 16 pm. Prior to the experiment, the microplates were kept for 1 hour next to the ZEBRABOX (Viewpoint Life Sciences) at 26 °C for acclimation after the transport. Measurement done in the ZEBRABOX consisted of four 15 minutes alternating cycles (white light, i.e. 100% stimulus / dark, i.e. no stimulus / white light / dark). Total distances swam (mm) by individual fish were measured with integration period of 60 seconds and processed in real time using the software Zebralab (Viewpoint Life Sciences). The threshold between “inactive” fish and “active” fish was set on 0.3 cm/sec. The sensitivity threshold of ZEBRABOX was set on 120.

2.5 Spontaneous movement and heartbeat

Embryos were handled as described above (2.2) and exposed to concentrations of 1 and 100 µg/L of substances alone and to the mixtures at two concentration levels (1 and 30 µg/L of each substance in the mixture). They were maintained at 26 °C until 21 hpf when randomized capture of videos was realized till 23 hpf. Crystallization dish containing 20 embryos (= one replicate) was acclimatized under the stereomicroscope for 30 sec and then 2 min video was captured. Positive control (1% v/v ethanol) was used to enhance the spontaneous movements (tail coils). Videos were then analyzed by DanioScope software (Noldus, Netherlands) and reported as spontaneous movements per minute.

The same embryos were used for the heart rate activity measurement during the 3 dpf starting at 72 hpf. Hatched larvae were first put in solution of 2.5% methylcellulose containing 50 mg/L of anesthetics MS-222 and placed in Petri dish. Larvae were acclimatized for 10 seconds under the stereomicroscope and 20 second videos were captured. Videos were analyzed using the DanioScope software and reported as beat per minute.

2.6 Zebrafish exposures for analysis of gene expressions

Embryos were handled as described above (2.3) and exposed to 1, 30 and 100 µg/L of pesticides assessed alone and two concentrations of the mixture of the three pesticides (1 and 30 µg/L, respectively of each compound). Hormone T3 (triiodothyronine) was used as positive control to assess the expression of thyroid-related genes. Dead or heavily malformed embryos/larvae were removed daily. Whole larvae were collected at 120 hpf into pools of 20 larvae per Eppendorf tube containing 500 µl *RNAlater*® and stored at -20 °C upon RNA extraction.

2.6.1 RNA isolation

The total RNAs were extracted using the SV Total RNA Isolation System Kit (Promega). Whole larvae were homogenized in 500 µl of RNA Lysis Buffer using glass beads (Sigma) in the FastPrep®-24 (M.P. Biomedicals, USA). Lysed samples were collected, centrifuged and mixed with 450 µl of 75% v/v ethanol and transferred on the spin column. Handling with spin columns was performed according to the manufacturer's instructions. Before the DNase incubation step, an additional centrifuge step was added (13500 rpm, 2 minutes) to dry the column. All RNA samples were treated with DNase I mixture (40 µl of Yellow core buffer, 5 µl of MnCl₂ and 5 µl of DNase I) for 15 minutes in water bath at 37 °C. The rest of the isolation was performed according to the manufacturer's instructions and purified RNAs were collected in 50 µl of Nuclease-Free water. The concentration and purity of collected RNA samples were verified spectrophotometrically at 260/280 and 260/230 nm wavelength with Nanodrop™. Purity of all the samples was between 2.0-2.1 for the ratio 260/280 and 1.8-2.2 for the ratio 260/230.

2.6.2 Reverse transcription

Reverse transcription phase was performed with GoScript™ Reverse Transcription System kit (Promega) according to the manufacturer's instructions using TProfessional Thermocycler (Biometra, Analytik Jena). Purified RNA (1 µg) was reversely transcribed in order to get final volume of 20 µl of cDNA, and it was stocked at -20 °C upon qPCR analysis.

2.6.3 Quantitative PCR

8 genes were selected to evaluate the effects of chosen pesticides and their mixture on the thyroid hormone system, mitochondrial metabolism, regulation of the cell cycle and oxidative stress defense. Three reference zebrafish genes were used in the analysis (*β-actin*, *efla* and *rpl13*). Functioning and references of used primers are shown in Table 2. Sequences, efficiencies, and accession numbers are presented in Supplementary Table S1. Primers were purchased from Elisabeth Pharmacon (Czech Republic).

QPCR was carried out with Brilliant III Ultra-fast SYBR Green QPCR Master Mix kit (Stratagene-Agilent) on the LightCycler® 480 (Roche). Each qPCR mix (20 µl) contained 5 µl of cDNA (10 times diluted), 2 µl of reverse and forward primer (2 µM) and 10 µl of SYBR from the kit and 3 µl of nuclease free water.

LightCycler was set according to the instructions of the SYBR kit manufacturer (Stratagene-Agilent). Briefly, the pre-incubation lasted 3 min at 95 °C, then the amplification consisted of

45 cycles with 1 cycle at 95 °C for 5 sec and 60 °C for 10 sec. The melting curve continued at 95 °C for 5 sec and at 65 °C for 1 min. Finally, the cooling step was performed at 40 °C for 30 sec. Melting curves of every reaction were analyzed to assess reaction specificity. The data were normalized to the mean of the Ct values for three reference genes, *β-act*, *elf1a* and *rpl13* and analyzed using the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). Results are shown as fold changes of exposed to control group.

Table 2

Primers used for qPCR analysis in zebrafish (*Danio rerio*) and their references.

Gene	Function	References
<i>β-act</i>	Reference gene	Dedeh <i>et al.</i> , 2015
<i>ef1a</i>	Reference gene	Gentès <i>et al.</i> , 2015
<i>rpl13</i>	Reference gene	Gentès <i>et al.</i> , 2015
<i>12S</i>	Mitochondrial metabolism	Arini <i>et al.</i> , 2015
<i>p53</i>	Regulation of the cell cycle/apoptose	Gentès <i>et al.</i> , 2015
<i>cat</i>	Oxidative stress defense	Lerebours <i>et al.</i> , 2009
<i>sod1</i>	Oxidative stress defense	Gentès <i>et al.</i> , 2015
<i>cyp26a1</i>	RA (retinoic acid) signaling pathway	Oliveira <i>et al.</i> , 2013
<i>dio2</i>	Thyroid hormone system	Yan <i>et al.</i> , 2012
<i>thra</i>	Thyroid hormone system	Yan <i>et al.</i> , 2012
<i>thrb</i>	Thyroid hormone system	Yan <i>et al.</i> , 2012

2.7 Chemical analysis and water quality

At the beginning and at end of the experiments, pH, dissolved oxygen and conductivity were controlled. Saturation by oxygen was always higher than 98%. pH varied between 7.6-7.9. Conductivity varied between 500 and 700 µS/cm with no difference between ISO medium and pesticide solutions. Stability of substances during the five-day experiment was verified by measuring their concentrations by LC-MS/MS analysis as described in detail in Supplementary Material S1. In brief, LC-MS/MS was performed with a Waters LC chromatograph (Waters, Manchester, U.K.) using Acquity BEH C18 column and gradient elution. Detection was performed on a Xevo TQ-S quadrupole mass spectrometer (Waters Manchester, U.K.) after ESI ionisation in a positive ion mode. The analytes were quantified using external calibration of metolachlor, MOA and MESA (0.1 – 500 µg/L in 20% of methanol) and the limit of detection (LOD, signal to noise ratio S/N>3) and quantification (LOQ, S/N>10) were 0.05 and 0.1 µg/L, respectively for each analyte.

2.8 Data analysis

Length of larvae was assessed with one-way ANOVA test. Sum of different observed malformations in all replicates of the three independent experiments were compared with

271 control animals using Fisher's exact test in software GraphPad Prism (Version 5, GraphPad
272 Software).

273 Different independent locomotion tests (independent experiments) were all assessed
274 statistically apart because of high variability in control fish. Final results are presented as
275 mean of fold changes with control of the three independent experiments.

276 Data of locomotion tests, spontaneous movements and heart rate were verified for normality
277 (Shapiro-Wilk test; $p > 0.01$) and homoscedasticity (Levene test; $p > 0.05$), and if confirmed,
278 ANOVA followed by Dunnet post-hoc test was used. In the other case, non-parametric
279 Kruskal-Wallis test with Mann-Whitney post-hoc test was used.

280 All data for the gene expression analysis were treated as described above (Shapiro-Wilk test;
281 $p > 0.01$; Levene test; $p > 0.05$). Eventual values higher or smaller than 3 standard deviations
282 were discarded. Data were log normalized prior to the analysis. ANOVA followed by Dunnet
283 post-hoc test was used. If normality or homoscedasticity were not confirmed, non-parametric
284 Kruskal-Wallis test with Mann-Whitney post-hoc tests were performed. All analyses were
285 performed using software Statistica 13.3 (StatSoft, version 13.2, USA).

286

287

3. Results

3.1. Exposure and chemical analysis

Stability of all three compounds during the five-day experiment was verified. Nominal and measured concentrations, as well as the stability are shown in Supplementary Table S2. The concentrations of compounds were stable with the maximum decline of 13,3% at 30 µg/L of S-metolachlor. Measured values corresponded to nominal ones in case of MOA and were slightly higher at S-metolachlor and MESA. To simplify the presentation, nominal concentrations are shown and discussed in the results and discussions sections.

3.2 Mortality, hatching success, length of fish and malformation results

No significant effects on mortality were observed in any concentration tested of S-metolachlor, MOA, MESA or mixture of the three compounds. All controls had mortality < 10% as preconized by OECD guideline (OECD, 2013). Survival in all tested conditions was between 97.2% and 100%. Similarly, no significant effects on hatching success were found in all conditions tested. Hatching success values were between 95% and 100% in all variants.

Observed developmental abnormalities and length of fish are presented in Table 3. Generally, no patterns or dose-response effects were observed after exposure to S-metolachlor, its two metabolites MOA and MESA or in the mixture. No effects on the length of fish were observed in any condition tested. The most frequently observed malformations were at swim bladder (non-inflation) and yolk sac (malabsorption). No statistically significant effects were detected after exposures to the lower concentrations (1 and 30 µg/L) of S-metolachlor, MOA and MESA or after exposure to the low concentration mixture (1 µg/L of each substance). Following statistically significant effects were observed: S-metolachlor induced non-inflation of the swim bladder (100 µg/L) and malabsorption of the yolk sac (300 µg/L) and MESA (300 µg/L) induced craniofacial malformations. In addition, mixture of the three substances at concentration of 30 µg/L per each substance induced spinal deformations. Morphological score endpoint which allows assessment of each fish individually revealed as most problematic concentrations of 100 and 300 µg/L of S-metolachlor, 100 µg/L of MESA and 30 µg/L of mixture.

317 **Table 3** Frequencies (%) of different types of developmental malformations, length of larvae (μm) and morphological score of zebrafish after
318 120 hours exposure to different concentrations of pesticide, metabolites and their mixtures. Each value represents mean $\pm$ standard deviation
319 from 3 independent experiments each based on N=60 embryos. *P < 0.05; **P < 0.01

		μm	%								Absolute value
Endpoint		Body length	Edemas	Craniofacial deformation	Spine deformation	Non-inflated gas bladder	Tail deformation	Necrosis	Yolk sac malabsorption	Cardiovascular toxicity	Morphologica l score
Concentration μg/L											
S-metolachlor	Control	3768 ± 103	1.1 ± 1.6	1.1 ± 1.6	0 ± 0	0.6 ± 0.8	0 ± 0	0.6 ± 0.8	1.1 ± 0.8	1.1 ± 1.6	0.9 ± 1.0
	1	3772 ± 118	0.6 ± 0.8	1.7 ± 1.4	1.1 ± 0.8	2.8 ± 2.8	0 ± 0	0.6 ± 0.8	3.9 ± 2.8	0 ± 0	1.6 ± 1.1
	30	3767 ± 131	1.7 ± 2.4	2.8 ± 3.9	1.1 ± 1.6	3.3 ± 1.4	0 ± 0	0 ± 0	3.3 ± 3.6	0 ± 0	1.8 ± 1.8
	100	3763 ± 123	1.7 ± 1.4	2.3 ± 3.2	0.6 ± 0.8	4.6 ± 1.6*	0 ± 0	0.6 ± 0.8	4.0 ± 0.8	0 ± 0	2.0 ± 1.1*
	300	3738 ± 131	1.7 ± 1.4	2.8 ± 1.6	1.7 ± 1.4	2.3 ± 0.8	0 ± 0	0.6 ± 0.8	5.8 ± 2.2*	0 ± 0	2.2 ± 0.8*
MOA	Control	3714 ± 136	1.1 ± 0.8	2.3 ± 2.1	0.6 ± 0.8	3.4 ± 2.4	0 ± 0	2.8 ± 4.0	2.3 ± 2.1	0 ± 0	1.9 ± 1.5
	1	3712 ± 130	2.3 ± 2.2	2.3 ± 2.2	1.8 ± 1.5	1.8 ± 2.5	0 ± 0	0.6 ± 0.8	1.8 ± 2.5	1.2 ± 1.7	1.8 ± 1.9
	30	3719 ± 160	1.1 ± 1.6	0.6 ± 0.8	0 ± 0	2.9 ± 1.6	0 ± 0	0.6 ± 0.8	1.7 ± 0	1.7 ± 2.4	1.3 ± 0.8
	100	3678 ± 169	3.5 ± 2.4	2.3 ± 1.6	2.9 ± 0.7	5.1 ± 2.4	0 ± 0	1.7 ± 2.4	2.8 ± 0.8	3.4 ± 4.8	3.0 ± 1.4
	300	3694 ± 163	1.1 ± 1.6	1.1 ± 1.6	1.7 ± 1.4	3.9 ± 0.7	0 ± 0	2.3 ± 3.2	2.9 ± 0.8	0.6 ± 0.8	2.1 ± 1.6
MESA	Control	3661 ± 140	0.6 ± 0.8	0.6 ± 0.8	1.1 ± 0.8	2.2 ± 2.1	0 ± 0	0 ± 0	1.7 ± 1.4	0 ± 0	0.7 ± 0.5
	1	3708 ± 112	0 ± 0	1.1 ± 1.6	0 ± 0	2.2 ± 2.1	0 ± 0	0 ± 0	0.6 ± 0.8	0 ± 0	0.6 ± 0.2
	30	3700 ± 146	0 ± 0	1.1 ± 1.6	0 ± 0	1.1 ± 1.6	0 ± 0	0.6 ± 0.8	0.6 ± 0.8	0 ± 0	0.6 ± 0.4
	100	3683 ± 118	1.1 ± 0.8	4.4 ± 5.2*	1.7 ± 1.4	3.3 ± 1.4	0 ± 0	1.1 ± 1.6	2.8 ± 0.8	0 ± 0	2.0 ± 1.7**
	300	3628 ± 127	0 ± 0	2.3 ± 2.1	0.6 ± 0.8	1.1 ± 0.8	0 ± 0	0 ± 0	1.7 ± 0	0 ± 0	0.9 ± 0.4
MIXTU RE	Control	3755 ± 149	0.7 ± 1.0	4.3 ± 2.6	0.6 ± 0.8	6.3 ± 4.4	0.6 ± 0.8	2.3 ± 1.7	6.9 ± 6.6	0.6 ± 0.8	3.2 ± 1.7
	1 (of each)	3763 ± 163	2.8 ± 0.8	2.9 ± 1.6	3.4 ± 2.8	4.5 ± 0.8	0.6 ± 0.8	1.1 ± 0.8	5.1 ± 2.4	0 ± 0	3.1 ± 1.1
	30 (of each)	3740 ± 193	3.1 ± 2.5	8.0 ± 1.0	4.6 ± 3.5*	6.7 ± 5.2	0 ± 0	3.0 ± 2.3	12.2 ± 8.6	1.9 ± 1.6	5.9 ± 2.0**

3.2 Light:dark locomotion test and heartbeat

None of the studied compounds or their mixtures caused any statistically significant effects on the swimming behavior. Detailed results are shown in Supplementary Material S2. Similarly, there were no effects observed at the heart rate of zebrafish larvae, detailed results are shown in Supplementary Material S3.

3.3 Spontaneous movements

Results of the spontaneous movements of the zebrafish embryos are presented in Figure 1. Environmental concentrations of S-metolachlor (1 µg/L) and of the mixture of the three substances (1 µg/L of each substance) significantly ($P=0.001$ and 0.022 respectively) reduced frequency of the tail movements. 0.5% solution of ethanol, used as positive control, significantly enhanced tail movements ($P=0.009$).

Figure 1. Spontaneous tail movements frequency per minute in zebrafish embryos exposed SM: S-metolachlor; MOA: Metolachlor oxanilic acid; MESA: Metolachlor ethanesulfonic acid; MIX: mixture of SM, MOA and MESA (concentration of 1 et 30 µg/L of each substance in the mixture); EtOH: ethanol; * $P < 0.05$, ** $P < 0.01$

3.5 Analysis of gene expression

Gene expression of eight pre-selected genes was analyzed. Results are shown in Figure 2 as fold changes of ratio between selected genes and housekeeping genes. Exposures to T3 was used as positive control for thyroid-related genes.

The expression of mitochondrially encoded 12S rRNA (*12S*) gene was not significantly upregulated or downregulated in comparison with the control for all the concentrations of all the pesticides tested (Fig.3a). *P53*, gene related to the regulation of the cell cycle and cell apoptosis, was significantly upregulated at 100 µg/L of MOA (1.8-fold, $P=0.005$) and 30 µg/L of MESA (2-fold, $P=0.022$). The mixture of SM, MOA and MESA in concentration of 1 µg/L of each pesticide (2-fold, $P<0.001$) induced significant upregulation of *p53* (Fig.3b). Oxidative stress defense associated genes catalase - *cat* (Fig.3c) and copper/zinc superoxide dismutase - *sod1* (Fig.3d) were not affected by any of the pesticides tested.

We also studied the genes associated to RA signaling pathway: gene coding for cytochrome P450, family 26, subfamily A, polypeptide 1 - *cyp26a1* (Fig.3e) and genes associated to thyroid metabolism disruption: iodothyronine deiodinase 2 - *dio2* (Fig.3f), thyroid hormone nuclear receptor α - *thra* (Fig.3g), and thyroid hormone nuclear receptor beta - *thrb* (Fig.3h). None of the four genes was deregulated after exposure to S-metolachlor. On the contrary, the metabolites and the mixture affected gene transcriptions. MOA significantly increased the expression of *dio2* (3.5-fold, $P=0.015$) and *thrb* (1.8-fold, $P=0.034$) in the highest concentration tested 100 µg/L. MESA significantly increased the mRNA expression level of *dio2* (5.6-fold, $P=0.002$), *thra* (2.1-fold, $P=0.004$) and *thrb* (1.9-fold, $P=0.003$) in the concentration of 1 µg/L. Upregulation of *thra* was also observed in the MESA concentration of 30 µg/L (2-fold, $P=0.027$). The mixture of the three substances (at low level - 1 µg/L of each substance) significantly upregulated the gene expression in all four genes - *cyp26a1* (2.3-fold, $P<0.001$), *dio2* (2.5-fold, $P=0.007$), *thra* (2.5-fold, $P=0.001$), *thrb* (2.2-fold, $P=0.001$). The T3 hormone significantly increased the expression of the mRNA transcription level of *cyp26a1* (4.2-fold, $P=0.001$) and *thra* (4.7-fold, $P<0.001$).

Figure 2. Expression levels of selected genes involved in mitochondrial metabolism (a), regulation of the cell cycle (b), oxidative stress defense (c, d), and thyroid metabolism disruption (e,f,g,h) associated genes in zebrafish larvae exposed for 5 days to SM: S-metolachlor; MOA: Metolachlor oxanilic acid; MESA: Metolachlor ethanesulfonic acid; MIX: mixture of SM, MOA and MESA; T3: triiodothyronine hormone; $P < 0.05$; $^{**}P < 0.01$; $^{***}P < 0.001$

4. Discussion

S-metolachlor and its metabolites are among the most commonly reported herbicides occurring in relatively high concentrations in European waterbodies (Accinelli et al., 2002; Cerejeira et al., 2003; Farlin et al., 2018; Kapsi et al., 2019; Meffe and de Bustamante, 2014; Vryzas et al., 2009). Despite of the overall importance, the toxicity of S-metolachlor on fish was commonly evaluated at environmentally non-relevant conditions and high concentrations (Quintaneiro et al., 2017) or using only the racemic mixture of metolachlor (Jin et al., 2011). More recently, toxicity of one of the metabolites - MOA, was assessed by Velisek et al. (2018a) on crayfish but the effects of S-metolachlor and its metabolites on fish remain poorly characterized. Our interest was mainly focused on thyroid disruption, due to findings of Jin et al. (2011) and Yang et al. (2015) who linked this type of disruption with some chloroacetanilide pesticides. Regarding the complexity of the thyroid system, an integrative approach combining multiple endpoints was used.

4.1 Effects on embryo-larval development

Few statistically significant induced malformations were detected in the present study but only at higher exposure levels. Spine deformations were induced in fish exposed to mixture of substances (30 µg/L of each of the three compounds, ~ 90 µg/L sum concentration) but, interestingly, no such effects were observed at any of the substances alone up to 300 µg/L. Apparently, concentration addition (CA) effects in the mixture of different compounds was a major driver beyond this observation (European Chemicals Agency, 2014). Correspondingly, Lydy *et al.*, (2004) discuss that, generally, pesticides of the same chemical class exert the CA effects. This is supported also by findings of Junghans et al. (2003) who reported the CA effects of the mixture of eight chloroacetanilide herbicides in the study with susceptible organism - algae. It should be noted that despite of the statistical significance, the frequency of spine malformation (4.6%) is rather low and might be of minor biological significance. Nevertheless, further experiments could eventually further explore the mechanism behind this observation.

Other significant malformations (craniofacial deformations, non-inflated swim bladder, yolk sac malabsorption) were induced by individual substances at higher concentrations (100 or 300 µg/L). The observed malformations could be linked to the disruption of thyroid signaling. Liu and Chan (2002) showed that thyroid disruption lead to retarded inflation of the swim bladder, retarded absorption of the yolk sac and retarded maturation of the gastrointestinal

system. Quintaneiro et al. (2017) described impairment of yolk sac absorption of 96 hpf zebrafish larvae exposed to high concentrations of S-metolachlor (45 mg/L).

In our study, no statistically significant mortality was observed for tested concentrations of all substances and the mixture. According to U.S. EPA Ecotox database the NOEL of S-metolachlor for bluegill is 1.5 mg/L and for rainbow trout 2.5 mg/L (measured at 96 hpf). Both concentrations are higher than our highest tested concentration of 300 µg/L. To our knowledge, no data are available for the effects of metabolites on fish. Recently, Velisek et al. (2018a) found no mortality of embryo-larval stages of crayfish exposed for 45 days up to 420 µg/L of MOA. No effect was found on the length of larvae or the hatching success of fish exposed to tested substances and the mixture. This finding is in concordance with 100 times higher LOEC (29.0 mg/L) of S-metolachlor for zebrafish embryos (96 hpf) for hatching success established by Quintaneiro et al. (2017). Thus, as it was expected, low environmentally relevant concentrations used in this study elicited no significant effects on traditional apical endpoints such as mortality, hatching success and length of zebrafish larvae and we further investigated other sublethal endpoints.

4.3 Effects on behavior

Use of zebrafish larvae in behavioral studies have multiple advantages for evaluation of neurotoxicity of pesticides as discussed by Pittman (2017). To our knowledge, this is the first study that investigated the effects of S-metolachlor and its metabolites on fish behavior although some studies suggested S-metolachlor effects on neurobehavior. Quintaneiro et al. (2017) exposed zebrafish embryos for 4 days to 25 mg/L of S-metolachlor and observed inhibited acetylcholinesterase activity. Villa et al. (2018) described decreased speed and distance travelled by Chironomous larvae upon metolachlor exposures (27.4 mg/L). Cook & Moore (2008) found that sublethal concentration of metolachlor (80 µg/L) altered the fighting behavior of crayfish. On the contrary, available study with MOA revealed no effect on the behavior of crayfish (activity, distance, speed) exposed up to 420 µg/L (Velisek et al., 2018a). Moreover, some pesticides of the same chemical class of chloroacetanilides were also reported to influence fish locomotion. Pretilachlor (1 mg/L) decreased feeding attempts and increased burst swimming reactions and buccal movements in adult fish *Clarias batrachus* (Soni and Verma, 2018). Acetochlor at 5 mg/L decreased spontaneous tail movements in zebrafish embryos (24 hpf) as well as total distance travelled, average speed and time of movement of larvae 6 dpf (Wang et al., 2019).

In the present study, S-metolachlor decreased spontaneous tail movements in zebrafish embryos, although this observation was recorded only in the lowest concentration (1 µg/L). Corresponding decrease was also observed after exposure to the lower concentration mixture (all three compounds, 1 µg/L each). The observed effects on spontaneous movements may be caused by S-metolachlor interfering with maternal T4 stock in yolk sac or its signaling via an $\alpha V\beta 3$ integrin as shown by Yonkers and Ribera (2008). The thyroid follicles start to secrete T4 hormone at 72 hpf (Porazzi et al., 2009), and nongenomic mechanism with maternal T4 influencing the neurodevelopment of zebrafish was previously reported (Yonkers and Ribera, 2008).

On the contrary, light:dark locomotion test didn't reveal any abnormal behavior. High biological variability in larvae movement could be linked to relatively wide time frame within the day (9 am – 4 pm), when the measurements were carried out. This was discussed by some authors who recommend specific more narrow frames like 11 am – 3 pm (Colwill and Creton, 2011) or 1 pm – 3:30 pm (MacPhail et al., 2009).

4.4 Effects on gene expression

Exposures to the metabolites appeared to have more pronounced effects on zebrafish larvae than the exposure to the parent compound. The highest tested concentration (100 µg/L) of MOA caused overexpression of genes *dio2*, *thrb* and *p53*. MESA induced overexpression of *dio2*, *thra* and *thrb* in fish exposed to low 1 µg/L, and *thra* and *p53* in fish exposed to 30 µg/L. The induction of *p53* indicates an activation of apoptotic processes or cell cycle arrest to protect the cells from additional detrimental effects (Ko and Prives, 1996). Genes *dio2*, *thra*, *thrb*, were proposed by Jin et al., (2011) as suitable biomarkers for thyroid system disruption in fish. Spaan et al., (2019) describe them as important genes (together with other deiodinases *dio1* and *dio3* and genes *hhex* and *NIS*) in the HPT axis. Deiodinase 2 is a selenoprotein that catalyzes the transformation of T4 to T3, thyroid hormones regulating the neurodevelopment. It is the most important deiodinase in zebrafish embryonic development as it controls the quantity of T3 hormone in the tissues (Walpita et al., 2009). Furthermore, deiodinases were found to affect the eye development of zebrafish (Houbrechts et al., 2016) and the overexpression of *dio2* has been linked with hypothyroidism (Orozco and Valverde-R, 2005). Thyroid hormone receptors α and β act as transcription factors for thyroid hormones. According to Walter et al. (2019), mRNA levels of deiodinases and thyroid hormone receptors vary throughout the zebrafish development, with expression peak of *dio2* and *thrb* around 72-96 hpf and following decrease at 120 hpf. Expression of *thra* is stable from 24 hpf

to 120 hpf. Furthermore, as described by Porazzi et al., (2009) and Raldúa et al., (2012), thyroid system in zebrafish is fully developed during 5 dpf.

Although MESA and MOA affected expressions of several genes, no changes were observed after exposure to S-metolachlor. Contradictory observations for S-metolachlor were found by Jin et al. (2011), where 10 µg/L caused overexpression of *dio2*, *thra* and *thrb* in juvenile medaka, only in females. These differences may be related to multiple factors such as exposure duration (14 days in Jin et al. (2011) in comparison to 5 days in our study) or use of racemic mixture of metolachlor by Jin et al. (2011), which has been found more toxic to non-target organisms than pure S-metolachlor (Liu et al., 2006; Ye et al., 2010). Use of different fish species in different life stage (1-month-old juveniles used by Jin et al. (2011) in comparison with embryos in the present study) is also of importance; differences in sensitivity of medaka and zebrafish have already been documented (Perrichon et al., 2014; Vignet et al., 2019).

The mixture of S-metolachlor and its two metabolites appeared to be most hazardous, and low environmental concentration (1 µg/L of each substance) altered transcription of several genes like *p53*, *dio2*, *thra*, *thrb* and *cyp26a1*. Gene *cyp26a1*, encoding for enzyme degrading retinoic acid, is very important in embryogenesis, and its overexpression can disturb all the retinoic acid signalling pathway and may lead to developmental malformations (Hu et al., 2008). Effects on gene expression observed at low concentration of the mixture were not confirmed at the high concentration. This mixture “low-dose effects” mechanism has been discussed in the field of thyroid, and generally endocrine, system disruption, where nonmonotonic responses and U-shaped dose-response curves have been documented (Melnick et al., 2002; Vandenberg et al., 2012).

The most known pesticides to exert thyroid dysfunction are organochlorine pesticides (Calsolaro et al., 2017). However, several chloroacetanilides have also been shown to disrupt the thyroid system including the above discussed metolachlor in adult and juvenile medaka (Jin et al., 2011), acetochlor (Jiang et al., 2015; Yang et al., 2015) and pretilachlor (Jiang et al., 2016) in zebrafish embryos, acetochlor in larval rare minnow (Li et al., 2009) or butachlor in adult rare minnow (Zhu et al., 2014). Similarly to the present work, the above-mentioned studies documented changes in gene expressions of deiodinases and thyroid nuclear receptors and used lower concentrations below hundreds of micrograms. Comparatively, Li et al., (2009) and Yang et al., (2015) also mentioned a non-sigmoidal dose response curve.

No changes were observed in the expression of the next three genes: *12S*, *sod1* and *cat*. Gene *12S* is associated with mitochondrial metabolism thus suggesting no effect on metabolic homeostasis (Arini et al., 2015). *Sod1* and *cat* are important genes managing some parts of the chain of the reactive oxygen species (ROS) in cell, scavenging respectively the superoxide radicals and hydrogen peroxide (Seifried et al., 2007). From these findings we conclude that S-metolachlor, its two metabolites and their mixture did not induce any oxidative stress detectable by *sod1* and *cat* in the zebrafish embryos. Quintaneiro et al. (2017) did not found either an effect of S-metolachlor on the catalase activity in zebrafish embryos exposed up to 25 mg/L. In contrast, MOA seems to have an inhibiting effect on catalase activity of a water crustacean: crayfish exposed to low concentrations (4.2 µg/L) (Velisek et al., 2018a).

5. Conclusion

Our work brings important findings about the effects of environmentally relevant mixture of one of the most used herbicides S-metolachlor and its two metabolites on early life stages of zebrafish. We have observed specific malformations that could be linked to thyroid disruption as well as decreased spontaneous movements of zebrafish larvae. Interestingly, metabolites were more harmful than S-metolachlor to zebrafish larvae at the level of gene expression with apparently strong effects on genes involved in cell cycle regulation and thyroid-related signaling. Further detailed analyses are needed to fully elucidate the dose-response relationship. The present study brings new information highlighting the importance of metabolites and the pesticides mixtures as emphasized in the European Environmental Action Programme (European Parliament and Council, 2013).

References

- Accinelli, C., Vicari, A., Pisa, P.R., Catizone, P., 2002. Losses of atrazine, metolachlor, prosulfuron and triasulfuron in subsurface drain water. I. Field results. *Agronomie* 22, 399–411.
- Alt, B., Reibe, S., Feitosa, N.M., Elsalini, O.A., Wendl, T., Rohr, K.B., 2006. Analysis of origin and growth of the thyroid gland in zebrafish. *Dev. Dyn.* 235, 1872–1883.
- Arini, A., Gourves, P.Y., Gonzalez, P., Baudrimont, M., 2015. Metal detoxification and gene expression regulation after a Cd and Zn contamination: An experimental study on *Danio rerio*. *Chemosphere* 128, 125–133.
- Atwood, D., Paisley-Jones, C., 2017. Pesticides Industry Sales and Usage 2008 - 2012.
- Blaser, H.-U., Spindler, F., 1997. Enantioselective catalysis for agrochemicals. The case histories of (S)-metolachlor, (R)-metalaxyl and clozylacon. *Top. Catal.* 4, 275–282.
- Calsolaro, V., Pasqualetti, G., Niccolai, F., Caraccio, N., Monzani, F., 2017. Thyroid Disrupting Chemicals. *Int. J. Mol. Sci.* 18.
- Cerejeira, M., Viana, P., Batista, S., Pereira, T., Silva, E., Valério, M., Silva, A., Ferreira, M., Silva-Fernandes, A., 2003. Pesticides in Portuguese surface and ground waters. *Water Res.* 37, 1055–1063.
- CHMU (Czech Hydrometeorological Institute), 2018. Online water quality database [Database]. URL www.hydro.chmi.cz/pasporty (accessed 11.28.18).
- Colwill, R.M., Creton, R., 2011. Locomotor behaviors in zebrafish (*Danio rerio*) larvae. *Behav. Processes* 86, 222–229.
- Cook, M.E., Moore, P.A., 2008. The Effects of the Herbicide Metolachlor on Agonistic Behavior in the Crayfish, *Orconectes rusticus*. *Arch. Environ. Contam. Toxicol.* 55, 94–102.
- De Liguoro, M., Bona, M.D., Gallina, G., Capolongo, F., Gallochio, F., Binato, G., Di Leva, V., 2014. A monitoring of chemical contaminants in waters used for field irrigation and livestock watering in the Veneto region (Italy), using bioassays as a screening tool. *Environ. Sci. Pollut. Res.* 21, 3546–3557.
- Dedeh, A., Ciutat, A., Treguer-Delapierre, M., Bourdineaud, J.P., 2015. Impact of gold nanoparticles on zebrafish exposed to a spiked sediment. *Nanotoxicology* 9, 71–80.
- Dobšíková, R., Blahová, J., Modrá, H., Škorič, M., Svobodová, Z., 2011. The effect of acute exposure to herbicide Gardoprim Plus Gold 500 SC on haematological and biochemical indicators and histopathological changes in common carp (*Cyprinus carpio* L.). *ACTA VET* 80, 359–363.
- Elliott, S.M., VanderMeulen, D.D., 2017. A regional assessment of chemicals of concern in surface waters of four Midwestern United States national parks. *Sci. Total Environ.* 579, 1726–1735.
- European Chemicals Agency, 2014. Transitional Guidance on the Biocidal Products Regulation: Transitional Guidance on mixture toxicity assessment for biocidal products for the environment.
- European Parliament and Council, 2013. General Union Environment Action Programme to 2020 “Living well, within the limits of our planet.”
- Farlin, J., Gallé, T., Bayerle, M., Pittois, D., Köppchen, S., Krause, M., Hofmann, D., 2018. Breakthrough dynamics of s-metolachlor metabolites in drinking water wells: Transport pathways and time to trend reversal. *J. Contam. Hydrol.* 213, 62–72.
- Fauvelle, V., Belles, A., Budzinski, H., Mazzella, N., Plus, M., 2018. Simulated conservative tracer as a proxy for S-metolachlor concentration predictions compared to POCIS measurements in

- 567 Arcachon Bay. Mar. Pollut. Bull. 133, 423–427.
- 568 Gentès, S., Maury-Brachet, R., Feng, C., Pedrero, Z., Tessier, E., Legeay, A., Mesmer-Dudons, N.,
569 Baudrimont, M., Maurice, L., Amouroux, D., Gonzalez, P., 2015. Specific Effects of Dietary
570 Methylmercury and Inorganic Mercury in Zebrafish (*Danio rerio*) Determined by Genetic,
571 Histological, and Metallothionein Responses. Environ. Sci. Technol. 49, 14560–14569.
- 572 Glinski, D.A., Purucker, S.T., Van Meter, R.J., Black, M.C., Henderson, W.M., 2018. Analysis of
573 pesticides in surface water, stemflow, and throughfall in an agricultural area in South Georgia,
574 USA. Chemosphere 209, 496–507.
- 575 Götz, T., Böger, P., 2004. The Very-Long-Chain Fatty Acid Synthase Is Inhibited by
576 Chloroacetamides. Verlag der Zeitschrift für Naturforsch. 549–553.
- 577 Gustavsson, M., Kreuger, J., Bundschuh, M., Backhaus, T., 2017. Pesticide mixtures in the Swedish
578 streams: Environmental risks, contributions of individual compounds and consequences of
579 single-substance oriented risk mitigation. Sci. Total Environ. 598, 973–983.
- 580 Gutiérrez, I.B., Mesquita, A.F., Gonçalves, F.J.M., Marques, J.C., Gonçalves, A.M.M., 2019.
581 Biomarkers' responses of the benthic clam *Scrobicularia plana* to the main active ingredients (S-
582 metolachlor and Terbutylazine) of a common herbicide. Ecol. Indic. 96, 611–619.
- 583 Hayes, A.W., Li, R., Hoeng, J., Iskandar, A., Peistch, M.C., Dourson, M.L., 2019. New approaches to
584 risk assessment of chemical mixtures. Toxicol. Res. Appl. 3, 1–10.
- 585 Heydens, W.F., Lamb, I.C., Wilson, A.G.E., 2010. Chloracetanilides. Hayes' Handb. Pestic. Toxicol.
586 1753–1769.
- 587 Hladik, M.L., Hsiao, J.J., Roberts, A.L., 2005. Are Neutral Chloroacetamide Herbicide Degradates of
588 Potential Environmental Concern? Analysis and Occurrence in the Upper Chesapeake Bay.
589 Environ. Sci. Technol. 39, 6561–6574.
- 590 Houbrechts, A.M., Vergauwen, L., Bagci, E., Van houcke, J., Heijlen, M., Kulemeka, B., Hyde, D.R.,
591 Knapen, D., Darras, V.M., 2016. Deiodinase knockdown affects zebrafish eye development at
592 the level of gene expression, morphology and function. Mol. Cell. Endocrinol. 424, 81–93.
- 593 Hu, P., Tian, M., Bao, J., Xing, G., Gu, X., Gao, X., Linney, E., Zhao, Q., 2008. Retinoid regulation of
594 the zebrafish *cyp26a1* promoter. Dev. Dyn. 237, 3798–3808.
- 595 ISO, 1996. ISO 7346-2:1996 - Water quality -- Determination of the acute lethal toxicity of substances
596 to a freshwater fish [*Brachydanio rerio* Hamilton-Buchanan (*Teleostei*, *Cyprinidae*)] -- Part 1:
597 Static method; Part 2: Semi-static method.
- 598 Jarque, S., Piña, B., 2014. Deiodinases and thyroid metabolism disruption in teleost fish. Environ. Res.
599 135, 361–375.
- 600 Jiang, J., Chen, Y., Yu, R., Zhao, X., Wang, Q., Cai, L., 2016. Pretilachlor has the potential to induce
601 endocrine disruption, oxidative stress, apoptosis and immunotoxicity during zebrafish embryo
602 development. Environ. Toxicol. Pharmacol. 42, 125–134.
- 603 Jiang, J., Wu, S., Liu, X., Wang, Y., An, X., Cai, L., Zhao, X., 2015. Effect of acetochlor on
604 transcription of genes associated with oxidative stress, apoptosis, immunotoxicity and endocrine
605 disruption in the early life stage of zebrafish. Environ. Toxicol. Pharmacol. 40, 516–523.
- 606 Jin, Y., Chen, R., Wang, L., Liu, J., Yang, Y., Zhou, C., Liu, W., Fu, Z., 2011. Effects of metolachlor
607 on transcription of thyroid system-related genes in juvenile and adult Japanese medaka (*Oryzias*
608 *latipes*). Gen. Comp. Endocrinol. 170, 487–493.
- 609 Junghans, M., Backhaus, T., Faust, M., Scholze, M., Grimme, L.H., 2003. Predictability of combined
610 effects of eight chloroacetanilide herbicides on algal reproduction. Pest Manag. Sci. 59, 1101–
611 1110.

612 Kapsi, M., Tsoutsi, C., Paschalidou, A., Albanis, T., 2019. Environmental monitoring and risk
613 assessment of pesticide residues in surface waters of the Louros River (N.W. Greece). *Sci. Total*
614 *Environ.* 650, 2188–2198.

615 Ko, L.J., Prives, C., 1996. p53: Puzzle and paradigm. *Genes Dev.* 10, 1054–1072.

616 Lerebours, A., Gonzalez, P., Adam, C., Camilleri, V., Bourdineaud, J.-P., Garnier-Laplace, J., 2009.
617 Comparative analysis of gene expression in brain, liver, skeletal muscles, and gills of zebrafish
618 (*Danio rerio*) exposed to environmentally relevant waterborne uranium concentrations. *Environ.*
619 *Toxicol. Chem.* 28, 1271–1278.

620 Li, W., Zha, J., Li, Z., Yang, L., Wang, Z., 2009. Effects of exposure to acetochlor on the expression
621 of thyroid hormone related genes in larval and adult rare minnow (*Gobiocypris rarus*). *Aquat.*
622 *Toxicol.* 94, 87–93.

623 Liu, H., Ye, W., Zhan, X., Liu, W., 2006. A comparative study of rac- and S-metolachlor toxicity to
624 *Daphnia magna*. *Ecotoxicol. Environ. Saf.* 63, 451–455.

625 Liu, Y.-W., Chan, W.-K., 2002. Thyroid hormones are important for embryonic to larval transitory
626 phase in zebrafish. *Differentiation* 70, 36–45.

627 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
628 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402–8.

629 Love, D.R., Pichler, F.B., Dodd, A., Copp, B.R., Greenwood, D.R., 2004. Technology for high-
630 throughput screens: the present and future using zebrafish. *Curr. Opin. Biotechnol.* 15, 564–571.

631 Lydy, M., Belden, J., Wheelock, C., Hammock, B., Denton, D., 2004. Challenges in Regulating
632 Pesticide Mixtures. *Ecol. Soc.* 9.

633 MacPhail, R.C., Brooks, J., Hunter, D.L., Padnos, B., Irons, T.D., Padilla, S., 2009. Locomotion in
634 larval zebrafish: Influence of time of day, lighting and ethanol. *Neurotoxicology* 30, 52–58.

635 Mai, H., Gonzalez, P., Pardon, P., Tapie, N., Budzinski, H., Cachot, J., Morin, B., 2014. Comparative
636 responses of sperm cells and embryos of Pacific oyster (*Crassostrea gigas*) to exposure to
637 metolachlor and its degradation products. *Aquat. Toxicol.* 147, 48–56.

638 Meffe, R., de Bustamante, I., 2014. Emerging organic contaminants in surface water and groundwater:
639 A first overview of the situation in Italy. *Sci. Total Environ.* 481, 280–295.

640 Melnick, R., Lucier, G., Wolfe, M., Hall, R., Stancel, G., Prins, G., Gallo, M., Reuhl, K., Ho, S.-M.,
641 Brown, T., Moore, J., Leakey, J., Haseman, J., Kohn, M., 2002. Summary of the National
642 Toxicology Program's report of the endocrine disruptors low-dose peer review. *Environ. Health*
643 *Perspect.* 110, 427–31.

644 Müller, M.D., Poiger, T., Buser, H.-R., 2001. Isolation and Identification of the Metolachlor
645 Stereoisomers Using High-Performance Liquid Chromatography, Polarimetric Measurements,
646 and Enantioselective Gas Chromatography. *J. Agric. Food Chem.* 49, 42–49.

647 Munn, M.D., Gilliom, R.J., Moran, P.W., Nowell, L.H., 2006. Pesticide Toxicity Index for Freshwater
648 Aquatic Organisms, 2nd Edition.

649 OECD, 2013. Test No. 210: Fish early life stage toxicity test. OECD Guideline for the testing of
650 chemicals. [WWW Document]. URL [https://www.oecd-ilibrary.org/docserver/9789264203785-
651 en.pdf?expires=1568722812&id=id&accname=ocid56014292&checksum=A5DDCACDA2BC8
652 F0AAD663E4B7777B894](https://www.oecd-ilibrary.org/docserver/9789264203785-en.pdf?expires=1568722812&id=id&accname=ocid56014292&checksum=A5DDCACDA2BC8F0AAD663E4B7777B894) (accessed 9.9.16).

653 Oliveira, E., Casado, M., Raldúa, D., Soares, A., Barata, C., Na, B.P., 2013. Retinoic acid receptors'
654 expression and function during zebrafish early development. *J. Steroid Biochem. Mol. Biol.* 138,
655 143–151.

656 Orozco, A., Valverde-R, C., 2005. Thyroid Hormone Deiodination in Fish. *Thyroid* 15, 799–813.

657 Panzica-Kelly, J.M., Zhang, C.X., Danberry, T.L., Flood, A., DeLan, J.W., Brannen, K.C., Augustine-
658 Rauch, K.A., 2010. Morphological score assignment guidelines for the dechorionated zebrafish
659 teratogenicity assay. *Birth Defects Res. Part B - Dev. Reprod. Toxicol.* 89, 382–395.

660 Perrichon, P., Le Bihanic, F., Bustamante, P., Le Menach, K., Budzinski, H., Cachot, J., Cousin, X.,
661 2014. Influence of sediment composition on PAH toxicity using zebrafish (*Danio rerio*) and
662 Japanese medaka (*Oryzias latipes*) embryo-larval assays. *Environ. Sci. Pollut. Res.* 21, 13703–
663 13719.

664 Pittman, J., 2017. Zebrafish Neurotoxicity Models, in: Kalueff, A.V. (Ed.), *The Rights and Wrongs of*
665 *Zebrafish: Behavioral Phenotyping of Zebrafish*. Springer, pp. 207–219.

666 Porazzi, P., Calebiro, D., Benato, F., Tiso, N., Persani, L., 2009. Thyroid gland development and
667 function in the zebrafish model. *Mol. Cell. Endocrinol.* 312, 14–23.

668 Quintaneiro, C., Patrício, D., Novais, S.C., Soares, A.M.V.M., Monteiro, M.S., 2017. Endocrine and
669 physiological effects of linuron and S-metolachlor in zebrafish developing embryos. *Sci. Total*
670 *Environ.* 586, 390–400.

671 Quintaneiro, C., Soares, A.M.V.M., Monteiro, M.S., 2018. Effects of the herbicides linuron and S-
672 metolachlor on Perez's frog embryos. *Chemosphere* 194, 595–601.

673 Raldúa, D., Thienpont, B., Babin, P.J., 2012. Zebrafish eleutheroembryos as an alternative system for
674 screening chemicals disrupting the mammalian thyroid gland morphogenesis and function.
675 *Reprod. Toxicol.* 33, 188–197.

676 Rose, M.T., Cavagnaro, T.R., Scanlan, C.A., Rose, T.J., Vancov, T., Kimber, S., Kennedy, I.R.,
677 Kookana, R.S., Van Zwieten, L., 2016. Impact of Herbicides on Soil Biology and Function. *Adv.*
678 *Agron.* 136, 133–220.

679 Russell, W.M.S., Burch, R.L., Hume, C.W., 1959. *The Principles of Humane Experimental*
680 *Technique*. Methuen, London.

681 Ryberg, K.R., Vecchia, A. V., Gilliom, R.J., Martin, J.D., 2014. Pesticide Trends in Major Rivers of
682 the United States, 1992–2010.

683 Seifried, H.E., Anderson, D.E., Fisher, E.I., Milner, J.A., 2007. A review of the interaction among
684 dietary antioxidants and reactive oxygen species. *J. Nutr. Biochem.* 18, 567–579.

685 Sidoli, P., Lassabatere, L., Angulo-Jaramillo, R., Baran, N., 2016. Experimental and modeling of the
686 unsaturated transports of S-metolachlor and its metabolites in glaciofluvial vadose zone solids. *J.*
687 *Contam. Hydrol.* 190, 1–14.

688 Soni, R., Verma, S.K., 2018. Acute toxicity and behavioural responses in *Clarias batrachus*
689 (Linnaeus) exposed to herbicide pretilachlor. *Heliyon* 4, e01090.

690 Spaan, K., Haigis, A.-C., Weiss, J., Legradi, J., 2019. Effects of 25 thyroid hormone disruptors on
691 zebrafish embryos: A literature review of potential biomarkers. *Sci. Total Environ.* 656, 1238–
692 1249.

693 Thomas Poiger, M.D.M. & H.-R.B., 2002. Verifying the Chiral Switch of the Pesticide Metolachlor on
694 the Basis of the Enantiomer Composition of Environmental Residues 56, 300–303.

695 Tian, D., Mao, H., Lv, H., Zheng, Y., Peng, C., Hou, S., 2018. Novel two-tiered approach of
696 ecological risk assessment for pesticide mixtures based on joint effects. *Chemosphere* 192, 362–
697 371.

698 U.S. Environmental Protection Agency, 1992. ECOREF #334 [Database]. *Pestic. Ecotoxicity*
699 *Database*. URL <https://cfpub.epa.gov/ecotox/search.cfm> (accessed 1.8.19).

- Vandenberg, L.N., Colborn, T., Hayes, T.B., Heindel, J.J., Jacobs, D.R., Lee, D.-H., Shioda, T., Soto, A.M., vom Saal, F.S., Welshons, W. V, Zoeller, R.T., Myers, J.P., Myers, J.P., 2012. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocr. Rev.* 33, 378–455.
- Velisek, J., Stara, A., Zuskova, E., Kubec, J., Buric, M., Kouba, A., 2018a. Chronic toxicity of metolachlor OA on growth, ontogenetic development, antioxidant biomarkers and histopathology of early life stages of marbled crayfish. *Sci. Total Environ.* 643, 1456–1463.
- Velisek, J., Stara, A., Zuskova, E., Kubec, J., Buric, M., Kouba, A., 2018b. Effects of s-metolachlor on early life stages of marbled crayfish. *Pestic. Biochem. Physiol.*
- Vignet, C., Cappello, T., Fu, Q., Lajoie, K., De Marco, G., Clérandeau, C., Mottaz, H., Maisano, M., Hollender, J., Schirmer, K., Cachot, J., 2019. Imidacloprid induces adverse effects on fish early life stages that are more severe in Japanese medaka (*Oryzias latipes*) than in zebrafish (*Danio rerio*). *Chemosphere*.
- Villa, S., Di Nica, V., Pescatore, T., Bellamoli, F., Miari, F., Finizio, A., Lencioni, V., 2018. Comparison of the behavioural effects of pharmaceuticals and pesticides on *Diamesa zernyi* larvae (*Chironomidae*). *Environ. Pollut.* 238, 130–139.
- Vryzas, Z., Vassiliou, G., Alexoudis, C., Papadopoulou-Mourkidou, E., 2009. Spatial and temporal distribution of pesticide residues in surface waters in northeastern Greece. *Water Res.* 43, 1–10.
- Walpita, C.N., Crawford, A.D., Janssens, E.D.R., Van der Geyten, S., Darras, V.M., 2009. Type 2 Iodothyronine Deiodinase Is Essential for Thyroid Hormone-Dependent Embryonic Development and Pigmentation in Zebrafish. *Endocrinology* 150, 530–539.
- Walter, K.M., Miller, G.W., Chen, X., Yaghoobi, B., Puschner, B., Lein, P.J., 2019. Effects of thyroid hormone disruption on the ontogenetic expression of thyroid hormone signaling genes in developing zebrafish (*Danio rerio*). *Gen. Comp. Endocrinol.* 272, 20–32.
- Wang, H., Meng, Z., Zhou, L., Cao, Z., Liao, X., Ye, R., Lu, H., 2019. Effects of acetochlor on neurogenesis and behaviour in zebrafish at early developmental stages. *Chemosphere* 220, 954–964.
- Yan, W., Zhou, Y., Yang, J., Li, S., Hu, D., Wang, J., Chen, J., Li, G., 2012. Waterborne exposure to microcystin-LR alters thyroid hormone levels and gene transcription in the hypothalamic–pituitary–thyroid axis in zebrafish larvae. *Chemosphere* 87, 1301–1307.
- Yang, M., Hu, J., Li, S., Ma, Y., Gui, W., Zhu, G., 2015. Thyroid endocrine disruption of acetochlor on zebrafish (*Danio rerio*) larvae. *J. Appl. Toxicol.* n/a-n/a.
- Ye, J., Zhao, M., Liu, J., Liu, W., 2010. Enantioselectivity in environmental risk assessment of modern chiral pesticides. *Environ. Pollut.* 158, 2371–2383.
- Yonkers, M.A., Ribera, A.B., 2008. Sensory Neuron Sodium Current Requires Nongenomic Actions of Thyroid Hormone During Development. *J. Neurophysiol.* 100, 2719–2725.
- Zemolin, C.R., Avila, L.A., Cassol, G.V., Massey, J.H., Camargo, E.R., 2014. Environmental fate of S-Metolachlor - A Review. *Planta Daninha* 32, 655–664.
- Zhu, L., Li, W., Zha, J., Wang, M., Yuan, L., Wang, Z., 2014. Butachlor causes disruption of HPG and HPT axes in adult female rare minnow (*Gobiocypris rarus*). *Chem. Biol. Interact.* 221, 119–126.