Petrological and geochemical constraints on the origin of apatite
ores from Mesozoic alkaline intrusive complexes, Central High-
Atlas, Morocco

Muhammad Ouabid^{1,2,*}, Otmane Raji¹, Jean-Marie Dautria³, Jean-Louis Bodinier^{1,3},
Fleurice Parat³, Hicham El Messbahi⁴, Carlos J. Garrido², Youssef Ahechach^{1,5}

1. Mohammed VI Polytechnic University, Geology & Sustainable Mining, Hay Moulay Rachid, 43150,
Benguerir, Morocco

2. Instituto Andaluz de Ciencias de la Tierra, Consejo Superior de Investigaciones Científicas–
Universidad de Granada, Avenida de las Palmeras 4, 18100 Armilla, Granada, Spain

3. Géosciences Montpellier, Université de Montpellier, CNRS, Cc 60, Place Eugène Bataillon, 34095
Montpellier Cedex 05, France

4. Département de Géologie, Faculté Polydisciplinaire, Université Sidi Mohamed Ben Abdellah, Route
d'Oujda, P.O. Box 1223, Taza, Morocco

5. Département de Géologie, Faculté des Sciences, Université Moulay Ismail, P.O. Box 11201 Zitoune,
Meknès, Morocco

*Corresponding author: Muhammad Ouabid, muhaouabid@gmail.com

Abstract

Five main apatite deposits from the Moroccan Central High-Atlas—including Anemzi, Tirrhist-Inouzane, Tassent, Ait-Daoud-Toumliline and Tasraft— have been investigated through integrated field, petrographic, mineralogical, and geochemical studies. The apatite ore occurs as veins, few mm to several cm thick, spatially associated with syenite–quartz monzonite (dominant) to monzodiorite–gabbrodiorite bodies emplaced during the Atlas Jurassic-Cretaceous alkaline magmatic event. Gem-quality apatite crystals, up to 15 cm in size, are associated with albite, clinopyroxene (hedenbergite–augite–diopside), amphibole (hornblende–edenite–actinolite), K-feldspar, quartz, magnetite, titanite, epidote, prehnite, and calcite. Two types of apatite have been distinguished according to their halogen contents: F-rich and Cl-rich apatites. Both types are enriched in REE (up to 2 wt.% Σ REE), and have major and trace element contents consistent with a magmatic origin, corroborated by the chemical similarity with accessory apatite in the host alkaline intrusions, as well as with other reference igneous apatites. However, hydrothermal fluids percolating through country sedimentary rocks were also involved, as indicated by the striking Na enrichment observed in the Cl-rich apatites, likely inherited from Triassic evaporites, showing close spatial relationships with the alkaline intrusions. These fluids would also account for pervasive Na-metasomatism of alkaline magmatic rocks, as observed in the wall rocks of the apatite veins.

Keywords: Moroccan Central High-Atlas, alkaline magmatic event, vein-type apatite ores, F-rich and Cl-rich apatites, magmatic origin/hydrothermal fluids.

1. Introduction

Apatite is the most ubiquitous accessory phosphorus(P)-bearing mineral found in sedimentary, igneous and metamorphic rocks. It is also sometimes a major rock-forming mineral in some peculiar rock types: e.g. sedimentary phosphorites and some igneous rocks of alkaline, carbonatitic and anorthositic affinities (e.g., McConnell 1973; Glenn et al., 1994; Dymek and Owens 2001; Martin and Rakovan, 2013; Duchesne and Liégeois, 2015; Pufahl and Groa et al., 2017). Apatite is the only major potential resource of P used in the manufacturing of e.g., nitrogen-phosphorus-potassium (NPK) fertilizers, animal feed supplements, ceramics, pharmaceuticals, textiles and batteries. The sedimentary phosphorites make the largest contribution to the world production of phosphate, whereas igneous apatite rocks account for only ~ 10 % (e.g., Jasinski, 2016; Pufahl and Groa et al., 2017). The igneous phosphate ores are nevertheless economically valuable since they provide high quality phosphate concentrates with low contents of unwanted contaminants such as Cd, As, Pb, Si and Al, and they are rich in Rare Earth Elements (REE) (e.g., Webster and Piccoli, 2015; Chakhmouradian et al 2017). In terms of world resources, Morocco is the largest holder and producer of sedimentary phosphates, with reserves estimated to represent about ~ 76 % of world reserves (e.g., Jasinski, 2016, 2019; Pufahl and Groa et al., 2017; www.ocpgroup.ma). However, Morocco also bears igneous phosphate ores, especially in the High-Atlas mountain range. Over the last two decades, gem-quality apatite crystals well-appreciated by mineral collectors and known as "Moroccan High-Atlas Imilchil-Anemzi apatite" were made available on the international mineral market (e.g., <https://www.minfind.com/mineral-569173.html>; [3](https://www.mchminerals.com/fine-</p></div><div data-bbox=)

[mineral-specimens/apatite-morocco](#); [Rakovan, 2015](#)). Despite their economic and scientific interests, only one recent study ([Dumańska-Słowik et al., 2018](#)) was devoted to the Anemzi apatite, chiefly dealing with organic inclusions, and a detailed characterization of the apatite ores remains to be done. This paper provides the first description of the main apatite deposits in the High-Atlas region. We carried out an exhaustive petrographic, mineralogical, and geochemical study (major, volatile, and trace elements) of the apatites and associated minerals, and of the host lithologies. These data are used to identify suitable sources for the phosphate mineralization.

2. Geological setting

The High-Atlas (up to 4000 m elevation) constitutes a typical intra-continental Alpine orogen, forming a thrust chain resulting from the Africa–Eurasia convergence. Its growth started during the early Mesozoic with the reactivation of Pangea paleo-faults which delimited subsidence sedimentary basins, leading to the Atlantic Ocean opening at the NW Africa margin (e.g., [Stampfli and Borel, 2002](#); [Piqué et al., 2002](#); [Teixell et al., 2003](#); [Laville et al., 2004](#); [Frizon de Lamotte et al., 2008, 2009](#); [Domènech et al., 2015](#)). This chain includes several segments, such as the Western, Central and Eastern High-Atlas (Fig. 1a). In contrast with the two other segments, the Central High-Atlas corresponds to the deepest part of the orogen (Fig. 1b; e.g., [Teixell et al., 2003](#); [Frizon de Lamotte et al., 2008, 2009](#); [Gouiza et al., 2010](#); [Michard et al., 2011](#)). Several works have described the lithostratigraphical sequence of this belt (Fig. 1b–c; [Charrière et al., 2005, 2009](#); [Haddoumi et al., 2002, 2010](#); [Ettaki et al., 2007](#); [Frizon de Lamotte et al., 2008, 2009](#); [Cavin et al., 2010](#); [Saura et al., 2014](#); [Charrière and Haddoumi, 2016, 2017](#); [Cavallina et al., 2018](#)). The first Mesozoic sediments are Triassic red argillites and

87 evaporites (mostly halite and gypsum) forming more than 1000 m-thick series which
 88 deposited within subsident basins and which cover unconformably a leveled
 89 Precambrian-Paleozoic basement (e.g., [Du Dresnay, 1987](#); [Piqué et al., 2002](#); [Eddif et](#)
 90 [al., 2007](#); [Ouabid et al., 2017, 2020](#); [Boukerrou et al., 2018](#); [Karaoui et al., 2021](#)). These
 91 deposits are generally interbedded with tholeiitic lava flows linking to the Central Atlantic
 92 Magmatic Province (e.g., [Marzoli et al., 2019](#) and references therein). Subsequently, a
 93 regional marine transgression led to the deposit of a series of Early Jurassic carbonates
 94 (mostly limestones and marls). During the Middle Jurassic and Lower Cretaceous,
 95 conformable red beds of continental sandstone settled, indicating the filling of basins
 96 and a marine regression. Overall, these post-rift basins were controlled by sinistral
 97 strike-slip faults, and separated by Triassic diapiric anticlinal ridges with dominant NE
 98 trending (Fig. [1b–c](#); e.g., [Laville and Pique, 1992](#); [Saura et al., 2014](#)). In the Middle
 99 Cretaceous, a new marine carbonated platform sets up (e.g., [Froitzheim et al. 1988](#);
 100 [Cavallina et al., 2018](#) and references therein), and at the end of the Cretaceous, the
 101 sediments consist of terrigenous red beds, expressing the early manifestation of the
 102 Alpine orogeny (e.g., [Laville et al. 1977](#); [Froitzheim et al. 1988](#); [Ettachfini and Andreu,](#)
 103 [2004](#); [Ettachfini et al., 2005](#); [Michard et al., 2011](#); [Cavallina et al., 2018](#)). The Central
 104 High-Atlas Mesozoic ridges are often unconformably overlain by Paleocene non-marine
 105 red bed formation which draws the synclinal geometries attributed to Late
 106 Cretaceous–Cenozoic shortening and halokinesis (Fig. [1c](#); e.g., [Charriere et al., 2009](#);
 107 [Michard et al., 2011](#)). The famous Cenozoic alkaline-carbonatite magmatic episode (~
 108 45–35 Ma) that occurred at NW Central High-Atlas Tamazight area (Fig. [1b](#)) coincides

with the African–European collisional event (e.g., [Bouabdli et al., 1988](#); [Bernard-Griffiths et al., 1991](#); [Bouabdellah et al., 2010](#)).

In addition, the core of the Central High-Atlas anticlinal ridges is formed by an important Jurassic–Cretaceous alkaline to transitional magmatism between ~ 165 Ma and ~ 125 Ma (Fig. [1b–c](#); e.g., [Hailwood and Mitchell 1971](#); [Westphal et al., 1979](#); [Beraâouz and Bonin, 1993](#); [Rahimi et al. 1997](#); [Armando, 1999](#); [Haddoumi et al., 2010](#); [Michard et al., 2011, 2013](#); [Bensalah et al., 2013](#)). This magmatism occurs as numerous dykes, sills and laccoliths, either intrusive along faults and narrow anticlinal ridges or intercalated in the thick Mesozoic sedimentary series (Fig. [1b–c](#); e.g., [Armando, 1999](#); [Teixell et al., 2003](#)). Basaltic lava flows within wide synclines are also reported (e.g., [Guezal et al., 2011](#); [Bensalah et al., 2013](#); [Michard et al., 2013](#)). The magma emplacement was mainly controlled by faults coeval with halokinetic tectonic and moderate syn-schistous deformation, might link to the widespread exhumation resulting from the orogen during the Upper Jurassic–Early Cretaceous (e.g., [Laville and Pique, 1992](#); [Frizon de Lamotte et al., 2008, 2009](#); [Michard et al., 2011](#); [Saura et al., 2014](#); [Torres-Lopez et al., 2016](#); [Essaifi and Zayane, 2018](#)). Metamorphic aureole around the intrusions is almost absent at the contact with the country sedimentary rocks, which are mostly faulted (e.g., Fig. [1b–c](#)). The tectonic boundaries are mostly vertical/sub-vertical faults marked by brecciated formation composed of fragments of igneous intrusive and carbonate rocks. However, some rare normal contacts between the intrusions and the sedimentary sequence are observed with chilled marginal magmatic rocks with carbonates recrystallization that occurred at about 300–500 °C ([Lhachmi, 1992](#), [Armando, 1999](#); [Lhachmi et al., 2001](#); [Essaifi and Zayane, 2018](#)). The

intrusive complexes include almost all terms of alkaline magma differentiation series (e.g., [Armando, 1999](#); [Lhachmi et al., 2001](#); [Zayane et al., 2002](#); [Essaifi and Zayane, 2018](#)). The intrusions are made up to 60–70% of troctolites, gabbros, and monzogabbros, while the remaining volume is represented by differentiated rocks (mostly monzonites and syenites). Depside the fragmentation of the intrusive bodies into several blocks which hampers to infer more information about the primary relationships, the differentiated unit seems to be intrusive into the mafic unit (e.g., [Armando, 1999](#); [Lhachmi et al., 2001](#); [Essaifi and Zayane, 2018](#)). The evolution of the magma sequence from mafic to felsic members would be mainly linked to fractional crystallization, whereas crustal contamination during the magma ascent cannot be excluded (e.g., [Armando, 1999](#); [Lhachmi et al., 2001](#), [Zayane et al., 2002](#); [Essaifi and Zayane, 2018](#)). The magmatism is interpreted to be formed in a transpressional tectonic setting where the Central High-Atlas experienced a significant uplift and orogenesis during the Jurassic (e.g., [Laville and Pique, 1992](#)), or in an asthenosphere mantle upwelling context beneath the orogen with partial melting of enriched upper mantle components without significant crustal extension (e.g., [Frizon de Lamotte et al., 2009](#); [Michard et al., 2013](#); [Essaifi and Zayane, 2018](#)). From our field observations, the major apatite deposits of the Moroccan Central High-Atlas —enclosing Anemzi, Tirrhist-Inougzane, Tassent-Tarstaft, Tarstaft, and Ait Daoud-Toumliline deposits, Figs. [1b](#), [c](#)— are always spatially associated to these Jurassic-Cretaceous alkaline intrusions. The apatite ores form mm- to several cm-thick veins without any preferential trending and the current mining is done only in an artisanal way and the only economic opportunities are for mineral collectors. Unfortunately, the limited extension of the exposed apatite

veins hides more information about the lateral extensions, but the scraping of the apatite ores generated by the mineral collectors could vary between 2 and 40 m long. These apatite-producing localities are isolated and less abundant. The contacts between the apatite veins and their host magmatic rocks are mostly sharp without any clear zonation and the mineralogical compositions of the vein-type apatite ores and its host rocks from different localities are slightly variable, whereas the main components are always alkali feldspars, pyroxene, amphibole, quartz, prehnite, calcite, magnetite, epidote, and titanite (detail below).

3. Sampling and analytical methods

The five Moroccan Central High-Atlas apatite deposits cited above (Fig. 1) were investigated. For each deposit, we sampled apatite mineralized veins and their direct country magmatic rocks (see Table 1). Apatite crystals have been separated by hand, included in epoxy mounts and polished. These apatites, the gangue-forming minerals, and the country rocks were also studied in polished thin-sections. For petrographical observations, we used standard microscope Leica DM2700P in the Geo-AnalyticalLab of Geology and Sustainable Mining Department, Mohammed VI Polytechnic University (UM6P, Benguerir, Morocco).

Seven representative samples of the magmatic rocks hosting the apatite mineralization veins were selected for whole-rock geochemical analyses of major elements and some trace elements performed at the Instituto Andaluz de Ciencias de la Tierra (IACT-CSIC, Granada, Spain). The samples include Anemzi AP45 and AP54, Tirrhist-Inouzane AP10, Ait Daoud-Toumliline AP08 and AP09, Tassent AP27 and

Tasraft AP61 (Table 1). Rock powders were obtained using a jaw crusher and an agate ring mill. Loss on ignition (L.O.I) was determined by drying the samples at 900°C, and ranges from 0.07 to 5.86 wt.%. (Cl, Cr, Co, Ni, Cu, Zn, Ga, Zr, Sr, Y, Nb) The analyses were performed using a sequential spectrometer Bruker AXS S4 Pioneer equipped with three analyzers (LiF200, OVO-55, PET). For X-ray fluorescence (XRF) major element, Cr, Ni, Zr, Sr and Y analyses, rock powders (c. 1 g) were weighed with di-lithium tetraborate flux, and then the mixture was fused at 1000 °C for 15 min. Within run precision (% RSD), measured by repeated analyses of USGS BHVO-2 and AGV-2 reference materials as external standards was better than 1.5% for all the elements except P (2.7%) (Varas-Reus et al., 2018). The other minor elements and SO₃ were analysed in pressed powder pellets in an independent calibration. The analyses of certified reference materials (BIR-1) have been used as external standard and the accuracies are <5% for major elements and <11% for minor elements.

In situ analyzes of major and minor elements for minerals were carried out using the Cameca SX100 Electron Microprobe at the Geosciences Montpellier (Montpellier, France). Analyses were obtained using a beam size of 30–40 µm for apatite and 5 µm for the minerals of gangue and country rocks, an accelerating voltage of 15 kV and a probe current of 20 nA. The counting time was fixed at 20 s for major elements and 40 s for F, Cl, and S. Combination of natural—including fluorapatite (F, P), chlorapatite (Cl), orthoclase (K), albite (Al, Na), wollastonite (Si, Ca), forsterite (Mg, Fe), baryte (S), rhodochrosite (Mn)—and synthetic (TiO₂ and Cr₂O₃) standards are used for the calibrations. The apatite and the associated pyroxene and amphibole were analyzed for trace elements using the Laser Ablation Coupled Plasma-Mass Spectrometry (LA-ICP-

MS) at Geosciences Montpellier (AETE platform, OSU-OREME). Signals were acquired in Time Resolved Acquisition. Each analysis consisted of a pre-ablation pulse of 5 s and 2 min for the blank and 1 min for measurement. The laser was fired employing an energy density of 15 J/cm² at a frequency of 5 Hz and using a spot size of 50 µm. Reference material NIST612 glass (Pearce et al., 1997) was used as external standard and CaO and SiO₂ contents determined by electron probe were used as internal standards. Data were processed using the GLITTER software (Griffin et al., 2008). LA-ICP-MS data accuracy and precision were assessed using the obtained results from the BIR-1G glass and show good agreement with published values for this international reference material (Gao et al., 2002; Jochum et al., 2005, 2016).

4. Petrography

4.1. Anemzi

In the Anemzi veins, the apatites (~ 10–30 vol.% of the vein) occur as hexagonal prisms with variable sizes that can reach more than 6 cm length and more than 3 cm width (Fig. 2). They are present in two different types of gangue (Fig. 2a–b). The first one is white in color and it mainly consists of calcite, plagioclase (albite), K-feldspar, magnetite, and quartz (Figs. 2c–d). Calcite occurs as medium- to fine subhedral to anhedral crystals (~ 0.1–0.8 mm in size) and is the dominant gangue-forming mineral (up to 40 vol.%), filling the spaces between the other minerals. Albite and K-feldspar are abundant (up to 25 vol.%) euhedral crystals (~ 0.2–1 cm in size) with sericitized cores and fresh rims. Magnetite is omnipresent and occurs as red-brownish euhedral prismatic to anhedral crystals up to 0.5 cm in size. Quartz is less abundant, euhedral to

subhedral, and small in size (< 0.5 mm). The second gangue type is dark, and it is made almost exclusively of needle-shaped amphibole (mostly hornblende) with subordinate chlorite as a product of amphibole transformation (Figs. 2b, e). The most beautiful Moroccan apatite gems are extracted from veins within white gangue: they are characterized by their perfect euhedral habit, transparency, green-yellow color and glassy luster (Fig. 2a). In contrast, the apatites associated with dark gangue are cloudy crystals, pale green to grey in color, with little or no luster and enclosed many euhedral needles of amphibole similar to those present in the matrix veins (Figs. 2b, e).

The Anemzi apatite veins are mainly hosted in medium- to coarse-grained mesocratic to leucocratic magmatic rock which consists of 40 to 60 vol.% of albitic plagioclase ($An < 10\%$), 20–25 vol.% of K-feldspar and 2–8 vol.% of quartz (Fig. 2c). Chlorite and Fe-oxides are omnipresent: they result from the complete transformation of ferromagnesian minerals (i.e., pyroxene, amphibole) which are no longer present in the rock. Accessory minerals include rutile, apatite, and zircon. Calcite is also observed as secondary mineral phase.

4.2. Tirrhist-Inouzane

The Tirrhist-Inouzane veins provide gem-quality apatites (up to 15 vol.%) transparency and luster with similar color than apatites from the Anemzi region (Figs. 3a, b). The size of apatite crystals is variable but can reach 3 cm length. The gangue consists of clinopyroxene, albite, epidote, quartz, K-feldspar, and subordinate chlorite (Fig. 3c–d). Clinopyroxene occurs as euhedral to subhedral crystals (up to 1 cm in size), forming the main gangue component (up to 35 vol.%). Zonation of some clinopyroxene

crystals is highlighted by the alternance of dark and light green domains. Epidote is abundant (up to 20 vol.%) and it appears as subhedral to euhedral crystals (~ 0.2–1 cm in size). Albite (dominant), quartz and K-feldspar (rare) represent around 30 vol.% and occur as medium- to fine subhedral to anhedral crystals (~ 0.1–1 mm in size), filling the interstices among clinopyroxene, epidote, and apatite. These apatite veins crosscut medium-grained to pegmatitic leucocratic magmatic rock rich in albitic plagioclase (up to 85 vol.%) (Figs. 3a, e, f) and also contain clinopyroxene (5–10 vol.%), and subordinate (< 5 vol.%) quartz and K-feldspar. Magnetite (sometimes with ilmenite exsolutions), stilpnomelane, apatite and zircon are also observed. Chlorite, epidote and sericite occur as secondary mineral assemblage.

4.3. Tassent

The apatites (~ 10–20 vol.%) associated with the Tassent veins are elongated crystals up to 15 cm length and 4 cm wide (Figs. 4a, b). They are pale yellow to pale green with an opalescent luster. The gangue mineral assemblage consists mainly of prehnite and quartz (~ 60 vol.%), plus clinopyroxene and epidote (Figs. 4c, d). Prehnite and quartz occur as euhedral to anhedral crystals (up to 3 mm in size). Clinopyroxene appears as euhedral to subhedral destabilized crystals (up to 1 cm in size). Epidote is occasionally observed as yellow-greenish euhedral crystals (< 0.5 mm in size).

The apatite veins are hosted by medium-grained mesocratic to leucocratic magmatic rocks (Figs. 4a, e), which consists of albitic plagioclase (up to 60–70 vol.%), amphibole (15–20 vol.%), quartz (5–10 vol.%) and subordinate (< 5 vol.%) K-feldspar

and clinopyroxene. Titanite, apatite and zircon are the main accessory minerals, and epidote and chlorite occur as alteration products.

4.4. Ait Daoud-Toumliline

The apatites (10–20 vol.%) from the Ait Daoud Toumliline veins are gem-quality dark green, with moderate vitreous luster and characterized by an abundance of amphibole inclusions (Figs. 5a to 5c). Their size can reach 3 cm. They are found in a gangue composed mainly of amphibole, prehnite, quartz, epidote, magnetite, albite and K-feldspar (Fig. 5d–e). Amphibole, prehnite and quartz represent ~ 60–70 vol.% of the vein and they occur as euhedral to subhedral crystals (up to ~ 4 mm in size). Epidote is abundant and crystallized as subhedral to euhedral crystals (< 1 mm in size). Albite, K-feldspar, and magnetite are also observed as medium to fine grains.

The apatite ores crosscut medium- to coarse-grained country mesocratic to melanocratic pale magmatic rocks with a primary paragenesis associating albitic plagioclase (60–65 vol.%) and amphibole (20–35 vol.%) (Fig. 5f). Clinopyroxene is also present and is frequently associated with Fe-Ti oxides (magnetite and ilmenite). Quartz and K-feldspar are rare. Titanite, rutile, and apatite are the main observed accessory minerals, and epidote, chlorite, and prehnite occur as secondary phases.

4.5. Tasraft

Two types of apatite occur at Tasraft: (i) the first one is similar to the apatites found in Ait Daoud-Toumliline (up to 2 cm in size, dark green translucent crystals with moderate vitreous luster and often containing tiny needle-shaped amphibole inclusions,

Fig. 6a); (ii) the second one corresponds to fine crystals (less than 0.5 cm in size) of gem-quality apatite with a color, transparency and luster similar to those found in Tirrhist-Inouzane and Anemzi. The gangue of these fine apatite gems consists mainly of clinopyroxene, albite, quartz, K-feldspar, and titanite (Figs. 6b, c, d, e). As observed for Tirrhist-Inouzane, clinopyroxene (up to 45 vol.% of the vein) is often zoned and occurs as euhedral to subhedral crystals (up to 1 cm in size). Albite (dominant), K-feldspar, and quartz represent around 40 vol.% and occur as medium euhedral to anhedral crystals (~ 0.1–1 mm in size), interstitial between clinopyroxene and apatite. Titanite is present and occurs as green-yellowish to clear prismatic crystals (< 5 vol.%, up to 1 cm in size).

The host rocks of these apatite veins are coarse- to medium-grained leucocratic magmatic rocks containing up to 70–80 vol.% of albitic plagioclase, up to 10 vol.% of clinopyroxene and subordinate K-feldspar and quartz. Amphibole, magnetite, titanite, rutile, apatite and zircon are accessory minerals (Figs. 6b–c). Epidote, chlorite, and sericite also occur as secondary phases.

5. Geochemistry of magmatic host rocks

Whole-rock geochemical analyses for the magmatic facies directly hosting the Moroccan Central High-Atlas vein-type apatite ores are given in Table 2. In the TAS classification (Fig. 7a; Middlemost, 1994), these rocks have intermediate to felsic (dominant) signature with SiO₂ contents ranging from ~ 53 to 65 wt.% and plot in the fields for gabbrodiorite–monzodiorite and syenite–quartz monzonite (dominant). They belong mostly to the alkaline/sub-alkaline series with alkali (K₂O + Na₂O) content from ~ 4.6 to ~ 10 wt.%. All these facies have relatively large ranges in Fe₂O₃^{tot} (~ 2–10.3

wt.%), MgO (Mg# ~ 0.04–0.8), CaO (~ 1.7–10.6 wt.%), Al₂O₃ (14.2–16.5 wt.%), TiO₂ (~ 0.4–2.3 wt.%) and P₂O₅ (~ 0.03–0.9 wt.%). Almost all these rocks are peralkaline [mol. (Al₂O₃/CaO+Na₂O + K₂O) = ACNK = 0.51–0.94] with low K contents [K# = mol. (K₂O/Na₂O+K₂O) = 0.01–0.1], except for Anemzi samples that belong to the high-K serie (K#: 0.26– 0.36) (Fig. 7b, Table 2, e.g., Rickwood, 1989; Maniar and Piccoli, 1989). Compared to the available data from the literature for the same Central High-Atlas magmatic intrusions further away from the apatite veins (Figs. 7a–c; data from Armando, 1999; Lhachmi et al., 2001; Zayane et al., 2002; Essaifi and Zayane, 2018), the studied rocks have geochemical compositions similar to the more differentiated members and differ significantly from the more primitive members (mostly gabbros). Evaluation of the alteration effects using the discriminated diagram of Hughes (1973) shows that almost all these differentiated rocks plot toward the field of sodic series (Na₂O up to ~ 9 wt.%), suggesting that they might have been affected by general Na-metasomatism conditions (Fig. 7c). In terms of volatile element compositions, Cl and SO₃ content reaches, respectively, a maximum of 0.25 wt.% and 0.04 wt.% in the Ait Daoud-Toumliline samples and are lowest in the other samples (0.05–0.13 wt.% Cl and 0.01 wt.% SO₃) (Table 2). Besides, all investigated rock samples have moderate contents of MnO (0.03–0.12 wt.%), Co (53–155 ppm), Ni (29–133 ppm), Cu (17–55 ppm), Zn (14–50 ppm), Ga (18–47 ppm), Sr (213–407 ppm), and Y (15–103 ppm). However, the more felsic rocks have slightly high Zr (up to 748 ppm) and Nb (up to 120 ppm) concentrations, and low values of Cr (less than 32 ppm) (Table 2).

6. Mineralogy

6.1. Apatite

6.1.1. Major element composition

Table 3 shows representative major element analyses of the studied apatites. The two major constituents —i.e., CaO and P₂O₅— show rather little variations: CaO and P₂O₅ vary between 53 and 57 wt.% and between 39 and 44 wt.%, respectively. The F content of the Anemzi apatite associated with calcite, feldspars, and magnetite matrix (white gangue; see above) is quite homogenous with about 2 wt.% and low Cl content (0.67–0.90 wt.%). In contrast, the Anemzi apatites associated with dark gangue (amphibole matrix), are rather heterogeneous: their F contents decrease from core to rim (from ~ 1.9 to ~ 0.8 wt.%) while Cl increases (from ~ 0.6 to ~ 2.5 wt.%) (Table 3). All analyzed Tirrhist-Inouzane apatites have relatively homogeneous F and Cl contents (~ 1.9–2.4 and ~ 0.34–0.47 wt.%, respectively). Similarly, the apatites from Tassent and Tasraft (fine gems) have F concentrations between 1.7 and 2.8 wt.% and low to moderate Cl contents (0.5–1.47 wt.%). On the other hand, distinct apatite compositions have been yielded by amphibole-bearing apatites from both Tasraft and Ait Daoud-Toumliline: the first ones show compositions intermediate between those of Cl-rich apatites and those of Cl-F apatites (F contents vary between 0.47 and 1.47 wt.% and Cl between 3.44 and 1.5 wt.%); the second ones are Cl-rich apatites with high Cl (up to 3.3 wt.%) and low F (~ 0.6–1 wt.%) contents. If we considered all apatites analyses, they show a good linear trend and progressive compositional variations from the F-rich to Cl-rich apatites as shown by Fig 8a.

In all studied apatites, the element substitutions are always very low, and they vary between the different apatite deposits and even between the crystals of the same deposit (see Table 3 and Figs. 8b, c, d). The major substituent element is Si ($0.1 < \text{SiO}_2$ wt.% < 0.8). Na_2O contents can reach 0.55 wt.%, whereas FeO^{tot} and MgO never exceed 0.11 and 0.08 wt.%, respectively. The S contents are particularly variable: SO_2 varies between 0 and 0.57 wt.%. In detail, the amphibole-bearing apatites from Ait Daoud-Toumline and Tasraft show quite similar compositions with relatively high Na_2O and MgO contents (up to 0.57 wt.% and 0.9 wt.%, respectively), FeO^{tot} contents between 0.04 and 0.11 wt.% and low concentrations of SiO_2 (< 0.4 wt.%) and SO_2 (~ 0 –0.3 wt.%). Similar FeO^{tot} and SiO_2 abundances (0.04–0.10 wt.% and 0.18–0.40 wt.%, respectively) have been measured in Tassent apatite but with generally low Na_2O (0.06–0.29 wt.%) and SO_2 (< 0.04 wt.%) contents and no MgO . The apatites from Tirrhist-Inouzane and Tasraft (fine gems) have MgO below detection limit and they generally contain similar low Na_2O (≤ 0.08 wt.%) and FeO^{tot} (≤ 0.04 wt.%) concentrations, whereas SiO_2 (up to 0.76 wt.%) and SO_2 (0.30–0.57 wt.%) abundances are relatively high. The Anemzi apatites from both white and dark gangues have comparable Na_2O (0–0.15 wt.%), MgO (≤ 0.01 wt.%) and FeO^{tot} (0.02–0.09 wt.%) concentrations. However, these two types of apatite are distinguished by their SiO_2 and SO_2 contents: the apatites from the dark gangue have SiO_2 (0.12–0.32 wt.%) and SO_2 (0–0.04 wt.%) significantly lower than those of the white gangue apatites (0.62–0.70 and 0.01–0.42 wt.%, respectively).

In addition, the rim–core chemical variations (e.g., SiO_2 , FeO^{tot} , Na_2O , Cl, and F) in some apatite crystals of veins are shown in Figs. 8e–h and Table 3. Despite their

large size, almost all fluorapatites —e.g., Tirrhist-Inouzane and Anemzi (white gangue)— have no chemical zoning. In contrast, the Anemzi amphibole-bearing apatites (dark gangue) display F-rich cores and Cl-rich rims and the SO₂ contents decrease from core to rim while Na₂O contents increase (Table 3).

To further constrain the relationships between the apatites and their magmatic host rocks, the apatites in Tirrhist-Inouzane syenites have also been analyzed (see Table 3). These igneous apatite grains have similar compositions compared to the Tirrhist-Inouzane, Anemzi (white gangue), Tassent and Tasraft (fine gems) apatite crystals in veins in terms of F (2–2.3 wt.%), Cl (0.5–0.8 wt.%), Na₂O (0.1–0.19 wt.%), MgO (< 0.01) and FeO^{tot} (0.04–0.09 wt.%), and they are clearly distinguished from both Ait Daoud-Toumliline and Tasraft amphibole-bearing apatite gems in veins as shown by Figs. 8a–c and Table 3. On the other hand, SiO₂ and SO₂ abundances in the syenite apatites are similar to those found in the gemmy apatites in veins from Tassent and Anemzi (dark gangue) (SO₂ < 0.04 wt.%, SiO₂ ~ 0.35 wt.%) (Fig. 8d).

6.1.2. Trace element composition

The trace element abundances of studied apatites given in Table 3 show that these minerals can incorporate a wide range of trace elements and often in large quantities. The most abundant trace elements in the Central High-Atlas gem-quality apatites in veins are REE (mainly La, Ce and Nd), Sr, Y, V, Mn, Th and U, whereas Sc, Co, Rb, Zr, Nb, Ba and Pb contents are very low (< 6 ppm) (Fig. 9 and Table 3). The highest REE concentrations are observed in the Ait Daoud-Toumliline apatites (Σ REE: 18190–22587 ppm), with variable abundances in the Anemzi (white gangue) and

Tasraft amphibole-bearing apatites (ΣREE : 3846–17432 ppm), moderate contents in Tassent apatites (ΣREE : 13140–13554 ppm) and low to moderate values (ΣREE : 4823–8495 ppm) in Tirrhist-Inouzane, Anemzi (dark gangue) and Tasraft (fine gems) apatites. Figs. 10 and 11 show, respectively, their REE patterns and multi-element spider diagrams normalized to chondrite abundances from McDonough and Sun (1995). In all apatites, a pronounced negative Eu-anomaly can be observed ($0.11 < \text{Eu}/\text{Eu}^* < 0.63$; Fig. 10). Almost all analyses show consistent enrichment in light REE relative to heavy REE with $(\text{Ce}/\text{Yb})_n$ ratios ranging from 12 to 78.

The V contents of apatites vary greatly between the different deposits. They are very low (between 3 and 4 ppm) in the Tassent apatites and slightly higher (between 18 and 34 ppm) in the Tirrhist-Inouzane and high in Ait Daoud-Toumliline (55–77 ppm) (Fig. 9a). We also find these high values but with variability in both Tasraft apatite types: 3–128 ppm in the amphibole-bearing apatites and 96–146 ppm in the fine apatite gems. In Anemzi, the V contents of the apatites from the dark gangue are low but relatively constant (9–14 ppm), while these from the white gangue are surprisingly variable and they can even reach high values (0.17–128 ppm) as in Tasraft. The Mn contents are very variable in the Anemzi apatites from the white gangue (56–397 ppm), while they are relatively stable in those from the dark gangue (274–312 ppm) (Fig. 9b). These latter values are relatively close to that found in the Tassent apatites (222–241 ppm Mn). The two types of apatite from Tasraft have Mn contents significantly different: 63–96 ppm for the fine gem-crystals and 138–250 ppm for the Amp-bearing crystals. The Mn contents of the apatites from Tirrhist-Inouzane (140–148 ppm) and Ait Daoud-Toumliline (116–131 ppm) are homogeneous and quite similar. Strontium

concentrations are moderate to high in all studied apatites (Fig. 9c): 407–686 ppm in Anemzi white gangue, 673–792 ppm in Anemzi dark gangue, 763–781 ppm in Tirrhist-Inouzane, 720–730 ppm in Tassent, 382–764 ppm in Tasraft and 386–483 ppm in Ait Daoud-Toumliline. Yttrium concentration ranges from 230 ppm to 4118 ppm (Fig. 9e): 1567–1672 ppm in Tassent, 1454–1692 ppm in Ait Daoud-Toumliline, 230–4118 ppm in Anemzi (white gangue), 425–947 ppm in Anemzi (dark gangue), 358–696 ppm in Tirrhist-Inouzane and 240–979 ppm in Tasraft. Thorium concentrations are moderate to high and usually ranges from 76 ppm to 530 ppm (Fig. 9f): 90–227 ppm in Anemzi white gangue, 79–112 ppm in Anemzi dark gangue, 140–182 ppm in Tirrhist-Inouzane, 76–83 ppm in Tassent, 63–253 ppm in Tasraft and 183–253 ppm in Ait Daoud-Toumliline. Uranium concentrations vary generally from 1.5 to 26 ppm (Fig. 9g). Rather similar high U values (16–26 ppm) are found in the apatites from the Anemzi (white gangue), Tirrhist-Inouzane and Tasraft (fine gems). The lowest U contents (1.5–7 ppm) are found in Tassent and mostly in amphibole-bearing apatites from Anemzi (dark gangue) and in these from Ait Daoud-Toumliline and Tasraft.

In addition, if we considered all apatites analyses from all studied veins, they show a rough positive correlation between total REE and Na₂O from the less to more Cl-rich apatites as shown by Fig 12. Although some compositional variations can be observed among the different apatites in veins (Figs. 9, 10, 11, 12, Table 3), their REE and multi-trace element patterns overlap and are globally parallel with those of the magmatic host rock apatites investigated here (Figs. 10–11).

6.2. The gangue minerals

6.2.1. Clinopyroxene

Clinopyroxene is exclusively observed in the F-rich apatites veins (e.g., Figs. 3, 4 and 6). The crystals are euhedral, up to 3 cm in size and dark to light green in color, sometimes with well-marked zoning corresponding to an alternation of dark and light green domains. Dark and light green grains also occur in almost all magmatic rocks hosting the apatite veins: they appear as euhedral to subhedral grains (0.5 and 2 mm, Figs. 3–4, 6). Major and trace element abundances of clinopyroxenes are given in Tables 4–5. The two major constituents —i.e., SiO_2 and CaO — show little variations; the representative values are $\sim 49\text{--}54$ wt.% SiO_2 and $20\text{--}26$ wt.% CaO . In detail, all analyzed dark green clinopyroxene zones and crystals from both apatite-bearing veins and country alkaline rocks are similar and have hedenbergite to Fe-rich augite compositions ($\text{Wo}_{44\text{--}49}$, $\text{En}_{12\text{--}26}$, $\text{Fs}_{27\text{--}43}$, see Fig. 13a and Table 4; e.g., Morimoto et al., 1988) with low Mg# [$\text{Mg}/(\text{Mg} + \text{Fe}^{2+}) \sim 0.32\text{--}0.66$], high total FeO content ($\sim 16\text{--}23$ wt. %), and moderate to high Na_2O values (up to ~ 3 wt. %) and MnO (up to 1.4 wt. %). However, the light green zones and crystals, are mainly diopside in compositions ($\text{Wo}_{46\text{--}51}$, $\text{En}_{27\text{--}41}$, $\text{Fs}_{8\text{--}25}$) with high Mg# $\sim$ (up to 0.95), and low total FeO content ($\sim 5\text{--}14$ wt. %), MnO ($\sim 0.1\text{--}0.4$ wt. %) and Na_2O values (mostly < 2 wt. %). All these clinopyroxene types have similar variable compositional range, with, Al_2O_3 ($\sim 0.1\text{--}2.6$ wt. %), TiO_2 (mostly < 1 wt. %) and Cr_2O_3 ($0\text{--}0.07$ wt. %) contents.

Fig. 13b shows REE patterns normalized to chondrite values (McDonough and Sun, 1995) of clinopyroxenes from Tasraft and Tassent F-rich apatite veins, and

Tirrhist-Inouzane host syenites (Table 5). All clinopyroxene REE concentrations are 10–160 times enriched relative to chondrite abundances ($\Sigma\text{REE} \sim 80\text{--}147$ ppm). They have parallel REE patterns [with slight high values in light REE in clinopyroxene from Tasraft apatite veins with $(\text{La}_n/\text{Yb}_n)$ up to 1.3] and show strongly developed negative Eu anomaly ($\text{Eu}/\text{Eu}^* \sim 0.19\text{--}0.37$). General high depletion for Cu, Rb, Ba, Nb, Ta, Pb, Th, and U (< 0.07 ppm) is present for all analyzed clinopyroxenes. They have generally $\sim 3\text{--}12$ ppm Sc, $9\text{--}23$ ppm Co, $\sim 17\text{--}80$ ppm Sr, $\sim 26\text{--}98$ ppm Zn, and $\sim 23\text{--}166$ ppm V.

6.2.2. Amphibole

Amphibole occurs as main gangue mineral in Cl-rich apatites veins (e.g., Ait-Daoud-Toumliline; Fig. 5) and as inclusions in almost all the Cl-rich apatites (e.g., Ait-Daoud-Toumliline, Tasraft and Anemzi; Figs. 2, 5–6). It forms euhedral and acicular green to brown crystals up to 2 cm in size. Amphibole also occurs as euhedral to subhedral grains ($\sim 0.2\text{--}1$ mm in size) in magmatic host rocks with colors similar to that of amphibole from apatite veins (e.g., Figs. 4–5). Table 4 shows representative major element compositions of the amphiboles. All amphiboles from apatite veins and their host rocks are calcic with high Ca content ($\sim 1.67\text{--}2.10$ atoms per formula unit = a.p.f.u), Si ($\sim 6.87\text{--}7.78$ a.p.f.u) and Mg ($\text{Mg}\# \sim 0.60$ to 0.90) with moderate Fe (mostly $\sim 0.54\text{--}1.20$ a.p.f.u), low to moderate (Na+K) values ($0.18\text{--}0.90$ a.p.f.u) and low Al concentrations ($\sim 0.1\text{--}1$ a.p.f.u), characteristics of magnesiohornblende-actinolite and edinite association (Fig. 13c; e.g., Leake et al., 1997). The amphibole trace element abundances have been analyzed for Anemzi apatite veins (dark gangue) and are given in Table 5. The REE abundance in amphiboles range from 46 to 101 ppm. In general,

their chondrite-normalized REE patterns are remarkably constant (Fig. 13d): they all display a gentle shape, flat pattern ($\text{La}_n/\text{Yb}_n \sim 0.7\text{--}1$) with a pronounced negative Eu anomaly ($\text{Eu}/\text{Eu}^* \sim 0.4\text{--}0.5$) similar to those of clinopyroxenes (Fig. 13c). They also have 32–650 ppm Cr and 364–405 ppm V, and Sc, Co, Zn, Sr, Y, Zr, and Nb abundances are less than 68 ppm, with low values (< 2 ppm) for Cu, Rb, Ba, Ta, Pb, Th, and U (Table 5).

6.2.3. Feldspars

Feldspars are commonly present in all studied vein-type apatite ores. Table 6 displays representative feldspar analyses. In all apatite ores and magmatic host rocks (Figs. 2–6), albite is the most abundant feldspar: the crystals are euhedral to subhedral and of centimeter size (up to 5 cm). Their compositions are similar, and they correspond almost pure to albite ($\text{Ab} \sim 90$ to 100). K-feldspar is also present but less abundant than albite: the crystals are euhedral to subhedral in the veins, subhedral to anhedral in magmatic host rocks and with sizes varying between 0.1 and 1 cm. Their compositions are between Or_{94} and Or_{99} . Anorthoclase compositions ($\text{Ab} \sim 70\text{--}75$, $\text{An} \sim 1\text{--}3$, $\text{Or} \sim 23\text{--}29$) are exclusively found in the magmatic host rocks (see Table 6).

6.2.4. Other minerals

Magnetite is found in all apatite veins: the crystals are euhedral to subhedral with up to 1 cm in size. This oxide is also observed in magmatic host rock as subhedral to anhedral grains up to 2 mm in size (e.g., Fig. 3). Representative compositions are reported in Table 7. Analyses of magnetite indicate moderate to high FeO^{tot} contents of

about ~ 91–95 wt.% in the apatite veins. In the magmatic host rock, the magnetite composition varies from magnetite ($\text{FeO}^{\text{tot}} \sim 90\text{--}95$ wt.%) to Ti-magnetite [$\text{FeO}^{\text{tot}} \sim 87$ wt.% and $\text{TiO}_2 \sim 3$ wt.%], sometimes with ilmenite exsolutions ($\text{TiO}_2 \sim 45\text{--}54$ wt.% and $\text{FeO}^{\text{tot}} \sim 46\text{--}50$ wt.%) (Table 7).

Titanite is also present in both apatite veins and host rocks and the analyses are reported in Table 7. It is associated with apatite veins and forms greenish yellow euhedral crystals up to 2 cm in size; in magmatic host rocks, it occurs as dark brown to colorless euhedral to subhedral crystals (0.5–2 mm). Titanite from veins has relatively homogeneous contents in SiO_2 , TiO_2 , and CaO (~ 29.5, ~ 41.5 and 29 wt.% respectively), whereas in the host rocks the titanite compositions are more variable: 20–32 wt.% SiO_2 , 29–54 wt.% TiO_2 and 19–29 wt.% CaO. On the other hand, rutile is observed and analyzed exclusively in the magmatic host rock, forming small euhedral to subhedral dark-red grains (up to 0.2 mm in size) and its analyses are also given in Table 7.

In the apatite mineralized veins, epidote (pistachite) occurs as green to greenish yellow euhedral crystals up to 3 cm in size, whereas in the alkaline host rocks it appears as yellow subhedral to anhedral grains of small size (0.02–0.5 cm). As shown in Table 8, the epidote compositions are similar in veins and their host rocks: 36–38 SiO_2 wt.%, 18.50–22.70 Al_2O_3 wt.%, 21–25 CaO wt.%, with slight variation FeO^{tot} content (~ 10–18 wt.%).

Prehnite is observed in the Tassent and Ait Daoud-Toumliline apatite mineralized veins. It occurs as radiating fibrous aggregates and is commonly translucent with

colorless to white greenish color (up to 4 cm in size, e.g., Figs. 4–5). Prehnite can occur also in the magmatic host rocks where it appears both as veinlets and as microcavity filling products. Analyses of this calcic mineral (~ 28 wt.% CaO) are shown in Table 8 and are homogenous and similar in both mineralized veins and host alkaline rocks.

7. Discussion and conclusions

According to our field observations, and petrological, mineralogical, and geochemical study, all Moroccan Central High-Atlas vein-type apatite ores (mostly gem-quality) are hosted by differentiated igneous intrusions ranging from syenite–quartz monzonite (dominant) to monzodiorite–gabbrodiorite (Fig. 7 and Table 2). The spatial association with Mesozoic (Jurassic-Cretaceous) alkaline/transitional rocks is typical of phosphates derived from silicate alkaline–carbonatite complexes (e.g., Simandl and Paradis, 2018 and references therein). However, carbonatites have not been found in the studied area, although the ore bodies are not so far to the well-known Tamazeght Eocene carbonatite complex (see Fig. 1b).

Our new data lead us to distinguish two types of apatite ores: (i) high gem-quality F-rich apatites associated with clinopyroxene (diopside to hedenbergite-augite) and (ii) gemmy Cl-rich apatites mainly formed in an amphibole matrix (hornblende–edenite to actinolite) (Figs. 2–6, 8, Table 3). Both ore bodies are also associated with feldspars (albitic plagioclase dominant and orthoclase), quartz, magnetite, titanite, epidote, calcite, and prehnite. All studied apatites of both deposit types have similar REE and multi-trace element patterns which suggest that they could have a relatively close origin (Figs. 9, 10, 11, Table 3). A comparison between the Central High-Atlas apatites and those of

549 the well-known Sweden Kiruna-type iron ores which formed by late-stage of magmatic
 550 differentiation of felsic magmas (pink field in Fig. 10; data from Frietsch and Perdahl,
 551 1995) shows that the REE compositions of apatites from both deposits are similar with
 552 pronounced negative Eu anomaly. Conversely, they differ from the REE patterns of the
 553 worldwide carbonatite apatites from the Oka (Canada), Kaiserstuhl (Germany),
 554 Jacupiranga (Brazil), Fen (Norway), Sokli (Finland), Hörningsholmen-Alnö (Sweden),
 555 Siilinjärvi (Finland) and Loolekop-Phalaborwa (South Africa) complexes (green field in
 556 Fig. 10; data from Ingrid, 1998; Chen and Simonetti, 2013; Doroshkevich et al., 2018;
 557 Decrée et al., 2020) (Eu/Eu^* up to 0.63 in our apatites and around 1 in worldwide
 558 carbonatite apatites). The pronounced Eu negative anomaly exhibited by all the Central
 559 High-Atlas apatites studied strongly suggests that feldspar crystallized from phosphate-
 560 rich melt prior to apatite crystallization. Such negative Eu anomalies in apatite have also
 561 been reported in fenites ($\text{Eu}/\text{Eu}^* \sim 0.73\text{--}0.84$, e.g., Decrée et al., 2020) as a result of
 562 cooling and crystallization of carbonatitic and alkaline magmas in the crust and the
 563 presence of alkali-rich fluids that metasomatize the surrounding country rocks (e.g.,
 564 Elliott et al., 2018 and references therein). Although many of the mineralogical features
 565 observed in studied apatite veins (evidence of e.g., K-feldspar, albite, clinopyroxenes
 566 and calcic amphiboles and other accessory minerals, Figs. 2–6) and the extensive sodic
 567 metasomatism affecting the magmatic country rocks (Fig. 7) suggest that fenitization
 568 may have occurred during the crystallization of apatite-rich veins (e.g., Zharikov et al.,
 569 2007; Elliott et al., 2018). The pronounced Eu anomaly together with Sr, Y, and Mn
 570 contents (Fig. 14a–b; Belousova et al., 2002) strongly indicate that Central High-Atlas
 571 apatites are co-genetic and chemically similar to the magmatic apatites of granitoids of

Kiruna type iron ores (Fig. 14a–b). Indeed, the mineralogical and geochemical signature of the Central High-Atlas apatite vein-type ores may be related to host-alkaline magma crystallization and evolution rather than carbonatite magma. These observations are corroborated by apatite discrimination diagrams based on the concentrations of several trace elements in apatite (Figs. 14c–d; Mao et al. 2016), showing that the Central High-Atlas apatites have composition close to apatites from iron oxide copper gold (IOCG) ore deposits and unmineralized rocks (likely syenites) (Fig. 14c). Generally, the IOCG systems are clearly interpreted as origin from magmatic hydrothermal systems (mainly porphyries, e.g., Barton, 2009). In addition, the mineralogy, texture, alteration halos and geochemistry data also suggest that the Central High-Atlas vein-type apatite ores could be largely linked to the Central High-Atlas differentiated magmatic rocks (mostly syenites) (Figs. 9, 10, 11, 12, 14). Plagioclases in both veins and host-rocks indicate that both derived from sodic-rich melt or/and underwent general albitization (see above and Table 6, Fig. 7b–c). The clinopyroxene and amphibole observed in both units (veins and host rocks) refer both to Ca-rich minerals and support a common igneous origin (Table 4–5 and Fig. 13). Altogether, field observations and mineralogical and geochemical signatures of veins and host-rocks suggest that Moroccan Central High-Atlas apatite ores might be related to the magmatic processes that lead to the formation of differentiated rock (likely syenites), although hydrothermal/continental contamination may have also been involved.

The absolute age of these apatite deposits is elusive, but they are observed in veins crosscutting the Jurassic-Cretaceous (ca. 165–125 Ma) alkaline igneous rocks of the Central High-Atlas and are probably coeval or post-date the crystallization of the

magmatic intrusions. The geochemical observations suggest that the crystallization environment of the high quality F-rich apatite ores (e.g., Figs. 2–3) remained chemically stable and that the intensive parameters controlling their crystallization process did not fluctuate (Fig. 8, Table 3). In contrast, the Cl-rich apatite ores enclosing intense randomly distributed amphibole inclusions in apatite gems indicate slight fluctuation of magma and/or fluid compositions. The presence of low temperature minerals (albite, quartz, epidote, prehnite, K-feldspar, magnetite and calcite) in both apatite deposits and in their host alkaline rocks strongly suggest that crystallization occurred during late magmatic stages and that the main geochemical signatures of apatite are most probably related to crystallization of alkaline magma rich in P and incompatible elements such as F (e.g., Fig. 8). It is also important to note that interactions of evolved magma with country sedimentary formations rich in Cl, Na and Ca most probably participated to the Cl-Na enrichment of apatites cannot be ignored. In Fig. 15, we provide the suitable fluids involved during the formation of the Central High-Atlas apatite mineralization. In the igneous systems the apatite is initially enriched in F owing to the high apatite–melt partition coefficient for F relative to Cl (e.g., Harlov, 2015; Webster and Piccoli, 2015). Throughout progressive crystallization of the magmas, the F contents decrease due to the crystallization of F-rich apatite resulting in progressive Cl enrichment of apatite (e.g., Fig. 8a and Table 3), Anemzi amphibole-bearing apatites being an intermediate case between the purely F-rich (e.g., Anemzi white gangue, Tirrhist-Inouzane, and Tassent) and Cl-rich apatite deposits (e.g., Tasraft and Ait Daoud-Toumliline). As mentioned above, the genesis of the Cl-rich apatites could also result from contamination of magma by Cl-enriched hydrothermal fluids due to

percolation through Triassic evaporates (mainly halite) buried beneath the Moroccan Central High-Atlas belt (Fig. 1c). This might be related to halokinetic tectonic on the Central High-Atlas magmatic intrusions emplacement (e.g., [Michard et al., 2011](#); [Saura et al., 2014](#)) and a general Na-metasomatism/fenitization associated with the emplacement of the alkaline magmatic rocks directly hosting these vein-type apatite ores (see Fig. 7; e.g., [Zirner et al., 2015](#); [Essaifi and Zayane, 2018](#)). The presence of solid and fluid organic inclusions in Anemzi gemmy apatites ([Dumańska-Słowik et al., 2018](#)), also supports a contamination of magmas by country sedimentary rocks. In addition, the hydrothermal fluids could also percolate through Mg-rich rocks (likely gabbros) present in the same area as attested by the presence of MgO in Cl-apatites (e.g., Fig. 8, Table 3). Altogether, this study highlighted the role of magma/rock and fluid/rock interactions and the complex chemical reactions involved during magma emplacement and the genesis of phosphate ores deposits. Further detailed field, petrological, structural, geochemical, geochronological and isotope investigations are however necessary in order to better quantified the igneous and sub-solidus processes as well as the exchange reactions and environmental conditions that lead to igneous phosphate ores deposits and the crystallization of gem-quality apatite crystals.

Acknowledgments

We would like to thank Dr. Franco Pirajno for editorial handling, and Dr. Kathryn M. Goodenough for the reviews that helped to clarify and improve some parts of the manuscript. This paper was funded in the frame of a master agreement between the OCP Group and Mohammed VI Polytechnic University (UM6P, Morocco). The authors wish to thank Bassou Zayi (UM6P-Geology & Sustainable Mining & UM6P–OCP Geo-

Analytical Lab) and Christophe Nevado and Doriane Delmas (Géosciences Montpellier) for their help in sample and thin section preparations.

References

Armando, G., 1999. Intracontinental alkaline magmatism: geology, petrography, mineralogy and geochemistry of the Jbel Hayim massif (central high Atlas Morocco). *Mém. Géol. l'Université Lausanne* 31, 1–106.

Barton, M.D., 2009. IOCG deposits: A Cordilleran perspective. In *Proc. 11th Biennial SGA Meeting*, pp. 5–7.

Belousova, E.A., Griffin, W.L., O'Reilly, S.Y., and Fisher, N.I., 2002, Apatite as an indicator mineral for mineral exploration: Trace-element compositions and their relationship to host rock type: *Journal of Geochemical Exploration* 76, 45–69.

Bensalah, M.K., Youbi, N., Mata, J., Madeira, J., Martins, L., El Hachimi, H., Bertrand, H., Marzoli, A., Bellieni, G., Doblas, M., Font, E., Medina, F., Mahmoudi, A., Beraâouz, H., Miranda, R., Verati, C., De Min, A., Ben Abbou, M., Zayane, R., 2013. The Jurassic-Cretaceous basaltic magmatism of the Oued El Abid syncline (High Atlas, Morocco): physical volcanology, geochemistry and geodynamic implications. *J. Afr. Earth Sci.* 81, 60–81.

Beraâouz, E.H., Bonin, B., 1993. Magmatisme alcalin intra-continentale en contexte de décrochement : le massif plutonique mésozoïque de Tirrhist, Haut Atlas (Maroc). *Comptes Rendus l'Academie Sci. Paris* 317, 647–653.

- 661 Bernard-Griffiths, J., Fourcade, S., Dupuy, C., 1991. Isotopic study (Sr, Nd, O and C) of
 662 lamprophyres and associated dykes from Tamazert (Morocco): crustal
 663 contamination processes and source characteristics. *Earth Planet. Sci. Lett.* 103,
 664 190–199.
- 665 Bouabdellah, M., Hoernle, K., Kchit, A., Duggen, S., Hauff, F., Klugel, A., Lowry, D.,
 666 Beaudoin, G., 2010. Petrogenesis of the Eocene Tamazert Continental Carbonatites
 667 (Central High Atlas, Morocco): Implications for a Common Source for the Tamazert
 668 and Canary and Cape Verde Island Carbonatites. *Journal of Petrology* 51, 1655–
 669 1686.
- 670 Bouabdli, A., Dupuy, C., Dostal, J., 1988. Geochemistry of Mesozoic alkaline
 671 lamprophyres and related rocks from the Tamazert massif, High Atlas (Morocco).
 672 *Lithos* 22, 43–58.
- 673 Boukerrou, S., Nalini, H., Moreira, H., Maacha, L., Zouhair, M., Outhounjite, M.,
 674 Ouirouane, S., Hibti1, M., Touil, A., 2018. Geochronology and geochemistry of
 675 Ediacaran volcanic rocks of the Tighardine ore deposit formation (western High
 676 Atlas, Morocco), *Arab. J. Geosci.*, 11: 22.
- 677 Cavallina, C., Papinia, M., Moratti, G., Benvenuti, M., 2018. The late Mesozoic evolution
 678 of the Central High Atlas domain (Morocco): Evidence from the paleo-drainage
 679 record of the Adrar Aglalag syncline. *Sedimentary Geology* 376, 1–17.
- 680 Cavin, L., Tong, H., Boudad, L., Meister, C., Piuze, A., Tabouelle, J., Aarab, M., Amiot,
 681 R., Buffetaut, E., Dyke, G., Hua, S., Le Loeuff, J., 2010. Vertebrate assemblages

from the early Late Cretaceous of southeastern Morocco: An overview. J. Afr. Earth
Sci. 57, 391–412.

Chakhmouradian, A. R., Reguir, E. P., Zaitsev, A. N., Couëslan, C., Xu, C., Kynický, J.,
Mumin, A.H., Yang, P., 2017. Apatite in carbonatitic rocks: Compositional variation,
zoning, element partitioning and petrogenetic significance. Lithos, 274, 188–213.

Charrière, A., Haddoumi, H., 2016. Les “couches rouges” continentales jurassico-
crétacées des Atlas marocains (Moyen Atlas, Haut Atlas central et oriental): bilan
stratigraphique, paléogéographies successives et cadre géodynamique. Boletín
Geológico y Minero 127, 407-430.

Charrière, A., Haddoumi, H., 2017. Dater les couches rouges continentales pour définir
la géodynamique atlasique. Géologues 194, 29–32.

Charrière, A., Haddoumi, H., Mojon, P.O., 2005. Découverte de Jurassique supérieur et
d'un niveau marin du Barrémien dans les “couches rouges” continentales du Haut
Atlas central marocain : implications paléogéographiques et structurales. Comptes
Rendus Palevol 4, 385–394.

Charrière, A., Haddoumi, H., Mojon, P.O., Ferriere, J., Cuhe, D., Zili, L., 2009. Mise en
évidence par ostracodes et charophytes de l'âge paléocène des dépôts discordants
sur les rides anticlinales de la région d'Imilchil (Haut Atlas, Maroc); conséquences
paléogéographiques et structurales. Comptes Rendus Palevol. 8, 9–19.

- 701 Chen, W., Simonetti, A., 2013. In-situ determination of major and trace elements in
702 calcite and apatite, and U–Pb ages of apatite from the Oka carbonatite complex:
703 Insights into a complex crystallization history. *Chemical Geology*, 353, 151–172.
- 704 Decrée, S., Savolainen, M., Mercadier, J., Debaille, V., Höhn, S., Frimmel, H., Baele,
705 J.M., 2020. Geochemical and spectroscopic investigation of apatite in the Siilinjärvi
706 carbonatite complex: Keys to understanding apatite forming processes and
707 assessing potential for rare earth elements. *Applied Geochemistry*, 123, 104778.
- 708 Domènech, M., Teixell, A., Babault, J., Arboleya, M.L., 2015. The inverted Triassic rift of
709 the Marrakech High Atlas: A reappraisal of basin geometries and faulting histories.
710 *Tectonophysics* 663, 177–191.
- 711 Doroshkevich, A.G., Prokopyev, I.R., Izokh, A.E., Klemm, R., Ponomarchuk, A.V.,
712 Nikolaeva, I.V., Vladyskin, N.V., 2018. Isotopic and trace element geochemistry of the
713 Seligdar magnesiocarbonatites (South Yakutia, Russia): Insights regarding the
714 mantle evolution beneath the Aldan-Stanovoy shield. *Journal of Asian Earth*
715 *Sciences*, 154, 354–368.
- 716 Duchesne, J.C., Liégeois, J.P., 2015. The origin of nelsonite and high-Zr ferrodiorite
717 associated with Proterozoic anorthosite. *Ore Geology Reviews*, 71, 40–56.
- 718 Du Dresnay, R., 1987. Jurassic development of the region of the Atlas Mountains of
719 Morocco chronology, sedimentation and structural significance. In: Cornelius, C.D.,
720 Jarnas, M., Lehmann, E.P. (Eds.), *Geology and Culture of Morocco*. Earth Sci. Soc.
721 Libya, Eighteenth Field Conference, 1977, pp. 77–99

- 722 Dumańska-Słowik, M., Weselucha-Birczyńska, A., Heflik, W., Maksymiuk, W., Sikorska-
723 Jaworowska, M., 2018. Organic inclusions evidence, composition, and
724 cathodoluminescence behaviour for the formation conditions of fluorapatite from
725 Anemzi (Morocco). *J. Raman Spectrosc.*, 1–13.
- 726 Dymek, R.F., Owens, B.E., 2001. Petrogenesis of apatite-rich rocks (nelsonites and
727 oxide-apatite gabbro-norites) associated with massif anorthosites. *Economic Geology*
728 96, 797–815.
- 729 Eddif, A., Gasquet, D., Hoepffner, C., Levresse, G., 2007. Age of the Wirgane
730 granodiorite intrusions (Western High-Atlas, Morocco): new U-Pb constraints. *J. Afr.*
731 *Earth Sci.* 47, 227–231.
- 732 Elliott, H.A.L., Wall, F., Chakhmouradian, A.R., Siegfried, P.R., Dahlgren, S.,
733 Weatherley, S., Finch, A.A., Mark, M.A.W., Dowman, E., Dady, E., 2018.
734 Fenites associated with carbonatite complexes: a review. *Ore Geology Review* 93,
735 38–59.
- 736 Essaifi, A., Zayane, R., 2018. Petrogenesis and origin of the Upper Jurassic-Lower
737 Cretaceous magmatism in Central High Atlas (Morocco): Major, trace element and
738 isotopic (Sr-Nd) constraints. *J. Afr. Earth Sci.* 137, 229–245.
- 739 Ettachfini, E.M., Andreu, B., 2004. Le Cénomanién et le Turonien de la Plate-forme
740 Préafricaine du Maroc. *Cretaceous Research* 25, 277–302.
- 741 Ettachfini, E.M., Souhel, A., Andreu, B., Caron, M., 2005. La limite Cénomanién-
742 Turonien dans le Haut Atlas Central, Maroc. *Geobios* 38, 57–68.

- 743 Ettaki, M., Ibouh, H., Chellai, E.H., Milhi, A., 2007. Les structures "diapiriques" liasiques
744 du Haut-Atlas central, Maroc : exemple de la "ride" d'Ikerzi. *Africa Geoscience*
745 *Review* 14, 73–99.
- 746 Frietsch, R., Perdahl, J.A., 1995. Rare earth elements in apatite and magnetite in
747 Kiruna-type iron ores and some other iron ore types. *Ore Geol. Rev.* 9, 489–510.
- 748 Frizon de Lamotte, D., Leturmy, P., Missenard, Y., Khomsi, S., Ruiz, G., Saddiqi, O.,
749 Guillocheau, F., Michard, A., 2009. Mesozoic and Cenozoic vertical movements in
750 the Atlas system (Algeria, Morocco, Tunisia): An overview. *Tectonophysics* 475, 9–
751 28.
- 752 Frizon de Lamotte, D., Zizi, M., Missenard, Y., Hafid, M., El Azzouzi, M., Charriere, A.,
753 Maury, R.C., Taki, Z., Benammi, M., Michard, A., 2008. The Atlas system. In:
754 Michard, A., Saddiqi, O., Chalouan, A., Frizon de Lamotte, D. (Eds.), *Continental*
755 *Evolution: The Geology of Morocco*. Springer-Verlag, Heidelberg, 133–202.
- 756 Froitzheim, N., Stets J., Wurster P., 1988. Aspects of western High Atlas tectonics.
757 *Lecture Notes in Earth Sciences* 15, 219–244.
- 758 Gao, S., Liu, X., Yuan, H., Hattendorf, B., Günther, D., Chen, L., Hu, S., 2002.
759 Determination of forty two major and trace elements in USGS and NIST SRM
760 glasses by laser ablation-inductively coupled plasma-mass spectrometry. *Geostand.*
761 *Newsl.* 26 (2), 181–196.
- 762 Glenn, C.R., Follmi, K.B., Riggs, S.R., GBaturin, .N., Grimm, K.A., Trappe, J., Abed,
763 A.M., Galli-Garrison, C., Garrison, R.E., Ilyin, A.V., Jehl, C., Rohrllich, V., Sadaqah,

R.M.Y., Schidlowski, M., Sheldon, R.E., Siegmund., H., 1994. Phosphorus and phosphorites: Sedimentology and environments of formation. *Eclogae Geologicae Helvetiae* 87, 747–88.

Gouiza, M., Bertotti, G., Hafid, M., Cloetingh, S., 2010. Kinematic and thermal evolution of the Moroccan rifted continental margin: Doukkala-High Atlas transect. *Tectonics*, 29, TC5008.

Griffin, W.L., Powell, W.J., Pearson, N.J., O'Reilly, S.Y., 2008. GLITTER: data reduction software for laser ablation ICP-MS. In: Sylvester, P. (Ed.), *Laser Ablation-ICP-MS in the Earth Sciences*. Mineralogical Association of Canada Short Course Series 40, 204–207.

Guezal, J., El Baghdadi, M., Barakat A., Rais, J., 2011. Le magmatisme jurassique-crétacé de Béni-Mellal (Haut-Atlas central, Maroc) : géochimie et signification géodynamique. *Bulletin de l'Institut Scientifique, Rabat*, 33, 17–23.

Haddoumi, H., Charrière, A., Feist, M., Andreu, B., 2002. Nouvelles datations (Hautérivien supérieur-Barrémien inférieur) dans les “couches rouges” continentales du Haut Atlas central marocain; conséquences sur l'âge du magmatisme et des structurations mésozoïques de la chaîne Atlasique. *Comptes Rendues Palevol* 1, 259–266.

Haddoumi, H., Charriere, A., Mojon, P.O., 2010. Stratigraphie et sédimentologie des «Couches rouges» continentales du Jurassique-Cretacé du Haut Atlas central

(Maroc): implications paleogéographiques et géodynamiques. *Geobios* 43, 433–451.

Hailwood, E.A., Mitchell, J.G., 1971. Palaeomagnetic and radiometric dating results from Jurassic intrusions in south Morocco. *Geophysical Journal International*, 24, 351–364.

Harlov, D.E., 2015. Apatite: A fingerprint for metasomatic processes. *Elements*, 11(3), 171–176.

Hughes, C.J. 1973. Spilites, keratophyres, and the igneous spectrum. *Geological Magazine*, 109(06): 513–527. doi:10.1017/ S0016756800042795.

Ingrid, H.K., 1998. Rare earth elements in sövitic carbonatites and their mineral phases. *Journal of Petrology*, 39(11-12), 2105–2121.

Jasinski, S.M., 2016. Mineral commodity summaries 2016. US Geol. Surv., 124–125. http://minerals.usgs.gov/minerals/pubs/commodity/phosphate_rock/mcs-2016-phosp.pdf.

Jasinski, S.M., 2019. Mineral commodity summaries 2019. US Geol. Surv., 122–123.

Jochum, K. P., Weis, U., Schwager, B., Stoll, B., Wilson, S. A., Haug, G. H., Andreae, M. O., and Enzweiler, J., 2016. Reference values following ISO guidelines for frequently requested rock reference materials, *Geostandards Geoanalytical Research*, 40, 333–350.

- 803 Jochum, K.P., Willbold, M., Raczek, I., Stoll, B., Herwig, K., 2005. Chemical
 804 characterisation of the USGS reference glasses GSA-1G, GSC-1G, GSD-1G, GSE-
 805 1G, BCR-2G, BHVO-2G and BIR-1G using EPMA, ID-TIMS, ID-ICPMS and LA-
 806 ICPMS. *Geostandards Geoanalytical Research* 29, 285–302.
- 807 Karaoui, A., Breitzkreuz, C., Karaoui, B., Yajoui, Z., Mahmoudi, A., Zanetti, A., Langone,
 808 A., 2021. The Ediacaran volcano-sedimentary succession in the Western Skoura
 809 inlier (Central High Atlas, Morocco): facies analysis, geochemistry, geochronology
 810 and geodynamic implications. *Int. J. Earth Sci.* [https://doi.org/10.1007/s00531-021-](https://doi.org/10.1007/s00531-021-01997-y)
 811 01997-y.
- 812 Laville E., Lesage J.L., Séguret, M., 1977. Géométrie, cinématique (dynamique) de la
 813 tectonique atlasique sur le versant Sud du Haut Atlas Marocain. Aperçu sur les
 814 tectoniques hercyniennes et tardi-hercyniennes. *Bulletin de la Société Géologique*
 815 *de France* 7, 499–523.
- 816 Laville, E., Pique, A., 1992. Jurassic penetrative deformation and Cenozoic uplift in the
 817 central High Atlas (Morocco): a tectonic model. *Structural and Orogenic inversions.*
 818 *Geol. Rundsch* 81, 157–170.
- 819 Laville, E., Pique, A., Amghar M., Cherroud, M., 2004. A restatement of the Mesozoic
 820 Atlasitic Rifting (Morocco). *J. Afr. Earth Sci.* 38, 145–153.
- 821 Leake, B.E., Wooley, A.R., Arps, C.E.S., Brich, W.D., Gilbert, M.C., Grice, J.D.,
 822 Hawthorne, F.C., Kato, A., Kish, H.J., Krivovichev, V.G., Linthout, K., Laird, J.,
 823 Mandarino, J.A., Maresch, W.V., Nickel, E.H., Rock, N.M.S., Schumacher, J.C.,

- 824 Smith, D.C., Stephenson, N.C.N., Ungaretti, L., Whittaker, E.J.W., Youzhi, G., 1997.
825 Nomenclature of amphiboles: report of the Subcommittee on Amphiboles of the
826 International Mineralogical Association, Commission on New Minerals and Mineral
827 Names. *Am. Mineral* 82, 1019–1037.
- 828 Lhachmi, A., Lorand, J.P., Fabries, J., 2001. Petrologie de l'intrusion alcaline
829 mésozoïque de la région d'Anemzi, Haut Atlas Central, Maroc. *J. Afr. Earth Sci.* 32,
830 741–764.
- 831 Lhachmi, A., 1992. Pétrologie de la série alcaline et des roches magmatiques associées
832 de la région d'Anemzi (Haut Atlas central. Maroc). Doctorat du Muséum National
833 d'Histoire Naturelle, France, Paris, 156p.
- 834 Maniar, P.D., Piccoli P.M., 1989. Tectonic discrimination of granitoids. *Geol Soc Am Bull*
835 101, 635–643.
- 836 Mao, M., Rukhlov, A.S., Rowins, S.M., Spence, J., Coogan, L.A., 2016. Apatite trace
837 element compositions: a robust new tool for mineral exploration. *Economic Geology*,
838 111(5), 1187–1222.
- 839 Martin, R., Rakovan, J., 2013. The geology of apatite occurrences. In *Apatite: The great*
840 *pretender*, ed. J. Rakovan, G. A. Staebler, and D. A. Dallaire, 21–27. Mineral
841 monograph 17. Denver, CO: Lithographie.
- 842 Marzoli, A., Bertrand, H., Youbi, N., Callegaro, S., Merle, R., Reisberg, L., Chiaradia, M.,
843 Brownlee, S., Jourdan, F., Zanetti, A., Davies, J., Cuppone, T., Mahmoudi, A.,
844 Medina, F., Renne, P.R., Bellieni, G., Crivellari, S., El Hachimi, H., Bensalah, M.K.,

- 845 Meyzen, C.M., Tegner, C., 2019. The Central Atlantic magmatic province (CAMP) in
846 Morocco. *Journal of Petrology* 60, 945–996.
- 847 McConnell, D., 1973. Apatite. Its Crystal Chemistry, Mineralogy, Utilization, and
848 Geologic and Biologic Occurrences. Springer, New York, 111p.
- 849 McDonough, W.F., Sun, S.S., 1995. The composition of the Earth. *Chemical Geology*
850 120, 223–253.
- 851 Michard, A., Frizon de Lamotte, D., Hafid, M., Charriere, A., Haddoumi, H., Ibouh, H.,
852 2013. Comment on “The Jurassice Cretaceous basaltic magmatism of the Oued El-
853 Abid syncline (High Atlas, Morocco): physical volcanology, geochemistry and
854 geodynamic implications” by Bensalah et al. *J. Afr. Earth Sci.* 81, 60–81. *J. Afr.*
855 *Earth Sci.* 88, 101–05.
- 856 Michard, A., Ibouh, H., Charriere, A., 2011. Syncline-topped anticlinal ridges from the
857 high Atlas: a moroccan conundrum, and inspiring structures from the syrian arc,
858 Israel. *Terra Nova* 23, 314–323.
- 859 Middlemost, E.A.K., 1994. Naming materials in magma/igneous rock system. *Earth Sci.*
860 *Rev.* 37, 215–224.
- 861 Morimoto, N., Fabries, J., Ferguson, A.K., Ginzburg, I.V., Ross, M., Seifert, F.A.,
862 Zussman, J., Aoki, K., Gottardi, G., 1989. Nomenclature of pyroxenes.
863 Subcommittee on pyroxenes. *American Mineralogist* 73, 1123–1133.

- 864 Ouabid M., Garrido, C.J., Ouali, H., Harvey, J., Hidas, K., Marchesi, C., Acosta-Vigil, A.,
865 Dautria, J.M., El Messbahi, H., Román-Alpiste, M.J., 2020. Late Cadomian rifting of
866 the NW Gondwana margin and the reworking of Precambrian crust – evidence from
867 bimodal magmatism in the early Paleozoic Moroccan Meseta. *International Geology*
868 *Review*, doi: 10.1080/00206814.2020.1818301.
- 869 Ouabid, M., Ouali, H., Garrido, C.J., Acosta-Vigil, A., Román-Alpiste, M.J., Dautria, J.M.,
870 Marchesi, C., Hidas, K. 2017. Neoproterozoic granitoids in the basement of the
871 Moroccan Central Meseta: correlation with the Anti-Atlas at the NW paleo-margin of
872 Gondwana. *Precambrian Research* 299, 34–57.
- 873 Pearce, N.J., Perkins, W.T., Westgate, J.A., Gorton, M.P., Jackson, S.E., Neal, C.R.,
874 Chenery, S.P., 1997. A compilation of new and published major and trace element
875 data for NIST SRM 610 and NIST SRM 612 glass reference materials. *Geostand.*
876 *Newsl.* 21, 115–144.
- 877 Piqué, A., Tricart, P., Guiraud, R., Laville, E., Bouaziz, S., Amrhar, M., Aït Ouali, R.,
878 2002. The Mesozoic–Cenozoic Atlas belt (North Africa), *Geodin. Acta* 15, 159–208.
- 879 Pufahl, P.K., Groat, L.A., 2017. Sedimentary and Igneous Phosphate Deposits:
880 Formation and Exploration: An Invited Paper. *Economic Geology* 112, 483–516.
- 881 Rahimi, A., Saidi, A., Bouabdelli, M., Beraaouz, E.H., Rocci, G., 1997. Crystallization
882 and fractionation of the post-Liassic intrusive series of Tasraft (central High Atlas,
883 Morocco). *Comptes Rendus de l'Académie des Sciences*, 324, 197–203.

- 884 Rakovan, J., 2015. Connoisseur's Choice: Fluorapatite, Acushnet Quarry, Bristol
885 County, Massachusetts. *Rocks & Minerals*, 90 (3), 244–259.
- 886 Rickwood, P.C., 1989. Boundary lines within petrologic diagrams, which use oxides of
887 major and minor elements. *Lithos* 22, 247–263.
- 888 Saura, E., Verges, J., Martín-Martín, J.D., Messenger, G., Moragas, M., Razin, P.,
889 Grelaud, C., Joussiaume, R., Malaval, M., Homke, S., Hunt, D.W., 2014. Syn- to
890 post-rift diapirism and minibasins of the Central High Atlas (Morocco): the changing
891 face of a mountain belt. *J. Geol. Soc. Lond.* 171, 97–105.
- 892 Simandl, G.J., Paradis, S., 2018. Carbonatites: related ore deposits, resources,
893 footprint, and exploration methods. *Applied Earth Science* 127, 123–152.
- 894 Stampfli, G.M., Borel, G.D., 2002. A plate tectonic model for the Paleozoic and
895 Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic
896 isochrons. *Earth and Planetary Science Letters* 196, 17–33.
- 897 Teixell, A., Arboleya, M.L., Julivert, M., 2003. Tectonic shortening and topography in the
898 central High Atlas (Morocco). *Tectonics*, 22(5), 1051.
- 899 Torres-Lopez, S., Casas, A.M., Villalín, J.J., El Ouardi, H., Moussaid, B., 2016.
900 PreCenomanian vs. Cenozoic folding in the High Atlas revealed by palaeomagnetic
901 data. *Terra Nova* 28, 110–119.
- 902 Varas-Reus, M.I., Garrido, C.J., Marchesi, C., Bosch, D., Hidas, K., 2018. Genesis of
903 ultra-high pressure garnet pyroxenites in orogenic peridotites and its bearing on the

- 904 compositional heterogeneity of the Earth's mantle. *Geochimica et Cosmochimica*
905 *Acta*, 232, 303–328.
- 906 Webster, J.D., and Piccoli, P.M., 2015. Magmatic Apatite: A powerful, yet deceptive,
907 mineral. *Elements*, 11, 177–182.
- 908 Westphal, M., Montigny, R., Thuizat, R., Bardon, C., Bossert, A., Hamzeh, R., Rolley,
909 J.P., 1979. Paléomagnétisme et datation du volcanisme permien, triasique et
910 crétacé du Maroc. *Can. J. Earth Sci.* 16, 2150–2164.
- 911 Zayane, R., Essaifi, A., Maury, R.C., Pique, A., Laville, E., Bouabdelli, M., 2002.
912 Cristallisation fractionnée et contamination crustale dans la série magmatique
913 jurassique transitionnelle du Haut Atlas central (Maroc). *C.R. Geoscience* 334, 97–
914 104.
- 915 Zharikov, V.A., Pertsev, N.N., Rusinov, V.L., Callegari, E., Fettes, D.J., 2007.
916 Metasomatism and metasomatic rocks. In: Recommendations by the IUGS
917 Subcommision of the Systematics of Metamorphic Rocks. British Geological Survey.
- 918 Zirner, A.L., Marks, M.A., Wenzel, T., Jacob, D. E., Markl, G., 2015. Rare earth
919 elements in apatite as a monitor of magmatic and metasomatic processes: The
920 Ilímaussaq complex, South Greenland. *Lithos*, 228, 12–22.

Figure captions

Fig. 1. (a) Sketch map showing the location of Central High-Atlas (CHA) segment in the north Morocco. **(b)** Geological map and **(c)** NW-SE cross-section of the Central High-Atlas [adapted from [Teixell et al. \(2003\)](#), [Michard et al. \(2011\)](#) and references therein], showing the spatial relationships among the Jurassic-Cretaceous alkaline magmatism and the Mesozoic sedimentary series, and location of the studied apatite deposits.

Fig. 2. (a–b) Hand specimen and **(c–e)** microscopic photographs of the Anemzi apatite ores showing gangue- and host rock-forming minerals: **(a, c, d)** white and **(b, e)** dark gangue types. Apatite (Ap), plagioclase (Pl), K-feldspar (Kfs), calcite (Cal), magnetite (Mag), quartz (Qz), amphibole (Amp).

Fig. 3. (a, b) Hand specimen and **(c, e, f, d)** microscopic photographs of the Tirrhist-Inouzane apatite mineralization showing gangue- and host rock-forming minerals. Clinopyroxene (Cpx), epidote (Ep), stilpnomelane (Stp).

Fig. 4. (a, b) Hand specimen and **(c, d, e)** microscopic photographs of the Tassent apatite mineralization showing gangue- and host rock-forming minerals. Prehnite (Prh), titanite (Ttn).

Fig. 5. (a, b, c) Hand specimen and **(d, e, f)** microscopic photographs of the Ait Daoud-Toumliline apatite mineralization showing gangue- and host rock-forming minerals.

Fig. 6. (a) Amphibole-bearing apatite gem in mount, **(b)** hand specimen and **(c, d, e)** microscopic photographs of the Tasraft apatite mineralization (fine gems), showing gangue- and host rock-forming minerals.

Fig. 7. Chemical classification for the lithologies hosting the Central High-Atlas vein-type apatite deposits. **(a)** TAS classification diagram (Middlemost, 1994). **(b)** K_2O vs. SiO_2 diagram of Rickwood (1989). **(c)** (K_2O+Na_2O) vs. $[K_2O/(K_2O+Na_2O)]*100$ plot (after Hughes, 1973). Literature data compiled from Armando (1999), Lhachmi et al. (2001), Zayane et al. (2002) and Essaifi and Zayane (2018).

Fig. 8. (a–d) Major element concentrations in Central High-Atlas apatites. **(e–h)** Results of electron microprobe analytical profiles, and trends within images of apatite crystals corresponding to locations of these profiles.

Fig. 9. Compositional variation diagrams for the most abundant trace elements in Central High-Atlas apatites.

Fig. 10. REE patterns normalized to chondrites (McDonough and Sun, 1995) for the Central High-Atlas apatites. Worldwide carbonatite apatite data are from Ingrid (1998), Chen and Simonetti (2013), Doroshkevich et al. (2018) and Decrée et al. (2020). Iron ore apatite data are from Frietsch and Perdahl (1995).

Fig. 11. Spider diagrams normalized to chondrites (McDonough and Sun, 1995) for the Central High-Atlas apatites.

Fig. 12. ΣREE (ppm) versus Na_2O content (wt.%) in apatite from veins and host rocks.

Fig. 13. Classification and REE patterns normalized to chondrites (McDonough and Sun, 1995) of studied clinopyroxenes and amphiboles: (a, b) clinopyroxenes and (c, d) amphiboles.

Fig. 14. (a-b) Trace element discrimination diagrams for studied Central High-Atlas apatites, compared to those of reference rock types (Belousova et al., 2002). Carbonatite field from Ingrid (1998), Chen and Simonetti (2013), Doroshkevich et al. (2018) and Decrée et al. (2020). Field of carbonatite related fenite apatites is plotted using data from Decrée et al. (2020). **(c-d)** Apatite metallogenic discrimination diagrams of Mao et al. (2016). **DP1-1** [= $-0.06461\log\text{Mn} - 1.56\log\text{Sr} + 2.609\log\text{Y} + 0.3631\log\text{La} - 1.766\log\text{Ce} + 0.6243\log\text{Eu} - 3.642\log\text{Dy} + 0.7086\log\text{Yb} - 1.178\log\text{Pb} + 0.4161\log\text{Th} + 0.963\log\text{U} + 6.589$]. **DP1-2** [= $0.2073\log\text{Mn} - 1.035\log\text{Sr} + 15.1\log\text{Y} + 4.995\log\text{La} - 5.804\log\text{Ce} + 0.1741\log\text{Eu} - 8.771\log\text{Dy} - 4.326\log\text{Yb} + 2.022\log\text{Pb} - 0.6719\log\text{Th} + 0.02096\log\text{U} - 10.45$]. **DP4-1** [= $5.379\cdot\log\text{V} + 1.0285\cdot\log\text{Mn} + 1.0004\cdot\log\text{Sr} - 0.0447\cdot\log\text{Ce} + 3.169\cdot\log\text{Eu} - 5.412\cdot\log\text{Yb} - 0.3302\cdot\log\text{Pb} - 0.1080\cdot\log\text{Th} + 5.385$].

Fig. 15. A schematic model for the two main Central High-Atlas vein-type apatite ores.

Tables

Table 1. Selected samples of the studied apatite deposits for mineralogical and geochemical analysis.

Table 2. Whole-rock XRF analyses of alkaline rocks directly hosting the Central High-Atlas vein-type apatite ores.

Table 3. Representative major and trace element compositions of the studied Central High-Atlas apatites.

Table 4. Representative compositions (wt.%) of the clinopyroxenes and amphiboles in the Moroccan Central High-Atlas apatite mineralized veins and magmatic host rocks.

Table 5. Trace element compositions (ppm) of the studied clinopyroxenes and amphiboles.

Table 6. Representative compositions (wt.%) of the feldspars in the studied Central High-Atlas apatite ores and their host alkaline rocks.

Table 7. Representative compositions (wt.%) of the magnetite, ilmenite, titanite and rutile from studied apatite veins and their host alkaline rocks.

Table 8. Representative compositions (wt.%) of the epidote and prehnite associated the apatite ores and their host alkaline rocks.