Assessing the potential risk and relationship between microplastics and phthalates in surface seawater of a heavily human-impacted metropolitan bay in northern China

Yandong Liu, Zhaozhao Li, Isabel Jalón-Rojas, Xiao Hua Wang, Erick Fredj, Dahai Zhang, Lijuan Feng, Xianguo Li

► To cite this version:

Yandong Liu, Zhaozhao Li, Isabel Jalón-Rojas, Xiao Hua Wang, Erick Fredj, et al.. Assessing the potential risk and relationship between microplastics and phthalates in surface seawater of a heavily human-impacted metropolitan bay in northern China. *Ecotoxicology and Environmental Safety*, 2020, 204, pp.111067. 10.1016/j.ecoenv.2020.111067 . hal-03442135

HAL Id: hal-03442135

<https://hal.science/hal-03442135>

Submitted on 25 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Assessing the potential risk and relationship between microplastics**
2 **and phthalates in surface seawater of a heavily human-impacted**
3 **metropolitan bay in northern China**
4

5 Yandong Liu ^a, Zhaozhao Li ^a, Isabel Jalón-Rojas ^{b, c}, Xiao Hua Wang ^c,
6 Erick Fredj ^d, Dahai Zhang ^a, Lijuan Feng ^a, Xianguo Li ^{a*}
7

8 ^a Key Laboratory of Marine Chemistry Theory and Technology (Ocean University of China),
9 Ministry of Education, Qingdao 266100, China

10 ^b UMR5805 EPOC, CNRS, OASU, Université de Bordeaux, Pessac, 33615, France

11 ^c The Sino-Australian Research Consortium for Coastal Management, School of Science, UNSW
12 Canberra, Canberra, ACT, 2600, Australia

13 ^d Jerusalem College of Technology, Jerusalem, Israel
14
15
16

17 *Corresponding author:

18 Key Laboratory of Marine Chemistry Theory and Technology (Ocean University of China),
19 Ministry of Education, Qingdao 266100, China

20 Tel: +86-532-66782215; Fax: +86-532-66782540

21 Email address: lixg@ouc.edu.cn (X. Li)
22

Abstract: The impacts of microplastics (MPs) and phthalates (PAEs), a class of MP-associated contaminants, on the marine environment are not thoroughly understood despite concern over their adverse effects on humans and ecosystems. Field studies linking MPs and PAEs in seawater have not yet been reported. We investigate for the first time the correlation between MPs contamination and the presence of PAEs in the surface seawater of Jiaozhou Bay (JZB), a semi-enclosed metropolitan bay in northern China heavily impacted by human activity. The abundance of MPs, dominated by polyethylene and polyethylene terephthalate mostly smaller than 2 mm, ranged between 24.44 items/m³ and 180.23 items/m³, with the majority being black and transparent fibers and fragments. Concentrations of PAEs varied from 129.96 ng/L to 921.22 ng/L. Relatively higher abundances of MPs and higher concentrations of PAEs were generally found in areas near riverine inputs and sewage treatment plants. There was a strong correlation between PAEs concentration and MPs abundance, suggesting that they are closely linked. In a risk assessment combining PAEs and MPs, the risk quotients (RQs) indicated that the ecological risk of di-n-butyl phthalate in JZB was relatively high ($0.046 < RQ < 0.516$); the risk of the other PAEs were low. The overall ecological hazard index (HI) of PAEs was low to medium ($0.098 < HI < 0.897$). The risk of MPs pollution in JZB, as indicated by Pollution Load Index (PLI), was moderate ($PLI_{JZB} = 11.76$), and mainly due to polyvinyl chloride.

Key words: Microplastics; phthalates; risk assessment; surface seawater; Jiaozhou Bay.

1. Introduction

Plastics and their products are widely used because of their convenience, light weight, and other advantageous properties (Andrés et al., 2014). However, the overuse and improper disposal of plastics have caused severe environmental problems (Rochman et al., 2013), because they can persist in the environment for tens to hundreds of years (Thompson et al., 2009). Large quantities of microplastics (MPs, size below 5mm) are released and transported into the ocean through rivers, sewage, and the atmosphere (Barnes et al., 2009). They originate either from intentional manufacturing (such as in facial cleansers, toothpastes, and plastic raw material; primary origin) or subsequent fragmentation of large plastics via mechanical, photochemical, and biological degradation (secondary origin) (Wright et al., 2013). MPs in marine systems are dispersed by waves, tides and ocean currents. They may sink through the water column (Cincinelli et al., 2019) and be incorporated into sediments (Klein et al., 2015; Tsang et al., 2017) or be assimilated by marine organisms (Kang et al., 2015) and hence transferred into the whole food web (Van et al., 2015), including marine mammals (Lusher et al., 2015a).

MPs easily adsorb heavy metals (Wang et al., 2017) and persistent organic pollutants (Guo et al., 2012a; Mai et al., 2018) due to their small particle size and high specific surface area, increasing their potential toxicity. Many harmful additives are also added during the production of plastics, such as plasticizers, flame retardants, and antioxidants (Hermabessiere et al., 2017). When MPs are immersed in seawater, they can release some of these poisonous substances (Hahladakis et al., 2018) and act as

vectors for pollutant transfer into the marine biota (Zhang et al., 2018a).

Plasticizers are polymer additives used to make plastics flexible, lightweight, and durable. Phthalic acid esters or phthalates (PAEs) have been the most commonly used plasticizers in the plastics industry for decades. PAEs as plasticizers are not chemically bonded with the plastic matrices, and are easily released into the surrounding environment (Dargnat et al., 2009). According to a report by the Information Handling Services (IHS) Chemistry, the amount of PAEs used to produce plasticizers worldwide was nearly 5.9 million tons in 2014 and is expected to rise to about 6.7 million tons by 2019 (Zhang et al., 2018c). Because of the large quantity and widespread application, PAEs are prevalent in the environment, and have been widely found in air, sea, and land environments (Wang et al., 2008; Zhang et al., 2018b). In addition, PAEs have been detected in some aquatic organisms, such as fish (Adeogun et al., 2015) and algae (Babu and Wu, 2010). PAEs are endocrine disruptive to animals and humans, and may be mutagenic, teratogenic, and carcinogenic (Guo et al., 2012b). Six PAEs, dimethyl phthalate (DMP), diethyl phthalate (DEP), di-n-butyl phthalate (DBP), butyl benzyl phthalate (BBP), bis(2-Ethylhexyl) phthalate (DEHP), and di-n-octyl phthalate (DnOP), were listed as environmental priority pollutants by the United States Environmental Protection Agency in 2013 (USEPA, 2013).

Jiaozhou Bay (JZB) is a typical semi-enclosed shallow bay in Qingdao, China, which forms a complex and diverse ecosystem with the surrounding land environment (Fu et al., 2007). JZB is also an important fishery and aquatic base, a sea-salt

production base, and a shipping and tourism economic zone in north China (Gao et al., 2003). With the rapid urbanization and industrialization along the JZB coast, large amounts of industrial waste and anthropogenic pollutants are discharged into JZB directly or via several rivers (Shi et al., 2011). The bay has a low water exchange, with an average residence time of 80 days, ranging from less than 20 days in the deep channel up to 120 days in the shallower northwest area (Liu et al., 2004). This long residence time is favorable for the accumulation of pollutants with a high ecological risk. Owing to its special and distinct location and features, JZB has been and continues to be strongly impacted by anthropogenic activities, providing a good system for the study of MPs and related plasticizers.

Although Zheng et al. (2019) measured MPs in JZB seawater and sediment samples, the level of PAEs has not yet been determined, and no study has evaluated the combined risk of MPs and PAEs using observation data. The influence of environmental parameters, such as water level, water temperature, pH, dissolved oxygen, salinity and conductivity, on the distribution and abundance of MPs and PAEs in semi-enclosed metropolitan bays has also not been reported. Therefore, the objectives of this study are: 1) to determine the level of PAEs in JZB surface seawater and the correlation, if any, between MPs and the environmental parameters; 2) to evaluate the ecological risk of MPs and PAEs. This study also provides a preliminary of the chemical additives in plastics that could be used to trace the source of MPs in the environment, and acts as a reference for the study of the environmental effects of and means of reducing MPs.

2. Materials and methods

2.1. Study area and sample collection

Located southeast of Shandong Peninsula and west of the Yellow Sea, JZB (35°55' N–36°18' N, 120°04' E–120°23' E) is surrounded by Qingdao, a metropolitan city in northern China (Fig. 1). The seawater area of JZB is about 370 km², with an average water depth of 7 m (Sun and Sun, 2015). The significant wave height of the bay is generally less than 5 m, with typical semidiurnal tides (Chen et al., 2012). Several seasonal rivers discharge water, waste, and sediments into the bay including the Dagu, Moshui, Licun, Haibo, and Yang Rivers (Dai et al., 2006; Liang et al., 2015). Several sewage treatment plants are also located on coastline of the bay, including the Tuandao, Haibo, and Licun sewage plants (Fig. 1).

MPs samples were collected from surface seawater at 22 sites (Fig. 1) using a seawater intake system on board the *R/V Yuan Jian 1* on 22 August, 2018 (Stations A1–A6), 10 September, 2018 (Stations C1–C9) and 2 January, 2019 (Stations D1–D7). All samples were collected at low tide. Table S1 in the Supplementary Information provides details of the sampling sites and environmental water parameters. An onboard flowmeter was used to measure the volume of pumped water. Typically, 300 L of seawater was filtered at each collection site using an online system with a 32µm steel sieve (Zhao et al., 2014). The retained particulate material was immediately back-flushed with Milli-Q water (Millipore Co., USA) into a 500 mL glass bottle and fixed in 2.5% formalin (Lattin et al., 2004).

For the PAEs analysis, seawater samples were collected at each station using a clean stainless-steel bucket and placed in clean 4L amber-glass bottles, with roughly 2 mL of 100 mg/L sodium azide added to each bottle to inhibit microbial activity (Gong et al., 2019). One field duplicated sample was randomly collected and examined at every three stations to check the representativeness of sampling. The water samples were stored in the dark at 4°C and processed as soon as possible. The water temperature (WT), pH, dissolved oxygen (DO), salinity (Sal.) and conductivity (Cond.) were measured in the field using an HQ40d multimeter (Hach, Loveland, CO, USA). The water level (WL) data were read from the sonar system on the ship.

2.2. Sample pretreatment and analysis

For the MPs measurement, samples were first oxidized to remove organic matter and then filtered. A stereoscopic microscope was used to examine the sample filters for MPs. When these were found, the polymer types were identified with a micro-Fourier transform infrared spectroscope (μ -FT-IR, PerkinElmer Spectrum Spotlight 400, PerkinElmer, USA). For detailed information about the procedure, readers should refer to the Supplementary Information Section S1.

Six USEPA priority PAEs (DMP, DEP, DBP, BBP, DEHP and DnOP) were identified and quantified in our study. Each seawater sample was filtered, the PAEs solid-phase extracted and then quantified with a GC-MS system (Agilent 7890A-5975C, Agilent Technologies Inc., California, USA). An internal calibration method was used for quantification. Detailed information can be found in Supplementary Information Section S2.

2.3. *Quality assurance (QA) and quality control (QC)*

Nitrile gloves and cotton lab coats were worn throughout the field work and laboratory experiments. All containers were non-plastic, and were cleaned with Milli-Q water. Blank membrane controls were run in the field and in the laboratory as background checks. Between each sample test, anhydrous ethanol was used to wipe the crystal on the μ -FT-IR to avoid cross-contamination (Zheng et al., 2019).

Non-plastic labware was also used for the PAEs analysis, and was cleaned three times with n-hexane before use to prevent any contamination. Detailed procedures and results of the QA/QC for the PAEs analysis, including blanks, spiked-matrix recoveries, duplicated sample analysis, and method detection limit (MDL), are described in Supplementary Information Section S3.

2.4. Risk assessment related to MPs and PAEs

Since MPs and PAEs are recent marine pollutants, there is still no systematic and standardized model specifically developed for assessing their potential ecological risk. Therefore, we adopted here the pollution load index (PLI) method (Tomlinson et al., 1980) for the risk assessment of MPs; this is based on both the abundance and the chemical components of the MPs (Xu et al., 2018). The risk evaluation of the PAEs was based on the risk quotient (RQ) and potential hazard index (HI) methods, following the guidance from the European Commission (EC, 2003). Detailed descriptions of the assessment methods can be found in Supplementary Information Section S4.

2.5. *Data analysis*

The abundance of MPs in seawater samples was expressed as items/m³. All PAEs concentrations were expressed as ng/L. The spatial distributions of the MPs and PAEs were processed using Kriging by Surfer (version 15.0). All statistical analyses were processed using SPSS (version 22.0). Statistical significance was considered at the 0.05 alpha level. The figures were drawn using Origin (version 9.1). Pearson correlation was performed to study the relationships between MPs, PAEs, and environmental water parameters using Matlab.

3. Results and discussion

3.1. MPs

3.1.1. MPs abundance in surface seawater

MPs were detected at all sampling stations (Fig. 2a, Table S1), with abundance ranging from 24.44 to 180.23 items/m³. The mean abundance was 80.46 ± 44.82 items/m³ with a median value of 65.34 items/m³. These values were compared with those found in surface waters in systems around the world, including JZB (Zheng et al., 2019), in order to calibrate the contamination level of MPs in JZB (Fig. 3). This comparison took into account the sampling method used in each study: bulk sampling (used in the present study) or trawling-net sampling. The abundance of MPs observed in this study was very similar to that reported by Zheng et al. (2019) in JZB (20–120 items/m³). Of the other studies using the bulk sampling method, a relatively higher abundance was recorded in the Yangtze Estuary, China (1675.8–6598.8 items/m³, Zhao et al., 2014), in the South China Sea (799–4339 items/m³, Cai et al., 2018), and

in the Incheon/Kyeonggi coastal region of South Korea (328–2876 items/m³, Chae et al., 2015). In contrast, significantly lower abundances of MPs were reported in many studies that used trawling-net sampling, for instance the Mediterranean Sea (0.15 items/m³; de Lucia et al., 2014), the Seto Inland Sea (0.39 items/m³; Isobe et al., 2014), and the East China Sea (0.029–0.305 items/m³; Zhao et al., 2014). In these studies, the lowest abundance of MPs in surface waters was four orders of magnitude lower than the highest for the bulk sampling method. Our results show that MPs were present in the surface seawater of JZB but, compared with other documented systems, the pollution level was still relatively low for a semi-enclosed bay heavily affected by human activity.

Spatially, the highest abundance of MPs, 180.23 items/m³, was found at Station D1 near the Tuandao sewage treatment plant; the lowest abundance, 24.44 items/m³, was observed at Station D4 near the mouth of the bay. The abundance was therefore up to seven times higher at sites near estuaries and sewage treatment plants than at the other stations in the bay. For example, the MPs abundance at Stations C6 and C7, close to the Lichun River and Haibo River estuaries, ranged from 98.82 to 159.42 items/m³, and the abundance at D1 and C7, close to the Tuandao and Haibo sewage treatment plants, ranged from 159.42 to 180.23 items/m³. These results suggest that MPs from rivers and sewage treatment plants were the major contributors to MPs in JZB. This agrees with previous studies that found rivers to be one of the major sources of marine MPs pollution (Schmidt et al., 2017), while estuaries are also potential MPs hot spots (Wright et al., 2013). Municipal sewage treatment plants have

long been considered important sources of MPs (Xu et al., 2019). Significant amounts of MPs still enter the environment from these despite the high removal rates of MPs in current sewage treatment (Murphy et al., 2016).

The distribution of MPs is not only related to human activity, but also closely related to specific geographical locations (Lusher et al., 2014) and ocean currents. The overall level of MPs in the surface seawater of JZB decreased gradually from east to west. The presence of residual currents, such as the clockwise circulation in the eastern region (Li et al., 2014; Yuan et al., 2019), and the low water exchange rate of the bay (Liu et al., 2004) favor the trapping of MPs near the sources. On the other hand, mixing processes combined with strong, flood-dominant tidal currents (Gao et al., 2014) tend to disperse the MPs to different regions of the bay. Clearly, the hydrodynamics plays a significant role in the abundance, distribution, and trapping of MPs. A detailed study of the transport mechanisms that determine the trajectories and distribution patterns of the MPs (e.g. Jalón-Rojas et al, 2019a, b) is a subject of our on-going modelling of JZB.

The D-series of samples were collected in a different season to the A- and C-series. Even though seasonal variations have been frequently reported (e.g. Pelamatti et al., 2019), this seems not to be the case in the present study. The MPs abundances were in a similar range at Stations A1–A6 (26.67–133.33 items/m³, average 66.85 items/m³), C1–C9 (40.00–159.42 items/m³, average 88.52 items/m³), and D1–D7 (24.44 –180.23 items/m³, average 81.77 items/m³) (distribution in Fig. 2, data in Table S1). While significant differences were observed between the different

geographical locations (within the same A, C or D series of samples), the variations between sampling times (between A, C and D series of samples) were relatively small.

3.1.2. MPs characteristics in surface seawater

MPs are generally classified into six categories according to their size: < 0.5 mm, 0.5–1 mm, 1–2 mm, 2–3 mm, 3–4 mm, and 4–5 mm. Fig. 4a shows that, except at Station D6, the surface seawater of JZB mainly contained MPs of size less than 2 mm, accounting for up to 71.80% of the total MPs (see Fig. S1a in Supplementary Information Section). In contrast, MPs with a size of 4–5 mm were the least abundant (7.72%). This dominance of smaller MPs is similar to that in many previous observational studies (e.g. Zhao et al., 2014; Zheng et al., 2019). In general, large plastics are gradually broken up into smaller pieces or particles in the marine environment. As a result, the number of small pieces of plastic increase as the size decreases. The toxicity of MPs particles is related to particle size: the smaller the size, the more toxic they are to marine zooplankton (Jeong et al., 2016).

Based on their morphological characteristics, MPs were classified into fibers, fragments, films, and granules. Typical MP samples are shown in Fig.S2 of Supplementary Information Section. Fibers and fragments were the main shapes here, accounting for 75.62% and 20.82% of the total MPs, respectively (Fig.S1b in Supplementary Information Section). MPs at Stations A4, A5, C3, and D4 were all fibers (Fig. 4b). Films were detected in five samples, accounting for 1.32% of the total MPs abundance. Granules were found only in three samples, accounting for 2.24% of

the abundance. Microbeads used in personal-care exfoliating products were not detected in this study.

Fishery activities in JZB contributed most to the presence of fibrous MPs, since plastic nets and ropes are the main fishing tools. Urban sewage has long been considered a major source of fibrous plastics, mainly from laundry (Zhao et al., 2015). The high presence of fibers supports the argument in Section 3.1.1 that the relatively high abundance of fiber MPs at the study sites may come, to a large extent, from domestic sewage. Although the use of plastic shopping bags has been restricted, they remain a source of film MPs (Sruthy and Ramasamy, 2017). Plastic granules from raw/unprocessed plastic materials, regarded as primary MPs (Law et al., 2010), were only detected in a few samples in our study area, consistent with previous studies in the East China Sea and Yangtze River Estuary (Zhao et al., 2014).

Colored MPs are more likely to be ingested by organisms, as they are more similar to their prey (Abayomi et al., 2017; Wang et al., 2017). Different colors of MPs were observed in the samples, including black, white, transparent, and colored ones. Black was the dominant color found in our study area (Fig. 4c), accounting for 51.17% of the total MPs (Fig. S1c), followed by colored MPs, including red, blue and green, accounting for 3.30~31.51% of the MPs. Transparent and white MPs, accounting for 12.45% and 4.87%, respectively, were also observed. These observations are consistent with a previous study by Zheng et al. (2019) that also observed large number of black fibers.

The color of MPs is usually derived from their original plastic products, but it

can also be changed because of photodegradation and other processes. Consequently, residence times of MPs in the water may be a controlling factor in the degree of color alteration. For example, some blue and green lines might lose most of their original colors and appear white (Zhang et al., 2017). In our study area, white MPs had a relatively higher abundance at Stations C6 (19%) and C7 (17%), which are close to the estuaries of the Licun and Haibo Rivers and municipal sewage treatment plants (Fig. 1). The waste-water treatment processes and the potential weathering undergone during the transport by rivers probably had a bleaching effect on the MPs, resulting in an increased proportion of lightly colored MPs in the effluent. Xu et al. (2019) also found that the abundance of light-colored MPs (white, transparent, yellow, yellow-brown) accounted for the highest proportion in sewage waters (58.54% and 65.38% in the influent and effluent, respectively). White or lightly colored MPs (e.g. discolored yellow, transparent) usually form a higher proportion in surface water (Campanale et al., 2020) due to their higher mass production and their subsequent improper disposal (Wang et al., 2020).

Eight different polymer types were identified in the surface seawater samples (Fig. 4d and Fig. S1d), including polyethylene (PE, accounting for 46.71% of the total abundance), polyethylene terephthalate (PET, 26.86%), rayon (14.80%), polypropylene (PP, 3.66%), polyvinyl chloride (PVC, 3.35%), polyamide (PA, 2.60%), polystyrene (PS, 1.09%), and polyester (0.92%). PE, PET, rayon, PP, and PVC were the dominant types of MPs (Fig. 5), with PE and PP were the most widely used in our daily life and industry (Klein et al., 2015), such as food packaging and

water pipeline. PE (with a density of 0.86–0.96 g/cm³) and PP (0.85–0.91 g/cm³) are readily transported by surface currents because of their lower specific densities compared to seawater (Jalón-Rojas et al., 2019a; Zheng et al., 2019). The fate of these MPs depends on biofouling or biological metabolism processes (Katija et al., 2017) which increase their density, causing their ultimate deposition onto the seafloor (Jalón-Rojas et al., 2019a).

3.2. PAEs

3.2.1. Concentration of PAEs in surface seawater

Fig. 2b and Table S1 summarize the PAEs concentrations in the JZB surface seawater at each site. All the PAEs, except DnOP, were detected in all samples (Table 1), with detection frequencies ranging from 41.38 to 100.00 %. The concentrations of Σ_6 PAEs (sum of the concentrations of the six PAEs) ranged from 129.96 to 921.22 ng/L, with an average of 342.01 ng/L. The concentration of DEHP was lower than the values reported for the Yellow River and Yangtze River in China (Sha et al., 2007; Wang et al., 2008), but significantly higher than that detected in the Tama River in Japan (Suzuki et al., 2001). In addition, DBP and DEHP were found to be much more abundant than any other four congeners (DMP, DEP, BBP and DnOP), with concentrations varying from < MDL to 248.03 ng/L (averaged at 158.09 ng/L, accounting for 46.22% of Σ_6 PAEs) and 60.42 to 617.18 ng/L (averaged at 140.31 ng/L, 41.02% of Σ_6 PAEs), respectively. This is consistent with environmental water samples from nine rivers and two seas in seven areas in China (Wu et al., 2013). DBP is widely used in cosmetics (Giulivo et al., 2016), and the DEHP came mainly from

plastics and the chemical industry (Simoneit et al., 2005). DBP and DEHP are also major PAEs in household garbage (in toys, plastic packaging materials, etc) (Gao and Wen, 2016). In summary, PAEs in JZB surface water mainly come from plastics, the chemical industry and household waste.

It is well documented that the octanol-water partitioning coefficient ($\lg K_{ow}$) of a PAE is closely related to its alkyl chain length: the longer the alkyl chain, the greater the octanol-water partitioning coefficient (as summarized in Table S2), which was proven by many previous studies including a recent correlation analysis (Li et al., 2020). In turn, the greater the octanol-water partitioning coefficient, the higher the lipophilicity (or fat-solubility) of PAEs, and the more difficult the degradation of PAEs by photolysis, hydrolysis, and biological processes (Gao and Wen, 2016). Therefore, it is not a surprise that the concentrations of DBP and DEHP (with relatively longer alkyl chains) in seawater are higher than those of the other PAEs. In addition, the higher the lipophilicity, the greater the threat to aquatic organisms (Dordio et al., 2011).

Legislation in China is continuously refining the standards for various pollutants in a wide range of wastes. However, MPs and PAEs discharge limits are not included in the current national standard for sewage discharge implemented by the Ministry of Environmental Protection of China (MEP, 2002a, b). In addition, there are still great challenges for removing organic pollutants in domestic sewage water, with biodegradation a key step. Technologies are still under development, which may be greatly benefited from identification of highly efficient degradation bacteria and

microbial communities. At most 18% of PAEs are removed by the current treatment equipment in sewage plants (Zhang et al., 2018b). Even if the sewage plants ensured strict observance of standards, the discharge of treated water would be still a significant source of PAEs.

Compared with other rivers, lakes, and oceans (Dargnat et al., 2009; Zhang et al., 2018b, c), the current PAEs concentration in JZB surface water is moderate to slightly low. However, with the rapid development of the marine economic zone along the coast of the Shandong Peninsula, pollution in JZB, including MPs and PAEs (and many other organic pollutants), is increasing markedly.

3.2.2. *Spatial distribution of PAEs*

The total PAEs concentrations in the east JZB, at Stations C6, C7, and D7 near the mouth of the Licun and Haibo Rivers, and at Stations D1 and C7 near the Tuandao and Haibo sewage treatment plants, were all relatively high (Fig. 2b, S3). The total PAEs concentration at Station D1 was as high as 921.22 ng/L. As for PAE congeners, the spatial variation trend of DBP and DEHP were similar to that of $\sum_6$ PAEs in seawater. This is not a surprise, since DBP and DEHP were the major components accounting for 87.24% of $\sum_6$ PAEs, as presented in Section 3.2.1.

In general, the PAEs concentrations at nearshore stations were higher than those in the middle of the bay, and the eastern-shore concentration higher than the western-shore. The higher concentrations of PAEs in the east probably come from surface runoff into JZB, with industries in Qingdao mainly located on the east coast. The Licun, Haibo, and several other rivers east of Qingdao are heavily polluted, with

very low flows, carrying a large amount of organic pollutants from various sources in the city into JZB. Several sewage plants, such as the Tuandao and Haibo sewage treatment plants, also discharge effluent directly into JZB. Many mariculture farms located in the northeastern waters close to the city also contributed to the high levels of PAEs in the region (Ting et al., 2017).

The seasonal variation in PAEs concentrations, reported in previous studies (e.g. Zeng et al., 2009), were much less than the variations between different sampling stations, similar to the case for MPs abundance discussed in Section 3.1.1. The PAEs' concentration ranges and mean values were of the same order of magnitude for Stations A1–A6 (188.51–391.62 ng/L, average 284.61 ng/L), C1–C9 (235.73–507.14 ng/L, average 347.54 ng/L) and D1–D7 (129.96–921.22 ng/L, average 384.10 ng/L).

3.3. Correlations between MPs, PAEs and environmental parameters.

As PAEs are used for plastics production, we investigated the correlation between the MPs abundance (items/m³) and PAEs concentrations (Σ_6 PAEs, ng/L) in JZB surface seawater to explore the possible relationship between the two pollutants. A significantly positive correlation was found ($r^2=0.745$, $p<0.01$, $n=22$; Fig. 6).

As discussed in Section 3.2, MPs and PAEs are correlated each other, both of which are the result of human activity. After they enter the marine system via either atmospheric deposition, surface runoff or riverine input, MPs are prone both to the adsorbing of waterborne contaminants and to the leaching of their plasticizers (Koelmans et al., 2016), leading to the strong correlation between MPs and PAEs found in JZB seawater. The adsorption of waterborne contaminants by MPs and the

release of additives are concurrent processes, and their relative contributions to PAEs levels are still a question to be resolved. It might be helpful to combine field studies with numerical simulations to establish the relationship between contaminant concentration and MPs abundance in specific samples, as well as to investigate the spread through the food chain of these contaminants.

Previous studies also found relationships between PAEs and MPs in a variety of cases. Baini et al. (2017) found correlations between four PAEs and MPs in neustonic samples from the northwestern Mediterranean Sea. Fossi et al. (2016) reported that PAEs were used as indirect (adsorbed contaminants) and plastic (component contaminants) tracers for MPs in the baleen-whale food chain. The present study reports their relationship in seawater samples, which further confirms that the migration behavior of organic pollutants, such as PAEs, is a possible indirect method to trace the source of MPs in the environment.

We also studied the correlations between MPs, PAEs, and various environmental water parameters (Fig. S4). Water level (WL) was significantly negatively correlated with MPs and PAEs ($r = -0.532$, $p < 0.05$; $r = -0.512$, $p < 0.05$, respectively). This suggests that WL is an important environmental factor in the distribution of MPs and PAEs. It could be that shallower waters are characterized by shorter residence times and slower currents, but the transport mechanisms of MPs in JZB need to be investigated in depth. It is interesting to note that pH, DO, Sal., and Cond. are strongly associated with WT (5.4–28.2°C), which is related to sampling time or season. However, there was no clear correlation between these parameters and MPs

abundance, neither with PAEs concentration.

3.4. Assessment of ecological risk

3.4.1 MPs risk assessment

The MPs polymer risk index H in JZB surface water varied from I to IV, with an uneven distribution (Fig. 2C). These high values were mainly due to the presence of high-risky polymers such as PVC, one of the most widely used plastics (Ye et al., 2017). Once PVC enters the marine environment, it can release carcinogenic monomers and other pollutants, impacting the wildlife (Green et al., 2016). Unreacted monomers and polymeric impurities in plastic products can also be released into the environment, threatening human health and ecosystem health (Araújo et al., 2002).

The MPs pollution load in JZB based on the Pollution Load Index (PLI) is given in Fig. 2d. Fifteen sites were moderately polluted, the other seven mildly polluted. According to this result, the MPs pollution risk in JZB surface seawater is not dramatic. The whole area is characterized by a medium level of pollution risk (Fig. 2c, d) with a PLI_{JZB} value of 11.76, which falls into risk category II.

3.4.2 Potential ecological risk related to PAEs

We assessed the ecological risk of the four main PAEs in JZB surface seawater. The risk quotient (RQ) of DMP, DEP, and DEHP were lower than 0.1 at the 22 sampling stations, except at Stations C7, C8, D1 and D7, indicating an overall low ecological risk. Relative higher RQs were found for DBP ($0.1 < RQ < 1$, except at Stations D4 and D7; Fig. S5), which indicates that DBP is a medium ecological risk. These results demonstrate that PAEs are becoming a potential threat to the ecology of

JZB, which deserves further study.

The hazard index (HI) values, ranging between 0.098 and 0.897, indicate medium-to-low ecological risk in JZB surface seawater. The HI value at Station D1 (0.897) was significantly higher than elsewhere, indicating that areas near sewage treatment plants are generally at a higher level of ecological risk. Higher HI levels were also observed at Station C6 (0.572) and C7 (0.588), which may be related to riverine input.

The risk assessment presented here has some limitations. For instance, it is difficult to determine the PNEC (Predicted No Effect Concentration) for each PAE congener because the environmental toxicology and ecotoxicological information on these pollutants are very limited. Further assessment of the ecological risks of these emerging pollutants is needed to provide more toxicity data (Gong et al., 2019).

4. Conclusions

The present study offers novel insights into the occurrence, distribution, correlation, and potential ecological risks of microplastics (MPs) and phthalates (PAEs) in surface seawater of a semi-enclosed bay heavily impacted by human activity, Jiaozhou Bay (JZB) in northern China.

The abundance of MPs in the bay was relatively low compared to other similar areas. The MP and PAE pollution was relatively severe near the estuaries and sewage plants, suggesting that rivers and sewage plants were the main sources of MPs and PAEs. A strong correlation between the abundance of MPs and concentration of PAEs

was observed in surface seawater.

The ecological risk assessment of PAEs, based on risk quotients, indicated that PAEs, presently at low-to-medium level, are becoming a potential threat to the JZB ecosystem. The risk-assessment model, based on MPs concentrations and polymer chemical hazards showed a high risk related to the presence of hazardous polymers such as PVC. The risk of MPs pollution in the bay was evaluated as medium.

Further studies on the impact of MPs on the transport of organic pollutants in marine environment are important in providing basic data for a source-to-sink study on pollutants in the ocean and a theoretical basis for the control of MPs pollution, as is more comprehensive modeling to help understand MP transport dynamics in a highly urbanized estuarine environment such as JZB.

Acknowledgments

This work was financially supported by the Fundamental Research Funds for the Central Universities in the Ocean University of China (No. 201861015) and a JZB scoping grant from Sino-Australian Research Centre for Coastal Management at UNSW Canberra. The μ -FT-IR test was carried out by the research team of Professor Chengjun Sun and the GC-MS detection by Dr. Bin Han and Professor Li Zheng (all at the Ecological Center of the First Institute of Oceanography, Ministry of Natural Resources, China), to whom we owe our great thanks. Our thanks also goes to Dr. Peter McInyre of..... for his help in English editing. This is publication No. 76 of the Sino-Australian Research Consortium for Coastal Management at UNSW Canberra,

485 Australia.

486

487 **References**

488 Abayomi, O.A., Range, P., Al-Ghouti, M.A., Obbard, J.P., Almeer, S.H., Ben-Hamadou, R., 2017.

489 Microplastics in coastal environments of the Arabian Gulf. *Mar. Pollut. Bull.* 124 (1),

490 181-188.

491 Adeogun, A.O., Ibor, O.R., Omogbemi, E.D., Chukwuka, A.V., Adegbola, R.A., Adewuyi, G.A.,

492 Arukwe, A., 2015. Environmental occurrence and biota concentration of phthalate esters in

493 Epe and Lagos Lagoons, Nigeria. *Mar. Environ. Res.* 108, 24-32.

494 Andrés, C., Fidel, E., J Ignacio, et al., 2014. Plastic debris in the open ocean. *Proc. Natl. Acad. Sci.*

495 U. S. A. 111 (28), 10239-10244.

496 Araújo, P.H.H., Sayer, C., Poco, J.G.R., Giudici, R., 2002. Techniques for reducing residual

497 monomer content in polymers: a review. *Polym. Eng. Sci.* 42, 1442-1468.

498 Babu, B., Wu, J.T., 2010. Production of phthalate esters by nuisance freshwater algae and

499 cyanobacteria. *Sci. Total Environ.* 408, 4969-4975.

500 Bagaev, A., Khatmullina, L., Chubarenko, I., 2018. Anthropogenic microlitter in the Baltic Sea

501 water column. *Mar. Pollut. Bull.* 129 (2), 918-923.

502 Baini, M., Martellini, T., Cincinelli, A., Campani, T., Minutoli, R., Panti, C., Finoia, M.G., Fossi,

503 M.C., 2017. First detection of seven phthalate esters (PAEs) as plastic tracers in superficial

504 neustonic/planktonic samples and cetacean blubber. *Anal. Methods.* 9 (9), 1512-1520.

505 Barnes, D. K. A.; Galgani, F.; Thompson, R. C.; Barlaz, M., 2009. Accumulation and

506 fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. B*, 364 (1526),

1985-1998.

Cai, M., He, H., Liu, M., Li, S., Tang, G., Wang, W., Huang, P., Wei, G., Lin, Y., Chen, B., Hu, J.,
Cen, Z., 2018. Lost but can't be neglected: Huge quantities of small microplastics hide in the
South China Sea. *Sci. Total Environ.* 633, 1206-1216.

Campanale, C., Stock, F., Massarelli, C., Kochleus, C., Bagnuolo, G., Reifferscheid, G., Uricchio,
V. F., 2020. Microplastics and their possible sources: The example of Ofanto River in
southeast Italy. *Environ. Pollut.* 258, 113284.

Chae, D.H., Kim, I.S., Kim, S.K., Song, Y.K., Shim, W.J., 2015. Abundance and distribution
characteristics of microplastics in surface seawaters of the Incheon/Kyeonggi Coastal Region.
Arch. Environ. Contam. Toxicol. 69 (3), 269-278.

Chen, B., Zhang, Y., Liu, J., Kong, X., 2012. Tidal current dynamic characteristic and its relation
with suspended sediment concentration in Jiaozhou Bay. *Adv. Mar. Sci.* 30 (1), 24-35 (in
Chinese).

Chen, M., Jin, M., Tao, P., Wang, Z., Xie, W., Yu, X., Wang, K., 2018. Assessment of
microplastics derived from mariculture in Xiangshan Bay, China. *Environ. Pollut.* 242,
1146-1156.

Cheung, P.K., Fok, L., Hung, P.L., Cheung, L.T.O., 2018. Spatio-temporal comparison of
neustonic microplastic density in Hong Kong waters under the influence of the Pearl River
Estuary. *Sci. Total Environ.* 628-629, 731-739.

Cincinelli, A., Martellini, T., Guerranti, C., Scopetani, C., Chelazzi, D., Giarrizzo, T., 2019. A
potpourri of microplastics in the sea surface and water column of the Mediterranean
Sea. *Trends Anal. Chem.* 110, 321-326.

529 Dai, J.C., Song, J.M., Zheng, G.X., 2006. Analysis of sedimentary environmental evolution of the
530 Jiaozhou Bay. *Adv. Mar. Sci.* 24 (3), 397-406 (in Chinese).

531 Dargnat, C., Blanchard, M., Chevreuil, M., Teil, M.J., 2009. Occurrence of phthalate esters in the
532 Seine River estuary (France). *Hydrol. Process.* 23, 1192-1201.

533 de Lucia, G.A., Caliani, I., et al., 2014. Amount and distribution of neustonic micro-plastic off the
534 western Sardinian coast (Central-Western Mediterranean Sea). *Mar. Environ. Res.* 100,
535 10-16.

536 Desforbes, J. W., Galbraith, M., Dangerfield, N., Ross, P. S., 2014. Widespread distribution of
537 microplastics in subsurface seawater in the NE Pacific Ocean. *Mar. Pollut. Bull.* 79(1), 94-99.

538 Dordio, A. V., Belo, M., Teixeira, D. M., Carvalho, A. P., Dias, C. M. B., Picó, Y., Pinto, A. P.,
539 2011. Evaluation of carbamazepine uptake and metabolism by *Typha* spp., a plant with
540 potential use in phytotreatment. *Bioresour Technol.* 102 (17), 7827-7834.

541 EC, 2003. European Commission Technical Guidance Document in Support of Commission
542 Directive 93/67/EEC on Risk Assessment for New Notified Substances and Commission
543 Regulation (EC) No 1488/94 on Risk Assessment for Existing Substance, Part II. Brussels,
544 Belgium.

545 Fossi, M. C., Marsili, L., Bani, M., et al., 2016. Fin whales and microplastics: The Mediterranean
546 Sea and the Sea of Cortez scenarios. *Environ. Pollut.* 209, 68-78.

547 Fu, M., Li, Z., Gao, H., 2007. Distribution characteristics of nonylphenol in Jiaozhou Bay of
548 Qingdao and its adjacent rivers. *Chemosphere.* 69 (7), 1009-1016.

549 Gao, D. W., Wen, Z. D., 2016. Phthalate esters in the environment: A critical review of their
550 occurrence, biodegradation, and removal during wastewater treatment processes. *Sci. Total*

551 Environ. 541, 986-1001.

552 Gao, G. D., Wang, X. H., Bao, X. W., 2014. Land reclamation and its impact on tidal dynamics in
553 Jiaozhou Bay, Qingdao, China. Estuar. Coast. Shelf Sci. 151, 285-294.

554 Gao, S.Q., Lin, Y.A., Jin, M.M., 2003. Transport and distribution of nutrients in anchovy
555 spawning ground to the southern waters of Shandong Peninsula. Acta Oceanol. Sin. 4,
556 533-546.

557 Giulivo, M., de Alda, M. L., Capri, E., Barceló, D., 2016. Human exposure to endocrine disrupting
558 compounds: Their role in reproductive systems, metabolic syndrome and breast cancer. A
559 review. Environ. Res. 151, 251-264.

560 Gong, J., Lin, C., Xiong, X., Chen, D., Chen, Y., Zhou, Y., Wu, C., Du, Y., 2019. Occurrence,
561 distribution, and potential risks of environmental corticosteroids in surface waters from the
562 Pearl River Delta, South China. Environ. Pollut. 251, 102-109.

563 Green, D.S., Boots, B., Sigwart, J., Jiang, S., Rocha, C., 2016. Effects of conventional and
564 biodegradable microplastics on a marine ecosystem engineer (*Arenicola marina*) and
565 sediment nutrient cycling. Environ. Pollut. 208, 426-434.

566 Guo, X.Y., Wang, X.L., Zhou, X.Z., Kong, X.Z., Tao, S., Xing, B.S., 2012a. Sorption of four
567 hydrophobic organic compounds by three chemically distinct polymers: role of chemical and
568 physical composition. Environ. Sci. Technol. 46 (13), 7252-7259.

569 Guo, Y., Zhang, Z., Liu, L., Li, Y., Ren, N., Kannan, K., 2012b. Occurrence and profiles of
570 phthalates in foodstuffs from China and their implications for human exposure. J. Agric.
571 Food Chem. 60 (27), 6913-6919.

572 Hahladakis, J.N., Velis, C.A., Weber, R., Iacovidou, E., Purnell, P., 2018. An overview of

573 chemical additives present in plastics: migration, release, fate and environmental impact
574 during their use, disposal and recycling. J. Hazard. Mater. 344, 179-199.

575 Hermabessiere, L., Dehaut, A., Paul-Pont, I., Lacroix, C., Jezequel, R., Soudant, P., Dufols, G.,
576 2017. Occurrence and effects of plastic additives on marine environments and organisms: a
577 review. Chemosphere. 182, 781-793.

578 Isobe, A., Kubo, K., Tamura, Y., Kako, S., Nakashima, E., Fujii, N., 2014. Selective transport of
579 microplastics and mesoplastics by drifting in coastal waters. Mar. Pollut. Bull. 89 (1-2),
580 324-330.

581 Isobe, A., Uchida, K., Tadashi, T., Lwasaki, S., 2015. East Asian seas: A hot spot of pelagic
582 microplastics. Mar. Pollut. Bull. 101 (2), 618-623.

583 Jalón-Rojas, I., Wang, X. H., Fredj, E., 2019a. On the importance of a three-dimensional approach
584 for modelling the transport of neustic microplastics. Ocean Science, 15(3), 717-724.

585 Jalón-Rojas, I., Wang, X. H., Fredj, E., 2019b. A 3D numerical model to track marine plastic
586 debris (TrackMPD): sensitivity of microplastic trajectories and fates to particle dynamical
587 properties and physical processes. Mar. Pollut. Bull. 141, 256-272.

588 Jeong, C.B., Won, E.J., Kang, H.M., Lee, M.C., Hwang, D.S., Hwang, U.K., Zhou, B., Souissi, S.,
589 Lee, S.J., Lee, J.S., 2016. Microplastic size-dependent toxicity, oxidative stress induction,
590 and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*). Environ.
591 Sci. Technol. 50 (16), 8849-8857.

592 Kang, J.H., Kwon, O.Y., Shim, W.J., 2015. Potential threat of microplastics to zooplanktivores in
593 the surface waters of the southern sea of Korea. Arch. Environ. Contam. Toxicol. 69 (3),
594 340-351.

595 Katija, K., Choy, C.A., Sherlock, R.E., Sherman, A.D., Robison, B.H., 2017. From the surface to
 596 the seafloor: how giant larvaceans transport microplastics into the deep sea. *Sci. Adv.* 3,
 597 e1700715.

598 Klein, S., Worch, E., Knepper, T.P., 2015. Occurrence and spatial distribution of microplastics in
 599 river shore sediments of the Rhine-Main area in Germany. *Environ. Sci. Technol.* 49,
 600 6070-6076.

601 Koelmans, A. A., Bakir, A., Burton, G. A., Janssen, C. R., 2016. Microplastic as a vector for
 602 chemicals in the aquatic environment: critical review and model-supported reinterpretation of
 603 empirical studies. *Environ. Sci. Technol.* 50 (7), 3315-3326.

604 Lattin, G.L., Moore, C.J., Zellers, A.F., Moore, S.L., Weisberg, S.B., 2004. A comparison of
 605 neustonic plastic and zooplankton at different depths near the Southern California shore. *Mar.*
 606 *Pollut. Bull.* 49 (4), 291-294.

607 Law, K.L., Morét-Ferguson, S., Maximenko, N.A., Proskurowski, G., Peacock, E.E., Hafner, J.,
 608 Reddy, C.M., 2010. Plastic accumulation in the North Atlantic subtropical gyre. *Science*. 329
 609 (5996), 1185-1188.

610 Liang, S.K., Pearson, S., Wu, W., Ma, Y.J., Qiao, L.L., Wang, X.H., Li, J.M., Wang, X.L., 2015.
 611 Research and integrated coastal zone management in rapidly developing estuarine harbours: a
 612 review to inform sustainment of functions in Jiaozhou Bay, China. *Ocean Coast. Manag.* 116,
 613 470-477.

614 Li, P., Li, G.X., Qiao, L.L., Chen, X.E., Shi, J.H., Gao, F., Wang, N., Yue, S.H., 2014. Modeling
 615 the tidal dynamic changes induced by the bridge in Jiaozhou Bay, Qingdao, China. *Cont.*
 616 *Shelf Res.* 84, 43-53.

617 Li, Y., Yan, H., Liu, Q., Li, X., Ge, J., Yu, X., 2020. Accumulation and transport patterns of six
 618 phthalic acid esters (PAEs) in two leafy vegetables under hydroponic conditions.
 619 Chemosphere. 249, 126457.

620 Liu, Z., Wei, H., Liu, G.S., Zhang, J., 2004. Simulation of water exchange in Jiaozhou Bay by
 621 average residence time approach. Estuar. Coast. Shelf Sci. 61 (1), 25-35.

622 Lusher, A.L., Burke, A., O'Connor, I., Officer, R., 2014. Microplastic pollution in the Northeast
 623 Atlantic Ocean: validated and opportunistic sampling. Mar. Pollut. Bull. 88 (1-2), 325-333.

624 Lusher, A. L.; Hernandez-Milian, G.; O'Brien, J.; Berrow, S.; O'Connor, I.; Officer, R., 2015a.
 625 Microplastic and macroplastic ingestion by a deep diving, oceanic cetacean: the True's
 626 beaked whale *Mesoplodon mirus*. Environ. Pollut. 199, 185-191.

627 Lusher, A.L., Tirelli, V., O'Connor, I., Officer, R., 2015b. Microplastics in Arctic polar waters: the
 628 first reported values of particles in surface and sub-surface samples. Sci. Rep. 5, 14947.

629 Mai, L., Bao, L.J., Shi, L., Liu, L.Y., Zeng, E.Y., 2018. Polycyclic aromatic hydrocarbons
 630 affiliated with microplastics in surface waters of Bohai and Huanghai Seas, China. Environ.
 631 Pollut. 241, 834-840.

632 MEP, 2002a. Discharge Standard of Pollutants for Municipal Wastewater Treatment Plant
 633 (GB18918-2002). China Environment Press, Beijing.

634 MEP, 2002b. Surface Water Quantity Standards (GB8938-2002). China Environment Press,
 635 Beijing.

636 Murphy, F., Ewins, C., Carbonnier, F., Quinn, B., 2016. Wastewater treatment works (WwTW) as
 637 a source of microplastics in the aquatic environment. Environ. Sci. Technol. 50 (11),
 638 5800-5808.

639 Pelamatti, T., Fonseca-Ponce, I. A., Rios-Mendoza, L. M., Stewart, J. D., Marin-Enriquez, E.,
 640 Marmolejo-Rodriguez, A. J., Hoyos-Padilla, E. M., Galvan-Magana, F. G., Gonzalez-Armas,
 641 R., 2019. Seasonal variation in the abundance of marine plastic debris in Banderas Bay,
 642 Mexico. *Mar. Pollut. Bull.* 145, 604-610.

643 Rochman, C.M., Browne, M.A., Halpern, B.S., Hentschel, B.T., Hoh, E., Karapanagioti, H.K.,
 644 Rios-Mendoza, L.M., Takada, H., Teh, S., Thompson, R.C., 2013. Policy: Classify plastic
 645 waste as hazardous. *Nature*. 494 (7436), 169-171.

646 Schmidt, C., Krauth, T., Wagner, S., 2017. Export of plastic debris by rivers into the sea. *Environ.*
 647 *Sci. Technol.* 51(21), 12246-12253.

648 Sha, Y., Xia, X., Yang, Z., Huang, G.H., 2007. Distribution of PAEs in the middle and lower
 649 reaches of the Yellow River, China. *Environ. Monit. Assess.* 124, 277-287.

650 Shi, J., Li, G., Wang, P., 2011. Anthropogenic influences on the tidal prism and water exchanges
 651 in Jiaozhou Bay, Qingdao, China. *J. Coast. Res.* 27(1), 57-72.

652 Simoneit, B. R., Medeiros, P. M., Didyk, B. M., 2005. Combustion products of plastics as
 653 indicators for refuse burning in the atmosphere. *Environ. Sci. Technol.* 39 (18), 6961-6970.

654 Sruthy, S., Ramasamy, E.V., 2017. Microplastic pollution in Vembanad Lake, Kerala, India: the
 655 first report of microplastics in lake and estuarine sediments in India. *Environ. Pollut.* 222,
 656 315-322.

657 Sun, S., Sun, X.X., 2015. *The Theory and Practice of Gulf Ecosystem-A Case Study for Jiaozhou*
 658 *Bay.* Science Press, Beijing, pp. 12-235 (in Chinese).

659 Suzuki, T., Yaguchi, K., Suzuki, S., Suga, T., 2001. Monitoring of phthalic acid monoesters in
 660 river water by solid-phase extraction and GC-MS determination. *Environ. Sci. Technol.* 35,

661 3757-3763.

662 Thompson, R.C., Moore, C.J., Vom Saal, F.S., Swan, S.H., 2009. Plastics, the environment and
663 human health: current consensus and future trends. *Philos. Trans. R. Soc. B.* 364 (1526),
664 2153-2166.

665 Ting, J. U., Wei, G. E., Chao, C. H. A. I., 2017. Pollution and ecological risk assessment of
666 polybrominated diphenyl ethers in the sediments of Jiaozhou Bay. *Environ. Chem. (China)*.
667 36 (4): 839-848.

668 Tomlinson, D.L., Wilson, J.G., Harris, C.R., Jeffrey, D.W., 1980. Problems in the assessment of
669 heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer Meeresun.*
670 33, 566-575.

671 Tsang, Y. Y., Mak, C. W., Liebich, C., Lam, S. W., Sze, E. T., Chan, K. M., 2017. Microplastic
672 pollution in the marine waters and sediments of Hong Kong. *Mar. Pollut. Bull.* 115 (1-2),
673 20-28.

674 USEPA, 2013. Electronic Code of Federal Regulations, Title 40-Protection of Environment, Part
675 423d Steam Electric Power Generating Point Source Category. Appendix A to Part 423-126,
676 Priority Pollutants.

677 Van Cauwenberghe, L.; Claessens, M.; Vandegehuchte, M. B.; Janssen, C. R., 2015. Microplastics
678 are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicolamarina*) living in natural
679 habitats. *Environ. Pollut.* 199, 10-17.

680 Wang, G., Lu, J., Tong, Y., Liu, Z., Zhou, H., Xiayihazi, N., 2020. Occurrence and pollution
681 characteristics of microplastics in surface water of the Manas River Basin, China. *Sci. Total*
682 *Environ.* 710, 136099.

683 Wang, J., Peng, J., Tan, Z., Gao, Y., Zhan, Z., Chen, Q., Cai, L., 2017. Microplastics in the surface
684 sediments from the Beijiang River littoral zone: composition, abundance, surface textures and
685 interaction with heavy metals. *Chemosphere* 171, 248-258.

686 Wang, P., Wang, S.L., Fan, C.Q., 2008. Atmospheric distribution of particulate- and gasphase
687 phthalic esters (PAEs) in a Metropolitan City, Nanjing, East China. *Chemosphere*. 72,
688 1567-1572.

689 Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics on
690 marine organisms: a review. *Environ. Pollut.* 178, 483-492.

691 Wu, X.L., Hong, H.J., Liu, X.T., Guan, W.B., Meng, L.X., Ye, Y., Ma, Y.Q., 2013. Graphene
692 dispersive solid-phase extraction of phthalate acid esters from environmental water. *Sci.*
693 *Total Environ.* 444, 224-230.

694 Xu, P., Peng, G., Su, L., Gao, Y., Gao, L., Li, D., 2018. Microplastic risk assessment in surface
695 waters: a case study in the Changjiang Estuary, China. *Mar. Pollut. Bull.*, 133, 647-654.

696 Xu, X., Jian, Y., Xue, Y., Hou, Q., Wang, L., 2019. Microplastics in the wastewater treatment
697 plants (WWTPs): Occurrence and removal. *Chemosphere*. 235, 1089-1096.

698 Ye, L., Qi, C., Hong, J., Ma, X., 2017. Life cycle assessment of polyvinyl chloride production and
699 its recyclability in China. *J. Clean. Prod.* 142, 2965-2972.

700 Yuan, Y., Jalón-Rojas, I., Wang, X. H., Song, D., 2019. Design, construction, and application of a
701 regional ocean database: A case study in Jiaozhou Bay, China. *Limnology and Oceanography:*
702 *Methods*, 17(3), 210-222.

703 Zeng, F., Wen, J., Cui, K., Wu, L., Liu, M., Li, Y., Lin, Y., Zhu, F., Ma, Z., Zeng, Z., 2009.
704 Seasonal distribution of phthalate esters in surface water of the urban lakes in the subtropical

city, Guangzhou, China. *J. Hazard. Mater.* 169, 719-725.

Zhang, H., Zhou, Q., Xie, Z., Zhou, Y., Tu, C., Fu, C., Mi, W., Ebinghaus, R., Christie, P., Luo, Y., 2018a. Occurrences of organophosphorus esters and phthalates in the microplastics from the coastal beaches in north China. *Sci. Total Environ.* 616-617, 1505-1512.

Zhang, W., Zhang, S., Wang, J., Wang, Y., Mu, J., Wang, P., Lin, X., Ma, D., 2017. Microplastic pollution in the surface waters of the Bohai Sea, China. *Environ. Pollut.* 231, 541-548.

Zhang, Z. M., Zhang, H. H., Zhang, J., Wang, Q. W., Yang, G. P., 2018b. Occurrence, distribution, and ecological risks of phthalate esters in the seawater and sediment of Changjiang River Estuary and its adjacent area. *Sci. Total Environ.* 619, 93-102.

Zhang, Z. M., Zhang, H. H., Zou, Y. W., Yang, G. P., 2018c. Distribution and ecotoxicological state of phthalate esters in the sea-surface microlayer, seawater and sediment of the Bohai Sea and the Yellow Sea. *Environ. Pollut.* 240, 235-247.

Zhao, S., Zhu, L., Li, D., 2015. Microplastic in three urban estuaries, China. *Environ. Pollut.* 206, 597-604.

Zhao, S., Zhu, L., Wang, T., Li, D., 2014. Suspended microplastics in the surface water of the Yangtze Estuary System, China: first observations on occurrence, distribution. *Mar. Pollut. Bull.* 86 (1-2), 562-568.

Zheng, Y., Li, J., Cao, W., Liu, X., Jiang, F., Ding, J., Yin, X., Sun, C., 2019. Distribution characteristics of microplastics in the seawater and sediment: A case study in Jiaozhou Bay, China. *Sci. Total Environ.* 674, 27-35.

Zhu, L., Bai, H., Chen, B., Sun, X., Qu, K., Xia, B., 2018. Microplastic pollution in North Yellow Sea, China: Observations on occurrence, distribution and identification. *Sci. Total*

727 Environ. 636, 20-29.

728 Zhu, J.M., Zhang, Q., Li, Y.P., Tan, S.D., Kang, Z.J., Yu, X.Y., Lan, W.L., Cai, L., Wang, J.Z.,

729 Shi, H.H., 2019. Microplastic pollution in the Maowei Sea, a typical mariculture bay of

730 China. Sci. Total Environ. 658, 62-68.

731

Figure Captions

Fig. 1 The geographic location of JZB and sampling stations of surface seawater. The red dots represent water sampling sites, the red pentagram represents the location of Qingdao municipal government, the transparent pentagrams represent the locations of district governments, and the blue rhombuses represent the locations of sewage treatment plants.

Fig. 2 The abundance of MPs (a), concentration of Σ_6 PAEs (b), MPs' polymer risk index H (c) and pollution load index PLI (d) in JZB surface seawater.

Fig. 3 Comparison of MPs abundance in JZB surface water with previously reported data.

Fig. 4 Proportion of different sizes (a), shapes (b), colors (c), polymer types (d) of MPs from JZB surface seawater at each sampling site.

Fig. 5 Identification of typical MPs in JZB surface seawater using μ -FT-IR. Values in brackets indicate the matches of the spectra with the standards. Scale bar of the right of photograph is 100 μ m.

Fig. 6 Linear regression of MPs abundance and Σ_6 PAEs observed in JZB surface seawater.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

873 **Fig. 5**

874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898

899 **Fig. 6**

900