[Click here to view linked References](#)

Mantle xenolith-bearing phonolites and basanites feed the active volcanic ridge of Mayotte (Comoros archipelago, SW Indian Ocean)

Carole Berthod⁽¹⁾, Etienne Médard⁽¹⁾, Andrea Di Muro^(2,3), Théo Hassen Ali⁽¹⁾, Lucia Gurioli⁽¹⁾, Catherine Chauvel⁽³⁾, Jean-Christophe Komorowski⁽³⁾, Patrick Bachèlery⁽¹⁾, Aline Peltier^(2,3), Mhammed Benbakkar⁽¹⁾, Jean-Luc Devidal⁽¹⁾, Pascale Besson⁽³⁾, Anne Le Friant⁽³⁾, Christine Deplus⁽³⁾, Sophie Nowak⁽³⁾, Isabelle Thinon⁽⁴⁾, Pierre Burckel⁽³⁾, Samia Hidalgo⁽³⁾, Nathalie Feuillet⁽³⁾, Stephan Jorry⁽⁵⁾, Yves Fouquet⁽⁵⁾.

(1) Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans, Clermont-Ferrand, France,

(2) Observatoire Volcanologique du Piton de la Fournaise, Institut de physique du globe de Paris, F-97418 La Plaine des Cafres, France

(3) Université de Paris, Institut de physique du globe de Paris, CNRS, F-75005 Paris, France

(4) BRGM, French Geological Survey, 3 avenue Claude Guillemin, BP36009, 45 060 Orléans, Cédex 2, France,

(5) Ifremer, Unité Géosciences Marines, F-29280 Plouzané, France

Keywords: Mayotte; Basanite; phonolite; magmatic system; fractional crystallization.

Abstract

Since 2018, the submarine east flank of Mayotte Island (Comoros archipelago) is the site of a major and still ongoing eruption, located at 3.5 km depth bsl on a WNW-ESE volcanic ridge. Oceanographic cruises carried out to monitor this seismo-volcanic crisis allowed us to determine that this volcanic ridge is built by a bimodal sodic alkaline magmatic series that includes basanites and phonolites. A petrological study of dredged samples allowed us to image the magmatic system feeding the volcanic ridge and to determine the link between basanitic and phonolitic magmas. The magmatic system feeding the volcanic ridge comprises multiple levels of magma storage. Basanitic magmas generated at 80-100 km mantle depth are stored in one or more deep reservoir(s) (≥ 37 km) and then in shallower

basanitic and phonolitic lenses located close to the Moho interface before rising the surface. This study identifies three possible scenarios: (1) the deep basanitic magma rises directly and quickly to the surface from the deep mantle reservoir (as is currently happening 60 km offshore), (2) the basanitic magma stalls in a shallower reservoir near the Moho before resuming its ascent toward the surface and erupting as porphyritic basanite, (3) the basanitic magma stops and evolves to phonolite in these sub-crustal reservoirs. The phonolitic lavas are produced by approximately 80% fractional crystallization (34% clinopyroxene, 30% anorthoclase feldspar, 15.5% magnetite, 12.5% olivine, 5% apatite and 4% ilmenite) of a hydrous basanitic magma at mantle depths ($P > 0.6$ GPa) under reduced oxygen fugacity ($\sim$ FMQ-1). In this third scenario, the phonolitic magma might be reactivated by the arrival of a new batch of deeper basanitic magma.

1. Introduction

The deepest part (3.5 km bsl) of the eastern submarine flank of Mayotte is the site of a long lasting and still ongoing (as of January 2021) effusive volcanic activity that followed an intense volcano-tectonic crisis beginning in May 2018 (Berthod et al., 2021; Cesca et al., 2020; Feuillet et al., 2021; Lemoine et al., 2020). A series of oceanographic cruises organized by the French scientific community have led to the discovery of an active volcanic ridge characterized by basanitic and phonolitic volcanic cones and lava fields, and extending along 60 km from the deep distal eruptive site to the subaerial easternmost part of Mayotte island, Petite Terre (Feuillet, 2019; Feuillet et al., 2021; Fouquet and Feuillet, 2019; Jorry, 2019). Both subaerial and submarine phonolitic eruptive products carry mm to dm sized ultramafic xenoliths testifying to rapid ascent from mantle levels. Worldwide, most phonolitic magmas evolve and reside in shallow crustal reservoirs (e.g., Scaillet et al. 2008; Grant et al. 2013; Moussallam et al. 2013), and mantle xenolith-bearing phonolites are extremely rare (Dautria et al., 1983; Grant et al., 2013; Price and Green, 1972). Phonolites usually result from fractional crystallization of a basanitic parental magma (Ablay et al., 1998; Kyle et al., 1992; Le roex et al., 1990), although phonolitic melts can be generated directly by mantle melting (Laporte et al., 2014). Berthod et al. (2021) demonstrate that lavas erupted in 2018-2019 from the distal submarine site are basanitic and ascended rapidly from a large deep reservoir located in the lithospheric mantle. Slight chemical evolution of the erupted lavas over time was interpreted as an evidence of syn-eruptive mixing between the ascending basanite and a shallower, cooler and more differentiated magma possibly stored close to the Moho interface.

Long-standing activity on the Mayotte active volcanic ridge allows us to explore the origin of deep phonolites and their evolution at mantle level together with their link with the basanite volcanism. Are basanite and phonolites both primitive mantle melting products? Or do phonolites derive through fractional crystallization of the basanites?

In order to answer these questions, we performed a detailed petrological and geochemical investigation, including petrological analysis, thermobarometry and in situ and whole-rock geochemistry of the submarine volcanic products collected along the Mayotte submarine ridge during three successive oceanographic cruises (MAYOBS 1, MAYOBS 2 and MAYOBS 4) between May and July 2019 (Feuillet, 2019; Feuillet et al., 2019; Fouquet and Feuillet, 2019; Jorry, 2019), and compared them with recent (Holocene) subaerial products collected during two field campaigns on Petite Terre in 2018 and 2019.

2. Geological setting

The Comoros Archipelago, located in the Mozambique Channel between the northern tip of Madagascar and the eastern coast of Mozambique (Fig. 1a) consists of four islands: Mayotte, Anjouan, Moheli and Grande Comore, from east to west. These islands are interconnected by a series of submarine volcanic ridges and are associated with atolls and partially emerged reef platforms (Tzevahirtzian et al. 2021). The overall archipelago has been recently interpreted as the NE-SW boundary between the Lwandle microplate and the Somalia plate, a boundary marked by a zone of active seismicity connecting the northern extremity of Madagascar in the east to the African coast in the west (Bertil and Regnault, 1998; Dziewonski et al., 1981; Ekström et al., 2012; Famin et al., 2020). This boundary may consist in a zone of broad deformation extending up to the northern half of Madagascar (Stamps et al., 2021) or in a more localized dextral transfer zone between the offshore branch of the East-African rift and Madagascar (Famin et al., 2020; Feuillet et al., 2021).

The origin of Comorian volcanism is a long-lasting debate (Bachelery and Hémond, 2016; Class et al., 2009; Claude-Ivanaj et al., 1998; Coltorti et al., 1999; Deniel, 1998a; Flower, 1973; Michon, 2016; Nougier et al., 1986; Pelleter et al., 2014; Strong, 1972; Thompson and Flower, 1971; Tzevahirtzian et al., 2021). According to the latest investigations, it is likely associated with lithospheric deformation possibly in relation with the East African Rift System (Courgeon et al., 2017; Lemoine et al., 2020; Michon, 2016; Nougier et al., 1986; Tzevahirtzian et al., 2021), rather than being the result of the interaction of a mantle plume

with oceanic lithosphere (Class et al., 2009, 2005, 1998; Class and Goldstein, 1997; Claude-Ivanaj et al., 1998; Deniel, 1998b; Emerick and Duncan, 1982; Hajash and Armstrong, 1972).

Mayotte, the easternmost part of the archipelago, is composed of a main volcanic island (Grande Terre) and a volcanic islet (Petite Terre), located 4 km east of Grande Terre and built on the modern barrier reef (Nehlig et al., 2013). Mayotte subaerial activity is subdivided into multiple phases of volcanism beginning in the southern part of Grande Terre (10.6 – 1.9 Ma) and then migrating toward the north (5 – 0.75 Ma) and the northeast (0.75 – Present) separated by periods of quiescence (Debeuf, 2004; Pelleter et al., 2014). Magmas emitted on Mayotte show a wider range of differentiation compared to the other islands of the Comoros archipelago, and they define two magmatic series ranging from strongly silica-undersaturated alkaline magmas in the older series to weakly alkaline magmas in the most recent ones (Fig. 2a).

Mayotte is considered the oldest island of Comoros archipelago with a maximum age of 20 Ma for the submarine onset of magmatic activity (Debeuf, 2004; Emerick and Duncan, 1982; Hajash and Armstrong, 1972; Michon, 2016; Nougier et al., 1986; Pelleter et al., 2014; Tzevahirtzian et al., 2021). Despite being the oldest island of the archipelago, Holocene volcanic activity has been documented both on land (Zinke et al., 2003) and on its submarine flanks. In May 2018, an intense seismo-volcanic crisis began on the distal part of a 60 km long NW-SE volcanic ridge that runs off the eastern submarine flank of Mayotte (Fig. 1b; Cesca et al. 2020; Lemoine et al. 2020). This WNW-ESE-oriented volcanic ridge extends from the site of the ongoing eruption in the east to Petite Terre Island in the west (Berthod et al., 2021; Feuillet et al., 2021), see Figure 1). Petite Terre is indeed characterized by the presence of very recent (200 to 4 ka; Zinke, (2003)) phonolitic phreatomagmatic activity with well-preserved maar structures, and strombolian mafic cones aligned on N140 fractures. Strombolian mafic cones are also present in the lagoon between Grande Terre and Petite Terre.

The presence of these recent volcanic structures along the east coast of Mayotte and the ongoing submarine eruption on the east flank of the island raises the question of the possible triggering of a future volcanic activity on the island itself. Indeed, ongoing volcanic seismicity is not only present below the active volcano, 60 km from the island, but the most significant seismic activity is currently recorded 5 – 15 km offshore the island of Petite Terre and below the volcanic ridge (Lemoine et al., 2020; ReVoSiMa, 2021).

Our first petrological study was focused on the lavas erupted at the active volcanic site, which represent an evolved basanite endmember (Berthod et al., 2021). The goal of this study

is to explore the most silicic (phonolite) endmember and the link between both endmembers to constrain the conditions of magma storage, evolution and ascent along the ridge. Our data contribute to image the magmatic system feeding the volcanic ridge east of Mayotte and to constrain the possible future volcanic hazards all along the Mayotte volcanic ridge.

3. Methods

1.1. Samples and dredges

Samples from the active submarine volcanic ridge on the East flank of Mayotte Island were recovered from water depths ranging from 3,200 to 1,000 m by the *R/V Marion Dufresne* in May, June and July 2019, during oceanographic cruises MAYOBS 1, 2 and 4 (Feuillet, 2019; Fouquet and Feuillet, 2019; Jorry, 2019). In addition to those realized on the eruptive site (MAY01-DR01, MAY02-DR08, MAY04-DR10, MAY04-DR11 and MAY04-DR12, Fig. 1b, Berthod et al., 2021), dredges have been performed on four strategic sites identified from high-resolution bathymetric survey of the entire volcanic ridge (Fig. 1 and Tab. 1). The dredged samples were recovered in the proximal part, 10 – 15 km east of Petite Terre Island, on and near a site informally nicknamed “Horseshoe” due to the horseshoe-shape of the submarine volcanic feature (MAY01-DR02, MAY02-DR06, MAY02-DR07 and MAY04-DR09). Three dredges collected samples from individual volcanic cones (MAY01-DR02, MAY02-DR06 and MAY04-DR09). One dredge (MAY02-DR07), was performed on a highly-reflective lava flow on the southeastern part of the “Horseshoe” site (Fig. 1b). Dredged samples were observed and described onboard. We selected unaltered samples with representative morphologies and textures, including quenched pillow rims, pillow cores, and samples containing xenoliths and crystals.

With the notable exception of MAY02-DR06, each dredge collected a very homogeneous set of rock types, a priori sourcing the same volcanic unit. Individual volcanic cones sampled by dredges MAY02-DR06 and MAY04-DR09, are built from lavas characterized by the presence of a chilled margin (1 – 2 cm thick), and an inward increase in vesicle density. A few crystals are observed in MAY02-DR06 and MAY04-DR09 samples. MAY01-DR02 and MAY02-DR07 samples display a glassy thick chilled margin (~ 4 cm), characterized by a rounded, cracked surface and sharp conchoidal fractures. The internal part presents a bedded aspect, which is delimited by elongated and flattened vesicles. These vesicles can reach several centimeters in length and are organized in the plane parallel to the surface of the

sample. Importantly, this facies is also observed in some samples from the MAY02-DR06 dredge.

Samples collected on Petite Terre island are also used for comparison and include a phonolitic glassy sample from the Vigie maar (MAY181215-14b, surrounding a large ultramafic xenolith - MAY181215-14a) and a mafic crystal-rich lithic fragment from a phonolitic pyroclastic deposit at Moya maar (MAY181214-02) that represents the island's basement. Angular, decametric to centimetric mantle xenoliths are found in pyroclastic phonolitic deposits from la Vigie (MAY181215-14a, MAY190911-4a, MAY190911-4b, MAY190911-4c, MAY190911-4d) and Dziani (MAY190911-6a) phonolitic maars on Petite Terre. Smaller xenoliths (cm to mm-sized) are also found in the phonolitic lava flow sampled by dredge MAY02-DR07 (MAY02-DR070207E, MAY02-DR070208E).

1.2. Bulk-rock analysis

Bulk rock compositions were obtained on samples distributed between four dredges and two sampling sites on Petite Terre (Fig. 1b). Major elements were analyzed by ICP-AES on a set of 7 samples at Laboratoire Magmas et Volcans (LMV, Clermont-Ferrand, France). Powdered samples were melted with LiBO_2 in a magnetic induction oven at 1100 °C for 5 min using graphite crucibles. The glass beads were then dissolved in a solution of deionized water and nitric acid (1 M) and diluted by a factor of 2000 to produce the solution analyzed by a HORIBA-Jobin-Yvon ULTIMA C ICP-AES. Analytical uncertainties ($\pm 2\sigma$) range between 1 and 3% except for Na_2O (7%) and P_2O_5 (10%). Four other samples (MAY181214-02, MAY181215-14b, MAY02-DR060201, and MAY02-DR060204) were analyzed at plateforme "Rayons X"- Université de Paris, with an X-ray fluorimeter Epsilon 3xl (Malvern-Panalytical) equipped with an Ag X-ray tube operating under He atmosphere, with 4 conditions during 120s: 5 kV– 60 μA without filter for the analysis of Na, Mg, Al and Si, 10 kV – 30 μA with a 7 μm titanium filter for the analysis of P, 12 kV – 25 μA with a 50 μm aluminium filter for the analysis of Ca, K and Ti, and 20 kV – 15 μA with a 200 μm aluminium filter during 120 s for the analysis of Mn and Fe. In order to avoid matrix and grain size effects, all samples were melted into "beads" prepared by mixing 0.1136 g of sample, 1.2312 g of fluxing agent ($\text{LiBO}_2/\text{Li}_2\text{B}_4\text{O}_7$) and 0.0187 g of non-wetting agent (LiBr) in a platinum crucible. The mixture was heated to 1050 °C for 25 minutes in a fusion instrument (LeNeo fluxer, Claisse). Calibration curves were obtained from identical beads of 14 geological reference materials (ACE, ANG, BCR-2, BEN, BHVO-2, BIR-1, BXN, DTN, FKN, GSN, MAN, Mica-Fe, Osh BO, UBN and BR24). The curves are perfectly linear over

the entire concentration range. Analytical uncertainties ($\pm 1\sigma$) are <5% for TiO₂, MnO and Fe₂O₃, 5% for MgO, SiO₂ and CaO, 10% for Al₂O₃, P₂O₅ and K₂O and 20% for Na₂O.

Trace elements analyses were performed by ICP-MS at Institut de physique du globe de Paris (IPGP, Paris, France). Around 50 mg of powdered rock samples were digested using a mix of 2 ml concentrated HNO₃ and 1 ml concentrated HF, heated in closed Teflon vessels at 110°C for 24 h. An additional 3 ml of concentrated HNO₃ was added to the samples after cooling and the mix was heated for another 24 h at 110 °C. Finally, 45 ml of ultrapure water was added to the samples after cooling and the solutions were sonicated for 4 hours. Samples were analyzed 24 to 48 hours later, after an additional 10x dilution with ultrapure water, using an inert introduction system on an Agilent 7900 ICP-MS. Calibration of rock samples was done against a BEN rock standard (Jochum et al., 2016). Analytical uncertainties are 6% or less for lithophile elements and 15% or less for chalcophile elements.

1.3. Electron microprobe analyses

Composition of minerals and glasses were analyzed using the CAMECA SXFive Tactis electron microprobe at LMV. We used an accelerating potential of 15 kV and a probe current of 15 nA (8 nA for glasses) for major and minor elements, with 10 s counting times. The beam was focused for minerals and defocused to a 20 µm diameter for glasses to avoid Na migration under the electron beam. Natural and synthetic mineral standards, including orthoclase (K, Al), albite (Na), wollastonite (Si, Ca), fayalite (Fe), forsterite (Mg), TiMnO₃ (Ti, Mn), NiO (Ni), Cr₂O₃ (Cr), and fluorapatite (P) were used for routine calibration.

4. Results

4.1. Whole-rock geochemistry

Major elements

Major element compositions are presented in Supplementary Tab. 1. Lavas from Mayotte volcanic ridge belong to the moderately silica-undersaturated trend identified for the Comoros magmas on Mayotte Island and at Karthala volcano on Grande Comore Island (Bachelery et al., 2016; Pelleter et al., 2014). The distribution of lava compositions is bimodal, with basanites sampled both in the distal (active eruptive site, Berthod et al., 2021) and proximal part of the ridge (MAY02-DR06, MAY04-DR09), and phonolites sampled in three dredges on the proximal part of the ridge (MAY01-DR02, MAY02-DR06, MAY02-DR07). Mafic

seamounts, with a SiO_2 content of 43.6 – 46.6 wt% and $\text{Na}_2\text{O}+\text{K}_2\text{O} = 6.2 - 7.9$ wt% plot in the basanite/tephrite field of the TAS diagram (Fig. 2a), and according to their normative olivine content should be classified as basanites. Their low MgO contents, between 4.5 and 6.4 wt%, and high FeO contents, 10.8 and 13.7 wt%, make them evolved basanites (Fig. 2b). Basanitic samples from the two dredges (MAY04-DR09 and MAY02-DR06) scattered along the ridge are very similar to lavas emitted at the 2018-2021 eruption site (Berthod et al. 2021, Fig. 2a). As already observed for the eruptive site (Berthod et al. 2021), evolved basanites of the volcanic ridge are characterized by an enrichment in P_2O_5 (0.9 – 2.0 wt%) compared to subaerial lavas from the Comoros archipelago (Supplementary Tab. 1). The basanitic sample collected on Petite Terre (MAY181214-02) display similar compositions with SiO_2 , $\text{Na}_2\text{O}+\text{K}_2\text{O}$, and MgO contents of 41.9 wt%, 2.9 wt%, and 15.4 wt%, respectively.

Conversely, lava flows sampled in the proximal ridge area (MAY01-DR02, MAY02-DR07) and north of this site (MAY02-DR06, Fig. 1b), fall within the phonolite field of the TAS diagram with SiO_2 contents ranging from 56.1 to 58.2 wt% and a $\text{Na}_2\text{O}+\text{K}_2\text{O}$ content of 11.9 – 13.0 wt% (Fig. 2a). Their low MgO and FeO contents (0.47 – 0.62 % and 6.2 – 6.5 %, respectively) reflect their differentiated character (Fig. 2b). The sample collected on Petite Terre (MAY181215-14b) also plots in the phonolite field with SiO_2 , $\text{Na}_2\text{O}+\text{K}_2\text{O}$ and MgO contents of 57.8 wt%, 5.6 wt%, and 1.2 wt%, respectively.

Only the largest mantle xenolith (MAY181215-14a) has been analyzed for bulk chemistry, and is characterized by a high MgO content of 46.7 wt% and a low alkali content of 0.36 wt%.

Trace elements

The chondrite normalized rare earth element (REE) patterns of the basanitic and phonolitic samples are distinct (Fig. 3a) with different concentrations and show a strong enrichment in light REE (LREE) compared to heavy REE (HREE). However, we note that phonolites are more fractioned in LREE while basanites are more fractioned in HREE. LREE/HREE ratios are lower in phonolites ($(\text{La}/\text{Yb})_{\text{C1}} = 17.45 - 22.97$) than in basanites ($(\text{La}/\text{Yb})_{\text{C1}} = 21.93 - 27.87$). The phonolite REE patterns crosscut those of the basanites in the middle to heavy REE range. In addition, contrary to the basanites, phonolites are characterized by negative europium anomalies.

Basanites and phonolites display different spider diagrams of incompatible trace elements normalized to chondrite concentrations (Fig. 3b). In particular, phonolites are characterized by depletions of TiO₂, Sr, and, for some of them, Ba, compared to basanites.

Phonolites are more enriched in incompatible elements than basanites (Fig. 3c). Indeed, Th content vary from 5 to 9 ppm in basanite and from 15 to 24 ppm in phonolite. With Co content of 24 – 50 ppm in basanite and of < 1 ppm in phonolite, all samples are characterized by a well-defined trend of decreasing Co with decreasing MgO content (Fig. 3d). This is true for all other compatible elements (Cr, Ni...). Scandium contents, ranging from 12 to 16 ppm, do not vary between basanites and phonolites (Fig. 3e). Barium content decrease from 1833 to 329 ppm with increasing SiO₂ content (Fig. 3f).

Phonolite samples collected on Petite Terre display compositions close to submarine phonolites with Th, Co, Sc and Ba of 19 ppm, 1 ppm, 15 ppm and 48 ppm, respectively (Fig. 3). The REE pattern is similar except for the Europium anomaly which is more pronounced. The basanitic sample is characterized by similar Th and Co content (4 ppm and 60 ppm, respectively) and higher Sc content (45 ppm) and a lower Ba content (200 ppm) compared to offshore basanites.

4.2. Petrology

The evolved basanites (MAY02-DR06 and MAY04-DR09) sampled at shallower water depth on the proximal part of the ridge and the mafic clast found scattered in the Moya phonolite maar (MAY181214-02) display a porphyritic texture with olivine and clinopyroxene crystals set in microcrystalline groundmasses (Fig. 4a, b, and c). These evolved basanites contain numerous glomerocrysts of olivine and clinopyroxene, with frequent normal zoning. Some clinopyroxene crystals also show sector-zoning (MAY02-DR06 samples). Olivine and clinopyroxene crystals are sometimes intergrown (Fig. 4b) indicating co-crystallization of these two phases. The groundmass consists of < 150 µm subhedral clinopyroxene microlites, which are occasionally zoned, and < 50 µm oxides. A few clinopyroxene glomerocrysts are characterized by three stages of growth including a rounded core surrounded by a rim with two stages of growth. The last stage is occasionally dendritic as observed in sample MAY02-DR060201m (Fig. 4a).

Dredged phonolites are characterized by a trachytic texture with < 200 µm iron-rich olivine (fayalite) microphenocrysts (Fig. 5). They have rounded (Fig. 5a) and/or skeletal (Fig.

5b) shapes and are intergrown with apatite and oxide crystals (Fig. 5c). The groundmass is composed of thin < 200 µm elongated alkali feldspars, < 100 µm apatite laths and < 50 µm euhedral oxide microlites (Fig. 5a, b, and c). There are two generation of feldspars in some samples (Fig. 5b), with rare microphenocrysts and elongated microlites. One sample (MAY02-DR070207) contains a few large (> 5 mm) rounded sanidine xenocrysts.

Glassy samples collected on land on Petite Terre show a similar trachytic texture and similar mineralogy. There, subhedral iron-rich olivine crystals are set in a groundmass containing < 50 µm feldspar microlites associated with small oxides (Fig. 5d).

Phonolitic samples (MAY02-DR07 offshore and MAY181215-14b on land) contain mantle xenoliths that range in size from a few mm (Fig. 4d) to more than 8 cm. Ultramafic xenoliths embedded in phonolites are mostly composed of large olivine and orthopyroxene crystals and small clinopyroxene and spinel crystals (Fig. 4d, e and f). Some of the orthopyroxene and clinopyroxene crystals contain exsolution lamellae (Fig. 4e and f). One of the xenoliths (MAY02-DR070208E) is crosscut by a vein containing phlogopite, pargasitic amphibole and a Ca-rich feldspar (Fig. 4d). The xenoliths are not in equilibrium with their host phonolite and are surrounded by a reaction rim (Fig. 4d). Reaction between olivine crystals and the phonolitic melt results in an overgrowth of new olivine enriched in iron, whereas reaction between orthopyroxenes and the phonolitic melt produces a thicker reaction-zone containing small olivine and oxide crystals.

4.3. Phase compositions and equilibria

Basanites

Basanites from dredge MAY02-DR06 are almost aphyric, and the small number of phenocrysts show evidence of disequilibrium. Olivine phenocrysts are normally zoned, with a sharp transition between a rounded Mg-rich core (Fo_{80.5–84.7}) and a rim enriched in Fe (Fo_{72.1–74.0}, Fig. 6a, Supplementary Tab. 2). According to Fe/Mg Kd, the rims are in equilibrium with the bulk rock composition whereas the cores are not, pointing to a xenocrystic/antecrystic origin for the cores. Clinopyroxene phenocrysts are also characterized by rounded cores (Fig. 6b, Supplementary Tab. 3), surrounded by a sieve-textured inner rim, and an outer rim with oscillatory zoning. Magnesian cores (Mg# 70.6 – 81.9) are generally enriched in Al (up to 8.8 wt% Al₂O₃) and Na (up to 1.33 wt% Na₂O), and are again not in equilibrium with the bulk rock composition. Rims (Mg# 67.6 – 76.4) have lower concentrations of Al (3.3 – 7.8%) and

Na (0.4 – 0.8 %). In one crystal, the Mg# 73 core contains inclusions of an aluminous spinel (spinel-hercynite solid solution) and amphibole (hastingsite). Oxide crystals are titanomagnetites with 62.8 – 69.3 % FeO and 12.6 – 19.3 % TiO₂ (Fig. 6c, Supplementary Tab. 4).

As for MAY02-DR06, olivine crystals in MAY04-DR09 samples are normally zoned, with a progressive compositional variation from Fo_{85.3} to Fo_{78.7} (Fig. 6a). The highest values indicate that olivine cores are not in Fe/Mg equilibrium with the bulk rock, pointing to partial accumulation of olivine and clinopyroxene, as suggested by the presence of glomerocrysts. Clinopyroxene crystals are also normally zoned, but with more discrete variations and some amount of oscillatory zoning, and Mg# variations from 85.5 down to values as low as 59.4 (Fig. 6b). A few crystals contain rounded cores corresponding to the most magnesian compositions. Also, most of the crystals are characterized by sector zoning. As described by Ubide et al., (2019), Neave et al. (2019) and Haddadi (2016), the hourglass sectors are depleted in minor elements (Ti, Al, Mn, Na, and in our case also Fe), and comparatively enriched in Mg compared to the prism sectors. Oxide crystals are titanomagnetites with 47.5 – 68.5 % FeO and 6 – 15.6% TiO₂ (Fig. 6c). A few chromium-rich spinels with Cr₂O₃ ranging from 12.5 to 21.8% also occur as inclusion in some clinopyroxene crystals.

The basanite sample from Petite Terre, MAY181214-02 is significantly enriched in large clinopyroxene and olivine crystals. Olivine crystals present a restricted compositional range (Fo_{77.7} – 86.6, Fig. 2b). Contrary to samples MAY02-DR06 and MAY04-DR09, olivine cores could be close to equilibrium with the bulk rock (Fe/Mg olivine Kd~0.38) but textural evidence again suggest significant accumulation. Clinopyroxene crystals in this sample are similar to those in MAY02-DR06 and MAY04-DR09 samples, with large variations of Mg# (from 41.8 to 87.9, Fig. 6b) and Al₂O₃ (2.7 – 12.2 wt%). Most clinopyroxene crystals are normally zoned, with a rather abrupt and partially rounded transition between Mg# ~85 cores and more Fe-rich rims (Mg#69-78), suggesting accumulation of Mg-rich antecrysts. One normally zoned crystal containing large apatite inclusions is less magnesium-rich, with a Mg# between 71 and 65.

Phonolites

Olivine crystals in submarine and subaerial phonolites are fayalites (Fo_{17.6} – 31.6 and Fo_{10.1} – 12.3 respectively) with 2.3 – 3.4 and 5.1 – 5.6 wt% MnO (Fig. 6a). They are intergrown with titanomagnetite (up to 18 wt% TiO₂) and apatite. Titanomagnetite crystals contain 72.1 –

78.4 wt% FeO, 10.4 – 18.3 wt% TiO₂, and 2.3 – 3.4 wt% Al₂O₃ (Fig. 6c). Alkali feldspars are mostly anorthoclase (Ab_{49.1 – 71.8}) with An and Or contents ranging from 3.0 to 23.8 wt% and from 10.1 to 44.1 % (Fig. 6d, Supplementary Tab. 5).

Except for P₂O₅ content, compositions of the glasses in most samples display a trend approximating a “liquid line of descent” (Fig. 2). The silica content in dredged phonolites glasses (MAY02-DR06, MAY01-DR02 and MAY02-DR07 dredges) varies from 56.8 to 60.5 % and MgO from 0.19 to 0.41 %. As SiO₂ increases, alkalis (Na₂O + K₂O) increase from 8.7 to 14.1 % and FeO content decreases from 6.4 to 4.4 wt%, respectively. Glasses from phonolites collected on Petite Terre Island present similar but more clustered compositions with SiO₂, MgO, alkalis and FeO contents of 57.6 – 60.1 %, 0.01 – 0.08 %, 14.4 – 15.1 %, and 4.5 – 4.8 %, respectively.

Xenoliths

Ultramafic xenoliths in phonolite lavas and pyroclasts are mantle fragments (harzburgite) dominated by Mg-rich olivine and orthopyroxene, with minor clinopyroxene and spinel. Fo_{90.2 – 93.1} olivine contain very little MnO (0.03 – 0.22 wt%) and low CaO concentrations (0-0.15 wt%), consistent with a mantle origin. Pyroxenes are Mg_{#91.3 – 94.9} clinopyroxene and Mg_{#88.6 – 95.1} orthopyroxene with CaO content of 17.1 – 24.8 % and 0.37 – 2.53 wt%, respectively. Oxides are chrome-spinel with Cr₂O₃ higher than 31% and FeO content ranging from 17.0 to 24.3 wt%. In one xenolith sample (MAY02-DR070208E), a vein contains a few crystals of bytownite plagioclase (An_{76.7 – 79.7}), pargasitic amphibole (Mg_# 92) and phlogopite mica (Mg_# 91.2 – 93.2).

5. Discussion

Both phonolites and basanites are present on the proximal part of the volcanic ridge and onshore, on Petite Terre Island. Phonolites and basanites collected offshore and on land display similar textural, petrological and geochemical features (Figs 2 – 6). Proximal basanites have a porphyritic texture with a microcrystalline groundmass made of clinopyroxene microlites and Fe-Ti oxides (Fig. 4a, b, and c), and glomerocrysts of zoned olivine and sector-zoned clinopyroxene. By contrast, the eastern part of the submarine ridge is only built from basanites characterized by an aphyric texture (ongoing eruptive site, Berthod et al., 2021).

All these samples belong to a moderately silica-undersaturated alkaline series (Fig. 2) which corresponds to the Karthala trend (Bachelery and Hémond, 2016). Also, geographically, all samples belong to a N140 alignment of numerous volcanic features (Fig. 1b, (Feuillet et al., 2021)). This 60 km long volcanic ridge runs on the eastern flank of Mayotte, from the site of a large active 2018-ongoing submarine eruption to the east, to Petite Terre Island to the west (Fig. 1b). The existence of this ridge raises two main scientific questions: (1) What is the origin of the phonolites, are they related to the basanites? (2) Is the entire volcanic ridge, on land and offshore, fed by the same plumbing system or are there numerous ascent pathways and magma storage zones? To answer these questions, it is necessary to understand how this compositional bimodality is achieved.

Since the phonolites contain mantle xenoliths (Fig. 4d, e and f), they were generated below the Moho, i.e., deeper than 17 km (Dofal et al., 2018). Most phonolites are generated by fractional crystallization in shallow crustal magma chambers (< 10 km, Grant et al. 2013 and references therein), and evidence for phonolitic melts at mantle depth is scarce (e.g., Price and Green 1972; Dautria et al., 1983; Grant et al. 2013). It has been demonstrated that phonolites can be directly produced by mantle melting (Irving and Price, 1981; Laporte et al., 2014; Loges et al., 2019). Such phonolites have low FeO concentrations (2 – 3 wt%) and are in equilibrium with high Mg# olivines (~Fo₉₀, Laporte et al. 2014). Phonolitic glass in equilibrium with mantle compositions have been analyzed in lherzolitic mantle xenoliths from La Grille on Grande Comore (Coltorti et al. 1999), and indeed show high Mg# (59 – 78) and low FeO concentrations (<3.1 wt%). The investigated phonolites of the Mayotte ridge, however, are highly evolved, with a significant enrichment in FeO (6.1 – 6.5 wt%) and are in equilibrium with low Mg# olivines (Fo_{13–28}), precluding an origin by direct mantle melting. The differentiated character of those phonolites would thus be better explained by fractional crystallization of a more primitive magma in a mantle reservoir.

5.1. Evidence of fractional crystallization

No primitive composition in equilibrium with a mantle lherzolite has been erupted along the ridge. This indicates the presence of one or more reservoirs in which the magma evolves between the source and the surface. According to our previous studies (Berthod et al. 2021; Feuillet et al. 2021), there are at least two levels of magma storage, one near 40 - 50 km depth in which the primitive magma differentiated to form the evolved basanites, and one upper level, possibly between 15 and 20 km where some more evolved magma is stored.

The evolution of igneous suites by fractional crystallization is characterized by decreasing compatible trace elements whereas incompatible elements increase and have constant ratios with each other (Kyle et al., 1992). According to our major and trace elements data, the fractionating assemblage likely includes clinopyroxene (decrease in CaO and Sc, Fig. 2c and 3e), feldspar (negative Eu anomaly, decrease in Ba and Sr, Fig. 3), magnetite and ilmenite (decrease in FeO, TiO₂, and Co, Figs 2 and 3), apatite (strong decrease in P, Supplementary Tab. 1) and possibly olivine (decrease in MgO, Ni and Co, Figs 2 and 3). There is no evidence for the crystallization of hydrous minerals (amphibole, mica), which usually produce negative anomalies in K₂O, Rb, and High Field Strength Elements like Nb, Ta, Zr and Hf (Matsui et al., 1977; Schnetzler and Philpotts, 1970).

The volcanism of the ridge extending from Petite Terre to the 2018 - 2021 eruption site is strongly bimodal, and no intermediate composition between evolved basanites and phonolites have been sampled so far despite three dredging campaigns and two sampling missions on land. This compositional gap commonly observed in tholeiitic and calc-alkaline series (e.g., Daly (1925), Grove and Donnelly-Nolan (1986), Reubi and Blundy (2009)) is rarely observed in alkaline series. This feature complicates petrological modelling, but does not in any way precludes an origin for the phonolites by differentiation of the basanites.

The possibility of a genetic link between basanites and phonolites was tested through a series of fractionation models: (1) Melts model, for major elements coupled with mass balance and (2) trace element correlation model.

5.2. Fractional crystallization tested with MELTS models and mass balance

Fractional crystallization models were run using the Rhyolite-MELTS software (Ghiorso and Gualda, 2015; Ghiorso and Sack, 1995; Gualda et al., 2012). We used the bulk composition of evolved basanite MAY01-DR0105 collected at the 2018-2021 eruption site, as parental melt, including 2.3 wt% H₂O as measured in olivine-hosted melt inclusions (Berthod et al. 2021). For each model, temperature decreased from 1200 to 700 °C, with a 10 °C step. Since no orthopyroxene and garnet crystals were found in Comoros lavas, they were excluded from the crystallizing assemblage. We tested a range of possible pressures from 0.4 to 1.2 GPa, H₂O concentrations from 1.0 wt% to 5.0 wt%, and a range of oxygen fugacities buffered between FMQ+1 and FMQ-2. Results of our Rhyolite-MELTS models are presented on Fig. 7. All models are mainly characterized by early crystallization of clinopyroxene, apatite and magnetite starting above 1100 °C. A second crystallization stage involves anorthoclase and

biotite with very late-stage leucite in some models. Early crystallization of olivine appears at low pressure and low oxygen fugacity. Ilmenite crystallizes in all models with $\text{H}_2\text{O} \leq 3.0 \%$ and generally appears early at high pressure and low oxygen fugacity. We observed that changes in initial H_2O concentration decreases the crystallization temperatures but only have a limited impact on the crystallization path. An increase in pressure produces a strong increase in FeO and Na_2O concentrations (Fig. 7). This is correlated with the involvement of more clinopyroxene, particularly early on, a decrease in magnetite and ilmenite contents, and absence of olivine. Variations in oxygen fugacity have a similar impact to pressure, with lower oxygen fugacity decreasing the amount of clinopyroxene and magnetite and resulting in higher FeO and Na_2O contents. Therefore, the exceptional FeO content in Petite Terre and submarine ridge phonolites (Fig. 7) may be a consequence of a deeper fractionation level compared to most phonolites (Grant et al., 2013; Moussallam et al., 2013; Scaillet et al., 2008) and/or particularly low oxygen fugacity. This observation is supported by the fact that the best fits to the phonolite composition are obtained at the highest pressures and lowest oxygen fugacities. The overall best model is obtained at 0.6 GPa with 2.3 wt% H_2O and an oxygen fugacity of FMQ-1 (Fig. 7), with a crystallizing assemblage of olivine (12.5 %), magnetite (15.5 %), apatite (5 %), ilmenite (4 %) and clinopyroxene (34 %) at high temperature, and anorthoclase (30%) below 1000 °C. Further crystallization after reaching the composition of MAY02-DR07 / MAY01-DR02 phonolites produces biotite, anorthoclase and leucite. Small amounts of fayalite crystallize in the lower oxygen fugacity models (FMQ-1 and lower). The presence of fayalite microlites in the phonolites is thus likely indicative of low oxygen fugacity, as confirmed by experimental studies (Giehl et al., 2013; Romano et al., 2020; Rutherford, 1969).

The phonolites from the ridge are characterized by an unusual enrichment in iron, compared to older phonolites from the main island of Mayotte (Debeuf, 2004; Pelleter et al., 2014), and most phonolites worldwide (Fig. 7). This iron enrichment, akin to the trend observed in the tholeiitic series (e.g. Carmichael 1964) has already been described in some lavas from the East African Rift (Barberi et al., 1971; Nash et al., 1969) and in the peralkaline rocks of the Gardar province in Greenland (Marks and Markl, 2003). Iron enrichment is usually explained by a late involvement of magnetite in the crystallization sequence (Chin et al., 2018; Luhr et al., 1984), possibly a consequence of low oxygen fugacity (e.g., Marks and Markl 2003, Nash et al. 1969) and/or low water content (Tatsumi and Suzuki, 2009). These

observations are in agreement with the Rhyolite-MELTS model, which suggests reducing conditions, and magma storage and evolution at upper mantle levels.

To confirm the Rhyolite-MELTS model, we performed a simple least-squares mass balance using major elements (Bryan et al. 1969). The calculation starts from the same composition (MAY01-DR0105 sample, Berthod et al. 2021) and ends with the composition of MAY01-DR020202 phonolite. We use microprobe analyses of phenocrysts and microlites from MAY01-DR0105 (olivine, clinopyroxene, magnetite, apatite), MAY02-DR08 (ilmenite) and MAY01-DR02 (anorthoclase and fayalite). The best model implies the crystallization of anorthoclase (50%), olivine (18 %), clinopyroxene (15 %), magnetite (8 %), apatite (5 %), and ilmenite (4 %).

5.3. Fractional crystallization tested with trace elements model

Trace element compositions of basanites found along the submarine ridge are similar to those reported for the 2018 - 2021 ongoing eruption (Fig. 3b, Berthod et al. 2021), and for the entire Comoros archipelago (Bachèlery and Hémond, 2016; Class et al., 1998; Claude-Ivanaj et al., 1998; Deniel, 1998b; Späth et al., 1996).

REE are strongly fractionated ($(La/Yb)_{pm} = 21.9 - 27.9$), indicating a deep peridotitic source with residual garnet, in agreement with previous studies on subaerial Comoros lava that propose a depth of partial melting near the spinel / garnet boundary, i.e. 80 – 100 km, (Bachèlery and Hémond, 2016; Class et al., 1998; Claude-Ivanaj et al., 1998; Deniel, 1998b; Späth et al., 1996) and our previous study of the ongoing 2018-2021 eruption (Berthod et al., 2021).

Phonolite and basanite lavas display similar REE enrichment but with significantly different patterns (Fig. 3a and b). In fact, phonolite REE patterns are less fractionated than the basanites ($(Gd/Y)_{pm} = 3.2 - 5.1$), which could suggest a different shallower source within the spinel stability field (Fig. 8a). In addition, if phonolites are derived from basanites by fractional crystallization, REE patterns should be more enriched than the mafic parental magma. To obtain similar patterns, the fractionated assemblage must include a REE-rich mineral that preferentially fractionate middle REE. All basanitic samples are characterized by an excess of phosphorus, also observed in all glass analyses (Fig. 8a), suggesting that apatite crystallization might play a role. In fact, apatite melt partition coefficients can reach high values (up to 21.7 for La, Luhr et al. 1984; Fujimaki 1986; Mahood and Stimac 1990;

Prowatke and Klemme 2006). Using these high partition coefficients and estimated amount of fractionated apatite from phosphorus concentration, we can reproduce the phonolite REE pattern (Fig. 8b, and Supplementary Tab. 6). Therefore, we suggest that REE ratios do not reflect a different source but are solely a consequence of apatite crystallization.

Phase proportions in the fractionated assemblage were estimated using the model proposed by Villemant et al. (1981). In their model, bulk partition coefficients in a monotonous fractionation series are calculated from the log/log plots of trace elements versus Th, taken as a perfectly incompatible reference element. In Mayotte samples, a few elements (Nb, Ta, Zr, Cs) are more incompatible than Th, so we used Cs, the most incompatible element, instead of Th. This is usually not done because of the high mobility of Cs, however, given the very young age of the investigated rocks, there is a very good correlation between Cs and Th ($r^2 = 0.972$) indicating that Cs mobility was negligible. We then selected a series of trace elements for which (1) the correlation with Th and Cs is good ($r^2 > 0.90$) or the value is constant, and (2) there is a good set of mineral/melt partition coefficient available (partition coefficients taken from Adam and Green (2006), Stix and Gorton (1990), Klemme et al. (2006) and Mahood and Stimac (1990)). We then performed a multiple regression analysis following Villemant et al. (1981) to extract the proportions of minerals in the fractionated assemblage, from the bulk partition coefficients and individual mineral partition coefficient. We only use minerals that have been observed in at least one dredged sample, i.e., olivine, clinopyroxene, magnetite, ilmenite, anorthoclase, and apatite. There is no break in slope in the log/log correlation, suggesting that the fractionating assemblage is relatively constant, at least within the resolution of our model, which is limited by the absence of intermediate compositions in the phonotephrite/tephriphonolite fields. HFSE are among the most incompatible elements, equally or more incompatible than Th, which preclude the crystallization of amphiboles or micas. Phase proportions in the crystallizing assemblage are dominated by feldspar (54 %), clinopyroxene (26 %), olivine (12 %), and apatite (8 %), with a negligible amount of magnetite. These results are in good agreement with the Rhyolite-MELTS model, and the mass-balance calculations.

5.4. Evidence of multiple levels of magma storage

Basanitic mantle reservoirs

As evidenced for lavas emitted by the 2018-2021 ongoing eruption on the eastern part of the submarine volcanic ridge (Fig. 1b), the low Mg# of the basanites indicates that they are not primary mantle melts (Berthod et al. 2021). This is supported by the presence of zoned olivine and clinopyroxene phenocrysts, which typically grow in dynamic reservoirs undergoing frequent replenishments and drainages (Fig. 4a, b, and c).

In addition, our barometric study of the clinopyroxene phenocrysts (Putirka, 2008) indicate multiple levels of magma storage beneath the volcanic ridge (Fig. 9). Since clinopyroxene compositions are neither in equilibrium with the bulk rocks nor in equilibrium with the residual glasses, the Putirka (2008) clinopyroxene-only geobarometer (equation 32a) is the most accurate geobarometer, with a standard error of estimate of 0.28 GPa (9 km). We estimate that the cores of clinopyroxene phenocrysts (Mg_{82-73}) in MAY02-DR06 basanite sample crystallized during a first phase of growth in a basanitic magma reservoir located at about 57 ± 9 km depth (Fig. 9). The location of this deep magma reservoir is in excellent agreement with seismicity location (ReVoSiMa, 2021) and is consistent with the previous geophysical studies (Cesca et al., 2020; Feuillet et al., 2021; Lemoine et al., 2020) and our previous geobarometric study on 2018-2021 erupted lavas which proposed a ≥ 37 km deep mantle reservoir (Berthod et al., 2021). Oscillatory zoned outer rims (Mg_{67-73}) crystallized during a second phase of growth in a shallower more differentiated basanitic magma chamber at an average depth of 18 ± 9 km (0.47 ± 0.28 GPa, Fig. 9).

Similar results have been obtained for sector-zoned clinopyroxene glomerocrysts in the MAY04-DR09 basanite sample. A few deep resorbed clinopyroxene cores (Mg_{83-8} , Fig. 9) crystallized at a pressure of 1.19 ± 0.28 GPa equivalent to a depth of 40 ± 9 km. For the sector-zoned rims, we only used the Al_2O_3 -rich prism sectors for barometry calculations, since hourglass sectors have been shown to produce erroneous pressure (Neave et al., 2015). Except for the resorbed cores, barometry indicate that clinopyroxene crystals ($\text{Mg}_{59-82.5}$) formed at intermediate pressures similar to those obtained for MAY02-DR06: 0.43 ± 0.28 GPa (16 ± 9 km, Fig. 9).

For the MAY181214-02 basanite sample collected on Petite Terre, no evidence was found for a deep fractionation level, and a mean depth of 17 ± 9 km (clinopyroxene $\text{Mg}_{47.9-67.5}$) has been calculated from phenocrysts and rims of the zoned clinopyroxene antecrysts (Fig. 9). Since MAY181214-02 ankaramite corresponds to an old lithic fragment collected on Petite Terre, our results suggest a stable depth for the shallower magmatic system feeding both on land and offshore part of the active volcanic ridge of Mayotte through time.

Our geobarometric results allow us to propose two different mantle storage levels: one deep mantle storage located below 40 km (Fig. 9) and another shallower storage at about 15 – 20 km, near the Moho (Moho depth estimated at ~17 km, Dofal et al. 2018). However, the large range of clinopyroxene compositions and zoning patterns preclude an identical transfer path for each magma pulse.

Phonolitic mantle reservoir

Phonolitic lavas contain clinopyroxene bearing harzburgite xenoliths (Fig. 4d, e and f). These fragments of mantle peridotite, incorporated into the phonolitic magma as it rose to the surface, indicate that they ascended from the mantle. In order to estimate the minimum depth of the phonolitic reservoir(s), we used a geothermobarometer (Putirka, 2008) on a series of clinopyroxene-orthopyroxene pairs in the mantle xenoliths (Fig. 4d, e and f). These geobarometric calculations also provide the maximum depth of the Moho below the investigated area. Temperatures were calculated using equation (36) of Putirka et al. (2008) with reported uncertainties of 45 °C (see Berthod et al., 2021 for discussion). For pressure, both equations (38) and (39) were tested against experiments performed at 1.00 GPa by Condamine and Médard (2014). Since equation (38) returned an average pressure of 1.02 GPa, whereas equation (39) returned a less accurate average pressure of 0.88, equation (38) was thus preferred for the calculation. It reproduces the experimental data with an average error (SEE) of 0.19 GPa. Calculations performed on clinopyroxene-orthopyroxene pairs in textural equilibrium and on exsolutions (Fig. 4d-f) give identical results. Assuming an average density of 2,890 kg.m⁻³ for the oceanic crust and 3,300 kg.m⁻³ for the lithospheric mantle, mantle xenoliths from phonolitic pyroclastic deposits from Petite Terre were equilibrated at the following depths (Fig. 9): 11.9 ± 3.4 km (MAY181215-14a), 12.3 ± 3.7 km (MAY190911-6a), 18.3 ± 5.4 km (MAY190911-4a), 14.5 ± 0.4 km (MAY190911-4b), 12.4 ± 2.9 km (MAY190911-4c) and 14.5 ± 2.1 km (MAY190911-4d). Clinopyroxene-orthopyroxene pairs were also analyzed in two more mantle xenoliths from phonolites of the submarine ridge and give similar equilibration depths of 15.2 ± 2.3 km (MAY02-DR070207) and 11.0 ± 1.3 km (MAY02-DR070208-xen, Fig. 9).

These depths are consistent with seismic data locating the Moho at about 17 km (Dofal et al., 2018), with the existence of numerous very low frequency seismic events between 10 and 30 km depths (Laurent et al., 2019, Feuillet et al., 2021), and with magnetotelluric sounding performed on the volcanic region (Darnet et al., 2020) which suggests the presence of magmatic conductive fluids at depth > 15 km. This is also coherent with the presence of an

evolved magma batch located above 17 ± 6 km (0.45 ± 0.19 GPa) intersected by the June 2019 lava flow at the active volcanic site (sample MAY02-DR08, Berthod et al., 2021). Finally, results from the Rhyolite-MELTS model confirm that differentiation occurred at about 20 km depth (0.6 GPa, Fig. 7).

Since similar minimum storage depths have been obtained for phonolites sampled on Petite Terre and on the intermediate part of the volcanic ridge. This raises the question as to whether these depths correspond to a single magma reservoir or to a zone made of several magma lenses. We suggest that the hundreds seamounts constituting the 60 km-long and 31 km-wide volcanic ridge (Fig. 1b) were probably emplaced between the last volcanic activity at Petite Terre (i.e. about 4000 years ago, Zinke et al. 2003) and the currently ongoing eruption. As shown by Castruccio et al. (2017), monogenetic volcanic fields are likely to be fed by ephemeral superficial magmas lenses rather than long-lived reservoirs.

We note that the evidence of multiple magma storage can also explain the bimodal volcanism. However, the minimum depth of the phonolitic storage obtained from the xenoliths is similar to the shallower basanitic reservoir (Fig. 9) located at about 15-20 km depth. Therefore, we can evaluate the possibility that phonolites are products of differentiation of basanite magmas in the shallower reservoirs located close to the MOHO seismic discontinuity interpreted as the Mantle-crust interface.

5.5. Integration in the geological context of Mayotte: possible eruption triggers?

Berthod et al. (2021) demonstrated that 50 % crystallization of a cumulate containing 80 % clinopyroxene and 20 % olivine is required to generate the basanite from a primary mantle melt in the deeper mantle reservoir (below 37 km). We then propose three scenarios for magma evolution beneath the volcanic ridge: (1) the basanitic magma rises directly and rapidly to the surface from the deep magma reservoir as evidenced by the petrological characteristics of the ongoing eruption (Berthod et al. 2021) and seismological data (Cesca et al. 2020; Feuillet et al. 2021; Lemoine et al. 2020). (2) zoned clinopyroxene phenocrysts in some basanitic lavas at proximal seamounts MAY02-DR06 and MAY04-DR09 and on Petite Terre indicate that the basanitic magma stalls at the Moho interface before resuming its ascent toward the surface. (3) the basanitic magma stops and evolves to phonolite in a reservoir below the Moho. In this third scenario, the phonolitic magma might be reactivated by the arrival of a deeper and more primitive magma sourced from a deep reservoir > 30 km. The

absence of phenocryst in the phonolites might be explained by an increase in temperature induced by the self-mixing reactivation (Couch et al. 2001).

Our three different models show that the phonolitic lavas can be produced by approximately 80 % fractional crystallization of a hydrous basanitic magma at mantle depths ($P > 0.6$ GPa) under reduced oxygen fugacity ($\sim$ FMQ-1). The total volume of MAY02-DR07 lava flow is estimated at 0.45 km^3 , therefore, the volume of basanite necessary to generate this phonolitic lava flow is at least 1.6 km^3 . Taking into account the 50 % crystallization in the deeper mantle reservoir, the eruption of 0.46 km^3 of phonolitic magmas thus requires intrusion of 4.6 km^3 of primary mantle melts.

Since 2018, Mayotte is the site of unprecedented and unexpected volcanic and seismic activities (Feuillet et al. 2021; Cesca et al. 2020; Lemoine et al. 2020). Following an oceanographic campaign, Feuillet et al. (2021) demonstrated that the hundreds of seismic events were associated with the development of a large volcanic edifice. As of January 2021, the volcano has risen 820 m above the 3,300 m deep seafloor with an erupted lava volume of $\sim 6.5 \text{ km}^3$ (Feuillet 2019; ReVoSiMa 2021) which represents by far the largest observed submarine eruption (Clague et al. 2011; Resing et al. 2011; Caress et al. 2012; Chadwick et al. 2016; Carey et al. 2018; Chadwick Jr et al. 2018) and the third largest mafic eruption of the last two millennia (Thordarson and Self 1993; Thordarson et al. 2001; Oppenheimer et al. 2018; Patrick et al. 2020). Berthod et al. 2021 demonstrated that this ongoing eruption is fed by a deep (≥ 37 km) and large ($\geq 10 \text{ km}^3$) mantle reservoir of evolved, alkaline magma having experienced extensive crystallization (50%).

This volcanic edifice is located on a N140 volcanic ridge that runs off the eastern flank of Mayotte and whose subaerial expression is Petite Terre Island (Feuillet et al. 2021, Fig. 1b), a ridge composed of hundreds of volcanic edifices (Fig. 1b). Large phonolitic lava flows are present on this ridge, about 10-15 km East of Petite Terre (Fig. 1b). Pyroclastic phonolitic edifices are also present on the submarine ridge and on Petite Terre. If phonolites derive by fractional crystallization from basanitic lavas similar to those feeding the ongoing eruption, and the structure of the plumbing system is similar all along the ridge, we suggest that the entire volcanic ridge could be fed by the same magmatic system. Moreover, geological mapping of the more than 300 submarine cones and vents clearly show that the zone of the Horseshoe 5-15 km from Petite Terre is the area with the highest density of individual eruptive vents per km^2 of the entire ridge (Fig. 1b). The extensive diversity of eruptive vents, eruptive styles, both explosive and effusive, the magma bimodal composition, the large

erupted volumes and multiple superimposed eruptive cones and lava domes suggests that volcanism in the Horseshoe area has been fed by a long-lasting and extensive transcrustal magmatic system that forms the core of the magmatic system branching out to the West towards the Petite Terre recent Holocene volcanic centers as well as branching out to the East towards the rest of the chain. The results of our petrological analysis coupled with seismic data (Cesca et al. 2020; Lemoine et al. 2020; Feuillet et al. 2021) support the hypothesis that this transcrustal magmatic system is currently undergoing a massive reactivation that is feeding the current exceptional voluminous submarine eruption. In such a model, existing phonolitic magma lenses could be reactivated by reinjection of the basanitic magma feeding the ongoing eruption (Sparks et al. 1977). Although no evidence of mingling or mixing has been identified in the phonolites so far, the presence of reversed zoned olivine phenocrysts in the recently erupted basanite lavas (Berthod et al. 2021) suggest that mixing involving a small shallow differentiated magma reservoir located close to the Moho ($< 17 \pm 6$ km) did occur beneath the current eruption site. This indicate that if eruptible magma is still present in lenses of the transcrustal mush magmatic system located at Moho depth underneath the ridge, it could be reactivated by the arrival of a deeper and more primitive magma. Such a scenario would have major implications in terms of increased volcanic hazard for Mayotte on land. In fact, more than 300,000 people are living on Mayotte with about 70,000 in the capital Mamoudzou located within the Holocene phreatomagmatic explosive volcanic zone, and 30,000 people on Petite Terre Island which hosts the only international airport of the archipelago, the electric plant and other economical and industrial assets as well as most of the public administration.

6. Conclusion

Our petrological study shows that the hundreds of edifices, individual cones and lava flows forming the active volcanic ridge on the east flank of Mayotte Island are characterized by a bimodal chemical distribution of erupted magmas, with basanite and phonolite but no intermediate compositions. Trace element characteristics indicate that those lavas result from partial melting near the garnet – spinel transition zone, between 80 – 100 km, which is consistent with the structuration of lithosphere in the region. Basanitic magmas are first stored and evolve in a deep magma reservoir located between 35 and 57 km. Clinopyroxene barometry highlights the existence of a second shallower storage system at about 15 – 20 km, probably made of multiple lenses of basanitic and phonolitic magmas that form an extensive

and long-lasting transcrustal mush magmatic system. Basanitic magmas evolve to phonolites by about 81 % fractional crystallization in this shallow storage system.

Three scenarios are proposed: (1) the basanitic magma can rise directly and quickly to the surface from the deep magma reservoir (current ongoing activity, Berthod et al. 2021). (2) the basanitic magma stalls in a shallower reservoir near the Moho before resuming its ascent toward the surface. (3) the basanitic magma stops and evolves to phonolite in the shallower magma storage zone. In this third scenario, the phonolitic magma might be reactivated by the arrival of a new batch of deeper basanitic magma. If the entire volcanic ridge is fed by the same extensive plumbing system sourcing the transcrustal mush magmatic system below the Horseshoe area, 5 – 15 km East of Petite Terre, the current reinjection of basanitic magma feeding the ongoing voluminous eruption could reactivate the phonolitic reservoirs that fed the volcanic areas on Petite Terre and the volcanic ridge crossing the eastern flank of Mayotte, drastically increasing volcanic hazards for the island of Mayotte. This is of particular concern given that Berthod et al. (2021) have shown that the timescales that separated the last geochemical perturbation of eruptible magma in a shallow Moho-depth storage zone recorded by zoned crystals, and the arrival of the magma on the surface were on the range of 30 ± 4 days.

References

- Ablay, G.J., Carroll, M.R., Palmer, M.R., Martí, J., Sparks, R.S.J., 1998. Basanite–phonolite lineages of the Teide–Pico Viejo volcanic complex, Tenerife, Canary Islands. *J. Petrol.* 39, 905–936. <https://doi.org/10.1093/petroj/39.5.905>
- Adam, J., Green, T., 2006. Trace element partitioning between mica-and amphibole-bearing garnet lherzolite and hydrous basanitic melt: 1. Experimental results and the investigation of controls on partitioning behaviour. *Contrib. to Mineral. Petrol.* 152, 1–17. <https://doi.org/10.1007/s00410-006-0085-4>
- Bachèlery, P., Hémond, C., 2016. Geochemical and Petrological Aspects of Karthala Volcano, in: *Active Volcanoes of the Southwest Indian Ocean*. Springer, pp. 367–384.
- Bachèlery, P., Morin, J., Villeneuve, N., Soulé, H., Nassor, H., Ali, A.R., 2016. Structure and eruptive history of Karthala volcano, in: *Active Volcanoes of the Southwest Indian Ocean*. Springer, pp. 345–366.
- Barberi, F., Bizouard, H., Varet, J., 1971. Nature of the clinopyroxene and iron enrichment in alkalic and transitional basaltic magmas. *Contrib. to Mineral. Petrol.* 33, 93–107.
- Berthod, C., Médard, E., Bachèlery, P., Gurioli, L., Di Muro, A., Peltier, A., Komorowski, J.,

- Benbakkar, M., Devidal, J., Langlade, J., Besson, P., Boudon, G., Rose-Koga, E.,
 Deplus, C., Le Friant, A., Bickert, M., Nowak, S., Thinon, I., Burckel, P., Hidalgo, S.,
 Jorry, S., Fouquet, Y., Feuillet, N., 2021. The 2018-ongoing Mayotte submarine
 eruption: magma migration imaged by petrological monitoring. Minor revisions, in
 progress. *Earth Planet. Sci. Lett.*
- Bertil, D., Regnault, J.M., 1998. Seismotectonics of Madagascar. *Tectonophysics* 294, 57–74.
[https://doi.org/10.1016/S0040-1951\(98\)00088-2](https://doi.org/10.1016/S0040-1951(98)00088-2)
- Castruccio, A., Diez, M., Ghos, R., 2017. The influence of plumbing system structure on
 volcano dimensions and topography. *J. Geophys. Res. Solid Earth* 122, 8839–8859.
<https://doi.org/10.1002/2017JB014855>
- Cesca, S., Letort, J., Razafindrakoto, H.N.T., Heimann, S., Rivalta, E., Isken, M.P., Nikkhoo,
 M., Passarelli, L., Petersen, G.M., Cotton, F., Dahm, T., 2020. Drainage of a deep
 magma reservoir near Mayotte inferred from seismicity and deformation. *Nat. Geosci.*
 13, 87–93. <https://doi.org/10.1038/s41561-019-0505-5>
- Chin, E.J., Shimizu, K., Bybee, G.M., Erdman, M.E., 2018. On the development of the calc-
 alkaline and tholeiitic magma series: A deep crustal cumulate perspective. *Earth Planet.*
Sci. Lett. 482, 277–287. <https://doi.org/10.1016/j.epsl.2017.11.016>
- Class, C., Goldstein, S.L., 1997. Plume-lithosphere interactions in the ocean basins:
 constraints from the source mineralogy. *Earth Planet. Sci. Lett.* 150, 245–260.
- Class, C., Goldstein, S.L., Altherr, R., Bachèlery, P., 1998. The process of plume–lithosphere
 interactions in the ocean basins—the case of Grande Comore. *J. Petrol.* 39, 881–903.
<https://doi.org/doi.org/10.1093/etroj/39.5.881>
- Class, C., Goldstein, S.L., Shirey, S.B., 2009. Osmium isotopes in Grande Comore lavas: a
 new extreme among a spectrum of EM-type mantle endmembers. *Earth Planet. Sci. Lett.*
 284, 219–227. <https://doi.org/10.1016/j.epsl.2009.04.031>
- Class, C., Goldstein, S.L., Stute, M., Kurz, M.D., Schlosser, P., 2005. Grand Comore Island:
 A well-constrained “low $^3\text{He}/^4\text{He}$ ” mantle plume. *Earth Planet. Sci. Lett.* 233, 391–409.
<https://doi.org/10.1016/j.epsl.2005.02.029>
- Claude-Ivanaj, C., Bourdon, B., Allègre, C.J., 1998. Ra–Th–Sr isotope systematics in Grande
 Comore Island: a case study of plume–lithosphere interaction. *Earth Planet. Sci. Lett.*
 164, 99–117. [https://doi.org/10.1016/S0012-821X\(98\)00195-2](https://doi.org/10.1016/S0012-821X(98)00195-2)
- Coltorti, M., Bonadiman, C., Hinton, R.W., Siena, F., Upton, B.G.J., 1999. Carbonatite
 metasomatism of the oceanic upper mantle: evidence from clinopyroxenes and glasses in
 ultramafic xenoliths of Grande Comore, Indian Ocean. *J. Petrol.* 40, 133–165.

- <https://doi.org/10.1016/j.epsl.2014.04.027>
- Condamine, P., Médard, E., 2014. Experimental melting of phlogopite-bearing mantle at 1 GPa: Implications for potassic magmatism. *Earth Planet. Sci. Lett.* 397, 80–92. <https://doi.org/10.1016/j.epsl.2014.04.027>
- Couch, S., Sparks, R.S.J., Carroll, M.R., 2001. Mineral disequilibrium in lavas explained by convective self-mixing in open magma chambers. *Nature* 411, 1037–1039. <https://doi.org/10.1038/35082540>
- Courgeon, S., Jorry, S.J., Jouet, G., Camoin, G., BouDagher-Fadel, M.K., Bachèlery, P., Caline, B., Boichard, R., Révillon, S., Thomas, Y., 2017. Impact of tectonic and volcanism on the Neogene evolution of isolated carbonate platforms (SW Indian Ocean). *Sediment. Geol.* 355, 114–131. <https://doi.org/10.1016/j.sedgeo.2017.04.008>
- Daly, R.A., 1925. The geology of Ascension island, in: *Proceedings of the American Academy of Arts and Sciences*. JSTOR, pp. 3–80.
- Darnet, M., Wawrzyniak, P., Tarits, P., Hautot, S., D’Eu, J.F., 2020. Mapping the geometry of volcanic systems with magnetotelluric soundings: Results from a land and marine magnetotelluric survey performed during the 2018–2019 Mayotte seismovolcanic crisis. *J. Volcanol. Geotherm. Res.* 406, 107046. <https://doi.org/10.1016/j.jvolgeores.2020.107046>
- Dautria, J.M., Girod, M., Rahaman, O., 1983. The upper mantle beneath eastern Nigeria: inferences from ultramafic xenoliths in Jos and Biu volcanics. *J. African Earth Sci.* 1, 331–338.
- Debeuf, D., 2004. *Étude de l’évolution volcano-structurale et magmatique de Mayotte (archipel des Comores, océan Indien)*. Université de la Réunion.
- Deniel, C., 1998a. Geochemical and isotopic (Sr, Nd, Pb) evidence for plume–lithosphere interactions in the genesis of Grande Comore magmas (Indian Ocean). *Chem. Geol.* 144, 281–303. [https://doi.org/10.1016/S0009-2541\(97\)00139-3](https://doi.org/10.1016/S0009-2541(97)00139-3)
- Deniel, C., 1998b. Geochemical and isotopic (Sr, Nd, Pb) evidence for plume–lithosphere interactions in the genesis of Grande Comore magmas (Indian Ocean). *Chem. Geol.* 144, 281–303. [https://doi.org/10.1016/S0009-2541\(97\)00139-3](https://doi.org/10.1016/S0009-2541(97)00139-3)
- Dofal, A., Fontaine, F.R., Michon, L., Barruol, G., Tkalcic, H., 2018. Crustal structure variation across the southwestern Indian Ocean from receiver functions determined at Ocean-Bottom Seismometers, in: *AGU Fall Meeting 2018*. AGU.
- Dziewonski, A.M., Chou, T., Woodhouse, J.H., 1981. Determination of earthquake source parameters from waveform data for studies of global and regional seismicity. *J.*

- Geophys. Res. Solid Earth 86, 2825–2852.
- Ekström, G., Nettles, M., Dziewoński, A.M., 2012. The global CMT project 2004–2010: Centroid-moment tensors for 13,017 earthquakes. *Phys. Earth Planet. Inter.* 200, 1–9.
- Emerick, C.M., Duncan, R.A., 1982. Age progressive volcanism in the Comores Archipelago, western Indian Ocean and implications for Somali plate tectonics. *Earth Planet. Sci. Lett.* 60, 415–428. [https://doi.org/10.1016/0012-821X\(82\)90077-2](https://doi.org/10.1016/0012-821X(82)90077-2)
- Famin, V., Michon, L., Bourhane, A., 2020. The Comoros archipelago: a right-lateral transform boundary between the Somalia and Lwandle plates. *Tectonophysics* 789, 228539. <https://doi.org/10.1016/j.tecto.2020.228539>
- Feuillet, N., 2019. MAYOBS1 cruise, RV Marion Dufresne. <https://doi.org/10.17600/18001217>
- Feuillet, N., Jorry, S., Rinnert, E., Thinon, I., Fouquet, Y., 2019. MAYOBS cruises, RV Marion Dufresne. <https://doi.org/10.18142/291>
- Feuillet, N., Jorry, S.J., Crawford, W., Deplus, C., Thinon, I., Jacques, E., Saurel, J.M., Lemoine, A., Paquet, F., Satriano, C., Aiken, C., Foix, O., Kowalski, P., Laurent, A., Rinnert, E., Cathalot, C., Donval, J.P., Guyader, V., Gaillot, A., Scalabrin, C., Moreira, M., PelRer, A., Beauducel, F., Grandin, R., Ballu, V., Daniel, R., Pelleau, P., Gomez, J., Besançon, S., Geli, L., Bernard, P., Bachelery, P., Fouquet, Y., BerRl, D., Lemarchand, A., Woerd, J. Van der, 2021. Birth of a large volcanic edifice through lithosphere-scale diking offshore Mayotte (Indian Ocean). *EarthArXiv*. <https://doi.org/10.31223/X5B89P>
- Flower, M.F.J., 1973. Evolution of basaltic and differentiated lavas from Anjouan, Comores Archipelago. *Contrib. to Mineral. Petrol.* 38, 237–260.
- Flower, M.F.J., 1972. Petrology of volcanic rocks from Anjouan, Comores Archipelago. *Bull. Volcanol.* 36, 238–250.
- Fouquet, Y., Feuillet, N., 2019. MAYOBS4 cruise, RV Marion Dufresne. <https://doi.org/10.17600/18001238>
- Fujimaki, H., 1986. Partition coefficients of Hf, Zr, and REE between zircon, apatite, and liquid. *Contrib. to Mineral. Petrol.* 94, 42–45.
- Ghiorso, M.S., Gualda, G.A.R., 2015. An H₂O–CO₂ mixed fluid saturation model compatible with rhyolite-MELTS. *Contrib. to Mineral. Petrol.* 169, 1–30. <https://doi.org/10.1007/s00410-015-1141-8>
- Ghiorso, M.S., Sack, R.O., 1995. Chemical mass transfer in magmatic processes IV. A revised and internally consistent thermodynamic model for the interpolation and extrapolation of liquid-solid equilibria in magmatic systems at elevated temperatures and

- pressures. *Contrib. to Mineral. Petrol.* 119, 197–212.
<https://doi.org/10.1007/BF00307281>
- Giehl, C., Marks, M., Nowak, M., 2013. Phase relations and liquid lines of descent of an iron-rich peralkaline phonolitic melt: an experimental study. *Contrib. to Mineral. Petrol.* 165, 283–304.
- Grant, T.B., Milke, R., Pandey, S., Jahnke, H., 2013. The Heldburg Phonolite, Central Germany: Reactions between phonolite and xenocrysts from the upper mantle and lower crust. *Lithos* 182, 86–101. <https://doi.org/10.1016/j.lithos.2013.09.012>
- Grove, T.L., Donnelly-Nolan, J.M., 1986. The evolution of young silicic lavas at Medicine Lake Volcano, California: Implications for the origin of compositional gaps in calc-alkaline series lavas. *Contrib. to Mineral. Petrol.* 92, 281–302.
<https://doi.org/10.1007/BF00572157>
- Gualda, G.A.R., Ghiorso, M.S., Lemons, R. V, Carley, T.L., 2012. Rhyolite-MELTS: a modified calibration of MELTS optimized for silica-rich, fluid-bearing magmatic systems. *J. Petrol.* 53, 875–890. <https://doi.org/10.1093/petrology/egr080>
- Hajash, A., Armstrong, R.L., 1972. Paleomagnetic and radiometric evidence for the age of the Comores Islands, west central Indian Ocean. *Earth Planet. Sci. Lett.* 16, 231–236.
[https://doi.org/10.1016/0012-821X\(72\)90195-1](https://doi.org/10.1016/0012-821X(72)90195-1)
- Irving, A.J., Price, R.C., 1981. Geochemistry and evolution of Iherzolite-bearing phonolitic lavas from Nigeria, Australia, East Germany and New Zealand. *Geochim. Cosmochim. Acta* 45, 1309–1320. [https://doi.org/10.1016/0016-7037\(81\)90224-6](https://doi.org/10.1016/0016-7037(81)90224-6)
- Jochum, K.P., Weis, U., Schwager, B., Stoll, B., Wilson, S.A., Haug, G.H., Andreae, M.O., Enzweiler, J., 2016. Reference values following ISO guidelines for frequently requested rock reference materials. *Geostand. Geoanalytical Res.* 40, 333–350.
<https://doi.org/10.1111/j.1751-908X.2015.00392.x>
- Jorry, S.J., 2019. MAYOBS2 cruise, RV Marion Dufresne.
<https://doi.org/10.17600/18001222>
- Klemme, S., Günther, D., Hametner, K., Prowatke, S., Zack, T., 2006. The partitioning of trace elements between ilmenite, ulvospinel, armalcolite and silicate melts with implications for the early differentiation of the moon. *Chem. Geol.* 234, 251–263.
<https://doi.org/10.1016/j.chemgeo.2006.05.005>
- Kyle, P.R., Moore, J.A., Thirlwall, M.F., 1992. Petrologic evolution of anorthoclase phonolite lavas at Mount Erebus, Ross Island, Antarctica. *J. Petrol.* 33, 849–875.
<https://doi.org/doi.org/10.1093/petrology/33.4.849>

- Laporte, D., Lambart, S., Schiano, P., Ottolini, L., 2014. Experimental derivation of nepheline syenite and phonolite liquids by partial melting of upper mantle peridotites. *Earth Planet. Sci. Lett.* 404, 319–331. <https://doi.org/10.1093/etroj/40.1.133>
- Le Bas, M.J., Streckeisen, A.L., 1991. The IUGS systematics of igneous rocks. *J. Geol. Soc. London.* 148, 825–833.
- Le roex, A.P., Cliff, R.A., Adair, B.J.I., 1990. Tristan da Cunha, South Atlantic: geochemistry and petrogenesis of a basanite-phonolite lava series. *J. Petrol.* 31, 779–812. <https://doi.org/10.1093/etrology/31.4.779>
- Lemoine, A., Briole, P., Bertil, D., Roullé, A., Foumel, M., THINON, I., Raucoules, D., Michele, M. de, Valty, P., 2020. The 2018-2019 seismo-volcanic crisis east of Mayotte, Comoros islands: seismicity and ground deformation markers of an exceptional submarine eruption. <https://doi.org/10.31223/osf.io/d46xj>
- Loges, A., Schultze, D., Klügel, A., Lucassen, F., 2019. Phonolitic melt production by carbonatite Mantle metasomatism: evidence from Eger Graben xenoliths. *Contrib. to Mineral. Petrol.* 174, 93. <https://doi.org/10.1007/s00410-019-1630-2>
- Ludden, J.N., 1977. The mineral chemistry and origin of xenoliths from the lavas of Anjouan, Comores Archipelago, western Indian Ocean. *Contrib. to Mineral. Petrol.* 64, 91–107.
- Luhr, J.F., Carmichael, I.S.E., Varekamp, J.C., 1984. The 1982 eruptions of El Chichón Volcano, Chiapas, Mexico: mineralogy and petrology of the anhydrite-bearing pumices. *J. Volcanol. Geotherm. Res.* 23, 69–108. [https://doi.org/10.1016/0377-0273\(84\)90057-X](https://doi.org/10.1016/0377-0273(84)90057-X)
- Mahood, G.A., Stimac, J.A., 1990. Trace-element partitioning in pantellerites and trachytes. *Geochim. Cosmochim. Acta* 54, 2257–2276. [https://doi.org/10.1016/0016-7037\(90\)90050-U](https://doi.org/10.1016/0016-7037(90)90050-U)
- Marks, M., Markl, G., 2003. Ilímaussaq ‘en miniature’: closed-system fractionation in an agpaite dyke rock from the Gardar Province, South Greenland (contribution to the mineralogy of Ilímaussaq no. 117). *Mineral. Mag.* 67, 893–919. <https://doi.org/10.1180/0026461036750150>
- Matsui, Y., Onuma, N., Nagasawa, H., Higuchi, H., Banno, S., 1977. Crystal structure control in trace element partition between crystal and magma. *Bull. Minéralogie* 100, 315–324.
- McDonough, W.F., Sun, S.-S., 1995. The composition of the Earth. *Chem. Geol.* 120, 223–253. [https://doi.org/https://doi.org/10.1016/0009-2541\(94\)00140-4](https://doi.org/https://doi.org/10.1016/0009-2541(94)00140-4)
- Michon, L., 2016. The volcanism of the Comoros archipelago integrated at a regional scale, in: *Active Volcanoes of the Southwest Indian Ocean*. Springer, pp. 333–344. https://doi.org/10.1007/978-3-642-31395-0_21

- 943 Moussallam, Y., Oppenheimer, C., Scaillet, B., Kyle, P.R., 2013. Experimental phase-
 944 equilibrium constraints on the phonolite magmatic system of Erebus Volcano,
 945 Antarctica. *J. Petrol.* 54, 1285–1307. <https://doi.org/10.1093/petrology/egt012>
- 946 Nash, W.P., Carmichael, I.S.E., Johnson, R.W., 1969. The mineralogy and petrology of
 947 Mount Suswa, Kenya. *J. Petrol.* 10, 409–439.
- 948 Neave, D.A., MacLennan, J., Thordarson, T., Hartley, M.E., 2015. The evolution and storage
 949 of primitive melts in the Eastern Volcanic Zone of Iceland: the 10 ka Grímsvötn tephra
 950 series (ie the Saksunarvatn ash). *Contrib. to Mineral. Petrol.* 170, 21.
 951 <https://doi.org/10.1007/s00410-015-1170-3>
- 952 Nehlig, P., Lacquement, F., Bernard, J., Caroff, M., Deparis, J., Jaouen, T., Pelleter, A.A.,
 953 Perrin, J., Prognon, C., Vittecoq, B., 2013. Notice de la carte géologique de Mayotte.
 954 BRGM/RP-61803-FR, 135 p., 45 ill., 1 ann.
- 955 Nougier, J., Cantagrel, J.M., Karche, J.P., 1986. The Comores archipelago in the western
 956 Indian Ocean: volcanology, geochronology and geodynamic setting. *J. African Earth Sci.*
 957 5, 135–144. [https://doi.org/10.1016/0899-5362\(86\)90003-5](https://doi.org/10.1016/0899-5362(86)90003-5)
- 958 Pelleter, A.-A., Caroff, M., Cordier, C., Bachèlery, P., Nehlig, P., Debeuf, D., Arnaud, N.,
 959 2014. Melilite-bearing lavas in Mayotte (France): An insight into the mantle source
 960 below the Comores. *Lithos* 208, 281–297. <https://doi.org/10.1016/j.lithos.2014.09.012>
- 961 Price, R.C., Green, D.H., 1972. Lherzolite nodules in a “mafic phonolite” from north-east
 962 Otago, New Zealand. *Nat. Phys. Sci.* 235, 133–134.
 963 <https://doi.org/10.1038/physci235133a0>
- 964 Prowatke, S., Klemme, S., 2006. Trace element partitioning between apatite and silicate
 965 melts. *Geochim. Cosmochim. Acta* 70, 4513–4527.
- 966 Putirka, K.D., 2008. Thermometers and Barometers for Volcanic Systems. *Rev. Mineral.*
 967 *Geochemistry* 69, 61–120. <https://doi.org/10.2138/rmg.2008.69.3>
- 968 Reubi, O., Blundy, J., 2009. A dearth of intermediate melts at subduction zone volcanoes and
 969 the petrogenesis of arc andesites. *Nature* 461, 1269–1273.
 970 <https://doi.org/10.1038/nature08510>
- 971 ReVoSiMa, 2021. Bulletin n°21 de l’activité sismo-volcanique à Mayotte, du 1 au 31 janvier
 972 2021. <https://doi.org/ISSN:2680-1205>
- 973 Romano, P., Scaillet, B., White, J.C., Andújar, J., Di Carlo, I., Rotolo, S.G., 2020.
 974 Experimental and thermodynamic constraints on mineral equilibrium in pantelleritic
 975 magmas. *Lithos* 376, 105793. <https://doi.org/10.1016/j.lithos.2020.105793>
- 976 Rutherford, M.J., 1969. An experimental determination of iron biotite-alkali feldspar

- equilibria. *J. Petrol.* 10, 381–408.
- Scaillet, B., Pichavant, M., Cioni, R., 2008. Upward migration of Vesuvius magma chamber over the past 20,000 years. *Nature* 455, 216–219. <https://doi.org/10.1038/nature07232>
- Schiano, P., Bourdon, B., Clocchiatti, R., Massare, D., Varela, M.E., Bottinga, Y., 1998. Low-degree partial melting trends recorded in upper mantle minerals. *Earth Planet. Sci. Lett.* 160, 537–550. [https://doi.org/10.1016/S0012-821X\(98\)00109-5](https://doi.org/10.1016/S0012-821X(98)00109-5)
- Schnetzer, C.C., Philpotts, J.A., 1970. Partition coefficients of rare-earth elements between igneous matrix material and rock-forming mineral phenocrysts—II. *Geochim. Cosmochim. Acta* 34, 331–340. [https://doi.org/10.1016/0016-7037\(70\)90110-9](https://doi.org/10.1016/0016-7037(70)90110-9)
- Späth, A., Roex, A.P. Le, Duncan, R.A., 1996. The geochemistry of lavas from the Comores Archipelago, Western Indian Ocean: petrogenesis and mantle source region characteristics. *J. Petrol.* 37, 961–991. <https://doi.org/10.1093/petrology/37.4.961>
- Stamps, D.S., Kreemer, C., Fernandes, R., Rajaonarison, T.A., Rambolamanana, G., 2021. Redefining East African Rift System kinematics. *Geology*.
- Stix, J., Gorton, M.P., 1990. Variations in trace element partition coefficients in sanidine in the Cerro Toledo Rhyolite, Jemez Mountains, New Mexico: Effects of composition, temperature, and volatiles. *Geochim. Cosmochim. Acta* 54, 2697–2708.
- Strong, D.F., 1972. Petrology of the island of Moheli, western Indian Ocean. *Geol. Soc. Am. Bull.* 83, 389–406.
- Tatsumi, Y., Suzuki, T., 2009. Tholeiitic vs calc-alkalic differentiation and evolution of arc crust: constraints from melting experiments on a basalt from the Izu–Bonin–Mariana Arc. *J. Petrol.* 50, 1575–1603. <https://doi.org/10.1093/petrology/egp044>
- Thompson, R.N., Flower, M.F.J., 1971. One-atmosphere melting and crystallization relations of lavas from Anjouan, Comores Archipelago, Western Indian Ocean. *Earth Planet. Sci. Lett.* 12, 97–107. [https://doi.org/10.1016/0012-821X\(71\)90060-4](https://doi.org/10.1016/0012-821X(71)90060-4)
- Tzevahirtzian, A., Zaragosi, S., Bachèlery, P., Biscara, L., Marchès, E., 2021. Submarine morphology of the Comoros volcanic archipelago. *Mar. Geol.* 432. <https://doi.org/10.1016/j.margeo.2020.106383>
- Ubide, T., Mollo, S., Zhao, J., Nazzari, M., Scarlato, P., 2019. Sector-zoned clinopyroxene as a recorder of magma history, eruption triggers, and ascent rates. *Geochim. Cosmochim. Acta* 251, 265–283. <https://doi.org/10.1016/j.gca.2019.02.021>
- Zinke, J., Reijmer, J.J.G., Thomassin, B.A., Dullo, W.-C., Grootes, P.M., Erlenkeuser, H., 2003. Postglacial flooding history of Mayotte lagoon (Comoro archipelago, southwest Indian Ocean). *Mar. Geol.* 194, 181–196. [https://doi.org/10.1016/S0025-3227\(02\)00705-](https://doi.org/10.1016/S0025-3227(02)00705-)

Figure Captions:

Fig. 1: a) Location of Comoros archipelago in the Mozambique Channel. b) Geological map of the active submarine volcanic ridge showing the location of our samples offshore and on land. Ages are from (Debeuf, 2004; Emerick and Duncan, 1982; Hajash and Armstrong, 1972; Nougier et al., 1986; Pelleter et al., 2014). DR labels correspond to the dredged samples.

Fig. 2: Geochemical signature of the volcanic ridge east of Mayotte. a) Classification of our samples using the TAS diagram after Le bas and Streckisen, (1991), b) FeO vs MgO diagram, c) CaO vs MgO diagram, d) Al₂O₃ vs MgO diagram. Blue arrows and green arrows respectively show the moderately silica-undersaturated (Karthala-type) and the highly silica-undersaturated (La Grille-type) trends for Mayotte lavas (from Bachèlery & Hémond 2016).

Fig. 3: Trace element variations in basanites and phonolites samples. a) Chondrite-normalized REE patterns, b) Spider diagrams of incompatible trace elements normalized to primitive-mantle concentrations (McDonough and Sun, 1995), c) V vs Th diagram, d) Co vs MgO diagram, e) Sc vs MgO diagram and f) Ba vs SiO₂ diagram.

Fig. 4: Textural and petrological features of basanite lavas sampled offshore on the submarine volcanic ridge (a) MAY02-DR060201m and b) MAY04-DR090104), and c) on Petite Terre Island (MAY181214-12). Mantle xenoliths in phonolites lavas: d) MAY02-DR070208E, e) MAY181215-14a, f) MAY190911-6a) contain large olivine and orthopyroxene crystals and smaller clinopyroxene and spinel crystals.

Fig. 5: Textural and petrological features of phonolite lavas sampled offshore on the active submarine volcanic ridge a) MAY02-DR0701v, b) MAY02-DR060204, c) MAY01-DR0202, and d) on Petite Terre Island (MAY181215-14b). Phonolites have a trachytic texture with laths of alkali feldspar (Afs), and skeletal to subhedral olivine (Fa) microphenocrysts intergrown with apatite (Ap) and oxide (Ox).

Fig. 6: Mineral compositions analyzed in the sampled rocks. a) olivine, b) clinopyroxene and orthopyroxene, c) Fe-Ti oxides, d) feldspar. Our data are compared to minerals in Comoros lavas (Coltorti et al., 1999; Flower, 1972; Ludden, 1977; Schiano et al., 1998; Späth et al., 1996). The two parallel clinopyroxene trends correspond to sector zoning.

Fig. 7: Rhyolite-MELTS models using MAY01-DR0105 basanite as parental magma. Models were run to test if the phonolites can be derived by deep fractional crystallization of the basanites. For each model, temperature decreases from 1200 to 700 °C, with a 10 °C step. We tested a-c) a range of possible pressures from 0.4 to 1.2 GPa, d-f) H₂O concentrations from 1.0 wt% to 5.0 wt%, and g-i) a range of oxygen fugacities buffered at FMQ+1, NNO~FMQ+0.5, FMQ, FMQ-1 and FMQ-2.

Fig. 8: Nature and depth of the source. REE elements could indicate two different sources for basanitic and phonolitic magmas. However, we suggest that REE elements used for this interpretation are impacted by apatite crystallization. b) Model REE patterns obtained by 80% fractional crystallization of a cumulate containing 3-4 wt% apatite (assuming REE are perfectly incompatible in all other phases) compared with phonolite patterns using different partition coefficients (Fujimaki, 1986; Luhr et al., 1984; Mahood and Stimac, 1990; Prowatke and Klemme, 2006).

Fig. 9: Barometric results plotted on a N140 cross section parallel to the volcanic ridge and compared to seismic events (ReVoSiMa, 2021) and magnetotelluric interpretation (presence of melt, Darnet et al., 2020). The latitude and longitude position for each sample is derived from the dredges location. Clinopyroxene barometry from equation 32a of (Putirka, 2008), which has a typical error of 0.28 GPa (9 km). Clinopyroxene/Orthopyroxene barometry in mantle xenoliths from equation (38) of Putirka (2008).

Fig. 10: Cartoon illustrating the magmatic system feeding the active volcanic ridge of Mayotte. The magma formed between 80 – 100 km of depth and then rises until it reaches a deep magma chamber (< 35 km). Using whole-rock geochemistry, Berthod et al., (2021) suggested that the magma have undergone fifty percent of crystallization in this reservoir (twenty percent of olivine and eighty percent of clinopyroxene). Following an internal or external trigger, the eruption begins, and magma rises to the surface. From this stage, we propose three scenarios: In the first scenario the magma rises directly and quickly to the

surface generating the aphanitic basanites (Berthod et al., 2021). In the second, the basanitic magma stalls at the Moho interface before resuming its ascent toward the surface. Finally, in the last scenario the basanitic magma stops and the depth of the Moho and evolves to phonolite with 81% of crystallization (anorthoclase (30%), olivine (12.5 %), magnetite (15.5 %), apatite (5 %), ilmenite (4 %) and clinopyroxene (34 %)) in a Moho reservoir. * Dofal et al., (2018).

Table Captions:

Tab. 1: Dredges and on land sample's locations

Figure 2

[Click here to access/download;Figure;Figure 2.png](#)

Figure 5

[Click here to access/download;Figure;Figure 5.png](#)

Figure 6

[Click here to access/download;Figure;Figure 6.png](#)

Figure 7

[Click here to access/download;Figure;Figure 7.png](#)

Figure 8

Figure 9

<i>Dredges</i>	<i>Oceanographic cruise</i>	<i>DOI : 10.18142/291</i>
DR02	MAYOBS 1	10.17600/18001217
DR06	MAYOBS 2	10.17600/18001222
DR07	MAYOBS 4	10.17600/18001238
DR09	MAYOBS 4	10.17600/18001238

On land sample		X
MAY181214-02	Petite Terre	45.297496 E
MAY181215-14	Petite Terre	45.286375 E

Start dredging			End dredging		
<i>Latitude</i>	<i>Longitude</i>	<i>Depth</i>	<i>Latitude</i>	<i>Longitude</i>	<i>Depth</i>
12°50,44' S	45°22,89' E	1370 m	12°50,35' S	45°22,94' E	1280 m
12°46.27 S	45°23.86 E	1267 m	12°46.27 S	45°23.48 E	1429 m
12°52.35 S	45°24.46 E	1590 m	12°52.08 S	45°23.96 E	1585 m
12° 48.84' S	45°21.42' E	1125 m	12° 48.99' S	45°21.35' E	945 m

Y
12.783116 S
12.800904 S