Deep particle stocks following the summer bloom around the Kerguelen islands: Insights into diatoms physiological state, community structure and mortality modes

Karine Leblanc, Augustin Lafond, Veronique Cornet, Justine Legras, Marie Barbara, B. Quéguiner, Karine Lafond, Barbara Marie

► To cite this version:

Karine Leblanc, Augustin Lafond, Veronique Cornet, Justine Legras, Marie Barbara, et al.. Deep particle stocks following the summer bloom around the Kerguelen islands: Insights into diatoms physiological state, community structure and mortality modes. *Journal of Marine Systems*, 2021, 222, pp.103609. 10.1016/j.jmarsys.2021.103609 . hal-03450616

HAL Id: hal-03450616

<https://hal.science/hal-03450616>

Submitted on 26 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deep particle stocks following the summer bloom around the Kerguelen islands : insights into diatoms physiological state, community structure and mortality modes.

Leblanc^{1*} Karine, Lafond¹ Augustin, Cornet¹ Véronique, Legras¹ Justine, Barbara Marie², Quéguiner¹ Bernard

¹Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille, France

² Sorbonne Université, CNRS, Laboratoire d'Océanographie Microbienne, LOMIC, F-66650, Banyuls/mer, France.

*corresponding author : Karine Leblanc (karine.leblanc@univ-amu.fr)

Abstract

Particles located at the interface between the surface ocean layer and the top of the mesopelagic domain are the initial vector of the biological pump yet, their nature is still largely unknown. During the MOBYDICK cruise in the vicinity of the Kerguelen Islands (Indian sector of the Southern Ocean) we deployed a recently available device that allows to concentrate and collect deep particles over a predetermined layer of water. In this paper, we present the first detailed quantitative and qualitative data of the collected particles and individual planktonic cells, including their taxonomy, carbon and lipid content, as well as cell viability, in order to characterize the particle stocks present at depth. The cruise was carried out at the end of the summer bloom, a period characterized by declining stocks of biogenic materials associated with various mortality processes of planktonic organisms. Unexpectedly, the majority of the collected particles consisted of single empty diatom frustules, whereas fecal pellets and aggregates accounted for only a minor fraction. The occurrence of distinct mortality processes, from parasitic infection to mesozooplankton grazing, and of distinct silicification degrees as well as life-stages could be identified in relation to diatom taxa, evidencing distinct export modes to intermediate and deep layers within the diatom community.

Keywords

Biological carbon pump, diatoms, mortality modes, Southern Ocean, diversity, deep particle stocks

1. Introduction

The global carbon (C) export to the deep ocean is still poorly constrained and estimates range from 5 and 20 Pg C yr⁻¹ (Laws et al., 2000; Henson et al., 2011; Puigcorbé et al., 2020). Despite the overarching importance of this process in Earth climate regulation, the mechanisms responsible for the variability of carbon export are still largely unknown. Characterizing the composition of the particulate C flux to depth together with the intricate ecological processes that can alter particle sinking rates are crucial in this context. Until now, the focus has been placed mainly on large (>500 µm)

aggregates and fecal pellets, otherwise known as marine snow, as primary vectors of C to the ocean's interior, mainly because of their elevated sinking rate, ranging from 5 to 2,700 m d⁻¹ (Turner, 2002; Riley et al., 2012). Methodological challenges to accurately quantify the flux of diverse types of particles over the entire size-spectrum, from µm to mm, have been identified already in the 80s (Alldredge and Silver, 1988) but are still difficult to overcome today, as no single instrument is able to characterize and quantify the entire particle size range together with its species-specific composition (Durkin et al., 2015). Recently, it has been suggested that *in-situ* aggregate size is not a sufficient descriptor for inferring their sinking rates and that more information is needed on their excess density, age, composition and porosity (Iversen and Lampitt, 2020, Laurenceau-Cornec et al., 2020).

Furthermore, the taxonomic composition of phytoplankton, their biomineral content, and the physiological status of cells in the surface layer have been recognized as central in marine snow genesis (Alldredge and Gotschalk, 1990; Klaas and Archer, 2002, Laurenceau-Cornec et al., 2019). Until recently, the importance of single cells, and in particular small phytoplankton (Waite et al., 2000; Richardson and Jackson, 2007), was not recognized because their sinking rate was deemed too slow to lead to any significant contribution to C export. Yet, recent studies report the presence of small particles under the form of single cells down to 1,000 m and deeper (Di Tullio et al., 2000; Dall'Olmo and Mork, 2014; Agusti et al., 2015, 2020; Durkin et al., 2016) revealing a potential important contribution of single cells to deep C stocks. The mechanisms invoked to explain these new observations are usually aggregation/disaggregation processes during descent in the water column, or entrainment through the mixed layer pump, in which deep mixing brings surface cells to depth rapidly, but such processes have not clearly been demonstrated (Gardner et al., 1995; Stemmann et al., 2004; Richardson and Jackson, 2007; Close et al., 2013; Giering et al., 2014; Durkin et al., 2015; Leblanc et al., 2018). A recent study carried out in the North Atlantic however estimated that eddy driven subduction could contribute to as much as 50% of total POC export (Omand et al., 2015). Only a few studies have reported the importance of whole diatom cells and spores in sinking C flux (Kemp et al., 2000; Riaux-Gobin et al., 2006; Salter et al., 2007; Rembauville et al., 2015; Romero and Fischer, 2017), but the exact quantitative contribution of single cells to deep C stocks and fluxes remains unknown (Le Moigne, 2019), as direct C measurements associated to intact phytoplankton cells are time consuming and seldom undertaken (Assmy et al., 2013; Rembauville et al., 2015). It remains unclear whether this deep C stock in the form of small individual cells plays a role in increasing the particulate C flux either directly or through deep aggregation and/or repackaging processes (Lam and Marchal, 2015).

Similarly, fecal pellets are commonly examined in the large size fraction (at least > 50 µm) but the role of minipellets (3-50 µm), as initially described by Gowing and Silver, (1985), has been overlooked. Minipellets can be produced by many different heterotrophs but are often linked to Phaeodaria, a group of siliceous Rhizaria (Gonzalez, 1992). Minipellets have been shown to represent a flux equivalent to 11-49% of the C flux of larger fecal pellets in the Eastern Tropical Pacific Ocean (Gowing and Silver, 1985). This size-class of fecal pellets has been reported as extremely abundant in a series of field studies from the late 80s and early 90s conducted in the

88 Eastern Tropical (Gowing and Silver, 1985) and North Pacific (Gowing, 1986), in the
89 Atlantic (Riemann, 1989) as well as in the Southern Ocean along the Antarctic
90 peninsula (Gonzalez, 1992). These types of particles have been ignored for more than
91 30 years in most biological C pump studies, except for a temporal survey from the
92 Northeast Atlantic, where high summer POC fluxes at 3000 m were shown to follow
93 peaks in phaeodarian abundance (Lampitt et al., 2009). Recently, a renewed interest
94 emerged with the recognized importance of Phaeodaria's role on both the Si and C
95 cycle in the mesopelagic zone in the world ocean (Stukel et al., 2018; Biard et al.,
96 2018). Another recent study in the North Pacific estimated that this group accounted
97 for up to 10% of the total organic carbon of sinking particles (Ikenoue et al., 2019).
98 Hence, marine snow and particles in the small size-range (<50-100 μm) have until
99 recently not been in the spotlight, for obvious methodological issues, and their relative
100 contribution to deep C stocks and fluxes remains to be fully quantified in various
101 oceanic regimes.

102 In 2015, a new device, called the Bottle net (Aquatic BioTechnology™) has allowed for
103 methodological improvements in the collection of deep particle stocks. The device
104 enables a rapid, detailed sampling of depth-integrated concentrated material collected
105 over a very short time period (during a CTD upcast), allowing for taxonomical but also
106 for physiological rate measurements since collection time is equal to that of Niskin
107 samples (Agusti et al., 2015). Collection of particles between 2,000 and 4,000 m with
108 this device during the Malaspina circumnavigation program in the subtropical
109 oligotrophic ocean revealed the ubiquitous presence of fresh single cells in this deep
110 layer. Diatoms, which do not dominate in surface oligotrophic waters, were surprisingly
111 major contributors to microplankton cells at depth (81.5%) and a large proportion (18%)
112 were intact viable cells, implying fast sinking rates estimated to 124-732 m d^{-1} (Agusti
113 et al., 2015) comparable to aggregates and fecal pellets sinking rates. This device
114 allows for a detailed analysis of small (>20 μm) plankton cells and particles in
115 microscopy, while sediment traps designed to measure downward fluxes do not
116 adequately sample this small fraction, which is often merged in larger aggregates and
117 phytodetritus within the collection cups and for which individual C content quantification
118 is difficult. It is complementary to polyacrimide gel traps which also allow to study the
119 contribution of individual cells and chains to the particulate flux (Laurenceau-Cornec
120 et al., 2015).

121 A Bottle net was deployed during the MOBYDICK (Marine Ecosystem Biodiversity and
122 Dynamics of Carbon around Kerguelen: an integrated view) cruise in the vicinity of the
123 Kerguelen Islands in February-March 2018, a period coinciding with the demise of the
124 recurrent summer bloom. The study region is considered as a natural fertilization
125 laboratory (Blain et al., 2008), and several previous studies focused on the large
126 phytoplankton bloom resulting from the island's Fe-enrichment of the plateau area by
127 comparison to the the neighboring HNLC (High Nutrient Low Chlorophyll) region
128 (Cornet-Barthaux et al., 2007; Mosseri et al., 2008; Armand et al., 2008a; Quéguiner,
129 2013; Lasbleiz et al., 2014; Lasbleiz et al., 2016). Here we compare results from
130 phytoplankton nets (Phytonets) and Bottle net casts, while a companion paper
131 describes in similar details the surface diatom communities from CTD Niskin collection
132 (Lafond et al., 2020). We observed a dominant contribution of single cell diatoms (93
133 $\pm 6\%$) to the deep particle stocks but a very small contribution of intact diatom cells

(~0.3 %) to C content in the intermediate layer (125-500 m), together with a very small fecal pellet contribution, that was dominated by the minipellet size-class. Taxonomical analyses revealed distinct communities west of Kerguelen in the HNLC area compared to the island's fertilized plateau and its eastern flank. Differences in silicification degrees as well as distinct mortality/export processes linked to surface nutrient depletion and trophic interactions (such as parasitic infection or grazing by phaeodarians) were identified in the upper layer, leading to distinct contributions of major diatom taxa to deep suspended particles.

2. Material and methods

2.1. Study site

The MOBYDICK cruise took place between February 18th and March 27th, 2018 aboard the R/V Marion Dufresne south of the Kerguelen Islands (Southern Ocean). Four stations were investigated with repeated visits (Fig. 1, Table 1), including two so-called reference stations, M2 (sampled thrice) and M4 (sampled twice), corresponding respectively to the Fe-enriched plateau and the HNLC off-plateau area. To increase information on the spatial variability, two intermediate stations, M1 and M3 were additionally sampled, once and twice respectively. Bottom depths spanned from 520 m at M2 to 4,730 m at M4. According to Pauthenet et al. (2018), station M3 was located within the Polar Frontal zone at this time of the year, whereas the other stations were located in the POOZ (Permanently Open Ocean Zone) of the Antarctic zone, with M1 and M4 both situated very close to the Polar Front (Fig. 1).

Figure 1: Station map of the four sampling sites around the Kerguelen Plateau, with surimposed satellite-Chl a map (monthly composite for March 2018).

Table 1: Metadata and hydrographical conditions at stations sampled during the MOBYDICK expedition.

Station	Latitude/Longitude	Date	Bottom depth (m)	MLD (m)	Ze 1% (m)	H ₄ SiO ₄ :NO ₃ ⁻
M1	49.85°S ; 74.90°E	09/03/2018	2 723	63	89	0.27 ± 0.01
M2-1	50.62°S ; 72.00°E	26/02/2018	520	79	64	0.06 ± 0.02
M2-2		06/03/2018		73	61	0.08 ± 0.03
M2-3		16/03/2018		80	58	0.13 ± 0.01
M3-1	50.68°S ; 68.06°E	04/03/2018	1730	74	93	0.12 ± 0.04
M3-3		19/03/2018		96	105	0.20 ± 0.17
M4-1	52.60°S ; 67.20°E	01/03/2018	4 731	69	95	0.17 ± 0.01
M4-2		12/03/2018		96	101	0.22 ± 0.04

2.2. Phytonet sampling

A 35 µm mesh phytoplankton net (hereafter called Phytonet) was deployed at all sites. Vertical net hauls were consistently realized between 0 and 125 m depth. Once on board, the net was gently rinsed from the outside using a hose connected to surface seawater. The window of the plankton collector was rinsed with a squirtbottle filled with 0.2 µm filtered seawater to unclog all aggregates. The entire remaining volume (between 400-500 mL) was immediately transferred to a polycarbonate bottle, which was then subsampled for diversity and cellular activity.

2.3. Bottle net sampling

A Bottle net (Aquatic Biotechnology™) was mounted on the rosette and deployed at each site. The device is a modified PVC bottle, holding a vertical 20 µm-plankton mesh and a plankton collector, that can be opened and closed between two chosen layers, to collect particles during the upcast (Agusti et al., 2015). Variable sampling depths were covered at each site and revisit depending on bottom bathymetry and CTD-profiles (Table 1, Fig. 2) for a total of 20 Bottle net casts. The intermediate layer (between approximately 125 and 500 m) was sampled at least once at each site, together with deeper layers whenever possible. At station M4-2, two Bottle net casts (125-250 and 250-500 m) were combined for better comparison with the other stations. The Bottle net top opening is round-shaped and similar in size to a standard Niskin bottle (7.5 cm diameter) and the amount of water filtered by the 20 µm mesh varied between 0.1 m³ (for the 125-150 m cast) and 9.5 m³ (for the deep 1900-4000 m cast). Sampling with the CTD was performed at half speed (0.5 m/s) during the Bottle net operations. Given the large body of water filtered, the total amount of particles and cells collected in approximately 50 mL of seawater varied between 26 x 10⁶ and 28 x 10⁷, thus allowing for a complete diversity analysis on concentrated material compared to Niskin bottles that contain only a very low average particle concentration in deep waters.

Blanks. The design of the Bottle net unfortunately renders the closing lid vulnerable to lifting by the up and down pulling motion during the upcast if the rosette is stopped several times in the surface layer in order to close other Niskin bottles. In order to assess the contamination that might arise from plankton rich surface water seeping into the Bottle net during this stop-and-go ascent period, we carried out 5 blank casts (3 at M2 and 2 at M3), with the Bottle net closed during the entire cast. Blank samples were then counted for particles and compared to the average counts measured at both stations. At M2, the blank samples contained 8.7% of the average cell counts and was mostly comprised of *Corethron inerme* filaments which were accumulating in the surface layer, while at M3, blank casts represented 2.7% of the average cell counts. Despite this slight contamination from surface waters, we are confident that our subsequent analyses and comparison to surface communities allowed to filter out this signal and still give valid conclusions on the nature of deep particles and export-related processes.

Figure 2: Bottle net casts depths at the 4 different sampling sites from West to East, and for the different revisits (1,2,3). Bottom bathymetry is indicated in bold italic numbers. The green rectangle indicates the phytonet collection depth in the surface layer from 0 to 125 m.

Collection. Samples were collected from the Bottle net on deck, after careful rinsing of its plankton net with a squirt bottle filled with 0.2 μm filtered seawater. The collected volume varied between 48 and 70 mL of seawater. Each sample was transferred immediately into a polypropylene vial and very gently homogenized before several aliquots were subsampled for biogenic silica (BSi), microscopical identification, diatom viability and lipid content.

2.4. Microscopical analyses

2.4.1. *Taxonomical identification and counting.* Two samples for diversity were fixed with acidified lugol and pH-buffered formol, while a third sample was filtered onto a 25 mm 0.2 μ m polycarbonate filter, rinsed with milliQ water then dried at room temperature, for Scanning Electron Microscopy (SEM) analyses. Upon sample collection from both the Bottle net and the Phytonet, a small aliquot was systematically observed on board within 30 min of collection using a bright-field Nikon TE-200 inverted microscope to observe any trophic behavior or parasitic activity. Further cell counts were later performed in the ground-based laboratory on a Nikon TE-200 inverted microscope, while detailed species identification has been carried out on a Phenom-Pro benchtop scanning electron microscope at 10 kV using the untreated and uncoated dried filters. Species identification relied mostly on Priddle and Fryxell (1985) and Scott and Marchant (2005).

2.4.2. *Mortality processes.* In order to gain further insights into biological interactions and mortality processes in the diatom community, we carefully identified cell status: intact (e.g. cytoplasm still present and visually unaltered), empty (e.g. completely intact frustule with no cytoplasm visible), broken (e.g. with a clean break at the girdle band junction), crunched (e.g. frustule with a jagged break outside of the girdle junction, suppl. Fig. S1), or infected (e.g. frustule filled with small black cells, suppl. Fig. S2). While many mortality pathways can explain the presence of empty and broken frustules, crunched frustules are most likely due to handling by copepod or amphipod mandibles (Assmy et al., 2007). Infected cells were only observed at M2 and refer mainly to the infection of *Corethron inerme* (and to a lesser extent of *Proboscia inerme* and *Rhizosolenia* spp.). Life stages such as resting spores and resting cells (mainly of *Odontella weissflogii*), as well as different morphological/winter forms within some species (e.g. *Chaetoceros atlanticus*, *Eucampia antarctica*) were also identified and counted separately (suppl. Fig. S3).

2.4.3. *Diatom viability.* To further improve the physiological description of diatom cells, we used the SYTOX Green viability probe (Veldhuis et al., 2001) on all Phytonet samples and on 13 out of 20 Bottle net samples. Upon collection, 990 μ L of sample was placed in an Eppendorf PE 1.5 ml vial and spiked with 10 μ L of 500 μ L SYTOX Green Nucleic Acid Stain (S7020, Life Technologies™) 5 mM solution, for a final concentration of 5 μ M and gently agitated. Samples were incubated for 30 min in the dark at *in situ* sampling temperature (4°C), then transferred onto a Sedgewick Rafter graduated 1 mL chamber and immediately counted on board on a Zeiss imager A2 epifluorescence microscope, using an FITC filter cube (λ_{ex} : 479/39 nm, λ_{em} : 522/40 nm, 497 nm LP). Non-viable cells, which have lost membrane integrity, incorporate the probe and their nuclei display a green fluorescence, while viable cells with intact membranes only show chlorophyll *a* (Chl *a*) red autofluorescence.

2.4.4. *Lipid content.* The presence of lipids inside diatom cells was assessed using Nile Red labelling (Greenspan, 1985). Seawater samples were treated on board immediately after collection. Cells were resuspended in 1 ml HEPES buffer 0.1 M (pH 7) containing 2% glutaraldehyde, 10 mM CaCl₂ and 10 mM MgCl₂. After 1 h incubation at 4°C and in the dark, samples were again centrifuged, rinsed and resuspended in 1 ml HEPES buffer 0.1M. Samples were stored at 4°C until analyses at the laboratory. Nile Red was added to each sample at a final concentration of 5 μ g/mL sample,

vortexed for 20 s and incubated 5 min at room temperature. The sample was then mounted onto a glass slide and observed on a Zeiss Observer Z1 epifluorescence inverted microscope using a DS Red filter cube (λ_{ex} : 550/25 nm; beamsplitter 570 nm LP, λ_{em} : 605/70 nm). The percentage of each diatom taxon in the Bottle net samples containing lipid droplets was then determined by scanning a counting a graduated Sedgewick Rafter chamber.

2.4.5. Spore induction and germination experiments

At station M2, a larger amounts of resting spores and winter forms were observed for several taxa such as *Odontella weissflogii*, *Chaetoceros atlanticus*, *Eucampia antarctica* and *Proboscia inerme*, coinciding with a higher Si-limitation at this site with low Si/DIN (Dissolved Inorganic Nitrogen) surface ratios (0.05-0.11). Hence, this site was chosen for exploratory experiments on resting spore induction and germination processes. At the first visit (M2-1), a spore induction experiment was carried out using 6 X 500 μL aliquots collected from the 100-450 m Bottle net sample. All aliquots were resuspended in 15 mL low nutrient water (collected at 30°S during transit to the study area) and three were placed in a dark incubator (low nutrient dark treatment) while the three other were placed in a lit incubator (50% of incoming surface light) cooled with running surface water. After 20 days, samples were fixed with acidified lugol and stored at 4°C. Resting spores and winter stages were enumerated back at the laboratory in all samples in an Utermöhl sedimentation cuve.

On the second visit at this site (M2-2), a spore germination experiment was conducted on the most frequently observed resting spores (*Odontella weissflogii*). Two times 15 resting cells of *O. weissflogii* were isolated under the microscope onboard from the 100-150 m Bottle net samples and resuspended in 10 mL filtered low nutrient water collected underway at 30°S. Both samples were placed in a lit surface incubator with no nutrient addition for the light treatment and +20 μM Si and +1 μM P for the light+nutrient treatment. After 12 days of incubation, samples were fixed with acidified lugol and stored at 4°C. Resting spores and vegetative cells were enumerated back at the ground-based laboratory in all samples in an Utermöhl sedimentation cuve.

2.5. Biogenic Silica (BSi)

An aliquot of 10 mL was filtered for biogenic silica (BSi) onto a 47 mm 0.6 μm polycarbonate filter, which was rinsed with milliQ water and dried at 60 °C for 24 h. Analyses were carried out in the laboratory following the triple NaOH/HF extraction procedure (Ragueneau et al., 2005). This method allows to correct for possible BSi overestimation due to the dissolution of siliceous lithogenic material during the first leaching. Blank values, estimated from measurement of 8 independent samples were $19.3 \pm 14.2 \text{ nmol L}^{-1}$ for biogenic silica, $13.3 \pm 1.6 \text{ nmol L}^{-1}$ for lithogenic silica, and $1.7 \pm 1.0 \text{ nmol L}^{-1}$ for particulate aluminum.

2.6. Carbon content calculations

Carbon content was calculated for diatoms in all Bottle net samples based on intact and broken cells with visible cytoplasmic content, after size and shape measurements in light microscopy (following NF EN 166195, 2015). The carbon conversion formula

304 from biovolume (in μm^3) was derived from Eppley et al. (1970) modified by Smayda,
305 (1978):

306
$$\text{Log}_{10} \text{ C biomass (pg C)} = 0.76 \cdot \text{Log}_{10} (\text{Biovolume}) - 0.352$$

307

3. Results

3.1. Characterization of deep particle stocks

Integrated and absolute particle abundances are presented in Fig. 3 and Table 2 respectively. Within the intermediate layer (i.e. 125-500 m), M1 exhibited the lowest particle (part.) abundances (3.1×10^7 part. m^{-2}) while the HNLC station M4 exhibited the highest abundances (2.6×10^8 part. m^{-2}). In this intermediate layer, abundances decreased at all sites between the first and last visits which occurred 11 to 15 days later, paralleling the demise of the bloom. The intermediate layer at the plateau station M2 could only be sampled between 100-450 m (1st visit), 125-450 m (2nd visit) and 125-375 m (3rd visit) which renders the comparison with the other stations more difficult. Nevertheless, according to the 125-450 m layer sampled during the second visit, we can fairly assume that particle abundances were intermediate (1.1×10^8 part. m^{-2}), being higher than M1 but lower than M3/M4. When expressed in number of particles per cubic meter (Table 2), we observe a clear decreasing trend with depth at all stations, except at M3-1 where particle concentrations were higher within the 125-500 m layer (6.1×10^5 part. m^{-3}) compared to the 60-125 m layer (4.0×10^5 part. m^{-3}). In the deep layers (i.e. > 500 m), particle concentrations ranged between 5.0×10^4 part. m^{-3} at M1 to 1.9×10^5 part. m^{-3} at M4-2.

Figure 3: a. Integrated particles per m^{-2} for diatoms, other microplankton and fecal pellets+aggregated for all Bottle net casts at the four study sites. The integration depth is indicated on the left axis. b. Relative percent contribution of main lineages to the other microplankton groups (excluding diatoms) in all Bottle net. c. Relative percent contribution of different types of inert particles : aggregates, minipellets <50 μm , and three different types of fecal pellets (fp.).

Table 2 : Particle concentration (total cells + aggregates + fecal pellets) in particles m⁻³ for all Bottle net samples according to station and sampling interval.

Station	Integrated sampling depth (m)	Particle concentration (particle m ⁻³)
M1	60-125 m	468 032
	125-500 m	83 680
	500-2500 m	49 826
M2-1	100-450 m	499 935
M2-2	100-150 m	1 267 622
	125-450 m	331 988
M2-3	125-375 m	307 361
	125-375 m	236 546
M3-1	60-125 m	398 300
	125-500 m	606 904
	500-1500 m	151 480
M3-3	125-500 m	245 226
	200-1500 m	113 800
M4-1	125-150 m	2 523 000
	125-500 m	689 680
	150-1900 m	148 016
	1900-4000 m	135 767
M4-2	125-500 m	600 736
	500-1000 m	189 719

Regarding the type of particles collected, one of the most striking features was the dominance of diatom cells at all sites and depths (Fig. 3a). On average, diatom cells represented 93 ± 6 % ($n = 19$) of the total particle amount, while other microplankton groups only represented 5 ± 4 %. Surprisingly, particles such as fecal pellets and aggregates only contributed to 2 ± 3 % of the total particle amount. The relative abundance of microplankton other than diatoms was the highest at M1 reaching up to 16 % of the total particle amount within the 60-125 m layer. Aggregates, mostly large-sized (100-400 μm) were the most numerous at M2-3 (125-375 m layer) contributing to 14 % (1.0×10^7 part. m⁻²) of the total particle amount while they did not exceed 1 % at the other stations. Interestingly, those large aggregates were not observed within the same layer sampled 8 h later, suggesting a short-lived export event.

Among identified microplankton groups, excluding diatoms (Fig. 3b), tintinnids (35 ± 12 %) and dinoflagellates (34 ± 21 %) were dominant at most sites and depths. Silicoflagellates was the following most abundant group (16 ± 15 %). They were notably found to increase with depth, and were the highest for the two deepest casts at M4-1 (48 % at 150-1900 m, 60 % at 1900-4000 m) and at M1 (26% at 500-2500 m). Siliceous Rhizaria was the next most abundant group with a large diversity of species belonging to different lineages (e.g. mainly Nassellaria, Phaeodaria, and Taxopodia). Nassellaria were present at all stations (8 ± 6 %) with the most abundant species belonging to the Theoperidae, Plagoniidae and Artostrobiidae groups. A rarely mentioned Taxopodia, which only has one described species (*Sticholonche zanclea*) was also very frequent at all sites and depths (5 ± 5 %). Phaeodarians were also often present though in minor proportions (2 ± 2 %) and were mostly represented by several *Protocystis* species (e.g. *P. tridens*, *P. swirei*, *P. balfouri*, *P. harstoni*, *P. micropelcus*). Although minor contributors to the deep water particle stocks, they should play an important role in the production of minipellets (Fig. 3c, Fig. 4a).

Among inert particles (Fig. 3c), identifiable fecal pellets (round, ovoid or elongated) were negligible (10% on average for all casts) compared to aggregates and minipellets which both constituted 45% on average of the remaining particles (see suppl. Fig. S4 for pictures of different types of particles). Minipellets are typically < 50 µm wide and are known to be excreted by the *Protocystis* genus (Phaeodaria) (Gonzalez, 1992) that was very abundant in surface Phytonets (data not shown). The occurrence of Phaeodaria and their trophic behavior will be further discussed in another paper (Leblanc et al., in prep). Here, minipellets were mostly constituted of diatom fragments, mostly belonging to the species *Fragilariopsis kerguelensis* (Fig. 4a, suppl. Fig. S4b), while *Protocystis* spp. were repeatedly observed in the Phytonet samples with one or several whole ingested *F. kerguelensis* cells. Small aggregates (<100 µm) were the most dominant type of aggregates and were also tightly packed with crushed diatom debris, as well as coccoliths (*Emiliana huxleyi*) in some samples. Fecal pellets imaged by SEM revealed a very high content in biominerals, mostly diatom frustule debris (Fig. 4).

Figure 4: a. Phaeodarian minipellets, b. ovoid fecal pellet, c. long fecal pellet, d. zoom on a round fecal pellet content. e. Loose fecal pellet containing recognizable debris of *Fragilariopsis kerguelensis* (pink) and centric diatoms (turquoise). A few coccoliths are also visible (purple).

3.2. Diatom physiological state

The state of the diatom cells within the upper 0-125 m layer sampled by the Phytonet and the deeper layers sampled by the Bottle net is presented in Fig. 5. Within the upper 0-125 m layer, diatoms were mostly intact (62 ± 12 %), while the remaining frustules were broken (24 ± 11 %) or empty (9 ± 6 %). No large differences were observed between stations, except at M4-1 where more broken frustules were observed (43 %). At the reference plateau station M2, a parasitic infection event was observed at the three visits, affecting almost exclusively the species *Corethron inerme* and to some lesser degree *Proboscia* and *Rhizosolenia*. The percentage of infected frustules varied between 14 % (1st visit) to 9 % (3rd visit) within the upper layer. At depth, infected cells were only observed at the last visit (i.e. M2-3, 21 % of the cells) within the 125-375 m layer. Interestingly, those infected cells were associated with large aggregates and were not found in the following cast performed 8 hours later.

In Bottle net samples, the most conspicuous feature was the decreasing contribution of intact cells with depth, at all stations and visits. Within the intermediate layer (125-500 m), frustules were mostly empty (49 ± 5 %) or broken (26 ± 6 %), while intact cells were a minority (23 ± 8 %). In the deep layers located below 500 m, only few intact cells were still observed, contributing to 5 ± 1 % of the total cells. Crunched frustules indicative of mesozooplankton grazing were rare in most samples (< 3 %), although they were more frequent at M1, with a contribution of 5-6 % within the 60-2500 m layer.

Figure 5 : Observed state of diatom cells in the Phytonet samples (upper pannel) and in the Bottle net samples (lower pannel) at each site and revisit. Collection depth is indicated below each pie chart. See method section for “crunched” and “infected” definition.

Viability test

Results from the SYTOX Green labelling experiment showing the percentage of viable cells quantified on board immediately after sampling are presented in Table 3. The percentage of viable diatom cells decreased at all sites between revisits, in both Phytonets and Bottle nets. Within the upper 0-125 m layer, the percentage of viable diatoms was the highest at station M3 (64 % then 47%) indicating a good physiological state of the cells, while it was the lowest at the eastern station M1 (18 %) and above the plateau at M2, where a decreasing trend from 47 % (1st visit) to only 6 % (3rd visit) highlights the decaying stage of the diatom population, even though visually intact cells represented a much larger contribution (Fig. 5). In the intermediate layer at M2, viable diatoms in the Bottle nets closely followed that of the Phytonets (43, 25 and 8 %). At the HNLC station M4, viable diatoms represented only a small third of total cells (28% then 26%). A larger proportion of live cells (64%) was found in a subsurface layer (125-150 m) at station M4-1, and this proportion was still fairly elevated in the deeper casts upon the first visit, with 29 and 24 % of viable cells in the 150-1900 and 1900-4000 m casts respectively. On the second visit however; this proportion dropped substantially, with only 4 % viable cells between 250-500 m and none in the 500-1000 m layer, also suggesting the decay of the diatom population.

Table 3: Percentage live diatom cells within total diatom cells counted after SYTOX labelling at each site, revisit and integrated sampling depth for the surface phytonet samples and for Bottle net casts.

Station	Integrated sampling depth (m)	% live diatoms
M1	0-125	18
	125-500	11
	500-2500	1
M2-1	0-125	47
	100-450	43
M2-2	0-125	19
	125-450	25
M2-3	0-125	6
	125-375	8
M3-1	0-125	64
	125-500	20
M3-2	0-125	47
	125-500	12
M4-1	0-125	28
	125-150	64
	125-500	51
	150-1900	29
	1900-4000	24
M4-2	0-125	26
	250-500	4
	500-1000	0

3.3. Diatom community structure within surface and deep layers

Relative abundances

Diatom relative abundances from samples collected by both the Phytonet and the Bottle net are presented in Fig. 6. Results show a clear difference in diatom communities between the two stations located above the plateau (M2) and east of Kerguelen (M1) and the two stations located west of Kerguelen (M4 and M3). Within

the upper 125 m layer, the Fe-fertilized plateau was dominated by *Corethron inerme* (81 ± 12 %) and *Proboscia* spp. (mostly *P. alata*; 9 ± 3 %). At M2, the Phytonet contents closely matched the surface layer niskin observations (Lafond et al., 2020) showing increasing abundances at each revisit of extremely long *Corethron inerme* filaments (up to 7-8 mm long, up to 74 cells in one filament) in the mixed layer (~80 m). At M1, some *Corethron* cells (13 %) were again observed in the surface Phytonet, but this genus was not dominant, while *Fragilariopsis kerguelensis* (42 %) and centric species constituted the bulk of diatom diversity.

In the intermediate layer (125-500 m) at M2 and M1, the Bottle net samples reflected a very similar community, that was dominated by a mix of *Eucampia antarctica* (30 ± 23 %), *F. kerguelensis* (26 ± 16 %), and *Corethron* spp. (19 ± 7 %), although the importance of *E. antarctica* declined throughout the survey period, while the contribution of *C. inerme* increased with time at M2. Species belonging to the genus *Pseudo-nitzschia* (mostly *P. heimii*) were also present contributing up to 18 % of the abundances at the last visit (125-375 m layer) and up to 25% in the deeper cast at M1 (500-2500 m). These results suggest that M1 was partly influenced by the nearby plateau despite its higher bathymetry which is coherent with the water mass circulation (Park et al., 2014).

By contrast, the surface layer of the western stations M3/M4 was dominated at the first visits by *Chaetoceros atlanticus* (64 ± 12 %), progressively replaced by a mix of different taxa at the second visit: *Chaetoceros atlanticus* (20 ± 11 %), other centrics (14 ± 9 % ; mainly unidentified centrics), *Actinocyclus/Thalassiosira* spp. (13 ± 16 %), *Dactyliosolen antarctica* (9 ± 4 %), *Proboscia* spp. (9 ± 1 %), *Rhizosolenia* spp. (9 ± 1 %), and *Corethron* spp. (8 ± 4 %). Diatom communities were much less diverse within the intermediate and deep layers and largely dominated by *Chaetoceros atlanticus* (71 ± 11 %), followed by *F. kerguelensis* (11 ± 5 %). Other *Chaetoceros* spp., mostly *C. dictyota* were also non negligible contributors to abundances at M4 (8 ± 2 % at both visits).

Winter stages and resting spores

The heavily silicified *Chaetoceros atlanticus* was a key species at the western stations M3/M4. It occurred under several forms: long chains composed of vegetative cells, a solitary stage with sigmoidal setae, which is likely a resting cell form, and under the 'bulbosum' form, which is thought to be the resting spore stage for this species (suppl. Fig. S3). In the surface layer, the vegetative stage was dominant at M4/M3/M1 but the solitary sigmoidal stage was also present even though less abundant. Interestingly, the 'bulbosum' stage was never observed in the Phytonets but was abundant in the subsurface Bottle net samples at M4-1 (125-150 m), M4-2 (125-250 m), M3 (65-125 m) and M1 (65-125 m), suggesting a sporulation event occurring below the mixed layer, although we cannot exclude that the 'bulbosum' form was not retained by the 35 µm mesh size of the Phytonet. The 'bulbosum' stage was also present albeit at lower abundances in almost all the deepest Bottle net casts.

Figure 6: Relative contribution of diatom cells main taxa to abundance (comprising both intact and empty cells) in the Phytonet samples (upper pannel) and in the Bottle net samples (lower pannel) at each site and revisit. Collection depth is indicated below each pie chart.

Winter stages of another heavily silicified species *Eucampia antarctica* (suppl. Fig. S3), as small rectangular forms (Fryxell and Prasad, 1990), were quasi absent from the Phytonet samples, which could also be due to the mesh size exceeding their average size (15-30 µm), but were on the other hand very abundant in the subsurface casts starting below 100 m at M2 and M1 in the Bottle net samples. This small winter form was minor at M3 and quasi absent at M4.

Several *Odontella weissflogii* morphotypes were also observed during the cruise. The vegetative lightly silicified stage was quasi absent from all samples, while a rectangular form, much more silicified than the vegetative stage and with short apical horns was the dominant form. This stage has been termed as the resting spore stage of *O. weissflogii* (Scott and Marchant, 2005) but no complete description of this stage is available. We also observed a larger rounder stage with even shorter apical horns, and therefore hypothesize that the rectangular form may be a resting cell or transition stage towards the actual resting spore, which would be the round form (suppl. Fig. S3). Both forms were absent at station M4 and M3, while the resting stage was most abundant at M2 and M1 in the upper Bottle net casts. The round spore stage was on average 10 times less abundant than the rectangular stage and was found at the same sites.

The small *Proboscia inermis* winter stages and abnormal forms (10-25 µm) of this stage (suppl. Fig. 2 in Lafond et al., 2020), were the next contributors to winter/resting

stage but were much less abundant, even though present at all sites, with a dominance at M4.

Several forms of *Rhizosolenia* with heavily silicified otaria- and clasper-lacking forms, such as *Rhizosolenia polydactyla f. squamosa* are thought to be resting spore/stage employed for overwintering but there is still a debate whether these morphotypes are winter resting stage or actual resting spore (Armand and Zielinski, 2001 and refs therein). This stage was present but as a very minor contributor in Phytonets and in one shallow Bottle net.

Figure 7: Relative contribution of diatom cells main taxa to C biomass (comprising intact cells only) in the Phytonet samples (upper pannel) and in the Bottle net samples (lower pannel) at each site and revisit. Collection depth is indicated below each pie chart.

Relative carbon biomass

Taxon-specific contribution to carbon biomass of intact cells only is presented in Fig. 7. Above the Kerguelen Plateau at M2, carbon biomass follows the same trend as abundances, with *C. inerme* (77 ± 8 %) and *Proboscia* spp. (12 ± 6 %) being the main contributors within the upper 125 m layer. Below 125 m, *Eucampia antarctica* contributed to 69 % of C biomass at the first visit, before its biomass decreased significantly at the third visit, when it was replaced by *C. inerme* (67-83 %). *Odontella weissflogii* as resting cells and/or spores contributed to 10 % at the first visit, and decreased with time, while *F. kerguelensis* contributed to 29 % at the second visit.

At the eastern station M1, *Corethron* spp. (24 %) was also an important contributor in the upper 125 m layers, but other taxa also drove the C biomass: *Rhizosolenia* spp. (19 %, e.g. *R. simplex*, *R. curvata*), and other centrics (18 %, e.g. *Asteromphalus hookeri*, *Actinocyclus octonarius*, *Azpeitia tabularis*, *Coscinodiscus* spp., unidentified centrics etc.). Below the surface layer, key species contributing to C biomass were similar to M2 (i.e. *C. inerme*, *E. antarctica*, *F. kerguelensis*), although species belonging to *Actinocyclus/Thalassiosira* complex became important (17 ± 3 % within the three sampled layers) and the species *Dactyliosolen antarctica* dominated C biomass in the intermediate 125-500 m layer (41 %).

At the western stations M3/M4, the biomass of large centric species became dominant although they were numerically few. Indeed, within the upper 125 m layer, *Proboscia* spp. (mainly *P. alata*), *Rhizosolenia* spp. (e.g. *R. curvata*, *R. polydactyla f. polydactyla*), *Actinocyclus/Thalassiosira* spp. (e.g. *A. curvatulus*, *A. octonarius*, *T. lentiginosa*, *T. tumida*), and other centrics (e.g. *Asteromphalus hookeri*, and unidentified centrics) contributed together to 78 ± 8 % of C biomass. In the intermediate layer, *Chaetoceros atlanticus* (39 ± 21 %) was the main contributor to C biomass, followed by the large centrics *Corethron* spp. (22 ± 19 %, mainly *C. pennatum*), *Proboscia* spp. (11 ± 12 %, mainly *P. alata*), and *Dactyliosolen antarctica* (8 ± 6 %). A similar pattern extended to the deep layers with *Chaetoceros atlanticus* still driving C biomass, except within the deepest layer sampled during the cruise (M4-1, 1900-4000 m) where *Fragilariopsis kerguelensis* contributed up to 43 % of the C biomass.

3.4. Si and C concentrations and integrated stocks in the intermediate and deep layers

BSi and diatom C in Bottle nets

Average and integrated biogenic silica and diatom carbon concentrations over the Bottle net sampling layers are presented in Table 4. Average concentrations of diatom C measured from intact cell counts were expectedly highest in all subsurface casts, with the highest value of $556 \mu\text{g C m}^{-3}$ over 100-150 m at M2-2 (main contributor *E. antarctica*), followed by M4-1 over 125-150 m with $389 \mu\text{g C m}^{-3}$ (main contributor *C. atlanticus*). M1 and M3 subsurface casts over 60-125 m contained much less intact diatoms, with 128 and $74 \mu\text{g C m}^{-3}$ respectively. Diatom C content then decreased at all sites with depth and reached the lowest values ($1-6 \mu\text{g m}^{-3}$) in all casts extending to over 1000 m depth. The highest concentrations for the intermediate layer (between 125 and 500 m) were found at M2-1 and M4-1, but decreased upon revisits to the same stations. Integrated diatom C values ranged between 1 and 40 mg C m^{-2} over the considered sampling layer, the highest diatom C content was measured at M2-1 (40 mg C m^{-2}) over 100-450 m followed by M4-1 (33 mg C m^{-2}) over 125-500 m. The lowest value for the intermediate layer was measured at M1 (2 mg C m^{-2} over 125-500 m).

Outside the plateau, BSi concentrations in the intermediate layer were highest at M4 (46 mg Si m^{-3}), followed by M3 ($18-12 \text{ mg Si m}^{-3}$) and M1 (15 mg Si m^{-3}). At M2, it reached 49 mg Si m^{-3} at the third visit within the 125-375 m layer. Integrated concentrations were highest at M4-1 in the intermediate ($17,272 \text{ mg Si m}^{-2}$) meso- ($42,277 \text{ mg Si m}^{-2}$) and bathypelagic ($28,374 \text{ mg Si m}^{-2}$) suggesting an efficient export of Si in HNLC waters. At M2, integrated concentrations increased steadily between the

first (2,088 mg Si m⁻²) and last visit (12,168 mg Si m⁻²) although the integrated depth was lower, meaning the Si stock increased significantly at depth.

Table 4: Amounts of intact diatom C (from microscopical counts) and total biogenic silica (from chemical measurements) in Bottle net samples (>20 µm) calculated as an average concentration (per m⁻³) in the considered layer, and as integrated values over the sampling depth (per m⁻²). For comparison, integrated Si and C stocks measured from discrete Niskin samples (> 0.4 - 0.7 µm) in the surface layer and over comparable depths are indicated in grey.

		Bottle net				Niskin	
		[diatom C]	[BSi]	diatom C	ΣBSi	ΣBSi	ΣPOC
		µg C m ⁻³	mg Si m ⁻³	mg m ⁻²	mg Si m ⁻²	mg Si m ⁻²	mg C m ⁻²
M4-1	0-125 m					16 405	4 409
M4-1	125-500 m					6 753	5 747
M4-1	125-1000 m						10 840
M4-1	125-150 m	389		10			
M4-1	125-500 m	89	46.1	33	17 272		
M4-1	150-1900 m	7	24.2	13	42 277		
M4-1	1900-4000 m	2	13.5	4	28 374		
M4-2	0-125 m					19 537	4 164
M4-2	125-1000 m					12 801	11 256
M4-2	125-250 m	148		19			
M4-2	250-500 m	13	24.1	3	6031	5 193	
M4-2	500-1000 m	6		3			
M3-1	0-125 m					14 066	3 983
M3-1	125-1000 m					11 190	9 590
M3-1	60-125 m	74		5			
M3-1	125-500 m	38	18.0	14	6753		
M3-1	500-1500 m	2	4.8	2	4780		
M3-3	0-125 m					11 916	3 209
M3-3	125-1000 m					10 811	8 544
M3-3	125-500 m	14	12.0	5	4848		
M3-3	200-1500 m	3		4			
M2-1	0-125 m					11 706	5 270
M2-1	125-350 m					8 207	7 786
M2-1	100-450 m	115	6.0	40	2088		
M2-2	0-125 m					9 464	4 355
M2-2	125-450 m					6 410	5 649
M2-2	100-150 m	556		28			
M2-2	125-450 m	30	25.2	10	8179		
M2-3	0-125 m					8 034	5 986
M2-3	125-450 m					4 961	6 841
M2-3	125-375 m	76		19			
M2-3	125-375 m	35	48.7	9	12168		
M1	0-125 m					9 943	3 977
M1	125-1000 m					6 897	9 177
M1	60-125 m	128		8			
M1	125-500 m	5	15.0	2	5639		
M1	500-2500 m	1	5.6	1	11236		

Comparison of integrated Si stocks between Niskin and Bottle net samples

Biogenic silica and particulate organic carbon were measured chemically from discrete Niskin samples and integrated over depth layers identical to the Bottle net whenever possible for comparison. For BSi, this direct comparison over the exact same sampling layer was only possible on two CTD casts at M4-1 (125-500 m) and M2-2 (125-450 m). Results presented in Table 4 show that ΣBSi in the Bottle net samples ($\Sigma\text{BSi}_{\text{BN}}$) were higher (2.5 and 1.3 times) than ΣBSi in Niskin bottles ($\Sigma\text{BSi}_{\text{Niskin}}$) which was unexpected, considering that the Bottle net mesh size was 20 μm , while Niskin samples were filtered onto 0.4 μm . The Niskin and Bottle net mouth openings are very close (~ 7 cm), and should not lead to different collection efficiencies. In this study, we can not conclude on whether the Bottle net collection efficiency is comparable to that of Niskin samples or if differences were only an effect of filtration size (both filter pore size and small vs large filtered volume effect). In future deployments, BSi and POC from Niskin bottles should be filtered onto 20 μm in parallel for a direct comparison with the Bottle net samples.

Last, a comparison between diatom C estimated from microscopical counts of Bottle net samples and $\Sigma\text{POC}_{\text{Niskin}}$ over the same depth range was possible for 6 casts (except at M1) and showed that diatom C in Bottle net samples represented only 0.1 to 0.6 % of $\Sigma\text{POC}_{\text{Niskin}}$ over the 125-500 or 125-1000 m layer.

3.5. Diatom lipid content

The proportion of each diatom species containing lipid droplets (as stained with Nile Red) is presented in Fig. 8. *Fragilariopsis kerguelensis* appears to be the taxon that is most likely to sink out of the surface layer with lipids, and its contribution to total stained cell was major (70-80%) at M4-1 and M1. *Chaetoceros atlanticus*, mostly in the form of single cells undergoing a sigmoidal stage or as spores (*C. atlanticus bulbosum*) was the next major taxon containing lipids, particularly at M3. At M2, *Eucampia antarctica* as well as *Corethron inerme* were also often showing lipid droplets, their relative contribution being similar to the evolution of total frustule abundance over the three visits. Finally, *Odontella weissflogii* resting stages/spores were also important lipid contributors during the first two visits at M2.

Figure 8: Relative contribution of main diatom taxa to lipid content, based on enumeration of cells positively labelled with Nile Red. Taxa contributing to less than 3 % of lipid containing cells were pooled together in the category “other spp.”

3.6. Spore germination and induction experiments at M2

Spore germination. At M2-2, 2 X 15 resting cells were collected from a subsurface Bottle net opened between 100 and 150 m. After 12 days incubation, the aliquot incubated with both light and nutrients (+Si+P) showed a successful germination of all resting stages, and cell count amounted to 6,072 of actively dividing vegetative *Odontella weissflogii* cells (Table 5). Cells looked healthy and no empty frustules were found. In the light only treatment, the 15 initial cells yielded 3 times less vegetative cells, with a total of 2,048 cells. Some resting cells remained in the sample (10), and could be either the ungerminated initially isolated resting cells or new resting cells formed over the course of the incubation. In addition, 758 larger rounder forms of *Odontella* resting cells, suggesting transition towards a resting spore stage, were counted in this treatment only, accounting for another 1/3 of cells. Several abnormal valves and half vegetative/half resting spore cells were also observed (see suppl. Fig. S5). Assuming that all resting cells germinated in the first treatment, the total amount of >6,000 cells in 12 days is achieved with a growth rate of 0.5 d⁻¹. In the light only treatment, assuming that only 5 out of the 15 resting cells germinated (since 10 intact resting cells were still observed in the sample), the final amount of >2,000 cells is also achieved with a growth rate of 0.5 d⁻¹.

Table 5: Results of the spore germination experiment at M2-2 after 12 days incubation of 2 replicate samples containing 15 resting cells/spores of *Odontella weissflogii* in a light + Si + P and in a light only treatment.

Spore germination experiment	Light +Si +P		Light only	
	Live	Empty	Live	Empty
<i>Odontella weissflogii</i> vegetative	6072	0	2048	78
<i>Odontella weissflogii</i> resting cell	0	2	10	6
<i>Odontella weissflogii</i> round form	0	0	758	0
Total <i>Odontella weissflogii</i> cells	6072	2	2816	84

Spore induction. At M2-1, 6 X 500 µL aliquots from the 100-450 m Bottle net were incubated in low nutrient water (LNW) + light or in low nutrient water and dark conditions for 20 days. Results are summarized in Table 6. In the LNW +dark treatment, about 30% more *Odontella weissflogii* resting cells and round forms were observed at the end of the incubation, showing a higher probability to form spores in both nutrient and light deprived environments. A similar trend was observed for *Eucampia antarctica* winter forms, which were 3 times more abundant in this treatment as well.

Table 6: Results of the spore induction experiment at M2-1 after 20 days of incubation of 2X3 replicate samples collected from phytonets and incubated in low nutrient water (LNW) in surface light conditions and in the dark.

<i>Spore induction experiment</i>	LNW + light treatment		LNW + dark treatment	
	Live	Empty	Live	Empty
<i>Odontella weissflogii</i> resting cells	64	81	91	65
<i>Odontella weissflogii</i> resting spores (round form)	4	9	9	5
<i>Eucampia antarctica</i> winter form	99	305	297	500

4. Discussion

4.1. Nature of the deep particle stock

The number of identifiable individual diatom frustules collected with the Bottle net in deep waters around Kerguelen islands by the end of summer was not negligible (10^7 - 10^8 cells m^{-2}) and the deep diatom stock (below 125 m) was one order of magnitude higher than the stock present in the upper (0-125 m) layer (Lafond et al., 2020). It was also 3 orders of magnitude higher than the concentration of single phytoplankton cells measured on average by Agusti et al. (2015) in the 2,000-4,000 m layer in the subtropical Ocean. This is not surprising considering the large annual diatom bloom event occurring around Kerguelen Islands due to natural Fe-fertilization, which has been documented during previous cruises in the same area (Armand et al., 2008a; Lasbleiz et al., 2016) and the shallower sampling depths. The export of diatom cells at depth was also expected considering that the MOBYDICK cruise occurred at the end of summer during the demise of the bloom, as previously documented from sediment trap samples in the same region (Rembauville et al., 2015), from sediment trap material over the Crozet plateau region (Salter et al., 2012; Salter et al., 2007) or further into the Australian sector of the Antarctic Zone (Rigual-Hernández et al., 2016; Rigual-Hernández et al., 2015). On the other hand, the dominance of single diatom cells (93%) over any other type of particles such as fecal pellets, phytodetritus and aggregates was unexpected. We have no other elements of comparison regarding the contribution of aggregates and fecal pellets in the subtropical ocean, as these were not reported in Agusti's Bottle net study.

Given the size of the dominant taxa observed in our study (20-50 µm), it is unlikely that single cells sank to depth without being integrated to larger aggregates. It is possible that the Bottle net sampling could have promoted the disaggregation of phytodetritus during the upcast and water flow through the 20 µm net, potentially explaining the very low contribution of aggregates and fecal pellets. However, the state of cells observed in microscopy directly after sampling on board in each sample, the absence of large

mucus/TEP aggregates and the morphological shape of diatoms and other cells (frustules and setae intact) suggest that these particles were likely not aggregated at the time of sampling. The similarity of the taxonomic diatom composition in the Bottle nets compared to the above surface layer phytonets (Fig. 6) furthermore suggests a rapid export mode of diatoms, which has already been hypothesized in the same region with observations of similar taxonomic composition between the surface layer and sediment taps moored at 300 m (Blain et al., 2020). We therefore hypothesize that these cells sank rapidly out of the surface layer, included in loose phytodetritus or aggregates which disaggregated *in situ* prior to collection, due to shear stress in the water column or to bacterial mineralization of TEP/colloids holding aggregates together.

Clearly, these cells were not exported as fecal pellets, given for instance the state of *Chaetoceros atlanticus* sigmoidal stages with unbroken setae, particularly in the deep cast (1900-4000 m) at M4-2. Microscopic examinations also allowed to observe loose aggregates still holding together as well as intact tightly packed fecal pellets (suppl. Fig. S4) which suggests that the Bottle net sampling was not disrupting particles or aggregates in a major way.

On one occasion, at one of the two 125-375 m casts performed at M2-3, numerous large aggregates were observed (14 % of total particles, but representing practically the entire inert particle fraction) together with cells covered with loose organic material. Unfortunately, it was complicated to estimate their C biomass due to their heterogeneity (e.g. density, composition etc.), but their contribution to C biomass was likely not negligible for this cast, since most of the other particles were empty diatom frustules. Interestingly, those large aggregates were not observed in the exact same layer 8 hours later (decrease in abundance by a factor X100), indicating a very brief export event at a high sinking rate. This station was sampled few days after a strong storm event which could have promoted both their physical aggregation through vertical mixing and their rapid export. Their disappearance 8 hours later further suggests an export event in the form of loose aggregates, that were not cohesive enough to stay aggregated below the surface layer. This observation could reconcile our data showing mostly single cells in the the intermediate layers, with the mechanism of rapid export proposed by Blain et al. (2020), which would lead to a rapid flushing of the surface layer of diatoms in the form of aggregates, once reached a critical cell concentration and stickiness threshold value. In another study carried out in the Antarctic Zone, sediment trap data results suggested a fast and undisturbed transport of particles between 2000 and 3700, and sinking velocities of $210 \text{ m}^2 \text{ d}^{-1}$ of the major diatom taxa were estimated (Rigual-Hernández et al., 2015).

Phytodetrital aggregates were previously reported to contribute significantly to export fluxes in this region. During ANTARES I, a thick fluff layer containing high pigment levels (up to $13 \mu\text{g L}^{-1}$) was observed west of Kerguelen at more than 3,000 m depth (Riaux-Gobin et al., 1997), while during spring (KEOPS 2 cruise), phytodetrital aggregates were the most abundant type of particles collected by polyacrylamide gel sediment traps (Laurenceau-Cornec et al., 2015). A study of surface sediment samples in the same area during the spring bloom in 2005 (KEOPS cruise) also revealed a number of intact diatom frustule on the seafloor, suggesting an efficient transport mode

of single cells at depth (Armand et al., 2008b), but which could also be the result of an aggregation/sinking/disaggregation sequence.

Finally, due to the very short period of particle collection (during a CTD upcast), we cannot exclude that we missed other important pulse export events such as the one observed at M2-3, especially at the end of the productive season which is often characterized by an intense and brief export event called the "fall dump" (Kemp et al., 2000; Quéguiner, 2013).

4.2. Mortality modes

Parasitic infection

During the cruise, the diatom contribution to C biomass was the highest above the plateau at M2 (Table 4). An important development of long filaments of *Corethron inerme* occurred in the course of three successive visits over nearly a month, and resulted in a doubling of diatom C biomass over the 0-100 m layer. This species was also mixed with a few (<10%) other large-size tubular centrics such as *Rhizosolenia* spp. and *Proboscia* spp, which likely occupied the same ecological niche. These species, with frustules relatively thinner than most other diatom taxa present at the other sites, grew at the station most depleted in silicic acid (1.2 to 2.6 μM), which could indicate their adaptation to low Si levels. During live observations on board, parasitic infection of many cells was clearly visible, with some cells filled with small black parasites < 1 μm in size (suppl. Fig. S2). This was the only site (except a single occurrence at M4-2) where such parasitic infection was observed. Single cell sequencing of isolates picked on board and co-occurrence network analyses suggest that these parasites belong to the Syndiniales group, which was not previously known to infect diatoms (Sassenhagen et al., 2020). Interestingly, at M2-1, *Corethron* was only a minor contributor to the intermediate layer stocks, which was initially dominated by *Eucampia*. The relative contribution of *Corethron* at intermediate depth increased over time, probably reflecting the sinking of infected cells, as is visible on the third visit with 15% of infected *Corethron* cells observed in the 125-375 m layer. The percentage of viable cells in the surface samples decreased from 47 % on the first visit to 6 % on the last (Table 2), reflecting massive cell mortality, which is the likely result of this parasitic infection. The apparition of infected cells below the euphotic layer at M2-3 could be the result of subsequent sinking of cells losing their membrane integrity and their buoyancy, but could also be linked to a strong storm event occurring between M2-2 and M2-3 that disrupted the surface layer and induced mixing with the intermediate layer. It is however not clear from our results if the parasitic infection led to preferential sinking of *Corethron* empty cells, or if this was just the result of the temporal increase of *Corethron* abundance over the three visits and mixed layer disruption following the storm. It would seem more likely that infected cells are rather lysed and remineralized in the surface layer through the viral shunt than massively exported at depth. However, a recent study conducted in the North Atlantic indicated that viral infection of a ballasted biomineral containing phytoplankton such as *Emiliania huxleyi* appeared to stimulate vertical export flux and rather enhanced the biological C pump (Laber et al., 2018). Other types of infections were observed during live observations of net material on board, such as small swimming nanoflagellates cells

invading *Rhizosolenia* spp. cells, or parasites flowing out of a *Ceratium lineatum* cell, which suggests that parasitic infections were actively occurring over the plateau at the end of the productive season. Zoosporic and fungal parasitic infection, in addition to viral and bacterial infection, are still a large unknown in our comprehension of diatoms as well as other phytoplankton's bloom termination and are thought to be much more abundant than reported in the literature (Gutiérrez et al., 2016; Scholz et al., 2016). Si limitation has furthermore been shown to facilitate viral infection of diatoms in a highly productive coastal system (Kranzler et al., 2019). In our study, parasitic infection was also observed at the most Si limited station (Si:N ratios of 0.05-0.11) and we hypothesize that the thinner frustule together with nutrient limitation could also facilitate the parasitic infection of large cylindric centrics such as *Corethron*, *Rhizosolenia* and *Proboscia*.

Grazing processes by microzooplankton

The high contribution of both dinoflagellates and tintinnids in the other microplankton compartment and the dominance of small-sized (<100 µm) aggregates over fecal pellets in the Bottle net samples is also more consistent with the dominance of an active microbial food web reflecting the demise of the bloom rather than with a more developed mesozooplankton compartment and longer food chains. Interestingly, minipellets (<30 µm) produced by Rhizaria were a major part of the detrital stock in all Bottle net samples. Their origin was confirmed by direct observations on board (suppl. Fig. S4) of active grazing on diatoms by the phaeodarian group *Protocystis* spp. (mostly *P. swirei* and *P. tridens*). This group of phaeodarians is known to feed on bacteria, algae and also fecal pellets or organic matter aggregates. Numerous small-sized (80-160 µm) *Protocystis* were observed ingesting whole cells of *Fragilariopsis kerguelensis* and excreting minipellets comprised between 10-30 µm, filled with *F. kerguelensis* debris as well as with other undetermined cells. Such an active grazing by phaeodarians on diatoms associated to the production of minipellets has seldom been reported except in older studies in the Pacific Ocean (Gowing and Silver, 1985), and in the Southern Ocean in the Antarctic peninsula (Gonzalez, 1992) and in the Weddell Sea (Gowing, 1989). Despite their small numerical abundance, this group, together with other siliceous Rhizaria, could have an important contribution to the Si stocks measured in the Bottle net, given their elevated skeleton Si content (in our study 2 nmol Si cell⁻¹) compared to diatoms, and their ability to concentrate Si debris in minipellets (Nakamura and Suzuki, 2015 and references therein; Leblanc et al., in prep.). The ecological role of these phaeodarians is likely to be significant as evidenced by previous studies around the Antarctic peninsula (Gonzalez, 1992), but it has received little attention until very recently, when new studies based on DNA metabarcoding have revealed the global importance of Rhizaria (Stukel et al., 2018) and phaeodarians in the particle vertical export (Gutierrez-Rodriguez et al., 2019). In a survey from 1988 in the Scotia Sea and the Weddell-Scotia confluence, integrated minipellets reached 10⁶ m⁻² values and were 5 orders of magnitude more abundant than krill feces (Gonzalez, 1992). In the Weddell Sea, phaeodarians were in turn actively grazed by salps such as *Salpa thompsoni* (Gowing, 1989), thereby linking the microbial food web to higher trophic levels. Other phaeodarians such as *Phaeodina*

antarctica were also observed in our study and agglutinated both whole diatom valves and silicoflagellates skeletons, resulting in elevated particulate Si cell⁻¹. Hence Rhizaria, through their siliceous skeletons and fecal pellets, and specifically minipellets, appear as active contributors to the downward Si flux at the end of the productive season.

Grazing processes by mesozooplankton

Mesozooplankton abundance was on average lower during the spring cruise (58 to 1,249 x10³ ind. m⁻²) that took place in 2010 in the same area (KEOPSII), compared to this study (64 to 860 x10³ ind. m⁻², A. Delegrange pers. comm.) reflecting the lower food availability in the surface layer at the end of summer. Despite the relative small contribution of fecal pellets to the detrital deep stocks in the Bottle net samples, signs of active mesozooplankton grazing (probably by copepods which was the dominant zooplankton group or amphipods like *Themisto gaudichaudii* frequently observed from micronekton sampling) were clearly indicated by the presence of crunched frustules debris outside as well as inside fecal pellets. Potential prey selection was observed, as crunched frustules were only observed for *Fragilariopsis kerguelensis* and several centric diatoms such as *Thalassiosira lentiginosa*, *Azpeitia tabularis* and *Asteromphalus hookeri* (suppl. Fig. S1). Other dominant diatoms such as *Eucampia antarctica*, *Corethron inerme* or *Chaetoceros atlanticus* were never observed with crunch marks, even though they were observed in SEM in fecal pellets, suggesting different types of grazers or feeding behaviors. During the EIFEX Fe-fertilization experiment, Assmy et al. (2013) reported that the thick Si frustule of *F. kerguelensis* made it less palatable for mesozooplankton and crunched end cells were interpreted as a result of copepods being deterred by the high energy expenditure needed to crush these preys (Hamm et al., 2003). Yet, in our study, fecal pellets observed in SEM also contained a number of crushed (and sometimes whole) *F. kerguelensis* valves together with other similarly heavily silicified diatoms such as *Eucampia antarctica*, *Chaetoceros atlanticus*, *Corethron criophilum* as well as many small intact *Chaetoceros* resting spores indicating active grazing and ingestion (Fig. 4). It is possible that the high lipid content of these species (Fig. 8) made them more palatable for zooplankton upon entering the winter diapause period. The SEM observations of fecal pellets <100 µm and the diversity that is observed thanks to many identifiable frustules (Fig. 4) suggest that even diatoms with thick frustules are actively grazed by copepods and other mesozooplankton species. However, we were not able to identify whether the preferential downward pathway for *F. kerguelensis* was sinking as single cells or included in fecal pellets. Furthermore, pulverized frustule debris were commonly observed during on board examinations but could not be quantified. Improved techniques would be needed to isolate fecal pellets, quantify their relative Si vs. C contents together with a precise taxonomic identification of frustule remains. Clearly, the Si/C ratio of these fecal pellets should be disproportionnally high compared to the single cell flux, even if it was dominated by empty frustules, if one considers the hundreds of siliceous frustules tightly packed into each fecal pellet.

4.3. Life stages : resting spores and winter forms

Diatom resting stages were more frequently observed in the HNLC waters west of Kerguelen and over the plateau than at station M1, and were only observed in centric species. The winter stage/resting spore formation is usually preferentially observed in centric diatoms and in the coastal environments, possibly allowing cells that have sunk to the sediments to resurface upon the next mixing event (McQuoid and Hobson, 1996). At M4 and M3 a particular solitary stage of *Chaetoceros atlanticus*, with sigmoidal setae (which we hypothesize is a resting cell form) and in lesser abundance, the spore form (*C. atlanticus* f. *bulbosum*) were dominant. On the plateau area, the abundance of the *C. atlanticus* complex was minor and may have resulted from some eastward advection, mixing some of these cells with the local community.

The plateau, which was the most Si-limited region at the end of summer (1.2 to 2.6 μM H_4SiO_4 and H_4SiO_4 :total $\text{N}_{\text{mineral}}$ ratios as low as 0.05-0.11), was dominated by the winter stage of *Eucampia antarctica* in the process of sinking out of the surface layer following a previous bloom, as they were more abundant in intermediate layer Bottle nets than in the surface Phytonets. Our results are consistent with observations reported for the Crozet Islands diatom export study in a similar comparison of an Fe-fertilized plateau vs. an Fe-limited station further off in the open ocean (Salter et al., 2012). In this study, the contribution of *Eucampia antarctica* viable heavily silicified and well preserved winter stages to diatom export flux at depth on the plateau was dominant (up to 71% contribution), while the vegetative stage was almost entirely absent. As nitrate was non-limiting, the authors suggested that low dissolved Fe and silicic acid concentrations at the end of the growth season were plausible triggers for the development of the winter stage and its subsequent sinking, which was tightly correlated to the enhanced C flux measured on the plateau (Salter et al., 2012). Winter forms of *Eucampia antarctica* were also abundant on the sediment floor of the Kerguelen Plateau (Armand et al., 2008b), confirming the important role of this emblematic species to the C pump in Fe-fertilized Southern Ocean island areas.

This species also co-occurred with *Odontella weissflogii*, which were likely sporulating, as resting cells and spores were present in the shallow Bottle net samples and much less abundant at depth than *E. antarctica*. Microscopic observations on board revealed healthy looking spores (packed with lipids), which further confirms the recent sporulation and on-going sedimentation event.

The exploratory sporulation/germination experiments conducted at M2 strongly suggest that Si availability was the key trigger for spore formation, since the light treatment yielded 3 times less vegetative cells of *O. weissflogii* than the Si+light treatment (Table 5). Furthermore the number of abnormal valves and general state of germinated cells were a clear indication of Si limitation, while nitrate was still sufficient ($>21 \mu\text{M}$). In the sporulation experiment the light deprivation in low nutrient waters stimulated a 30 % increase in *O. weissflogii* resting stages, while *E. antarctica* winter stages increased by a factor of 3 (Table 6), which could reflect different species-specific responses to environmental triggers.

Small *Thalassiosira* and *Chaetoceros* spores ($<20 \mu\text{m}$) were abundant in Niskin bottle samples below the mixed layer at M2 and M1 (Lafond et al., 2020) but were neither

observed in Phytonets nor Bottle nets because of too large mesh sizes (35 and 20 μm). Even though in small abundances, they however could contribute up to 40% of diatom C biomass at these sites below the mixed layer, and are therefore potentially largely underestimated in the deep Bottle net casts, which do not allow sampling such small sized cells. These observations are congruent with previous reports of increased dominance of *Chaetoceros Hylaochaete* resting spores (CRS) in trap samples (Rembauville et al., 2016) and of dominance of CRS together with *E. antarctica* in sediments over the Kerguelen plateau (Armand et al., 2008b). New findings also show that viral infection of the bloom-forming genus *Chaetoceros* could induce mass formation of resting spores as a defense strategy (Pelusi et al., 2020), which is yet another mechanism that could lead to mass carbon export events by diatom spores.

4.4. Silicification degree and role in export

A schematic graph of diatom communities, Si:C stoichiometry and key processes both on and off-plateau is presented in Fig. 9. Clear differences over and outside the plateau for Si/C stoichiometrical ratios in particulate matter based on diatom carbon biomass were observed with the highest average Si:C_{diatoms} ratios from Niskin bottle at off-plateau stations (2.7 ± 0.7) and lowest on the plateau (0.6 ± 0.2) (Lafond et al., 2020).

Unfortunately we cannot produce a direct comparison from the Bottle net samples due to lacking POC data, but floristic fauna as well as microscopic observations confirmed the dominance of more heavily silicified species outside the Fe-fertilized plateau with a large dominance of the *Chaetoceros atlanticus* 'bulbosum' complex and *F. kerguelensis* at M4/M3 (Fig. 6). Frustule abundance was dominated by these two taxa, evidencing their elevated contribution to the downward Si flux. *F. Kerguelensis* is known to be the dominant species in sediment and main opal contributor of the summer POOZ (Permanently Open Ocean Zone), where it accounts for ~80% of the total diatom assemblage (Crosta et al., 2005) and its maximum occurrence is reported in waters 1-7°C (Jacques, 1983) which covers the temperature range measured during MOBYDICK (2-5°C). In our study region, the contribution of *F. kerguelensis* in the sediment varies between 30 and 75%, while *T. lentiginosa*, which was also abundant here, is the second most abundant taxon in Southern Ocean sediments with a reported range from 5-30% and has the same pattern of distribution in the sediments than *F. kerguelensis* (Crosta et al., 2005; Shukla et al., 2016). This is consistent with our findings showing that *F. Kerguelensis* is the dominant contributor to biomass in the deepest sample at M4 (Fig. 7), suggesting that it is the most efficient species at injecting C below the 2000 m horizon. On the plateau (M2), the more weakly silicified *Corethron inerme* was dominant in the surface layer, but the deep particle stock was also enriched with more heavily silicified species such as *E. antarctica* and *F. kerguelensis* even though in lesser numbers than west of Kerguelen.

It is notable that *F. kerguelensis* silicification degree, quantified with a silicification fluorescent probe (Lafond et al., 2020), was higher at M4/M3/M1 than at M2, reflecting the different ecological situations regarding Fe limitation. Hence, both taxonomic differences (mix of *Corethron* with *Eucampia* and *Fragilariopsis*) and lesser silicification degree of the dominant species can explain the lower Si/C_{diatom} ratios on the plateau.

Finally, *Eucampia* and *Fragilariopsis* were the dominant contributors to the Si flux while *Corethron inerme* was the main contributing species to C export, in particular upon the last visit at M2 (Fig. 7), again showing differences in the relative contribution of diatom species to Si and C export.

Figure 9: Conceptual schematic of the main results obtained above the Kerguelen Plateau at the naturally Fe-fertilized station (M2) and off-plateau in the HNLC station (M4). The blue curve on the left illustrates the vertical evolution of density with depth at station M2. Instruments used to sample the different water layers are displayed on the right handside. Only major taxa (>10% of total abundance) are represented here for clarity, and their relative contribution to abundance is indicated in black numbers.

5. Conclusions

The similarity of deep Bottle net samples with the overlying layer is not compatible with very slow sinking particles or suspended cells with no flux, and indicates rather fast export mechanisms to depth or intense mixing. The Bottle net does not allow to determine whether the collected material was part of the slow/fast sinking flux or a fraction of suspended particles, as can be done with a marine snow catcher for instance (Riley et al., 2012). A combination of such tools, together with gel traps (Lundsgaard, 1994) would allow to make progress on the characterization of the C attenuation curve in the meso- to bathypelagic ocean. *In situ* sinking velocities of aggregates and fecal pellets are still difficult to estimate due to the lack of data especially on their porosity and density, while size may not be the best predictor for this process (Iversen and Lampitt, 2020; Smith, 2014). Our results suggest a rapid export of diatom cells in the form of loose aggregates/phytodetritus, that reached intermediate and deep layers, but which were disaggregated prior to sampling, explaining the dominance of single cells in bottle net.

Our study shows that the concept of the Bottle net offers new investigative capabilities to the existing toolbox of instruments (McDonnell et al., 2015). The tools to be developed in this direction should prove a very useful complement to collect suspended and sinking particles in the ocean's interior. The Bottle net allowed to carry out a detailed study of diatom taxonomy, C biomass, physiological state of cells, trophic behaviour and life cycles on depth-integrated concentrated material which could not have been obtained from discrete deep Niskin samples, in which particle concentration is much too low for such analysis. In particular, it allowed to identify various mortality modes at the end of the final summer bloom. *Fragilariopsis kerguelensis*, which was previously considered as a very resistant species against grazers (Hamm et al., 2003; Assmy et al., 2013; Quéguiner, 2013), has been shown here to be actively grazed by mesozooplankton with a lot of crunched frustules, but also ingested by phaeodarian barely larger than the diatom cell length. High contents of crushed frustules debris observed in fecal pellets and minipellets also showed that this species is indeed palatable for various grazers. The Bottle net also allowed us to collect small-sized particles and to evidence that minipellets were the dominant fecal material in the water column at this stage of the season. Unfortunately, we could not estimate their contribution to the deep C stocks, even though previous studies have evidenced that it could be important.

Finally, the Bottle net could also help identify sporulation events linked to surface Si-limitation over the plateau and evidence that parasitic infection of diatoms could help inject both C and Si in the form of whole intact cells below the mixed layer, rather than be remineralized entirely in the surface layer. Much more emphasis needs to be placed on the end of bloom periods during field work, in order to be able to progress on identifying processes leading to the demise of the blooms and to understand how parasites and viruses may alter species succession and the fate of primary produced C. We also stress that live observations on board are absolutely necessary in identifying different trophic behaviours, such as infection by parasites that can be easily overlooked when examining fixed samples back at the ground-based laboratory.

Future sampling strategies in this region (and presumably on a global scale) will need to incorporate the biogeochemical and taxonomic characteristics of all components of vertical particle flux in a more integrated manner, allowing the intensity of the biological carbon pump to be more precisely related to environmental conditions. This strategic development proves to be crucial to provide elements for predicting the evolution of the biological carbon pump in these times of rapid climate change.

Acknowledgments

We thank B. Quéguiner, the PI of the MOBYDICK project, for providing us the opportunity to participate to this cruise, the chief scientist I. Obernosterer and the captain and crew of the R/V Marion Dufresne for their enthusiasm and support aboard during the MOBYDICK–THEMISTO cruise (doi:10.1BE00/1F00040G). This work was supported by the French oceanographic fleet ("Flotte océanographique française"), the French ANR ("Agence Nationale de la Recherche", AAPC 201B program, MOBYDICK Project number: ANR-1B-CE01-001G), and the French Research program of INSH-

1004 CNRS LEFE/CYBER (“Les enveloppes fluides et l’environnement” – “Cycles
1005 biogéochimiques, environnement et ressources”). We also thank Dr. D. Vaultot and I.
1006 Probert who gave us access to the Phenom-Pro at the Roscoff Biological Station. The
1007 authors wish to thank Frédéric Le Moigne for constructive discussions on this paper.

1008 **Declaration of interests**

1009 The authors declare that they have no known competing financial interests or personal
1010 relationships that could have appeared to influence the work reported in this paper.

1011

1012

1013

Supplementary Figures

1014 An overview of the diversity of all taxonomic groups found in Niskins, phytonet and Bottle nets can be
 1015 found here <https://plankton.mio.osupytheas.fr/mobydick-2018/>. An overview of the composition of fecal
 1016 pellets and aggregates in both light microscopy and SEM is available here
 1017 <https://plankton.mio.osupytheas.fr/mobydick-fecal-pellets/>

1018
 1019
 1020
 1021
 1022
 1023
 1024

Supl. Figure S1. Selection of images showing crunched siliceous diatom frustules (a to k) and Phaeodarian skeleton (l), indicating active mesozooplankton grazing. a. *Asteromphalus hookeri*, b-f. *Thalassiosira lentiginosa*, g-h. *Fragilariopsis kerguelensis*, j. *Proboscia* sp., k. *Membraneis* sp. l. skeleton of *Protocystis* sp. (Phaeodaria).

1025

1026

1027 **Suppl. Figure S2:** Images showing a parasitic infection of *Corethron inerme* colonies at station M2.
1028 Black arrows indicate infected cells.

1029

Suppl. Figure S3 : Morphological variations in two important species observed during the cruise. a to f. *Chaetoceros atlanticus bulbosum* complex. G to i: vegetative stage of *Odontella weissflogii*, rectangular resting cell/spore stage, round resting spore stage. J to l : *Eucampia antarctica* from vegetative colonial to small winter form.

Suppl. Figure S4: Different types of inert particles observed in the Bottle net samples. a) Minipellet (<50 μm) produced by *Protocystis* spp., b) aggregates <100 μm , c) aggregates 100-400 μm , d) spheroid fecal pellet, e) oval fecal pellet, f) cylindrical fecal pellet.

Suppl. Figure S5: *Odontella weissflogii* cells after 12 days incubation of resting cells in a light+Si+P treatment (a-b) and light only (c-e) treatment. Vegetative cells actively dividing (a-b), resting spore stage (c), half-vegetative half-spore like stage (d), abnormal valve due to the lack of Si (e). Scale bar: 25 μm.

References

- Agusti, S., Gonzalez-Gordillo, J.I., Vaque, D., Estrada, M., Cerezo, M.I., Salazar, G., Gasol, J.M., Duarte, C.M., 2015. Ubiquitous healthy diatoms in the deep sea confirm deep carbon injection by the biological pump. *Nat Commun* 6, 7608.
- Agustí, S., Krause, J.W., Marquez, I.A., Wassmann, P., Kristiansen, S., Duarte, C.M., 2020. Arctic (Svalbard islands) active and exported diatom stocks and cell health status. *Biogeosciences* 17, 35-45.
- Allredge, A.L., Gotschalk, C.C., 1990. The Relative Contribution of Marine Snow of Different Origins to Biological Processes in Coastal Waters. *Cont Shelf Res* 10, 41-58.
- Allredge, A.L., Silver, M.W., 1988. Characteristics, Dynamics and Significance of Marine Snow. *Prog Oceanogr* 20, 41-82.
- Armand, L.K., Cornet-Barthaux, V., Mosseri, J., Quéguiner, B., 2008a. Late summer diatom biomass and community structure on and around the naturally iron-fertilised Kerguelen Plateau in the Southern Ocean. *Deep-Sea Research Part II-Topical Studies in Oceanography* 55, 653-676.
- Armand, L.K., Crosta, X., Queguiner, B., Mosseri, J., Garcia, N., 2008b. Diatoms preserved in surface sediments of the northeastern Kerguelen Plateau. *Deep-Sea Research Part II-Topical Studies in Oceanography* 55, 677-692.
- Armand, L.K., Zielinski, U., 2001. Diatom species of the genus *Rhizosolenia* from Southern Ocean sediments: Distribution and taxonomic notes. *Diatom Res* 16, 259-294.
- Assmy, P., Henjes, J., Klaas, C., Smetacek, V., 2007. Mechanisms determining species dominance in a phytoplankton bloom induced by the iron fertilization experiment EisenEx in the Southern Ocean. *Deep-Sea Res Pt I* 54, 340-362.
- Assmy, P., Smetacek, V., Montresor, M., Klaas, C., Henjes, J., Strass, V.H., Arrieta, J.M., Bathmann, U., Berg, G.M., Breitbarth, E., Cisewski, B., Friedrichs, L., Fuchs, N., Herndl, G.J., Jansen, S., Kragefsky, S., Latasa, M., Peeken, I., Rottgers, R., Scharek, R., Schuller, S.E., Steigenberger, S., Webb, A., Wolf-Gladrow, D., 2013. Thick-shelled, grazer-protected diatoms decouple ocean carbon and silicon cycles in the iron-limited Antarctic Circumpolar Current. *Proceedings of the National Academy of Sciences of the United States of America* 110, 20633-20638.
- Biard, T., Krause, J.W., Stukel, M.R., Ohman, M.D., 2018. The Significance of giant Phaeodarians (Rhizaria) to Biogenic Silica Export in the California Current Ecosystem. *Global Biogeochemical Cycles*.
- Blain S., Quéguiner B., Trull T.W., 2008. The natural iron fertilization experiment KEOPS (KErguelen Ocean and Plateau compared Study): An overview. *Deep Sea Research Part II: Topical Studies in Oceanography*, 55 (5-7), 559-565.
- Blain, S., Rembauville, M., Crispi, O., Obernosterer, I., 2020. Synchronized autonomous sampling reveals coupled pulses of biomass and export of morphologically different diatoms in the Southern Ocean. *Limnology and Oceanography*.
- Close, H.G., Shah, S.R., Ingalls, A.E., Diefendorf, A.F., Brodie, E.L., Hansman, R.L., Freeman, K.H., Aluwihare, L.I., Pearson, A., 2013. Export of submicron particulate

1089 organic matter to mesopelagic depth in an oligotrophic gyre. Proceedings of the
1090 National Academy of Sciences 110, 12565-12570.

1091 Cornet-Barthaux, V., Armand, L., Queguiner, B., 2007. Biovolume and biomass
1092 estimates of key diatoms in the Southern Ocean. Aquat Microb Ecol 48, 295-308.

1093 Crosta, X., Romero, O., Armand, L.K., Pichon, J.J., 2005. The biogeography of major
1094 diatom taxa in Southern Ocean sediments: 2. Open ocean related species.
1095 Palaeogeogr Palaeocl 223, 66-92.

1096 Dall'Olmo, G., Mork, K.A., 2014. Carbon export by small particles in the Norwegian
1097 Sea. Geophys Res Lett 41, 2921-2927.

1098 DiTullio, G., Grebmeier, J., Arrigo, K., Lizotte, M., Robinson, D., Leventer, A., Barry,
1099 J., VanWoert, M., Dunbar, R., 2000. Rapid and early export of *Phaeocystis antarctica*
1100 blooms in the Ross Sea, Antarctica. Nature 404, 595-598.

1101 Durkin, C.A., Estapa, M.L., Buesseler, K.O., 2015. Observations of carbon export by
1102 small sinking particles in the upper mesopelagic. Mar Chem 175, 72-81.

1103 Durkin, C.A., Koester, J.A., Bender, S.J., Armbrust, E.V., 2016. The evolution of silicon
1104 transporters in diatoms. J Phycol 52, 716-731.

1105 Eppley, R.W., Reid, F.M.H., Strickland, J.D.H., 1970. The ecology of the plankton off
1106 La Jolla, California, in the period April through September, 1967. III. Estimates of
1107 phytoplankton crop, size, growth rate, and primary production. Bull. Scripps Inst.
1108 Oceanogr. 17, 33-42.

1109 Fryxell, G.A., Prasad, A.K.S.K., 1990. *Eucampia antarctica* var. *recta* (Mangin) stat.
1110 nov. (Biddulphiaceae, Bacillariophyceae): life stages at the Weddell Sea ice edge.
1111 Phycologia 29, 27-38.

1112 Gardner, W.D., Chung, S.P., Richardson, M.J., Walsh, I.D., 1995. The oceanic mixed-
1113 layer pump. Deep Sea Research Part II: Topical Studies in Oceanography 42, 757-
1114 775.

1115 Giering, S.L., Sanders, R., Lampitt, R.S., Anderson, T.R., Tamburini, C., Boutrif, M.,
1116 Zubkov, M.V., Marsay, C.M., Henson, S.A., Saw, K., Cook, K., Mayor, D.J., 2014.
1117 Reconciliation of the carbon budget in the ocean's twilight zone. Nature 507, 480-483.

1118 Gonzalez, H.E., 1992. Distribution and abundance of minipellets around the Antarctic
1119 peninsula. Implications for protistan feeding behaviour. Mar Ecol Prog Ser 90, 223-
1120 223.

1121 Gowing, M.M., 1986. Trophic biology of phaeodarian radiolarians and flux of living
1122 radiolarians in the upper 2000 m of the North Pacific central gyre. Deep Sea Research
1123 Part A. Oceanographic Research Papers 33, 655-674.

1124 Gowing, M.M., 1989. Abundance and feeding ecology of Antarctic phaeodarian
1125 radiolarians. Mar Biol 103, 107-118.

1126 Gowing, M.M., Silver, M.W., 1985. Minipellets: a new and abundant size class of
1127 marine fecal pellets. J Mar Res 43, 395-418.

1128 Greenspan, P., 1985. Nile red: a selective fluorescent stain for intracellular lipid
1129 droplets. The Journal of Cell Biology 100, 965-973.

1130 Gutiérrez, M.H., Jara, A.M., Pantoja, S., 2016. Fungal parasites infect marine diatoms
 1131 in the upwelling ecosystem of the Humboldt current system off central Chile.
 1132 *Environmental Microbiology* 18, 1646-1653.

1133 Gutierrez-Rodriguez, A., Stukel, M.R., dos Santos, A.L., Biard, T., Scharek, R., Vaultot,
 1134 D., Landry, M.R., Not, F., 2019. High contribution of Rhizaria (Radiolaria) to vertical
 1135 export in the California Current Ecosystem revealed by DNA metabarcoding. *The ISME*
 1136 *journal* 13, 964-976.

1137 Hamm, C.E., Merkel, R., Springer, O., Jurkojc, P., Maier, C., Prechtel, K., Smetacek,
 1138 V., 2003. Architecture and material properties of diatom shells provide effective
 1139 mechanical protection. *Nature* 421, 841-843.

1140 Henson, S.A., Sanders, R., Madsen, E., Morris, P.J., Le Moigne, F., Quartly, G.D.,
 1141 2011. A reduced estimate of the strength of the ocean's biological carbon pump.
 1142 *Geophys Res Lett* 38.

1143 Ikenoue, T., Kimoto, K., Okazaki, Y., Sato, M., Honda, M.C., Takahashi, K., Harada,
 1144 N., Fujiki, T., 2019. Phaeodaria: An Important Carrier of Particulate Organic Carbon in
 1145 the Mesopelagic Twilight Zone of the North Pacific Ocean. *Global Biogeochemical*
 1146 *Cycles* 33, 1146-1160.

1147 Iversen, M.H., Lampitt, R.S., 2020. Size does not matter after all: No evidence for a
 1148 size-sinking relationship for marine snow. *Prog Oceanogr* 189.

1149 Jacques G., 1983. Some ecophysiological aspects of the Antarctic phytoplankton. *Polar*
 1150 *Biology*, 2, 27-33.

1151 Kemp, A.E.S., Pike, J., Pearce, R.B., Lange, C.B., 2000. The "Fall dump"- a new
 1152 perspective on the role of a "shade flora" in the annual cycle of diatom production and
 1153 export flux. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2129-
 1154 2154.

1155 Klaas C., Archer D.E., 2002. Association of sinking organic matter with various types of
 1156 mineral ballast in the deep sea: Implications for the rain ratio. *Global Biogeochemical*
 1157 *Cycles*, 16 (4), 63-61 - 63-14.

1158 Kranzler, C.F., Krause, J.W., Brzezinski, M.A., Edwards, B.R., Biggs, W.P.,
 1159 Maniscalco, M., McCrow, J.P., Van Mooy, B.A., Bidle, K.D., Allen, A.E., 2019. Silicon
 1160 limitation facilitates virus infection and mortality of marine diatoms. *Nature*
 1161 *microbiology*, 1-8.

1162 Laber, C.P., Hunter, J.E., Carvalho, F., Collins, J.R., Hunter, E.J., Schieler, B.M., Boss,
 1163 E., More, K., Frada, M., Thamatrakoln, K., Brown, C.M., Haramaty, L., Ossolinski, J.,
 1164 Fredricks, H., Nissimov, J.I., Vandzura, R., Sheyn, U., Lehahn, Y., Chant, R.J., Martins,
 1165 A.M., Coolen, M.J.L., Vardi, A., DiTullio, G.R., Van Mooy, B.A.S., Bidle, K.D., 2018.
 1166 Coccolithovirus facilitation of carbon export in the North Atlantic. *Nat Microbiol* 3, 537-
 1167 547.

1168 Lafond, A., Leblanc, K., Legras, J., Cornet, V., Quéguiner, B., 2020. The structure of
 1169 diatom communities constrains biogeochemical properties in surface waters of the
 1170 Southern Ocean (Kerguelen Plateau). *J Marine Syst*, 103458.

1171 Lam, P.J., Marchal, O., 2015. Insights into particle cycling from thorium and particle
 1172 data. *Annual review of marine science* 7, 159-184.

1173 Lampitt, R.S., Salter, I., Johns, D., 2009. Radiolaria: Major exporters of organic carbon
1174 to the deep ocean. *Global Biogeochemical Cycles* 23, GB1010.

1175 Lasbleiz, M., Leblanc, K., Armand, L.K., Christaki, U., Georges, C., Obernosterer, I.,
1176 Quéguiner, B., King, G., 2016. Composition of diatom communities and their
1177 contribution to plankton biomass in the naturally iron-fertilized region of Kerguelen in
1178 the Southern Ocean. *FEMS microbiology ecology* 92, 653--676.

1179 Lasbleiz, M., Leblanc, K., Blain, S., Ras, J., Cornet-Barthaux, V., Hélias Nunige, S.,
1180 Quéguiner, B., 2014. Pigments, elemental composition (C, N, P, and Si), and
1181 stoichiometry of particulate matter in the naturally iron fertilized region of Kerguelen in
1182 the Southern Ocean. *Biogeosciences* 11, 5931-5955.

1183 Laurenceau-Cornec, E.C., Trull, T.W., Davies, D.M., Bray, S.G., Doran, J., Planchon,
1184 F., Carlotti, F., Jouandet, M.P., Cavagna, A.J., Waite, A.M., Blain, S., 2015. The
1185 relative importance of phytoplankton aggregates and zooplankton fecal pellets to
1186 carbon export: insights from free-drifting sediment trap deployments in naturally iron-
1187 fertilised waters near the Kerguelen Plateau. *Biogeosciences* 12, 1007-1027.

1188 Laurenceau-Cornec, E.C., Le Moigne, F.A., Gallinari, M., Moriceau, B., Toullec, J.,
1189 Iversen, M.H., Engel, A., De La Rocha, C.L., 2020. New guidelines for the application
1190 of Stokes' models to the sinking velocity of marine aggregates. *Limnology and*
1191 *Oceanography* 65, 1264-1285.

1192 Laws, E.A., Landry, M.R., Barber, R.T., Campbell, L., Dickson, M.-L., Marra, J., 2000.
1193 Carbon cycling in primary production bottle incubations: inferences from grazing
1194 experiments and photosynthetic studies using and in the Arabian Sea. *Deep Sea*
1195 *Research Part II: Topical Studies in Oceanography* 47, 1339-1352.

1196 Le Moigne, F.A.C., 2019. Pathways of Organic Carbon Downward Transport by the
1197 Oceanic Biological Carbon Pump. *Frontiers in Marine Science* 6.

1198 Leblanc, K., Queguiner, B., Diaz, F., Cornet, V., Michel-Rodriguez, M., Durrieu de
1199 Madron, X., Bowler, C., Malviya, S., Thyssen, M., Gregori, G., Rembauville, M.,
1200 Grosso, O., Poulain, J., de Vargas, C., Pujo-Pay, M., Conan, P., 2018. Nanoplanktonic
1201 diatoms are globally overlooked but play a role in spring blooms and carbon export.
1202 *Nat Commun* 9, 953.

1203 Lundsgaard, C., 1994. Use of high viscosity medium in studies of aggregates,
1204 Sediment Trap Studies in the Nordic Countries. *Symposium Proc., Mar. Biol. Lab.,*
1205 *Helsingør, Denmark*, pp. 141-152.

1206 McDonnell, A.M., Lam, P.J., Lamborg, C.H., Buesseler, K.O., Sanders, R., Riley, J.S.,
1207 Marsay, C., Smith, H.E., Sargent, E.C., Lampitt, R.S., 2015. The oceanographic
1208 toolbox for the collection of sinking and suspended marine particles. *Prog Oceanogr*
1209 133, 17-31.

1210 McQuoid, M.R., Hobson, L.A., 1996. Diatom Resting Stages. *J Phycol* 32, 889-902.

1211 Mosseri, J., Quéguiner, B., Armand, L., Cornet-Barthaux, V., 2008. Impact of iron on
1212 silicon utilization by diatoms in the Southern Ocean: A case study of Si/N cycle
1213 decoupling in a naturally iron-enriched area. *Deep Sea Research Part II: Topical*
1214 *Studies in Oceanography* 55, 801-819.

1215 Nakamura, Y., Suzuki, N., 2015. Phaeodaria: diverse marine cercozoans of world-wide
1216 distribution, *Marine Protists*. Springer, pp. 223-249.

1217 Omand, M.M., D'Asaro, E.A., Lee, C.M., Perry, M.J., Briggs, N., Cetinić, I.,
1218 Mahadevan, A., 2015. Eddy-driven subduction exports particulate organic carbon from
1219 the spring bloom. *Science* 348, 222-225.

1220 Park, Y.H., Durand, I., Kestenare, E., Rougier, G., Zhou, M., d'Ovidio, F., Cotté, C.,
1221 Lee, J.H., 2014. Polar Front around the Kerguelen Islands: An up-to-date
1222 determination and associated circulation of surface/subsurface waters. *Journal of*
1223 *Geophysical Research: Oceans* 119, 6575-6592.

1224 Pauthenet, E., Roquet, F., Madec, G., Guinet, C., Hindell, M., McMahon, C.R.,
1225 Harcourt, R., Nerini, D., 2018. Seasonal Meandering of the Polar Front Upstream of
1226 the Kerguelen Plateau. *Geophys Res Lett* 45, 9774-9781.

1227 Pelusi, A., De Luca, P., Manfellotto, F., Thamatrakoln, K., Bidle, K.D., Montresor, M.,
1228 2020. Virus-induced spore formation as a defense mechanism in marine diatoms. *New*
1229 *Phytol.*

1230 Priddle, J., Fryxell, G., 1985. Handbook of the common plankton diatoms of the
1231 Southern Ocean. British Antarctic Survey.

1232 Puigcorbé V., Masqué P., Le Moigne F.A.C., 2020. Global database of ratios of
1233 particulate organic carbon to thorium-234 in the ocean: improving estimates of the
1234 biological carbon pump. *Earth System Science Data*, 12 (2), 1267-1285.

1235 Quéguiner, B., 2013. Iron fertilization and the structure of planktonic communities in
1236 high nutrient regions of the Southern Ocean. *Deep Sea Research Part II: Topical*
1237 *Studies in Oceanography* 90, 43-54.

1238 Ragueneau, O., Savoye, N., Del Amo, Y., Cotten, J., Tardiveau, B., Leynaert, A., 2005.
1239 A new method for the measurement of biogenic silica in suspended matter of coastal
1240 waters: using Si:Al ratios to correct for the mineral interference. *Cont Shelf Res* 25,
1241 697-710.

1242 Rembauville, M., Blain, S., Armand, L., Quéguiner, B., Salter, I., 2015. Export fluxes in
1243 a naturally iron-fertilized area of the Southern Ocean – Part 2: Importance of diatom
1244 resting spores and faecal pellets for export. *Biogeosciences* 12, 3171-3195.

1245 Rembauville, M., Manno, C., Tarling, G.A., Blain, S., Salter, I., 2016. Strong
1246 contribution of diatom resting spores to deep-sea carbon transfer in naturally iron-
1247 fertilized waters downstream of South Georgia. *Deep Sea Research Part I:*
1248 *Oceanographic Research Papers* 115, 22-35.

1249 Riaux-Gobin, C., Hargraves, P.E., Neveux, J., Oriol, L., Vétion, G., 1997. Microphyte
1250 pigments and resting spores at the water-sediment interface in the Subantarctic deep
1251 sea (Indian sector of the Southern Ocean). *Deep Sea Research Part II: Topical Studies*
1252 *in Oceanography* 44, 1033-1051.

1253 Riaux-Gobin, C., Fontugne, M., Jensen, K.G., Bentaleb, I., Cauwet, G., Chrétiennot-
1254 Dinot, M.J., Poisson, A., 2006. Surficial deep-sea sediments across the polar frontal
1255 system (Southern Ocean, Indian sector): Particulate carbon content and microphyte
1256 signatures. *Mar Geol* 230, 147-159.

1257 Richardson, T.L., Jackson, G.A., 2007. Small phytoplankton and carbon export from
1258 the surface ocean. *Science* 315, 838-840.

1259 Riemann, F., 1989. Gelatinous phytoplankton detritus aggregates on the Atlantic deep-
1260 sea bed. *Mar Biol* 100, 533-539.

1261 Rigual-Hernández, A.S., Trull, T.W., Bray, S.G., Armand, L.K., 2016. The fate of
1262 diatom valves in the Subantarctic and Polar Frontal Zones of the Southern Ocean:
1263 Sediment trap versus surface sediment assemblages. *Palaeogeography,*
1264 *Palaeoclimatology, Palaeoecology* 457, 129-143.

1265 Rigual-Hernández, A.S., Trull, T.W., Bray, S.G., Closset, I., Armand, L.K., 2015.
1266 Seasonal dynamics in diatom and particulate export fluxes to the deep sea in the
1267 Australian sector of the southern Antarctic Zone. *J Marine Syst* 142, 62-74.

1268 Riley, J., Sanders, R., Marsay, C., Le Moigne, F.A., Achterberg, E.P., Poulton, A.J.,
1269 2012. The relative contribution of fast and slow sinking particles to ocean carbon
1270 export. *Global Biogeochemical Cycles* 26.

1271 Romero, O.E., Fischer, G., 2017. Shift in the species composition of the diatom
1272 community in the eutrophic Mauritanian coastal upwelling: Results from a multi-year
1273 sediment trap experiment (2003–2010). *Prog Oceanogr* 159, 31-44.

1274 Salter, I., Kemp, A.E.S., Moore, C.M., Lampitt, R.S., Wolff, G.A., Holtvoeth, J., 2012.
1275 Diatom resting spore ecology drives enhanced carbon export from a naturally iron-
1276 fertilized bloom in the Southern Ocean. *Global Biogeochemical Cycles* 26, n/a-n/a.

1277 Salter, I., Lampitt, R.S., Sanders, R., Poulton, A., Kemp, A.E.S., Boorman, B., Saw, K.,
1278 Pearce, R., 2007. Estimating carbon, silica and diatom export from a naturally fertilised
1279 phytoplankton bloom in the Southern Ocean using PELAGRA: A novel drifting
1280 sediment trap. *Deep Sea Research Part II: Topical Studies in Oceanography* 54, 2233-
1281 2259.

1282 Sassenhagen, I., Irion, S., Jardillier, L., Moreira, D., Christaki, U., 2020. Protist
1283 Interactions and Community Structure During Early Autumn in the Kerguelen Region
1284 (Southern Ocean). *Protist* 171, 125709.

1285 Scholz, B., Guillou, L., Marano, A.V., Neuhauser, S., Sullivan, B.K., Karsten, U.,
1286 Kupper, F.C., Gleason, F.H., 2016. Zoospore parasites infecting marine diatoms - A
1287 black box that needs to be opened. *Fungal Ecol* 19, 59-76.

1288 Scott, F.J., Marchant, H.J., 2005. Antarctic marine protists. Australian Biological
1289 Resources Study Canberra.

1290 Shukla, S.K., Cressin, J., Crosta, X., 2016. *Thalassiosira lentiginosa* size variation and
1291 associated biogenic silica burial in the Southern Ocean over the last 42 kys. *Marine*
1292 *Micropaleontology* 127, 74-85.

1293 Smayda, T.J., 1978. From phytoplankton to biomass. 6. Phytoplankton manual.

1294 Smith, H.E., 2014. The contribution of mineralising phytoplankton to the biological
1295 carbon pump in high latitudes. University of Southampton.

1296 Stemmann, L., Jackson, G.A., Gorsky, G., 2004. A vertical model of particle size
1297 distributions and fluxes in the midwater column that includes biological and physical
1298 processes—Part II: application to a three year survey in the NW Mediterranean Sea.
1299 *Deep Sea Research Part I: Oceanographic Research Papers* 51, 885-908.

1300 Stukel, M.R., Biard, T., Krause, J., Ohman, M.D., 2018. Large Phaeodaria in the
1301 twilight zone: Their role in the carbon cycle. *Limnology and Oceanography* 63, 2579-
1302 2594.

1303 Turner, J.T., 2002. Zooplankton fecal pellets, marine snow and sinking phytoplankton
1304 blooms. *Aquat Microb Ecol* 27, 57-102.

1305 Veldhuis, M., Kraay, G., Timmermans, K., 2001. Cell death in phytoplankton:
1306 correlation between changes in membrane permeability, photosynthetic activity,
1307 pigmentation and growth. *Eur J Phycol* 36, 167-177.

1308 Waite, A.M., Safi, K.A., Hall, J.A., Nodder, S.D., 2000. Mass sedimentation of
1309 picoplankton embedded in organic aggregates. *Limnology and Oceanography* 45, 87-
1310 97.

1311

1312