IMMUNO-METABOLIC SIGNATURES OF DENDRITIC CELLS ASSOCIATE WITH T-CELL RESPONSES IN MELANOMA PATIENTS

Juraj Adamik, Deena Maurer, Paul Munson, Alexis Combes, Philippe Pierre,
Matthew Krummel, Rafael J Argüello, Lisa Butterfield

► To cite this version:

Juraj Adamik, Deena Maurer, Paul Munson, Alexis Combes, Philippe Pierre, et al.. IMMUNO-METABOLIC SIGNATURES OF DENDRITIC CELLS ASSOCIATE WITH T-CELL RESPONSES IN MELANOMA PATIENTS. SICT, Nov 2020, Los Angeles, United States. pp.A325 - A325, 10.1136/jitc-2020-sitc2020.0299 . hal-03451694

HAL Id: hal-03451694

<https://hal.science/hal-03451694>

Submitted on 26 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Trial Registration Clinicaltrials. gov NCT01343043

Ethics Approval This study was approved by the appropriate institutional review boards and independent ethics committees.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0298>

299

IMMUNO-METABOLIC SIGNATURES OF DENDRITIC CELLS ASSOCIATE WITH T-CELL RESPONSES IN MELANOMA PATIENTS

¹Juraj Adamik*, ²Deena Maurer, ¹Paul Munson, ³Alexis Combes, ⁴Philippe Pierre, ³Matthew Krummel, ⁴Rafael Argüello, ¹Lisa Butterfield. ¹Parker Institute for Cancer Immunotherapy, University of California, San Francisco, San Francisco, CA, USA; ²University of Pittsburgh, Pittsburgh, PA, USA; ³University of California, San Francisco, San Francisco, CA, USA; ⁴Centre d'Immunologie de Marseille-Luminy, Marseille, France

Background The therapeutic efficacy of Dendritic cells (DC) vaccines remains low and there is an unmet need for more effective vaccine design to achieve durable clinical outcomes. Our study analyzed the transcriptomic and energetic metabolism profile of an adenoviral-based DC vaccine targeted against three commonly shared melanoma antigens: Tyrosinase, MART-1 and MAGE-A6 from 35 subjects enrolled in a Phase I study of autologous DC vaccines in late-stage melanoma.¹ To further investigate the immuno-metabolic features of monocyte-derived DC vaccines, we are employing a novel flow cytometry-based method, called SCENITH² to integrate functional metabolic states with multiparametric DC immune phenotypes.

Methods iDC were generated from HD and patient monocytes using GM-CSF+IL-4 for 5d. DC were matured (mDC) using IFN γ +LPS for additional 24 hrs. Tolerogenic DC (Tol DC) were generated using vitamin-D3 and dexamethasone. Seahorse[®] was used to measure DC metabolic profile. Cytek/Aurora spectral flow cytometry was used for multiparametric phenotypic and metabolic analysis by SCENITHTM.

Results Melanoma patient mDC used for autologous vaccine generation showed significantly altered metabolic gene signatures associated with enhanced oxidative phosphorylation (OXPHOS) and lipid metabolism pathways as compared to HD mDC. Furthermore, increased enrichment for mitochondrial respiration genes involved in the TCA cycle, electron transport chain and fatty acid oxidation (FAO) correlated with inferior tumor antigen-specific T cell responses and clinical outcome in patients. Seahorse analyses confirmed that HD and good outcome patient DC demonstrated the highest maturation-induced reduction in maximal oxygen consumption rate/OXPHOS and exogenous FAO. Interestingly, while the glycolytic rate of non-responding patient DC was the lowest, overall, we observe only a moderate increase in glycolytic capacity during DC maturation. SCENITH analysis showed that unlike monocytes, which are primarily glycolytic, differentiated mono-derived iDC and mDC utilize both glycolysis and mitochondrial respiration. Interestingly, under tolerogenic (Tol) differentiation conditions Tol iDC shift from glucose dependence into FAO and/or glutaminolysis while Tol mDC strongly depend on OXPHOS. Consistent with dependence on mitochondrial respiration, Tol mDC exhibit reduced HIF1 α levels together with enhanced p-AMPK:p-mTOR ratio. Additionally, we show that the altered metabolism of Tol mDC is linked to retention of CD14-monocyte antigen with reduced DC markers HLA-DR, CD86, CD206, CD11c, CD33, with increased PD-L1 and ILT3 expression. Furthermore, we show that unlike HD mDC, tolerogenic and melanoma patient-

derived mDC populations exhibit similar metabolic and immune characteristics.

Conclusions We demonstrate that metabolic profile of DCs is tightly associated to the immuno-stimulatory potential of DC vaccines from cancer patients. Using SCENITH, we linked phenotypic and functional metabolic changes associated to immune signatures that correspond to healthy and immuno-suppressed DC differentiation.

Ethics Approval The clinical trial reported was fully approved by the Univ. Pittsburgh PRC and IRB (PRO12010416, #09-021) and had FDA IND #15044 and NCT01622933.

REFERENCES

1. Butterfield LH, Vujanovic L, Santos PM, Maurer DM, Gambotto A, Lohr J, Li C, Waldman J, Chandran U, Lin Y, et al. Multiple antigen-engineered DC vaccines with or without IFN α to promote antitumor immunity in melanoma. *Journal for immunotherapy of cancer* 2019;7:113.
2. Argüello RJ, Combes AJ, Char R, Bousiquot E, Gigan JP, Camosseto V, Samad B, Tsui J, Yan P, et al. SCENITH: A flow cytometry based method for functional profiling energy metabolism with single cell resolution. *BioRxiv* 2020.03.10.985796; doi: <https://doi.org/10.1101/2020.03.10.985796>. *Cell Metabolism*. (Sneak and Peek: under review 10.2139/ssrn.3565001)

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0299>

300

FINAL ANALYSIS OF A PROSPECTIVE, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED PHASE IIB TRIAL OF TUMOR LYSATE, PARTICLE-LOADED, DENDRITIC CELL VACCINE IN STAGE III/IV MELANOMA: 36-MONTH ANALYSIS

¹Lexy Adams*, ¹Robert Chick, ¹Guy Clifton, ¹Timothy Vreeland, ¹Patrick McCarthy, ¹Anne O'Shea, ¹Phil Kemp Bohan, ¹Annelies Hickerson, ¹John Myers, ¹Jessica Cindass, ¹Diane Hale, ²Mark Faries, ³John Hynstrom, ⁴Adam Berger, ⁵James Jakub, ⁶Jeffrey Sussman, ⁷Montaser Shaheen, ⁸Thomas Wagner, ⁹George Peoples. ¹Brooke Army Medical Center, San Antonio, TX, USA; ²John Wayne Cancer Institute, Los Angeles, CA, USA; ³University of Utah, Salt Lake City, UT, USA; ⁴Rutgers Cancer Institute of New Jersey, Philadelphia, PA, USA; ⁵Mayo Clinic, Rochester, MN, USA; ⁶University of Cincinnati, Cincinnati, OH, USA; ⁷University of Arizona, Phoenix, AZ, USA; ⁸Orbis Health Solutions, Greenville, SC, USA; ⁹Cancer Vaccine Development Program, San Antonio, TX, USA

Background The tumor lysate, particle-loaded, dendritic cell (TLPLDC) vaccine is created ex vivo by loading autologous dendritic cells (DC) with yeast cell wall particles (YCWP) containing autologous tumor lysate, thus delivering tumor antigens to the DC cytoplasm via phagocytosis. TLPLDC then activates a robust T cell response against the unique antigens for each patient. The primary analysis of the prospective, randomized, multi-center, double-blind, placebo-controlled phase IIB trial in patients with resected stage III/IV melanoma showed TLPLDC improved 24-month disease-free survival (DFS) in the per-treatment (PT) analysis (patients completing the 6-month primary vaccine series). Here, we examine the secondary endpoint of 36-month DFS and overall survival (OS).

Methods Patients with resected stage III/IV melanoma were randomized 2:1 to TLPLDC vaccine or placebo (autologous DC loaded with empty YCWP). Treatments were given at 0, 1, 2, 6, 12 and 18 months. The protocol was amended to include patients receiving concurrent checkpoint inhibitors (CPIs) to follow changes in standard of care. The co-primary endpoints were 24-month DFS by intention-to-treat (IT) analysis and per-treatment (PT) analysis, with secondary endpoints including 36-month DFS and OS by ITT and PT analysis, pre-specified analysis by stage, and safety as measured by CTCAE v4.03.