Imprint of seasonality changes on fluvio-glacial dynamics across Heinrich Stadial 1 (NE Atlantic Ocean)

Wiem Fersi, Aurélie Penaud, Mélanie Wary, Samuel Toucanne, Claire Waelbroeck, Linda Rossignol, Frédérique Eynaud

► To cite this version:

Wiem Fersi, Aurélie Penaud, Mélanie Wary, Samuel Toucanne, Claire Waelbroeck, et al.. Imprint of seasonality changes on fluvio-glacial dynamics across Heinrich Stadial 1 (NE Atlantic Ocean). *Global and Planetary Change*, 2021, 204, pp.103552. 10.1016/j.gloplacha.2021.103552 . hal-03455758v1

HAL Id: hal-03455758

<https://hal.science/hal-03455758v1>

Submitted on 30 Nov 2021 (v1), last revised 15 Dec 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Imprint of seasonality changes on fluvio-glacial dynamics across Heinrich Stadial 1 (NE Atlantic Ocean)

**Wiem Fersi ^{(a)*}, Aurélie Penaud ^(a), Mélanie Wary ^(b), Samuel Toucanne ^(c), Claire
Waelbroeck ^(d), Linda Rossignol ^(e), Frédérique Eynaud ^(e)**

*(a) Univ Brest (UBO), CNRS, UMR 6538 Laboratoire Géosciences Océan (LGO), F-
29280 Plouzané, France*

*(b) Institut de Ciència i Tecnologia Ambientals (ICTA-UAB), Universitat Autònoma de
Barcelona, Bellaterra, Catalonia, Spain*

*(c) Ifremer, Laboratoire Géophysique et environnements Sédimentaires. F-29280
Plouzané, France*

(d) LOCEAN/IPSL, Sorbonne Université-CNRS-IRD-MNHN, UMR7159, Paris, France

*(e) Univ Bordeaux, CNRS, UMR 5805 Environnements et Paléoenvironnements
Océaniques et Continentaux (EPOC), F-33405 Talence, France*

*Corresponding author. Tel.: +33-298-498-741; fax: +33-298-498-760.

E-mail addresses: wiem.fersi@univ-brest.fr, aurelie.penaud@univ-brest.fr

Abstract

The northern Bay of Biscay has previously proven its great potential for recording the ‘*Fleuve Manche*’ paleoriver (i.e., the largest Pleistocene river in Europe) fluvio-glacial activity. In this study, new dinoflagellate cyst (dinocyst) analyses have been carried out at sub-centennial resolution in core MD13-3438 to reconstruct the deglacial history of the ‘*Fleuve Manche*’ paleoriver runoff coupled with European Ice Sheets (EIS) fluctuations across Heinrich Stadial 1 (HS1: 18.2–14.6 ka BP), a key extreme climatic event of the last glacial period. Prior to Heinrich Event (HE) 1 (16.7–14.6 ka BP), the onset of HS1 (18.2–16.7 ka BP) appears here marked by enhanced ‘*Fleuve Manche*’ paleoriver runoff, materialized by laminated deposits. Our work suggests a novel sub-centennial scale subdivision of the early HS1 (laminated) interval into 5 sub-phases when episodes of substantial fluvio-glacial delivery concomitant with warm summers alternate with episodes of moderate runoff associated with extended cold winters. We argue that multidecadal seasonal changes played a key role in the hydrological regime of western Europe during this HS1 interval, with the retreat of the southern limb of the EIS, and associated influx of meltwater and fluvio-glacial delivery, which were strongly influenced by those multidecadal changes in seasonality. Interestingly, our paleoclimatic record not only evidences the crucial role of seasonality in controlling climate and hydrological variations during HS1 but also shows a remarkable echo with reconstructions from the western Mediterranean Basin, highlighting common climate forcings at regional scale during the last deglaciation.

Keywords: *Heinrich Stadial 1; Dinoflagellate cysts; northern Bay of Biscay; ‘Fleuve Manche’ paleoriver; European Ice Sheets*

1. Introduction

The last glacial period was accompanied by millennial-scale abrupt climate shifts, portrayed in Greenland ice-cores as rapid transitions from cold atmospheric phases termed Greenland Stadials (GS) to warm atmospheric phases referred to as Greenland Interstadials (GI; e.g., Dansgaard et al., 1993; Rasmussen et al., 2014). Despite their original designation, these climate excursions had an impact across the globe (e.g., Voelker, 2002). In the North Atlantic Ocean, some GS were associated with massive iceberg surges mainly from the Laurentide Ice Sheet (LIS) *via* the Hudson Strait Ice Stream (e.g., Bond et al., 1992, 1993; Broecker et al., 1992, 1994; Hemming, 2004), identified in marine sediments as Ice Rafted Debris (IRD)-enriched layers (e.g., Bond et al., 1993; Broecker, 1994; Heinrich, 1988). These massive iceberg (and thus freshwater) surge events are known as Heinrich Events (HEs including HE1), with their corresponding stadial phases called Heinrich Stadials (HSs including HS1; Barker et al., 2009; Sanchez Goñi and Harrison, 2010). The associated huge freshwater releases resulted in large reductions of the Atlantic Meridional Overturning Circulation (AMOC; e.g., McManus et al., 2004; Stanford et al., 2006, 2011; Ng et al., 2018; Toucanne et al., 2021). Numerous studies demonstrated that Greenland Iceland and European Ice Sheets were also major contributors to the oceanic disturbances in the North Atlantic Ocean, especially when considering the surge sequencing along time (e.g., Bond et al., 1997, 1999; Grousset et al., 2000, 2001; Hemming et al., 2000, 2004; Knutz et al., 2001, 2007; Hall et al., 2006; Peck et al., 2006; Nygård et al., 2007; Toucanne et al., 2008, 2010, 2015). HS1, including the HE1 layer, occurred at the onset of the last deglaciation (~19-11 ka BP; Clark et al., 2012a), just before the abrupt Bølling-Allerød (B/A) warming event starting at ca. 14.7 ka BP (Rasmussen et al., 2014) and after the Last Glacial Maximum (LGM; Mix et al., 2001). Over the LGM, which was characterized by a large European Ice Sheet (EIS) including the British-Irish (BIIS) and Scandinavian (SIS) Ice Sheets, the '*Fleuve Manche*' paleoriver (Channel River) was one of the largest river systems that drained western Europe (e.g., Gibbard, 1988; Toucanne et al., 2009, 2010, 2015). This huge fluvial system included the French, Belgian and British rivers, and the merged German, Polish and Dutch rivers, on the exposed English Channel and North Sea Basin, respectively. Multiproxy studies conducted along the northwestern European margin and especially from the northern Bay of Biscay (e.g., Zaragosi et al., 2001; Auffret et al., 2002; Mojtahid et al., 2005, 2017; Eynaud et al., 2007, 2012; Penaud et al., 2009; Toucanne et al., 2009, 2010), identified recurrent phases of meltwater inputs at the onset of HS1 (between 18.3 and 17 ka BP). Materialized in sediments

as millimeter- to centimeter-scale laminations, they were attributed to the seasonal melting of the EIS and seasonal subsequent freshwater discharge from the '*Fleuve Manche*' paleoriver. Further works provided new evidence of a differential contribution from ice sheets to the laminated deposit, with a particularly large SIS/Baltic sourced part during the last deglaciation and the HS1 interval (Toucanne et al., 2015). Until now, palynological investigation of this laminated facies (Eynaud, 1999, 2007, 2012; Auffret et al., 2000; Zaragosi et al., 2001; Penaud et al., 2009) was performed at resolution varying between 70 and 250 years only, due to the strong dilution of palynomorphs in sediments. Such laminated facies, corresponding to exceptionally high sedimentation rates, appear as ideal candidates to increase the temporal resolution of marine records and thus improve our understanding of short-lived fluctuations in the regime of the '*Fleuve Manche*' paleoriver and associated EIS dynamics.

Our study thus constitutes the first detailed dinocyst study encompassing the HS1 interval in the Bay of Biscay with a special focus on the laminated facies deposited at the onset of HS1. Our main objective was to decipher the set and sequence of events that occurred during this period over the northern Bay of Biscay. Our high-resolution palynological study was conducted on core MD13-3438 and combined with micropaleontological, geochemical and sedimentological analyses available for the twin reference core MD95-2002. Our multiproxy approach led to:

- (1) the reconstruction of the coupled EIS and '*Fleuve Manche*' paleoriver dynamics across HS1;
- (2) the study of high frequency seasonal variability within the laminated deposit, providing the first reconstruction of the sub-centennial climate variability across HS1 in the NE Atlantic;
- (3) and the characterization of sea surface conditions over the northern Bay of Biscay using dinocyst quantifications (keeping in mind their potentialities and limits in the study area).

2. Environmental context

2.1. Location of the studied core

The Calypso long piston core MD13-3438 (47°27' N; 8°27' W; 2180 m water depth; 36 m long) and twin core MD95-2002 (47°27' N; 8°32' W; 2174 m water depth) were respectively collected during the VT 133 / MERIADZEK (Woerther, 2013) and MD101-IMAGES (Bassinot and Labeyrie, 1996) oceanographic cruises on board the R/V Marion Dufresne (Table 1).

These marine sedimentary archives were retrieved from the Meriadzek Terrace, northern Bay of Biscay, directly off the mouth of the '*Fleuve Manche*' paleoriver (Fig. 1a). Their recovery, on a structure lying 600 m above the abyssal plain, guarantees few disturbances from gravity processes despite the proximity of deep-sea turbidite systems (Auffret et al., 2000; Zaragosi et al., 2000, 2001). The late Quaternary sedimentation of this area was imprinted by supplies from the northwestern European drainage catchments *via* the '*Fleuve Manche*' paleoriver during glacial lowstands (Auffret et al., 2000; Bourillet et al., 2003; Mojtahid et al., 2005; Zaragosi et al., 2006; Eynaud et al., 2007; Toucanne et al., 2008, 2009).

At present, the water column is structured by the deep Labrador Sea Water (~1500-2000 m depth) characterized by salinity ranging from 35 to 35.5 psu (Cossa et al., 2004) and the intermediate warm and salty (35.7 psu) Mediterranean Outflow Water (800-1500 m depth). Down to 800 m depth, the modern European Slope Current carries warm and salty waters to the Nordic Seas (Berx et al., 2013). Surface waters over the study area correspond to the southward recirculation of the North Atlantic Current (NAC; Fig. 1a), i.e., the south-eastern branch of the subpolar North Atlantic gyre (e.g., Sutton and Allen, 1997; Daniault et al., 2016). At the study site, modern (i.e., pre-21st century) mean Sea Surface Temperature (SST) and Salinity (SSS) are 11.7±0.6 °C and 35.54±0.05 psu in winter, and 17.5±1.0 °C and 35.58±0.10 psu in summer (World Ocean Atlas, 2001; Conkright et al., 2002).

2.2. Climatic changes in the northern Bay of Biscay during the last 40 kyr with focus on HS1

A large number of studies have shown that core MD95-2002 (Fig. 1a) has archived the history of deglacial pulses and meltwater discharge from the EIS (Zaragosi et al., 2001; Ménot et al., 2006; Eynaud et al., 2007, 2012; Penaud et al., 2009; Toucanne et al., 2009, 2015; Fig. 1b).

Data acquired on MD95-2002 (Fig. 1b) indeed provided nearshore marine to terrestrial information (Table 1 for related references).

In the North Atlantic Ocean, HS can be recognized on the basis of *Neogloboquadrina pachyderma* abundances close to 100% (e.g., Broecker et al., 1992; Eynaud et al., 2009). At site MD95-2002, HS1 (blue band on Fig. 1b) is thus identified as the interval between 18.2–14.6 ka BP (Fig. 1b). Early HS1 is evidenced by repeated alternation of millimeter- to centimeter-scale deposits (i.e., laminae) of mud and Coarse Lithic Grains (CLG) (Zaragosi et al., 2001). This facies results from an intense activity of the ‘*Fleuve Manche*’ paleoriver in response to the substantial EIS retreat (e.g., Mojtahtid et al., 2005; Zaragosi et al., 2006; Eynaud et al., 2007; Toucanne et al., 2008). This laminated interval is associated with high concentrations of the freshwater micro-algae *Pediastrum* (Penaud et al., 2009), high values of the Ti/Ca-XRF ratio (i.e., detrital *versus* biogenic proxy for enhanced terrigenous supply; Toucanne et al., 2009, 2012, 2015) and of the Branched and Isoprenoid Tetraether (BIT) index (i.e., a proxy for the relative fluvial input of terrestrial organic matter in the marine environment; Ménot et al., 2006). All those tracers highlight strong fluvial inputs and terrestrial-sourced organic sediment advections over the northern Bay of Biscay. This period is concomitant with major EIS melting events, especially from the Baltic Ice Stream of the SIS, accompanied by seasonal spring-summer meltwater discharge into the Bay of Biscay (Zaragosi et al., 2001; Mojtahtid et al., 2005; Ménot et al., 2006; Eynaud et al., 2007, 2012; Penaud et al., 2009; Toucanne et al., 2008, 2009, 2015). Within the laminated interval, the concentrations of CLG (Fig.1b) include both, Laurentide and European-sourced IRDs and also fluvial lithic grains transported *via* the ‘*Fleuve Manche*’ paleoriver. Increased CLG concentrations then recorded between 16.7 and 15 ka BP (Fig.1b; Zaragosi et al., 2001) characterize the HE1 phase. This interval contains the ‘conventional’ HE1 layer marked by notable peaks of magnetic susceptibility and carbonates (low Ti/Ca values) at ~16 ka BP characterizing the typical LIS surge (Grousset et al., 2000; Toucanne et al., 2015; Fig.1b).

3. Material and methods

3.1. Stratigraphy of core MD13-3438

The age model of core MD13-3438 is wedged on the last updated chronostratigraphy of core MD95-2002 (Toucanne et al., 2015). This latter was built with the Clam software (Blaauw, 2010) by integrating 22 AMS- ^{14}C dates over the last 40 kyr with additional tie-points: (i) 4 AMS- ^{14}C dates tied from the neighbouring cores MD03-2690 and MD03-2692, and (ii) *N. pachyderma* abundances correlated with the NGRIP $\delta^{18}\text{O}$ signal. Modern reservoir age correction is estimated to about 352 ± 92 years. Prior to HS1 and during the Holocene, the reservoir age constantly averaged around 400 ± 200 ^{14}C years. During HS1, the B/A and the Younger Dryas, average reservoir ages were respectively estimated to about 970, 680 and 875 years with uncertainties of 200 years (1s) (Toucanne et al., 2015).

The age model of core MD13-3438 was established by the correlation of its Ti/Ca-XRF signal with that of its twin core MD95-2002 (Fig. 1b). The XRF analysis of core MD13-3438 was performed at Ifremer (Plouzané) using an Avaatech X-Ray Fluorescence core scanner at 1 cm-resolution. Tie points used to synchronize both Ti/Ca-XRF signals can be found in Table 2. The correlation is supported by planktonic foraminiferal data (counts of the total assemblage performed on the >150 μm sediment fraction and on >300 individuals for each sample at EPOC laboratory), especially the high *N. pachyderma* relative abundances delimiting the HS1 interval (Fig. 1b). Our 5 cm-sampling frequency for dinocyst analyses enables us to achieve a temporal resolution of about 18 to 186 years, with a mean resolution of 58 years in all the studied sections, and of about 29 years in the laminated sequence, characterized by sedimentation rate values around 185 cm/kyr (Fig. 1b). This resolution thus provides valuable new details on the deglacial climatic history of the northern European margin across the HS1 interval.

It is worth noting that recent studies showed that the rapid cooling marking the onset of HS1 in the North Atlantic took place at $17.48 \text{ ka} \pm 0.21 \text{ ka}$ (1σ) (Missiaen et al., 2019; Waelbroeck et al., 2019) while maximal *N. pachyderma* abundances (i.e., allowing identifying HS 1) in core MD95-2002 between 880-390 cm, and between 575-245 cm in core MD13-3438, start at around 18.2 ka BP (Fig. 1b). Our chronologies are likely too old by about 700 years at the onset of HS1 as a result of underestimated reservoir ages over the last deglaciation.

192

193 **3.2. Dinocyst analyses**

194 a. Laboratory procedure, dinoflagellate cyst identification and diversity indices

195 A total of 76 samples were analysed from the 18.4-14 ka BP interval encompassing HS1.
196 Palynological treatments were performed at EPOC laboratory (University of Bordeaux,
197 France) following a standard protocol described by de Vernal et al. (1999). Calibrated tablets
198 of known concentrations of *Lycopodium clavatum* spores were added to each sample before
199 chemical treatments in order to estimate palynomorph concentrations. Chemical treatments
200 include cold HCl (10 %), cold HF (40 and 70%) and sieving through single-use 10 µm nylon
201 mesh screens. For each sample, an average of about 400 specimens (minimal counts of 175
202 cysts) was achieved using a Leica DM 2500 microscope at ×630 magnification except for the
203 laminated sequence between 575 and 330 cm for which an average of 120 specimens
204 (minimal counts of 100 cysts) was obtained due to strong dilution.

205 Dinocyst ecology has been thoroughly described through the progressive development and
206 compilation of atlases of modern cyst distribution (Matthiessen, 1995; Rochon et al., 1999;
207 Marret and Zonneveld, 2003; Zonneveld et al., 2013; van Nieuwenhove et al., 2020; Marret et
208 al., 2020). For this study, taxonomic attribution and the grouping of some species were done
209 in accordance with those atlases. *Brigantedinium* spp. taxa include all spherical brown cysts
210 excluding *Dubridinium* spp. Other peridinioid cysts were grouped (i.e., *Quinquecuspsis* spp.,
211 *Lejeunecysta* spp., *Dubridinium* spp. and *Votadinium* spp.) as miscellaneous peridinioid cysts
212 (MPCs). Dinocyst assemblages were described with the relative abundances of each taxon
213 calculated on the basis of the total sum of specimens counted including unidentified taxa and
214 excluding pre-Quaternary cysts. Finally, the species richness and Margalef index (Harper,
215 1999), calculated using PAST v.1.75b (Hammer et al., 2001), have been used to estimate the
216 dinocyst assemblage diversity as an additional ecological indicator. The species richness
217 represents the number of different taxa identified within each studied sample. The Margalef
218 diversity index (Margalef, 1958) is defined by: Margalef's Index = $(S - 1) / \ln(N)$, where S
219 and N correspond to the total number of species (S) and of individuals (N) in the sample.

220

221 b. Dinoflagellate cyst ratios and fluvial-derived palynological tracers

222 A “Warm/Cold” (W/C) ratio (Table 3) was used to qualitatively address SST variations
223 (Turon and Londeix, 1988; Versteegh, 1994; Combourieu- Nebout et al., 1999; Eynaud et al.,
224 2016; Penaud et al., 2016). Also, the “Heterotrophic/Autotrophic” (H/A) ratio (Table 3) was

calculated. This ratio is often used to discuss changes in primary productivity (PP) since heterotrophic dinoflagellates have a strict heterotrophic strategy of nutrition, being indirectly related to food resources, especially diatoms (Wall et al., 1977; Lewis et al., 1990; Marret, 1994; Zonneveld et al., 1997, 2001, 2013). It is worth noting that a large part of gonyaulacoids cysts (phototrophic cyst producing taxa) may survive thousands of years in well oxygenated sediments but heterotrophic peridinioids are extremely sensitive and vulnerable to early diagenesis. Since availability of oxygen in the sediments is the most important diagenetic variable, it has been suggested that the amount of species-selective degradation (i.e., here calculated as the H/A ratio for instance) may also be related to bottom water oxygen concentration, itself related to the rate of deep-ocean ventilation (Zonneveld et al., 2008). Both PP and/or taphonomic issues have to be taken into account to fully understand the H/A ratio.

The “*Lingulodinium machaerophorum* / *Operculodinium centrocarpum*” (Lmac/Ocen) ratio was used to discuss continental *versus* oceanic influences at the core location (e.g., Penaud et al., 2020), *L. machaerophorum* being a taxon dominating (nearly monospecifically) in estuarine environments of the French Atlantic coast (Wall et al., 1977; Morzadec-Kerfourn, 1977; Ganne et al., 2016; Lambert et al., 2017). Finally, a new ratio “*L. machaerophorum* / *Islandinium minutum*” (Lmac/Imin) has been used in this study as a proxy for summer *versus* winter prevailing seasonality modes. *I. minutum* is abundant in polar regions where surface waters do not exceed 0 °C in winter (Zonneveld et al., 2013).

In addition, pre-Quaternary dinocysts and Non Pollen Palynomorphs (NPP) including freshwater micro-algae *Pediastrum* spp. coenobia, *Botryococcus* spp. and *Concentricystes* spp.) were counted on the same palynological slides and expressed in absolute concentrations (palynomorphs/cm³). It has been demonstrated that *Pediastrum* spp. freshwater micro-algae are related to strong river discharge in marine environments (Zaragosi et al., 2001; Lézine et al., 2005; Eynaud et al., 2007; Penaud et al., 2009). Also, the biostratigraphical study of Kaiser (2001, unpublished data), based on the identification of reworked dinocysts in core MD95-2002 revealed that they were derived from the second half of the Mesozoic (Late Jurassic) to the Early Tertiary (Miocene), then characterizing the *Manche*, Parisian Basin and South England geological formations. Therefore, concentrations of *Pediastrum* spp. micro-algae and pre-Quaternary dinocysts, together with the ratio of “Reworked” (Rd, pre-Quaternary) *versus* “Modern” (Md) dinocysts (i.e., Rd/Md ratio), constitute robust proxies to discuss ‘*Fleuve Manche*’ paleoriver discharge in the Bay of Biscay (e.g., Zaragosi et al., 2001; Kaiser, 2001; Eynaud et al., 2007; Penaud et al., 2009). The Rd/Md ratio, reworked cyst

and *Pediastrum* spp. concentrations, together with *L. machaerophorum* occurrences and the Lmac/Ocen ratio will be referred to as Fluvial-derived Palynological Tracers (FPT) allowing discussing fluvio-glacial delivery to the NW European margin.

c. Dinocyst-based quantitative reconstructions of sea-surface parameters

The Modern Analogue Techniques (MAT), run on the “R version 2.7.0” software (R Development Core Team, 2008; <http://www.r-project.org/>), was applied on dinocyst assemblages to estimate past quantitative sea-surface environmental parameters. The MAT consists in the comparison of fossil records with modern dinocyst assemblages from the most recent update of the standardized Northern Hemisphere “modern” dinocyst database, which includes the abundance of 71 different taxa and 1968 sites in relation to 17 modern environmental parameters (de Vernal et al., 2020). This method relies on the assumption that modern relationships between hydrographical parameters and dinocyst assemblages were still valid in the past (e.g., Guiot and de Vernal, 2007). Sources of uncertainties, quantified with the Root Mean Square Errors (RMSE), could derive from the lack of modern analogues corresponding to fossil assemblages (Guiot and de Vernal, 2007; de Vernal et al., 2020).

The quantification of hydrological parameters is based on a weighted average of the values obtained for the five best modern analogues, with the maximum weight being given to the statistically closest analogue. Regarding the threshold distance ($d_T=1.2$), analogues are i) good when the distance $d < d_T/2$, ii) acceptable when $d_T/2 < d < d_T$, and iii) poor when $d > d_T$ (de Vernal et al., 2005). The n=1968 database and its related environmental database, allows the reconstruction of summer and winter SST and SSS, mean annual Primary Productivity (PP), and Sea Ice Cover Duration (SIC_D), with RMSE of ± 1.8 °C for SST_{summer}, ± 1.2 °C for SST_{winter}, ± 2.1 psu for SSS_{summer}, ± 1.1 psu for SSS_{winter}, ± 138 gC m⁻² for PP_{annual}, ± 1.5 months year⁻¹ for SIC_D.

4. Palynological results

Based on a cluster analysis run on dinocyst taxa percentages with the Psimpoll program, 9 palynozones were identified in core MD13-3438. We labelled them according to the stratigraphic interval they match: LGM (595-578 cm), HS1-a as the onset of HS1 prior to HE1 – HS1-a being subdivided into 5 sub-palynozones termed HS1-a1 to HS1-a5 (578-330 cm), and HS1-b (330-270 cm) and HS1-c (270-240 cm) as the first and second phase of HE1, respectively. Palynological data are presented and discussed according to these palynozones. They are plotted *versus* depth (cm) in result figures (Figs. 2 to 4), to exhibit the regularly sampled data without considering the strong impact of sedimentation rates, and then *versus* age (Cal ka BP) in the discussion figures (Figs. 5 to 7).

4.1. Quaternary dinocyst and other palynomorph abundances and derived indices

a. General observations on MD13-3438 results

A total of 31 different Quaternary dinocyst taxa (autotrophic or heterotrophic) have been identified (Table 3) with a species richness of about 15 different taxa per slide varying between 7 taxa at 580 cm and 25 taxa at 240 cm (Fig. 2). The Margalef index strongly matches the species richness with a general increasing trend from the bottom to the top of the studied section (Fig. 2), diversity being the lowest during the end of the LGM. Quaternary dinocyst concentrations vary from about 1200 to 47600 cysts/cm³, with average values of 5200 cysts/cm³ (Fig. 3). Overall, dinocyst assemblages are dominated by heterotrophic taxa (mean value of 57%) including especially *Brigantedinium* spp. (30%) and miscellaneous peridinioid cysts (MPCs) (21%), as well as *Islandinium minutum* (3%), cysts of *Protoperidinium nudum* grouped with *Selenopemphix quanta* (1%) as well as *Echinidinium* spp. (2%; Fig. 3). The H/A ratio trend (Fig. 2) seems to be mainly explained by fluctuations in *Brigantedinium* spp. (Fig. 2). The lowest values of this ratio are observed across the LGM, followed by high to moderate values across HS1-a and by a decreasing trend from the start of HS1-b to the onset of the B/A. Autotrophic taxa (mean value of 43%) are dominated by the following species: *Operculodinium centrocarpum* (16%), *Bitectatodinium tepikiense* (9%), *Lingulodinium machaerophorum* (4%), cysts of *Pentapharsodinium dalei* (3%), *Nematosphaeropsis labyrinthus* (3 %), *Spiniferites lazus* (2%), *Spiniferites ramosus* (2%) *Spiniferites belerius* (2%) and *Spiniferites septentrionalis* (2%) (Fig. 3).

317

318 b. Detailed observations according to the 9 palynozones

319 Each of the 9 palynozones can be described in terms of Quaternary dinocyst percentages,
320 which they were statistically established from, but also in terms of other micropaleontological
321 indicators. They are labelled from LGM to B/A (Fig. 2, Fig. 3):

322 • **LGM**: This lowermost zone exhibits the highest dinocyst concentrations (10000 to 47000
323 cysts/cm³) explained by maximal values of *O. centrocarpum* occurrences (higher than 85%).
324 *N. pachyderma* percentages show relatively low values (lower than 22%).

325 • **HS1-a**: The transition between LGM and HS1-a is marked by an abrupt decrease of *O.*
326 *centrocarpum* percentages and the increase of heterotrophic taxa (*Brigantedinium* spp. and
327 MPCs) representing more than 60% of the total cyst assemblages until 330 cm. Also, this
328 interval is characterized by increasing *N. pachyderma* percentages. Five sub-zones have been
329 distinguished: Sub-zone **HS1-a1** corresponds to the first evidence of increasing FPT values
330 and shows higher percentages of *B. tepikiense*, *Brigantedinium* spp. as well as MPCs. In sub-
331 zone **HS1-a2**, *B. tepikiense* percentages decrease while a diversification of the assemblage is
332 observed, with especially higher relative abundances of *Spiniferites* spp., *L. machaerophorum*
333 and cysts of *P. dalei*, while Quaternary dinocyst concentrations are the highest of the whole
334 HS1-a interval. Importantly, concentrations of reworked cysts and of total freshwater micro-
335 algae show maximal values during this HS1-a2 sub-zone (Fig. 2). Dinocyst concentrations
336 reach minimal values (2500 cyst/cm³) within HS1-a3, HS1-a4 and HS1-a5, when important
337 increases of MPCs are also recorded. While **HS1-a3** is characterized by the dominance of
338 *Brigantedinium* spp., HS1-a4 and HS1-a5 are marked by maximal values of *L.*
339 *machaerophorum* (16%) and the highest Lmac/Ocen values. A second peak of freshwater
340 micro-algae concentrations associated with maximal values of Rd/Md ratio additionally
341 characterises HS1-a4 (Fig. 2). *I. minutum* abundances increase from the beginning of HS1-a5
342 at 360 cm concomitantly with decreasing FPT values (Fig. 2, Fig. 3).

343 • **HS1-b**: This interval is generally characterized by a strong drop of MPCs with low to near-
344 zero values persisting all the way up to the top of the studied interval, and the gradual decline
345 of *L. machaerophorum* percentages. Maximal values of *I. minutum* (maximum about 20%)
346 and *S. septentrionalis* (maximum about 6%) occur in this zone.

347 • **HS1-c**: This interval is characterized by higher percentages of *O. centrocarpum* (20%), *B.*
348 *tepikiense* (18%), and *Echinidinium* spp. (10%), while *I. minutum* and *L. machaerophorum*
349 strongly decrease and are nearly absent until the end of the studied sequence.

• **B/A:** This interval is marked by the significant drop of *N. pachyderma* percentages from ~95% to ~35%. Dinocyst assemblages are characterized by the quasi-total disappearance of MPCs as well as significant percentages of *N. labyrinthus* (~15%). The first part of the B/A interval is characterized by increasing percentages of *Brigantedinium* spp. and occurrences of *Impagidinium* spp., while the second part of B/A (lowest *N. pachyderma* abundances) is characterized by increasing *B. tepikiense* and cysts of *P. dalei* percentages.

4.2. Dinocyst-based sea-surface quantifications

a. General observations on MD13-3438 results

Our MAT-based reconstructions rely on one to five modern analogues (Fig. 4a). Overall (76 samples in total), 5 samples are scored as “good” (especially within the basal LGM interval and at the top of the B/A interval) and 45 samples are scored as “acceptable” and range between the threshold distance ($d_T=1.2$) and $d_T/2$ (Fig. 4a). Samples scoring as “poor” are more particularly found within HS1-b and HS1-c, intervals also characterized by fewer analogues. Fig. 4a shows that the d_T pattern is very close to that of the H/A ratio. Consequently, uncertainties in dinocyst quantifications are likely related with heterotrophic taxa occurrences and especially with the high abundances of the *Brigantedinium* spp. cysts. From the $n=1968$ modern dinocyst database (de Vernal et al., 2020), 30 modern analogues are selected as the best analogues during the MAT calculations. They are mainly distributed along the eastern and western northern coasts of Canada, eastern coasts of USA, along the eastern and western coasts of Greenland and in the northern Atlantic Ocean, as well as in the Norwegian and Arctic Seas (Fig. 4b). Boxes have been delimited in Fig. 4b to show the main locations of those best analogues across each time interval in Fig. 4a (color code at the right of the figure). Within the end of the LGM, best analogues are located in the western coasts of Canada and in the Norwegian Sea. Most of the modern analogues selected within HS1-a are located in the eastern coasts of USA and Canada. For HS1-b, analogues are located in the eastern and western coasts of Greenland and Svalbard. Analogues found within HS1-c are more diversified and located in the eastern coasts of USA, Canada and Greenland and in the North Sea, this latter representing the principal source of modern analogues for the upper B/A interval (Figs. 4a and b).

b. Dinocyst-based sea surface parameters estimates

382 Dinocyst-derived mean summer SSTs (Fig. 4a) vary between -0.9 and 29.6 °C (average of
383 17.3 °C) and mean winter SSTs (Fig. 4a) range between -1.8 and 18.5 °C (average of 7.3 °C).
384 SST_{summer} and SST_{winter} values show similar trends, with their highest values recorded within
385 HS1-a (SST_{summer} of 20 °C and SST_{winter} of 9 °C). These atypically elevated values, especially
386 for SST_{summer}, exceed modern mean SSTs for the northern Bay of Biscay (modern average
387 SST_{summer} of 17.5 and SST_{winter} of 11.7°C). This zone is characterized by strong occurrences
388 of heterotrophic taxa (*Brigantedinium* spp. and MPCs), as also highlighted by the high H/A
389 ratio (Fig. 2), and modern analogues have been found along the eastern coasts of USA in
390 subtropical areas also characterized by highly productive (strong fluvial discharge) conditions
391 (modern analogue value of PP_{annual} around 1800 gC m⁻²).

392 Dinocyst-derived SSS_{summer} and SSS_{winter} (Fig. 4a) show low salinities along the studied
393 section ranging between 25 and 34 psu (average of 30 psu), below mean average modern
394 values of about 35.6 psu over the northern Bay of Biscay. The lowest salinities are recorded
395 during the laminated interval HS1-a with SSS_{summer} values of 29 psu, consistent with strong
396 fluvial discharge such as observed today on the eastern coasts of USA where analogues were
397 selected (Figs. 4a, b). The highest SSS values (around 34 psu) are recorded in the B/A
398 interval.

399 Although quantifications are especially critical within HS1-b because of the lack of modern
400 analogues across this interval (i.e., only 3 over 11 levels provided parameter estimates based
401 on MAT), this latter appears characterized by maximal Sea Ice Cover duration (“SIC_D” of ca.
402 10 months per year; Fig. 4a) and the lowest SST_{summer} and SST_{winter} (ca. 5 °C and -2 °C,
403 respectively; Fig. 4a). Finally, reconstructed PP_{annual} appears the highest within HS1-a (950
404 gC m⁻²; Fig. 4a).

5. The Last Deglaciation on the northern Bay of Biscay

Our compilation of MD13-3438 multiproxy signals indicative of fluvial discharge, allows us to describe for the first time the consequence of those hydrographic events at sub-centennial scale over the period covering the LGM to the B/A in the northern Bay of Biscay. We superimpose to our new and high-resolution results some of those previously published and established at lower temporal resolution (but longer time-scale) from twin core MD95-2002, (Fig. 5; Fig. 6; Zaragosi et al., 2001), and highlight the additional information provided by our new palynological study.

5.1. The warm climatic conditions bracketing HS1

Although the LGM and B/A are not the main targeted periods of our study, with consequently few samples documenting those intervals in core MD13-3438, it is important to describe the climatic context bracketing the main period of focus HS1. We thus, in the following, have synthesized information consistent with earlier works (especially including those relying on longer time-scale and multiproxy MD95-2002 records) and their derived climatic interpretations.

a. The end of the LGM

This interval is characterized by extremely low abundances of *N. pachyderma* (Figs. 5 and 6) and the highest percentages of *O. centrocarpum* (Fig. 6), considered to be a tracer of the NAC (Turon, 1984; Eynaud et al., 2004, 2012; Penaud et al., 2008, 2009). It is also associated with relatively high SST_{winter} (~9°C; Fig. 6) and high SSS (~34 psu; Fig. 6; see also Eynaud et al., 2012, focused on LGM salinities derived from multiproxy records). Those relatively warm conditions were suggested to be induced by a significant penetration of the warm and salty NAC in the Bay of Biscay (Eynaud, 1999; Zaragosi et al., 2001; Eynaud et al., 2007, 2012; Penaud et al., 2009) and more generally a more vigorous NAC in the NE Atlantic Ocean (Weinelt et al., 1996; Rosell-Melé and Comes, 1999; Kucera et al., 2005; de Vernal et al., 2000, 2002, 2005, 2006; Caille et al., 2013; Wary et al., 2015).

In parallel, a slight increase of the ‘*Fleuve Manche*’ paleoriver activity is recorded (through higher concentrations of reworked palynomorphs – Zaragosi et al., 2001 – and slight increase of the BIT-index and C/N ratio – Ménot et al., 2006; Fig. 5), synchronously with the 19-ka meltwater pulse identified in Europe by significant melting of the BIIS in Ireland (K-MWP:

Kilkeel event; Clark et al., 2004; Fig. 7) and of the southern SIS (i.e., R4 event; Toucanne et al., 2015; Fig. 7), while summer insolation at 65°N increases (Berger and Loutre, 1991; Fig. 7). In core MD13-3438, the end of the LGM displays small amounts of terrestrial organic matter exported to the northern Bay of Biscay and moderate marine productivity as suggested by low to moderate values of inferred PP, moderate MPCs (Fig. 5) and the lowest values of *Brigantedinium* spp. cyst percentages (Fig. 3).

b. The onset of the B/A

The onset of this relatively warm interval is characterized by a remarkable decrease in polar foraminiferal *N. pachyderma* percentages (Figs. 5 and 6) accompanied by a significant increase of *N. labyrinthus* percentages (Fig. 6) and a brief incursion of *Impagidinium* spp. (Fig. 3). Also, increases of dinocyst concentrations are coeval with a moderate TOC/N ratio in core MD95-2002 (Ménot et al., 2006; Fig. 5) suggesting a moderate PP increase. Since 14.6 ka BP, warmer sea-surface conditions (high SST_{winter} close to modern ones and high W/C ratio; Fig. 6) are associated with a rapid SSS increase (about 34 psu; Fig. 6) highlighting a vigorous NAC with warm and salty Atlantic surface waters carried to the study area, consistent with a concomitant strong AMOC as depicted in various records from the whole North Atlantic Ocean (e.g. McManus et al., 2004; Ng et al., 2018; Fig. 7). Data acquired in the northern North Atlantic (Caulle et al., 2013; Wary et al., 2015), the western Iberian margin (Bard et al., 2000; Martrat et al., 2007; Eynaud et al., 2009; Hodell et al., 2013; Salgueiro et al., 2014; Naughton et al., 2016) and the Alboran Sea (Cacho et al., 1999; Martrat et al., 2014; Català et al., 2019) support this observation of warmer conditions at a regional scale. It appears accompanied by rapid forest development as documented in the western Mediterranean borderlands and NW Iberia at the onset of the B/A (Naughton et al., 2007, 2016; Fletcher and Sanchez-Goñi, 2008; Combourieu-Nebout et al., 2009; Camuera et al., 2019, 2021; Fig. 7) concomitant with increasing humidity, as also identified in speleothem records from southwestern Europe (Genty et al., 2006; Moreno et al., 2010; Jalut et al., 2010). An increase in seaward transfer of fluvially-derived sediments is reported directly north of the Meriadzek Terrace ca. 16-14 ka, indicating possible influence of the retreating Irish Ice Sheet (which is not connected to the ‘*Fleuve Manche*’ paleoriver) at that time (Toucanne et al., 2008). However, our data show no such evidence of deglacial meltwater fluxes at site MD13-3438, neither with FPT values (only a slight increase of *Pediastrum* spp. concentrations; Fig. 5) nor with the BIT-index (Ménot et al., 2006; Fig. 5). This lack of deglacial evidence on the Meriadzek Terrace may be explained by the major episode of sea-level rise referred to as

“Meltwater Pulse 1A” (MWP-1A, Fig. 7) which occurred at around ~14.6-14.3 ka BP. This led to a ~20 m sea-level rise in less than 500 years (Weaver et al., 2003; Deschamps et al., 2012; Lambeck et al., 2014; Fig. 7) probably responsible for the displacement of the ‘*Fleuve Manche*’ mouth about 300 km eastward (Toucanne et al., 2012, 2010), thus rapidly halting any ‘*Fleuve Manche*’ imprint on the sedimentary record of our study site.

5.2. A sub-centennial subdivision of HS1 in the northern Bay of Biscay

a. Early-HS1, laminated interval (HS1-a): 18.2-16.7 ka BP

General observations

Overall, the HS1-a interval is characterized by a huge increase in terrigenous sediment supplies in the study area, evidenced through higher values of the Ti/Ca-XRF ratio (Fig. 5) and higher sedimentation rates (mean values of about 200 cm/kyr at the study site) at site MD13-348, as well as through maximal values of the BIT-index (core MD95-2002; Ménot et al., 2006; Fig. 5) and low CLG concentrations mainly due to dilution by terrigenous sediment at site MD95-2002. This interval is also marked by a reduction of *O. centrocarpum* percentages from about 90% to 10 % and increasing percentages of *Spiniferites* spp. (i.e., *S. bentorii* and *S. lazus*), which point to a transition from a full-oceanic (end of the LGM) to a neritic to coastal influence (cf. Penaud et al., 2020 for modern ecological requirements of dinocyst taxa in the study area) with enhanced PP (Fig. 5). This is also confirmed by high values of the Lmac/Ocen ratio (Fig. 7) and, in general, by the substantial rise of FPT values. These indications support the progradation of a large outer-shelf delta on the margin at that time and huge delivery of the ‘*Fleuve Manche*’ sediment load close to the Meriadzek Terrace (Toucanne et al., 2012). Geochemical investigations suggested that the meltwater release pattern is mainly related to the Baltic Ice Stream of the SIS (R5 event; Toucanne et al., 2015; Fig. 7).

Enhanced proximal fluvio-glacial delivery from the ‘*Fleuve Manche*’ paleoriver was suggested to result from the seasonal influx of meltwater from the retreating EIS and responsible for the formation of the laminated facies (Zaragosi et al., 2001, 2006; Mojtahid et al., 2005; Ménot et al., 2006; Eynaud et al., 2007, 2012; Toucanne et al., 2008, 2009; Penaud et al., 2009; Fig. 7). Here, dinocyst-derived quantitative reconstructions further show a significant drop of SSS, especially of SSS_{summer} (mean values about 29 psu; Fig. 6), with a large seasonal SSS amplitude (>2 psu; Fig. 6), suggesting that the EIS melting occurred

mainly during the spring-summer season (Mojtahid et al., 2005; Toucanne et al., 2009, 2015; Eynaud et al., 2012; see Fig. 8 for a conceptual scheme of past hydrographic changes in the study area).

The HS1-a interval is characterized by a sustainable increase in the H/A ratio (fig. 7). Interestingly, this pattern matches the AMOC trend (cf. $^{231}\text{Pa}/^{230}\text{Th}$ values; McManus et al., 2004; Ng et al., 2018; Fig.7) with high values of heterotrophic cysts during the AMOC slowdown in the North Atlantic Ocean and conversely. This observation highlights the probable taphonomic issue linked to varying bottom water ventilation (Zonneveld et al., 2008). In our study, heterotrophics (Fig. 5) may be related to either increasing PP and/or better preservation of peridinioid cysts under low bottom-water oxygenation. Remarkably, benthic foraminiferal taxa, from the northern Bay of Biscay, indicated both a general eutrophication and a severe bottom water dysoxia in the sea floor (Mojtahid et al., 2017). Overall cold and dry conditions are observed in the southern and western Iberian Peninsula, with a strong reduction of the Mediterranean and temperate forests (Fletcher and Sanchez Goñi, 2008; Naughton et al., 2016; Camuera et al., 2021; Fig.7) and a notable increase of xerophytic taxa (i.e., steppic taxa) (Camuera et al., 2021; Fig.7).

Sub-centennial variability of fluvio-glacial discharge on northern Bay of Biscay

Our unprecedented high-resolution multiproxy study of the laminated facies shows a new sub-centennial-scale structure, following a multi-step scenario encompassing 5 intervals (Figs. 5 to 7; cf. subsection 4.1). Enhanced fluvial inputs were recorded within HS1-a2 corresponding to maximal terrestrial advection and HS1-a4 consisting to the strongest stratification level of the surface water column and the strongest westward shift of the freshwater front toward the ocean (Fig. 5). Interestingly, these intervals are characterized by lower and fluctuating *N. pachyderma* percentages and increased values of SST and seasonality reconstructed from dinocysts at site MD13-3438, testifying warm sea-surface conditions (Fig. 6). Decreased fluvial discharge marked the HS1-a1, HS1-a3 and HS1-a5 intervals, this latter appears characterized by the establishment of cold sea surface conditions in the northern Bay of Biscay within the latter.

The oscillating pattern of ‘*Fleuve Manche*’ paleoriver runoff illustrated here by the different phases identified in HS1-a finds a striking echo in the environmental fluctuations from arid to more humid conditions in the Iberian Peninsula between 18.4 and 16.4 ka BP (Camuera et al., 2021: Fig. 7), with phases of higher runoff matching those of increased humidity in southern

Europe. Furthermore, these observations are regionally supported by SST records from the Alboran Sea (Cacho et al., 1999, 2006; Martrat et al., 2014; Fig.7).

The role of multidecadal changes in seasonality on fluvio-glacial fluxes to the northern Bay of Biscay

The first conceptual model proposed by Mojtabid et al. (2005) suggested: (i) enhanced melting of EIS and surrounding glaciers, responsible for enhanced clay and CLG fluxes to the Bay of Biscay via the '*Fleuve Manche*' paleoriver during spring and especially summer seasons, and (ii) freezing SST and sea-ice cover establishment inhibiting EIS melting during winter seasons. While first studies attributed CLG-rich units to the seasonal calving of EIS icebergs onto the northern Bay of Biscay (Mojtabid et al., 2005; Zaragosi et al., 2006; Eynaud et al., 2007), Toucanne et al (2009) later proposed that the episodic rainout of these particular CLG originates instead from the '*Fleuve Manche*' paleoriver discharge of anchor-ice (i.e., ice attached to the riverbed) and of sediment-rich frazil-ice (Reimnitz and Kempama, 1987; Kempama et al., 2001).

Echoing the seasonal dynamic of the '*Fleuve Manche*' paleoriver, our data interestingly suggest warmer sea-surface conditions within HS1-a (mean SST values about ~9°C and high values of the W/C ratio; Figs. 6) associated with a significant temperature seasonality increase (> 12°C; Fig. 6), especially within periods of maximal fluvial discharge, namely HS1-a2 and HS1-a4. To further investigate this, we specifically created a 'summer *versus* winter index' as the ratio between two major dinocyst taxa occurring regionally and illustrating two opposite environmental dynamics: the estuarine taxon *L. machaerophorum* and the polar taxon *I. minutum* (Lmac/Imin ratio; Fig.7). We consider *L. machaerophorum*, whose highest abundances at present are encountered in areas under strong fluvial influence, as characteristics of the '*Fleuve Manche*' paleoriver discharge induced by European ice sheet retreat and melting during the deglaciation, i.e., which occurred preferentially during summer seasons according to earlier sedimentological studies cited above. In contrast, occurrences of the polar species *I. minutum* are associated with cold sea-surface conditions and expansion of temporary sea-ice cover (1-3 months/year, Fig. 6) over the studied interval in the northern Bay of Biscay (as it is at present at high latitudes), then mostly expressing a winter signal (Fig. 8).

The Lmac/Imin ratio (Fig. 7) matches the seasonality signal (Fig. 6). It shows high values within most of the HS1-a interval and particularly during maximal runoff periods HS1-a2 and HS1-a4. On the contrary, intervals characterized by lower fluvio-glacial inputs (HS1-a1, HS1-

a3 and HS1-a5) depict low L_{mac}/I_{min} values, highlighting the limited influence of summer seasons and the prevalence of cold winters (Fig. 8). We thus argue that multidecadal changes of the seasonality pattern played a substantial role on the EIS and ‘*Fleuve Manche*’ paleoriver sub-centennial dynamics during the last deglaciation, with release of river ice and meltwater via the ‘*Fleuve Manche*’ paleoriver favoured during episodes with prevailing summer mode, and reversely formation of anchor- and frazil-ice in the ‘*Fleuve Manche*’ paleoriver favoured during episodes of winter prevailing mode (see Fig. 8 for a conceptual scheme of past hydrographic changes in the study area).

Such an enhanced warm summer-prevailing mechanism could then explain the EIS melting, as far as in the northern European lowlands, in an ‘apparent’ long cold climate stadial (Fig. 8). HS1 is generally seen as a very cold and dry interval in the North Atlantic Ocean and across Europe (Bard et al., 2000), marked by overall light NGRIP $\delta^{18}O$ values, especially during HE1 (Fig. 7). The strong changes in SST seasonality observed in the northeastern Atlantic Ocean, and weakly expressed in high latitudes, may then support the hypothesis that Greenland temperatures mainly represent winter temperatures (Denton et al., 2005, 2010; Buizert et al., 2014, 2018). However, it is worth noting the occurrence of a plateau of slightly lighter NGRIP $\delta^{18}O$ values over the HS1-a interval (Fig. 7) arguing for slightly warmer atmospheric conditions.

b. Late-HS1, namely HE1 layer (HS1-b and HS1-c): 16.7–14.6 ka BP

In our study, at 16.7 ka BP, the onset of HE1 phase is characterized by the disappearance of laminae deposits as well as a drastic drop of sedimentation rates (from 156 to 41 cm/kyr; Fig. 5) and in the BIT-index (Ménot et al., 2006; Fig. 5). Those changes are associated with a substantial decline of FPT values including the L_{mac}/O_{cen} ratio. All these observations point to a considerable decrease of the ‘*Fleuve Manche*’ paleoriver runoff. Also, high values of $C_{37:4}$ alkenones (i.e., a biomarker derived from haptophyte algae, and used as a proxy for low salinity water associated with icebergs; Ménot et al., 2006) are synchronous with high CLG concentrations (Fig. 6). The local hydrology is thus strongly impacted by both Laurentide- and European-derived icebergs at that time (Grousset et al., 2000). Increasing percentages of the polar species *I. minutum* (Fig. 6), recognized as a good tracer of cold surface waters seasonally to quasi-permanently covered with sea-ice (Zonneveld et al., 2013; Radi et al., 2013), are synchronous with low reconstructed SST_{winter} (Fig. 6) indicating cold sea-surface conditions (Fig. 6).

Dinocyst assemblages enabled us to determine two sub-zones, HS1-b and HS1-c, with a 120 years study resolution on the entire HE1 interval:

HS1-b (16.7–15.6 ka BP; 85 years study resolution) is characterized by the drastic decrease of fluvial inputs (low FPT values, Fig. 5; decreasing L_{mac}/O_{cen} ratio, Fig. 7). Maximal *I. minutum* (Fig. 6) and *S. septentrionalis* (Fig. 3) occurrences are recorded at 16.4 ka BP, depicting the coldest conditions (Figs. 6 and 7). Low *O. centrocarpum* percentages additionally suggest a still weakened NAC, consistently with $^{231}Pa/^{230}Th$ record which suggests that a weakened AMOC state persisted for over a millennium (~16.5–15 ka BP) in the North Atlantic Ocean (Ng et al., 2018; Fig. 7). These cold surface conditions recorded on the northern Bay of Biscay may have been responsible for a weaker EIS melting and are consistent with records from the IRD belt indicating a widespread cooling associated with a major calving episode of the LIS (Stanford et al., 2011; Hodell et al., 2017). This interval is indeed marked by a notable peak of magnetic susceptibility (Figs. 5 and 6) and significant CLG concentrations (core MD95-2002; Figs. 5 and 6) accompanied by minimal values of the Ti/Ca-XRF ratio at 16 ka BP (Fig. 5) indicative of carbonate-rich CLG (Auffret et al., 1996; Toucanne, 2009;). This may correspond to the ‘cemented marls’ of Hemming (2004) interpreted as the LIS (i.e., Hudson Strait) iceberg delivery to the North Atlantic Ocean (Toucanne et al., 2015).

Extremely cold conditions at our study site (Figs. 6, 7 and 8) are synchronous with extremely cold boreal temperatures (lightest NGRIP $\delta^{18}O$ values of the entire HE1 phase; Fig. 7) and with the coldest and most arid environments documented in marine palynological sequences from the Alboran Sea (Combourieu-Nebout et al., 2002, Fletcher and Sanchez Goñi, 2008; Fig7) and off the Iberian Peninsula (Turon et al., 2003; Naughton et al., 2007, 2009, 2016). The dryness over the Mediterranean borderlands increased (Morellón et al., 2009; Camuera et al., 2019; Fig. 7) in relation with the southward migration of the polar front (Eynaud et al., 2009).

HS1-c (15.6–14.6 ka BP; 170 years study resolution) is characterized by higher percentages of *B. tepikiense* and *P. dalei* cysts, currently observed in surface sediments from the subpolar North Atlantic basin (Harland, 1983; de Vernal et al., 1992; Dale, 1996; Rochon et al., 1999), and also clearly accompanied by both decreasing *I. minutum* percentages and CLG concentrations (Figs. 5 and 6) as well as increasing *O. centrocarpum* percentages (Fig. 3). This suggests warm sea-surface conditions (Fig. 6) with increased thermal seasonal amplitudes (Figs. 6, 7 and 8). The retreat of winter sea-ice marks the end of the HS1 interval in the northeastern Atlantic Ocean. Increasingly warm conditions on the northern Bay of

637 Biscay are consistent with an intensified NAC and increasing SSTs as detected at regional
638 scale (e.g., Caille et al., 2013; Wary et al., 2015; Naughton et al., 2016; Hodell et al., 2013;
639 Cacho et al., 1999; Martrat et al., 2014; Català et al., 2019). Also, warmer and more humid
640 conditions are observed in the southern and western Iberian Peninsula (Fletcher and Sanchez
641 Goñi, 2008; Naughton et al., 2009, 2016; Camuera et al., 2021; Fig.7) that may be attributed
642 to the northward displacement of the polar front (Cayre et al., 1999; Naughton et al., 2009,
643 2016).

6. Conclusion

The high-resolution palynological investigation of the last deglaciation in core MD13-3438 (northern Bay of Biscay) highlights significant climatic and paleoenvironmental changes related to both the proximal European Ice Sheets (EIS) and the '*Fleuve Manche*' paleoriver dynamics. Dinocyst-based quantitative reconstructions provide an evaluation of past hydrographical changes. Seven short-scale sub-phases within the HS1 interval were identified for the first time on the northern Bay of Biscay:

HS1-a (i.e., laminated interval; 18.2–16.7 ka BP) is generally characterized by marked multidecadal changes in the seasonality range, i.e., warm summers resulting in enhanced EIS melting and '*Fleuve Manche*' paleoriver runoff, causing the largest drop of SSS and cold winters resulting in freezing conditions and lower '*Fleuve Manche*' paleoriver activity. Fluvial-derived palynological tracers indicate that this interval is subdivided into five sub-phases. Two maximal meltwater episodes (HS1-a2, 17.8–17.6 ka BP and HS1-a4, 17.4–17 ka BP) are detected, with substantial erosional processes in the '*Fleuve Manche*' basin and strong stratification of the water column. Both events correspond to strong seasonality phases dominated by summer-prevailing modes. Fluvio-glacial discharge of the '*Fleuve Manche*' paleoriver, although significant, is comparatively moderate during HS1-a1 (18.2–17.8 ka BP), HS1-a3 (17.6–17.4 ka BP) and HS1-a5 (17–16.7 ka BP), and possibly result from a decrease of the seasonality that could have limited the EIS melting at that time. Indeed, our new data suggest winter-prevailing modes at that time.

HS1-b (i.e., first phase of HE1; 16.7–15.6 ka BP) includes the Laurentide-sourced IRDs recorded at ~16 ka BP. This interval is characterized by a substantial decrease of proximal fluvial inputs and cold winters. Cooling sea-surface conditions are recorded with seasonal winter sea-ice cover occurrences and the lowest reconstructed SSTs.

HS1-c (i.e., second phase of HE1; 15.6–14.6 ka BP) is characterized by warmer sea-surface conditions, suggesting the advection of Atlantic surface waters conveyed by the NAC and leading to the transition to the warm B/A.

Our reconstructed sub-centennial variability suggests the crucial role of multidecadal seasonal changes, especially within the early HS 1 interval, for fluvio-glacial dynamics and the melting of the EIS. Furthermore, our records are in agreement with marine and terrestrial sequences from the western Mediterranean basin suggesting common regional forcings acting on the multi-scale climate variability across the last deglaciation.

7. Acknowledgments

This work was supported by the French projects: ANR IDEGLACE, INSU RISCC, INSU ICE-BIO-RAM and by the European Research Council ERC grant ACCLIMATE/n° 339108. This work results from regional, national and international collaborations, between LGO laboratory (Brest University,), Ifremer (GM), EPOC laboratory (Bordeaux University,), LSCE and the Universitat Autònoma of Barcelona. We received funding from the CG29 (*Conseil Général du Finistère*, 29) and financial support from LGO and EPOC laboratories. This work was also supported by ISblue project, Interdisciplinary graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French government under the program "Investissements d'Avenir". We thank Mikael Rovere and Patrice Woerther for their assistance onboard the R/V Marion Dufresne, Muriel Georget (EPOC) and Pierre-Olivier Coste (LGO) for their help and laboratory assistance. Mélanie Wary's contribution was also supported through funding from the Spanish Ministry of Science, Innovation and Universities, through the "María de Maeztu" program for Units of Excellence (MDM-2015-0552).

8. Table caption

Table 1: Datasets acquired on both cores MD95-2002 and MD13-3438 with their corresponding references.

Table 2: List of tie points defined by aligning the Ti/Ca-XRF signal of the well-dated core MD95-2002 with that of the studied core MD13-3438.

Table 3: List of identified autotrophic and heterotrophic dinocyst taxa according to their ecological preferences: cold surface-waters (in blue; C) and warm surface-waters (in orange; W). The Warm/Cold qualitative temperature index is based on these W and C species.

9. Figure captions

Figure 1: Paleogeographic reconstruction of western Europe during the LGM (~ 20 ka BP) modified from Toucanne et al. (2015) and records of the past fluvial activity of the 'Fleuve Manche' paleoriver. **a)** The yellow star corresponds to the location of the study core MD13-

3438 and of the core MD95-2002. Black lines represent the extension of the Northern Hemisphere ice caps: (1) the European Ice Sheet (EIS) including the Scandinavian (SIS) and British-Irish Ice Sheets (BIIS) (Boulton et al., 2001; Clark et al., 2012b; Ehlers et al., 2011), with their main fluvio-glacial paths (black arrows), as well as the Alpine Ice Sheet; and (2) the Laurentide Ice Sheet (LIS) with main fluvio-glacial path (white arrow). The bold white arrow identifies the drainage of the '*Fleuve Manche*' paleoriver. The North Atlantic surface circulation (red arrows for the warmer North Atlantic Current (NAC) and its branches) is also shown with the return flow pathway of the deep waters (blue arrows). White shading indicates the extent of the EIS and LIS. The trajectory of the icebergs from the LIS are represented with light blue arrows and white arrows indicate the main supply sources of freshwater to the North Atlantic. **b)** Interval from 40 to 5 ka BP: July insolation at 65°N (Berger and Loutre, 1991), Greenland $\delta^{18}\text{O}$ record (GICC05; Svensson et al., 2008, in black), Ti/Ca- XRF data and *N. pachyderma* abundances of core MD13-3438 (blue) compared with the MD95-2002 dataset (in red) including: *N. pachyderma* abundances (Zaragosi et al., 2001), coarse lithic grain concentrations (CLG; 10^3 grains/g dry sed.; Zaragosi et al., 2001), magnetic susceptibility of bulk sediment (U_{em}; Grousset et al., 2000), fluvial input proxies (number of laminae per cm, concentrations of freshwater micro-algae *Pediastrum* spp.; Zaragosi et al., 2001) and the branched and isoprenoid tetraether (BIT) index (Ménot et al., 2006). Light grey band represents the LGM (23–19 ka BP) while blue band allows visualizing HS1 in both cores MD95-2002 and MD13-3438 (18.4–14 ka BP). Dark grey band represents the conventional HE1 layer occurring at ~16 ka BP.

Figure 2: Palynological data against depth (cm) with taxonomical indices (species richness, black; Margalef index, grey), percentages of the most abundant taxa *Brigantedinium* spp., ratio between Heterotrophic and Autotrophic (H/A) dinocysts (grey), percentages of *L. machaerophorum*, ratio between *L. machaerophorum* and *O. centrocarpum* (grey), concentrations of reworked dinocysts (blue) and of total freshwater microalgae (dark grey line: including *Botryococcus* spp. (purple), *Concentricystes* spp. (pink) and *Pediastrum* spp. (green)). The reworked (pre-Quaternary) vs. modern (in situ) cyst ratio (Rd/Md) is highlighted in grey. Horizontal lines delineate the nine palynozones / climatic subdivisions discussed in the manuscript. LGM: Last Glacial Maximum; HS1-a1 to HS1-a5: Laminated interval; HS1-b and HS1-c: Heinrich Event 1; B/A: Bølling/Allerød.

Figure 3: Diagram of dinocyst assemblages (i.e., major taxa with values higher than 2% at

least once in palynological assemblages) *versus* depth (cm) for core MD13-3438, compared with the relative abundances of the planktonic foraminiferal species *N. pachyderma* and with total dinocyst concentrations (black line) including autotrophic (black dotted line) and heterotrophic (grey dotted line) taxa (cysts/cm³). Red and black stars indicate AMS-¹⁴C dates obtained from cores MD95-2002 and MD03-2690 (Eynaud et al., 2012; Mojtahid et al., 2005; Toucanne et al., 2008; 2015; Zaragosi et al., 2006; Zaragosi et al., 2001) respectively, and grey star refers to the NGRIP tie point used for the MD95-2002 age model. These ages are here projected as equivalent depths for core MD13-3438 using the twins Ti/Ca-XRF signals of both MD95-2002 and MD13-3438 cores (Fig. 1b). Palynozones/climatic subdivisions were based on dinocyst clustering established by Psimpoll program. Horizontal lines delineate the nine palynozones discussed in the manuscript (same as in Fig. 2). High percentages of dinocyst taxa are shown by orange vertical arrows for long-lasting periods and by dotted orange vertical arrows for short periods.

Figure 4: a) Dinocyst-based environmental parameters reconstructed for core MD13-3438 with the Modern Analog Technique (MAT) using the n=1968 modern dinocyst database (de Vernal et al., 2020) *versus* depth: SST and SSS for the summer (red dotted line) and winter (blue dotted line), PP_{annual} (black dotted line), and Sea Ice cover Duration (months/year; grey dotted line). Number of analogues found for each assemblage (minimum of 0 and maximum of 5 allowed with the MAT; dark red). Distances between fossil MD13-3438 assemblages and their “closest” modern analogue (i.e., D_{min}). Threshold distance value (d_T=1.2; red line) for quantification robustness: good analogues if D_{min} is between 0 (perfect analogue) and d_T/2 (orange line); acceptable analogues if D_{min} is between 0.6 and d_T=1.2, and poor analogues (i.e. caution with quantifications) if D_{min} > d_T (de Vernal et al., 2005). Horizontal lines delineate the nine palynozones / climatic subdivisions discussed in the manuscript (same as in Fig. 2 and 3). **b)** Map showing the geographical distribution of the closest analogues (i.e., corresponding to D_{min}) selected by the MAT for the studied section of core MD13-3438 (environmental dataset also from de Vernal et al., 2020). Analogues have been grouped according to geographical areas (color code for boxes; also repeated in Fig. 4a for source area of closest analogues in each palynozone).

Figure 5: Deglacial records of the ‘*Fleuve Manche*’ paleoriver activity on the northern Bay of Biscay between 18.4 and 14 ka BP. For core MD13-3438 (in blue): *N. pachyderma* percentages, sedimentation rates (cm/kyr), Ti/Ca-XRF ratio, freshwater microalgae

Pediastrum spp. concentrations (10^3 algae/cm³), reworked dinocyst concentrations (10^3 cysts/cm³), *L. machaerophorum* percentages and concentrations (10^3 cysts/cm³), reconstructed PP_{annual} (10^3 g C m⁻²) and peridinioid cyst concentrations (10^3 cysts/cm³). For core MD95-2002 (in red): *N. pachyderma* percentages, coarse lithic grain concentrations CLG (10^3 grain/g dry sed.), number of laminae (laminae/cm), Reworked *versus* Modern dinocyst ratio (Rd/Md, shaded red), *L. machaerophorum* percentages (shaded red) (Zaragosi et al., 2001), magnetic susceptibility of bulk sediment (U em; Grousset et al., 2000), Ti/Ca-XRF ratio (Toucanne et al., 2015), branched and isoprenoid tetraether (BIT index; Ménot et al., 2006) and Carbon/Nitrogen ratio (C/N; Ménot et al., 2006). Yellow bands highlight periods of reduced runoff (HS1-a1, HS1-a3 and HS1-a5) and orange bands correspond to the periods of maximal fluvio-glacial discharge. Blue and pink bands highlight the HE1 phase (HS1-b, including the conventional HE1 layer at ~16 ka BP, and HS1-c, respectively). Horizontal lines delineate the nine palynozones / climatic subdivisions discussed in the manuscript (same as in Fig. 2, 3 and 4).

Figure 6: Reconstructed hydrological parameters derived from dinocyst assemblages of the core MD13-3438 (in blue, this study) compared with data from MD95-2002 (in red; Zaragosi et al., 2001). *N. pachyderma* percentages, coarse lithic grain concentrations CLG, magnetic susceptibility of bulk sediment (U em ; Grousset et al., 2000), and number of laminae are compiled with relative abundances of the main dinocyst species, percentages of C37:4 among C37 alkenones (red, Ménot et al., 2006), Warm/Cold dinocyst ratio (W/C, This study), dinocyst-derived sea-surface parameters of MD13-3438 (Sea Surface Temperatures: SST_{winter} in blue, SST_{summer} in orange; Sea Surface Salinity: SSS_{summer-winter}; Seasonality (SST_{summer} - SST_{winter})). Yellow bands highlight periods of reduced runoff (HS1-a1, HS1-a3 and HS1-a5) and orange bands correspond to the periods of maximal fluvio-glacial discharge. Blue and pink bands highlight the HE1 phase (HS1-b and HS1-c, respectively). Horizontal lines delineate the nine palynozones / climatic subdivisions discussed in the manuscript (same as in Fig. 2, 3, 4 and 5).

Figure 7: Synthesis of main information regarding regional paleoclimatic reconstructions for the last deglaciation and our climate subdivisions especially within HS1.

Composite Relative Sea Level (RSL) curve (Lambeck et al., 2014) with identified K-MWP at 19 ka BP and MWP-1A at ~14.6 ka BP, July Insolation curve at 65°N (Berger and Loutre, 1991), $\delta^{18}\text{O}$ NGRIP record (GICC05; Svensson et al., 2008), ²³¹Pa/²³⁰Th signal (Ng et al.,

2018), *N. pachyderma* percentages (blue, this study; red for the core MD95-2002 (Zaragosi et al., 2001)), H/A ratio (blue, this study; has been also translated and reversed for comparison), Runoff events (R) 4 and 5 (core MD95-2002, Toucanne et al., 2015), *L. machaerophorum* versus *O. centrocarpum* (Lmac/Ocen ratio, This study), *L. machaerophorum* versus *I. minutum* (Lmac/Imin ratio, This study), dinocyst-derived sea-surface seasonality (SSTsummer–SSTwinter) and three-point moving average of Xerophyte and Mediterranean forest percentages from southern Spain (Camuera et al., 2021) as well as temperate forest percentages from the Alboran Sea (Fletcher and Sanchez Goñi, 2008). The proposed comparison respects dating uncertainties and strong reservoir age impacts at that time may be responsible for generating temporal offsets between records.

Dashed grey lines correspond to original values, whereas continuous lines correspond to smoothed data. Yellow bands highlight periods of reduced runoff (HS1-a1, HS1-a3 and HS1-a5) and orange bands correspond to periods of maximal fluvio-glacial discharge. Blue and pink bands highlight the HE1 phase (HS1-b, including the conventional HE1 layer at ~16 ka, and HS1-c, respectively).

Figure 8: Conceptual model illustrating hydrological conditions and processes involved for fluvio-glacial dynamics across the HS1 interval in the northern Bay of Biscay.

10. Appendix caption

Appendix: Exhaustive table of palynological results (percentages, concentrations and reconstructed parameters)

11. References

- Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., et al., 2006. The Greenland ice core chronology 2005, 15-42 kyr. Part 1: constructing the time scale. *Quat. Sci. Rev.* 25, 3246–3257.
- Auffret, G.A., Boelaert, A., Vergnaud-Grazzini, C., Muller, C., Kerbrat, R., 1996. Identification of Heinrich layers in core KS 01 North-Eastern Atlantic (46°N, 17°W), implications for their origin. *Mar. Geol.* 131, 5–20.
- Auffret, G., Zaragosi, S., Voisset, M., Droz, L., Loubrieu, B., Pelleau, P., Savoye, B., Bourillet, J.-F., Baltzer, A., Bourquin, S., Dennielou, B., Coutelle, A., Weber, N.,

845 Floch, G., 2000. First observations on the morphology and recent sedimentary
846 processes of the Celtic Deep Sea Fan [Premieres observations sur la morphologie et
847 les processus sedimentaires recents de l'Eventail celtique]. *Oceanologica Acta* 23,
848 109–116.

849 Auffret, G., Zaragosi, S., Dennielou, B., Cortijo, E., Van Rooij, D., Grousset, F., Pujol, C.,
850 Eynaud, F., Siegert, M., 2002. Terrigenous fluxes at the Celtic margin during the last
851 glacial cycle. *Mar. Geol.* 188, 79–108.

852 Bard, E., Rostek, F., Turon, J.L., Gendreau, S., 2000. Hydrological impact of Heinrich events
853 in the subtropical Northeast Atlantic. *Science* 289, 1321–1324.

854 Barker, S., Diz, P., Vautravers, M. J., Pike, J., Knorr, G., Hall, I. R., Broecker, W. S., 2009.
855 Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature* 457,
856 1097–1102.

857 Bassinot, F., Labeyrie, L., 1996. IMAGES I, MD101: A coring cruise of the R/V Marion
858 Dufresne in the North Atlantic Ocean and Norwegian Sea. Les rapports de campagnes
859 à la mer. Institut Paul-Emile Victor (IPEV).

860 Berger, A., Loutre, M.F., 1991. Insolation values for the climate of the last 10 million years.
861 *Quat. Sci. Rev.* 10, 297–317.

862 Berx, B., Hansen, B., Østerhus, S., Larsen, K. M., Sherwin, T., Jochumsen, K., 2013.
863 Combining in-situ measurements and altimetry to estimate volume, heat and salt
864 transport variability through the Faroe Shetland Channel. *Ocean Science Discussions*,
865 10.

866 Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of radiocarbon sequences.
867 *Quat. Geochronol.* 5, 512–518.

868 Bond, G.W., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S.,
869 Jantschik, R., Clasen, S., Simet, C., Tedesco, K., Klas, M., Bonani, G., Ivy, S., 1992.
870 Evidence for massive discharges of icebergs into the North Atlantic Ocean during the
871 last glacial period. *Nature* 360, 245–249.

872 Bond, G., Broecker, W., Johnsen, S., McManus, J., Labeyrie, L., Jouzel, J., Bonani, G., 1993.
873 Correlations between climate records from North Atlantic sediments and Greenland
874 ice. *Nature* 365, 143–147.

875 Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen,
876 H., Hajdas, I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic
877 Holocene and glacial climates. *Science* 278, 1257–1266.

878 Bond, G., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G., Johnson, S.,
879 1999. The north Atlantic's 1-2 kyr climate rhythm: relation to Heinrich events,
880 Dansgaard/Oeschger cycles and the little ice age. In: Clark, P.U., Webb, R.S.,
881 Keigwin, L.D. (Eds.), *Mechanisms of Global Climate Change at Millennial Time*
882 *Scales*. American Geophysical Union, Washington, D.C, 35–58.

883 Boulton, G.S., Dongelmans, P., Punkari, M., Broadgate, M., 2001. Paleoglaciology of an ice
884 sheet through a glacial cycle: the European ice sheet through the Weichselian. *Quat.*
885 *Sci. Rev.* 20, 591–625.

886 Bourillet, J.-F., Reynaud, J.-Y., Baltzer, A., Zaragosi, S., 2003. The 'Fleuve Manche': the
887 submarine sedimentary features from the outer shelf to the deep-sea fans. *J. Quat. Sci.*
888 18, 261–282.

889 Broecker, W., Bond, G., Klas, M., Clark, E., McManus, J., 1992. Origin of the northern
890 Atlantic's Heinrich events. *Clim. Dyn.* 6, 265–273.

891 Broecker, W.S., 1994. Massive iceberg discharges as triggers for global climate change.
892 *Nature* 372, 421–424.

893 Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Lecavalier, B. S., Kindler, P., et al., 2014.
894 Greenland temperature response to climate forcing during the last deglaciation.
895 *Science* 345, 1177–1180.

896 Buizert, C., Keisling, B. A., Box, J. E., He, F., Carlson, A. E., Sinclair, G., De Conto, R. M.,
897 2018. Greenland-wide seasonal temperatures during the last deglaciation. *Geophys.*
898 *Res. Lett.* 45, 1905–1914.

899 Cacho, I., Grimalt, J. O., Pelejero, C., Canals, M., Sierro, F. J., Flores, J. A., Shackleton, N.,
900 1999. Dansgaard-Oeschger and Heinrich event imprints in Alboran Sea
901 paleotemperatures. *Paleoceanography* 14, 698–705.

902 Cacho, I., Shackleton, N., Elderfield, H., Sierro, F. J., Grimalt, J. O., 2006. Glacial rapid
903 variability in deep-water temperature and $\delta^{18}\text{O}$ from the Western Mediterranean Sea.
904 *Quat. Sci. Rev.* 25, 3294–3311.

905 Camuera, J., Jimenez-Moreno, G., Ramos-Román, M.J., García-Alix, A., Toney, J.L.,
906 Anderson, R.S., Jimenez-Espejo, F., Bright, J., Webster, C., Yanes, Y., 2019.
907 Vegetation and climate changes during the last two glacial-interglacial cycles in the
908 western Mediterranean: a new long pollen record from Padul (southern Iberian
909 Peninsula). *Quat. Sci. Rev.* 205, 86–105.

910 Camuera, J., Jiménez-Moreno, G., Ramos-Román, M. J., García-Alix, A., Jiménez-Espejo, F.
911 J., Toney, J. L., Anderson, R. S., 2021. Chronological control and centennial-scale

912 climatic subdivisions of the Last Glacial Termination in the western Mediterranean
 913 region. *Quat. Sci. Rev.* 255, 106814.
 914 Català, A., Cacho Lascorz, I., Frigola Ferrer, J. I., Pena, L. D., Lirer, F., 2019. Holocene
 915 hydrography evolution in the Alboran Sea: A multi-record and multiproxy
 916 comparison. *Clim. Past.* 15, 927–942.
 917 Caille, C., Penaud, A., Eynaud, F., Zaragosi, S., Roche, D. M., Michel, E., Boulay, S.,
 918 Richter, T., 2013. Sea-surface hydrographical conditions off South Faeroes and within
 919 the North-Eastern North Atlantic through MIS 2: The response of dinocysts. *J. Quat.*
 920 *Sci.* 28, 217–228.
 921 Cayre, O., Lancelot, Y., Vincent, E., 1999. Paleoceanographic reconstructions from
 922 planktonic foraminifera off the Iberian Margin: temperature, salinity, and Heinrich
 923 events. *Paleoceanography* 14, 384–396.
 924 Clark, P.U., McCabe, A.M., Mix, A.C., et al. 2004. Rapid rise of sea level 19,000 years ago
 925 and its global implications. *Science* 304, 1141–1144.
 926 Clark, P. U., Shakun, J. D., Baker, P. A., et al., 2012a. Global climate evolution during the
 927 last deglaciation. *Proc. Natl. Acad. Sci.* 109, 1134–1142.
 928 Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Jordan, C., Sejrup, H.S., 2012b. Pattern and
 929 timing of retreat of the last British-Irish Ice Sheet. *Quat. Sci. Rev.* 44, 112–146.
 930 Combourieu Nebout, N., Londeix, L., Baudin, F., Turon, J. L., Von Grafenstein, R., 1999.
 931 Quaternary marine and continental paleoenvironments in the western Mediterranean
 932 (Site 976, Alboran Sea): palynological evidence. In *Proceedings of the Ocean Drilling*
 933 *Program. Scientific Results* 161, 457–468.
 934 Combourieu-Nebout N., Turon, J. L., Zahn, R., Capotondi, L., Londeix, L., Pahnke, K., 2002.
 935 Enhanced aridity and atmospheric high-pressure stability over the western
 936 Mediterranean during the North Atlantic cold events of the past 50 ky. *Geology* 30,
 937 863–866.
 938 Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U.,
 939 Marret, F., 2009. Rapid climatic variability in the west Mediterranean during the last
 940 25 000 years from high resolution pollen data. *Clim. Past.* 5, 503–521.
 941 Conkright, M. E., Locarnini, R. A., Garcia, H. E., O'Brien, T. D., Boyer, T. P., Stephens, C.,
 942 Antonov, J. I., 2002. *World Ocean Atlas 2001: Objective analyses, data statistics, and*
 943 *figures: CD-ROM documentation.*

944 Cossa, D., Cotté-Krief, M. H., Mason, R. P., Bretaudeau-Sanjuan, J. J. M. C., 2004. Total
 945 mercury in the water column near the shelf edge of the European continental margin.
 946 *Marine Chemistry* 90, 21–29.

947 Daniault, N., Mercier, H., Lherminier, P., Sarafanov, A., Falina, A., Zunino, P., et al., 2016.
 948 The northern North Atlantic Ocean mean circulation in the early 21st century.
 949 *Progress in Oceanography* 146, 142–158.

950 Dale, B., 1996. Dinoflagellate cyst ecology: modeling and geological applications. In: J.
 951 Jansonius and D.C. McGregor (Editors), *Palynology: principles and applications*, Vol.
 952 3. AASP Foundation, Salt Lake City, 1249–1275.

953 Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer,
 954 C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjörnsdottir, A. E., Jouzel, J., Bond, G.,
 955 1993. Evidence for general instability of past climate from a 250-kyr ice-core record.
 956 *Nature* 364, 218–220.

957 de Vernal, A., Harland, R., 1992. Quaternary organic-walled dinoflagellate cysts of the North
 958 Atlantic Ocean and adjacent seas: ecostratigraphy and biostratigraphy. In “Neogene
 959 and Quaternary Dinoflagellate Cyst of the North Atlantic Ocean and Adjacent Seas:
 960 Ecostratigraphy and Biostratigraphy” (M. J. Head and J. H. Wrenn, Eds.), pp. 289–
 961 328. American Association of Stratigraphic Palynologists Foundation, Dallas.

962 de Vernal, A., Henry, M., Bilodeau, G., 1999. Technique de préparation et d’analyse en
 963 micropaléontologie. *Les Cahiers du GEOTOP* vol. 3, Université du Québec à
 964 Montréal, Montréal, Canada.

965 de Vernal, A., Hillaire-Marcel, C., Turon, J.-L., Matthiessen, J., 2000. Reconstruction of sea-
 966 surface temperature, salinity, and sea-ice cover in the northern North Atlantic during
 967 the last glacial maximum based on dinocyst assemblages. *Can. J. Earth Sci.* 37, 725–
 968 750.

969 de Vernal, A., Hillaire-Marcel, C., Peltier, W.R., Weaver, A.J., 2002. The structure of the
 970 upper water column in the northwest North Atlantic: modern vs. Last Glacial
 971 Maximum conditions. *Paleoceanography* 17, 1050.

972 de Vernal, A., Eynaud, F., Henry, M., Hillaire-Marcel, C., Londeix, L., Mangin, S.,
 973 Matthiessen, J., Marret, F., Radi, T., Rochon, A., Solignac, S., Turon, J.-L., 2005.
 974 Reconstruction of sea-surface conditions at middle to high latitudes of the Northern
 975 Hemisphere during the Last Glacial Maximum (LGM) based on dinoflagellate cyst
 976 assemblages. *Quat. Sci. Rev.* 24, 897–924.

977 de Vernal, A., Rosell-Melé, A., Kucera, M., Hillaire-Marcel, C., Eynaud, F., Weinelt, M.,
978 Dokken, T., Kageyama, M., 2006. Comparing proxies for the reconstruction of LGM
979 sea-surface conditions in the northern North Atlantic. *Quat. Sci. Rev.* 25, 2820–2834.

980 de Vernal, A., Radi, T., Zaragosi, S., Van Nieuwenhove, N., Rochon, A., Allan, E., et al.,
981 2020. Distribution of common modern dinoflagellate cyst taxa in surface sediments of
982 the Northern Hemisphere in relation to environmental parameters: The new n= 1968
983 database. *Mar. Micropaleontol.* 101796.

984 Denton, G.H., Alley, R.B., Comer, G.C., Broecker, W.S., 2005. The role of seasonality in
985 abrupt climate change. *Quat. Sci. Rev.* 24, 1159–1182.

986 Denton, G.H., Anderson, R.F., Toggweiler, J.R., Edwards, R.L., Schaefer, J.M., Putnam,
987 A.E., 2010. The last glacial termination. *Science* 328, 1652–1656.

988 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., et al., 2012.
989 Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago. *Nature*
990 483, 559–564.

991 Ehlers, J., Gibbard, P.L., Hughes, P.D., 2011. *Quaternary Glaciations: Extent and*
992 *Chronology*, vol. 15. Elsevier, Amsterdam, pp. 1126.

993 Eynaud, F., 1999. *Kystes de dinoflagellates et evolution paléoclimatique et paléohydrologique*
994 *de l'Atlantique Nord au cours du Dernier Cycle Climatique du Quaternaire*, Ph.D.
995 thesis, 291 pp. Univ. of Bordeaux 1, Bordeaux, France.

996 Eynaud, F., Turon, J. L., Duprat, J., 2004. Comparison of the Holocene and Eemian
997 palaeoenvironments in the South Icelandic Basin: dinoflagellate cysts as proxies for
998 the North Atlantic surface circulation. *Rev. Palaeobot. Palynol.* 128, 55–79.

999 Eynaud, F., Zaragosi, S., Scourse, J.D., Mojtahid, M., Bourillet, J.F., Hall, I.R., Penaud, A.,
1000 Locascio, M., Reijonen, A., 2007. Deglacial laminated facies on the NW European
1001 continental margin: the hydrographic significance of British Ice sheet deglaciation and
1002 Fleuve Manche paleoriver discharges. *Geochem. Geophys. Geosyst.* 8, Q06019.

1003 Eynaud, F., De Abreu, L., Voelker, A., Schönfeld, J., Salgueiro, E., Turon, J. L., et al., 2009.
1004 Position of the Polar Front along the western Iberian margin during key cold episodes
1005 of the last 45 ka. *Geochem. Geophys. Geosyst.* 10, Q07U05.

1006 Eynaud, F., Malaizé, B., Zaragosi, S., de Vernal, A., Scourse, J., Pujol, C., et al., 2012. New
1007 constraints on European glacial freshwater releases to the North Atlantic Ocean.
1008 *Geophys. Res. Lett.* 39.

1009 Eynaud, F., Londeix, L., Penaud, A., Sanchez-Goni, M. F., Oliveira, D., Desprat, S., Turon, J.
1010 L., 2016. Dinoflagellate cyst population evolution throughout past interglacials: Key

1011 features along the Iberian margin and insights from the new IODP Site U1385 (Exp
1012 339). *Glob. Planet. Chang.* 136, 52–64.

1013 Fensome, R. A., Williams, G. L., 2004. The Lentin and Williams index of fossil
1014 dinoflagellates, 2004 edition. In: AASP Foundation Contributions Series 42, pp. 909.

1015 Fletcher, W.J., Sanchez Goñi, M.F., 2008. Orbital- and sub-orbital-scale climate impacts on
1016 vegetation of the western Mediterranean basin over the last 48,000 yr. *Quat. Res.* 70,
1017 451–464.

1018 Ganne A., Leroyer C., Penaud A., Mojtahid M., 2016. Present-day palynomorph deposits in
1019 an estuarine context: the case of the Loire Estuary. *Journal of Sea Research* 118, 35–
1020 51.

1021 Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K.,
1022 Chkir, N., Hellstrom, J., Wainer, K., Bourges, F., 2006. Timing and dynamics of the
1023 last deglaciation from European and North African $\delta^{13}\text{C}$ stalagmite profiles of
1024 comparison with Chinese and South Hemisphere stalagmites. *Quat. Sci. Rev.* 25,
1025 2118–2142.

1026 Gibbard, P. L., 1988. The history of the great northwest European rivers during the past three
1027 million years. *Philosophical Transactions of the Royal Society of London. B,*
1028 *Biological Sciences* 318, 559–602.

1029 Grousset, F.E., Pujol, C., Labeyrie, L., Auffret, G., Boelaert, A., 2000. Were the North
1030 Atlantic Heinrich events triggered by the behaviour of the European ice sheets?
1031 *Geology* 28, 123–126.

1032 Grousset, F.E., Cortijo, E., Herve, L., Richter, T., Burdloff, D., Duprat, J., Weber, O., 2001.
1033 Zooming in on Heinrich layers. *Paleoceanography* 16, 240–259.

1034 Guiot, J., de Vernal, A., 2007. Transfer functions: methods for quantitative paleoceanography
1035 based on microfossils. In: Hillaire-Marcel (Ed.), *Proxies in Late Cenozoic*
1036 *Paleoceanography. Developments in Marine Geology* 1. Elsevier, pp. 523–563.

1037 Hall, I.R., Moran, S.B., Zahn, R., Knutz, P.C., Shen, C.-C., Edwards, R.L., 2006. Accelerated
1038 drawdown of meridional overturning in the late-glacial Atlantic triggered by transient
1039 pre-H event freshwater perturbation. *Geophys. Res. Lett.* 33, L16616.

1040 Hammer, Ø., Harper, D. A., Ryan, P. D., 2001. PAST: Paleontological statistics software
1041 package for education and data analysis. *Palaeontologia electronica* 4, 9.

1042 Harland, R., 1983. Distribution maps of Recent dinoflagellate cysts in bottom sediments from
1043 the North Atlantic Ocean and adjacent seas. *Palaeontology* 26, 321–387.

1044 Harper, D. A., 1999. Numerical palaeobiology: computer-based modelling and analysis of
1045 fossils and their distributions. John Wiley & Sons Inc.

1046 Harrison, S. P., Goñi, M. S., 2010. Global patterns of vegetation response to millennial-scale
1047 variability and rapid climate change during the last glacial period. *Quaternary Science*
1048 *Reviews* 29, 2957–2980.

1049 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast Atlantic
1050 Ocean during the past 130,000 years. *Quat. Res.* 29, 142–152.

1051 Hemming, S.R., Bond, G.C., Broecker, W.S., Sharp, W.D., Klas-Mendelson, M., 2000.
1052 Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ ages of individual hornblende grains for varying Laurentide
1053 sources of iceberg discharges 22,000 to 10,500 yr B.P. *Quat. Res.* 54, 372–383.

1054 Hemming, S.R., 2004. Heinrich events: massive late Pleistocene detritus layers of the North
1055 Atlantic and their global climate imprint. *Rev. Geophys.* 42, RG1005, 1–43.

1056 Hodell, D., Crowhurst, S., Skinner, L., Tzedakis, P. C., Margari, V., Channell, J. E., et al.,
1057 2013. Response of Iberian Margin sediments to orbital and suborbital forcing over the
1058 past 420 ka. *Paleoceanography* 28, 185–199.

1059 Hodell, D. A., Nicholl, J. A., Bontognali, T. R., Danino, S., Dorador, J., Dowdeswell, J. A., et
1060 al., 2017. Anatomy of Heinrich Layer 1 and its role in the last deglaciation.
1061 *Paleoceanography* 32, 284–303.

1062 Ivanovic, R. F., Gregoire, L. J., Burke, A., Wickert, A. D., Valdes, P. J., Ng, H. C., et al.,
1063 2018. Acceleration of northern ice sheet melt induces AMOC slowdown and northern
1064 cooling in simulations of the early last deglaciation. *Paleoceanography and*
1065 *Paleoclimatology* 33, 807–824.

1066 Jalut, G., Turu i Michels, V., Dedoubat, J.J., Otto, T., Ezquerro, J., Fontugne, M., Belet, J.M.,
1067 Bonnet, L., García de Celis, A., Redondo-Vega, J.M., Vidal-Romaní, J.R., Santos, L.,
1068 2010. Palaeoenvironmental studies in NW Iberia (Cantabrian range): vegetation
1069 history and synthetic approach of the last deglaciation phases in the western
1070 Mediterranean. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 297, 330–350.

1071 Kaiser, J., 2001. Caractérisation palynologique des flux terrigènes Manche-Golfe de
1072 Gascogne au cours du Dernier Maximum Glaciaire et du réchauffement Holocène.
1073 Maîtrise des Sciences de l'Environnement, Univ. Bordeaux 1, Bordeaux, France, 30
1074 pp.

1075 Kempama, E.W., Reimnitz, E., Barnes, P.W., 2001. Anchor-ice formation and ice rafting in
1076 southwestern Lake Michigan, U.S.A. *Journal of Sedimentary Research* 71, 346–354.

1077 Knutz, P.C., Austin, W.E.N., Jones, E.J.W., 2001. Millennial-scale depositional cycles related
1078 to British Ice Sheet variability and North Atlantic paleocirculation since 45 yr B. P.,
1079 Barra Fan, U. K. margin. *Paleoceanography* 16, 53–64.

1080 Knutz, P.C., Zahn, R., Hall, I.R., 2007. Centennial-scale variability of the British Ice Sheet:
1081 implications for climate forcing and Atlantic meridional overturning circulation during
1082 the last deglaciation. *Paleoceanography* 22, PA1207.

1083 Kucera, M., Weinelt, M., Kiefer, T., Pflaumann, U., Hayes, A., Weinelt, M., et al., 2005.
1084 Reconstruction of sea-surface temperatures from assemblages of planktonic
1085 foraminifera: multi-technique approach based on geographically constrained
1086 calibration data sets and its application to glacial Atlantic and Pacific Oceans. *Quat.*
1087 *Sci. Rev.* 24, 951–998.

1088 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., Sambridge, M., 2014. Sea level and global ice
1089 volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National*
1090 *Academy of Sciences* 111, 15296–15303.

1091 Lambert C., Vidal M., Penaud A., Combourieu-Nebout N., Lebreton V., Ragueneau O.,
1092 Gregoire G., 2017. Modern palynological record in the Bay of Brest (NW France):
1093 Signal calibration for palaeo-reconstructions. *Rev. Palaeobot. Palynol.* 244, 13–25.

1094 Lewis, J., Dodge, J. D., Powell, A. J., 1990. Quaternary dinoflagellate cysts from the
1095 upwelling system off shore Peru, Hole 686B, ODP leg 112. *Proceedings of ODP*
1096 *Scientific Results*, 112, 323–327.

1097 Lézine, A.M., Duplessy, J.C., Cazet, J.P., 2005. West African monsoon variability during the
1098 last deglaciation and the Holocene: evidence from fresh water algae, pollen and
1099 isotope data from core KW31, Gulf of Guinea. *Palaeogeogr. Palaeoclimatol.*
1100 *Palaeoecol* 219, 225–237.

1101 Marcott, S. A., Clark, P. U., Padman, L., Klinkhammer, G. P., Springer, S. R., Liu, Z., et al.,
1102 2011. Ice-shelf collapse from subsurface warming as a trigger for Heinrich events.
1103 *Proceedings of the National Academy of Sciences* 108, 13415–13419.

1104 Margalef, R., 1958. Temporal succession and spatial heterogeneity in phytoplankton.
1105 *Perspectives in marine biology* 323–349.

1106 Marret, F., 1994. Distribution of dinoflagellate cysts in recent marine sediments from the east
1107 Equatorial Atlantic (Gulf of Guinea). *Rev. Palaeobot. Palynol.* 84, 1–22.

1108 Marret, F., Zonneveld, K.A.F., 2003. Atlas of modern organic-walled dinoflagellate cyst
1109 distribution. *Rev. Palaeobot. Palynol.* 125, 1–200.

1110 Marret, F., Scourse, J., Austin, W., 2004. Holocene shelf-sea seasonal stratification dynamics:
 1111 A dinoflagellate cyst record from the Celtic Sea, NW European shelf. *The Holocene*
 1112 14, 689-696.

1113 Marret, F., Bradley, L., de Vernal, A., Hardy, W., Kim, S. Y., Mudie, P., et al., 2020. From
 1114 bi-polar to regional distribution of modern dinoflagellate cysts, an overview of their
 1115 biogeography. *Marine Micropaleontology* 159, 101753.

1116 Martrat, B., Grimalt, J. O., Shackleton, N. J., de Abreu, L., Hutterli, M. A., Stocker, T. F.,
 1117 2007. Four climate cycles of recurring deep and surface water destabilizations on the
 1118 Iberian margin. *Science* 317, 502–507.

1119 Martrat, B., Jimenez-Amat, P., Zahn, R., Grimalt, J. O., 2014. Similarities and dissimilarities
 1120 between the last two deglaciations and interglaciations in the North Atlantic region.
 1121 *Quat. Sci. Rev.* 99, 122–134.

1122 Matthiessen, J., 1995. Distribution patterns of dinoflagellate cysts and other organic-walled
 1123 microfossils in recent Norwegian-Greenland Sea sediments. *Mar. Micropaleontol.* 24,
 1124 307–334.

1125 McManus, J.F., Francois, R., Gherardi, J.M., Keigwin, L.D., Drown-Leger, S., 2004. Collapse
 1126 and rapid resumption of Atlantic meridional circulation linked to deglacial climate
 1127 changes. *Nature* 428, 834–837.

1128 Ménot, G., Bard, E., Rostek, R., Weijers, W.H., Hopmans, E.C., Schouten, S., Sinninghe
 1129 Damsté, J.S., 2006. Early reactivation of European rivers during the last deglaciation.
 1130 *Science* 313, 1623–1625.

1131 Missiaen, L., Waelbroeck, C., Pichat, S., Jaccard, S. L., Eynaud, F., Greenop, R., Burke, A.
 1132 2019. Improving North Atlantic marine core chronologies using ^{230}Th normalization.
 1133 *Paleoceanography and paleoclimatology* 34, 1057–1073.

1134 Mix, A.C., Bard, E., Schneider, R., 2001. Environmental processes of the ice age: land,
 1135 oceans, glaciers (EPILOG). *Quat. Sci. Rev.* 20 (4), 627—657.

1136 Mojtahid, M., Eynaud, F., Zaragosi, S., Scourse, J., Bourillet, J.F., Garlan, T., 2005.
 1137 Palaeoclimatology and palaeohydrography of the glacial stages on Celtic and
 1138 Armorican margins over the last 360 000 yrs. *Marine Geology* 224, 57–82.

1139 Mojtahid, M., Toucanne, S., Fentimen, R., Barras, C., Le Houedec, S., Soulet, G., et al., 2017.
 1140 Changes in northeast Atlantic hydrology during Termination 1: Insights from Celtic
 1141 margin's benthic foraminifera. *Quaternary Science Reviews* 175, 45–59.

1142 Morellón, M., Valero-Garcés, B., Vegas-Vilarrúbia, T., González-Sampériz, P., Romero, Ó.,
 1143 et al., 2009. Late glacial and Holocene palaeohydrology in the western Mediterranean
 1144 region: The Lake Estanya record (NE Spain). *Quat. Sci. Rev.* 28, 2582–2599.

1145 Moreno, A., Stoll, H.M., Jimenez-Sanchez, M., Cacho, I., Valero-Garcés, B., Ito, E.,
 1146 Edwards, L.R., 2010. A speleothem record of rapid climatic shifts during last glacial
 1147 period from Northern Iberian Peninsula. *Glob. Planet. Chang.* 71, 218–231

1148 Morzadec-Kerfourn, M.-T., 1977. Les kystes de Dinoflagellés dans les sédiments récents le
 1149 long des côtes bretonnes. *Rev. Micropaleontol.* 20, 157–166.

1150 National Oceanographic Data Centre (NODC), 2001. World Ocean Atlas.
 1151 http://www.nodc.noaa.gov/OC5/WOD01/pr_wod01.html.

1152 Naughton, F., Sanchez Goñi, M.F., Desprat, S., Turon, J.-L., Duprat, J., Malaizé, B., Joli, C.,
 1153 Cortijo, E., Drago, T., Freitas, M.C., 2007. Present-day and past (last 25 000 years)
 1154 marine pollen signal off western Iberia. *Mar. Micropaleontol.* 62, 91–114.

1155 Naughton, F., Sanchez Goñi, M.F., Kageyama, M., Bard, E., Cortijo, E., Desprat, S., Duprat,
 1156 J., Malaizé, B., Joli, C., Rostek, F., Turon, J.-L., 2009. Wet to dry climatic trend in
 1157 north western Iberia within Heinrich events. *Earth Planet. Sci. Lett.* 284, 329–342.

1158 Naughton, F., Goñi, M. S., Rodrigues, T., Salgueiro, E., Costas, S., Desprat, S., et al., 2016.
 1159 Climate variability across the last deglaciation in NW Iberia and its margin.
 1160 *Quaternary International* 414, 9–22.

1161 Ng, H. C., Robinson, L. F., McManus, J. F., Mohamed, K. J., Jacobel, A. W., Ivanovic, R. F.,
 1162 Gregoire, L.J., Chen, T., 2018. Coherent deglacial changes in western Atlantic Ocean
 1163 circulation. *Nature communications* 9, 1–10.

1164 North Greenland Ice Core Project, m., 2004. High-resolution record of Northern Hemisphere
 1165 climate extending into the last interglacial period. *Nature* 431, 147–151.

1166 Nygård, A., Sejrup, H.P., Hafliðason, H., Leksens, W.A.H., Clark, C.D., Bigg, G.R., 2007.
 1167 Extreme sediment and ice discharge from marine-based ice streams: new evidence
 1168 from the North Sea. *Geology* 35, 395–398.

1169 Peck, V.L., Hall, I.R., Zahn, R., Elderfield, H., Grousset, F., Hemming, S.R., Scourse, J.D.,
 1170 2006. High resolution evidence for linkages between NW European ice sheet
 1171 instability and Atlantic Meridional Overturning Circulation. *Earth Planet. Sci. Lett.*
 1172 243, 476–488.

1173 Penaud, A., Eynaud, F., Turon, J.L., Zaragosi, S., Marret, F., Bourillet, J.F., 2008. Interglacial
 1174 variability (MIS 5 and MIS 7) and dinoflagellate cyst assemblages in the Bay of
 1175 Biscay (North Atlantic). *Mar. Micropaleontol.* 68, 136–155.

1176 Penaud, A., Eynaud, F., Turon, J. L., Zaragosi, S., Malaizé, B., Toucanne, S., Bourillet, J. F.
1177 2009. What forced the collapse of European ice sheets during the last two glacial
1178 periods (150 ka BP and 18 ka cal BP)? Palynological evidence. *Palaeogeogr.*
1179 *Palaeoclimatol. Palaeoecol.* 281, 66–78.

1180 Penaud, A., Eynaud, F., Voelker, L., Helga, A., Turon, J. L., 2016. Palaeohydrological
1181 changes over the last 50 ky in the central Gulf of Cadiz: complex forcing mechanisms
1182 mixing multi-scale processes. *Biogeosciences* 13, 5357–5377.

1183 Penaud, A., Ganne, A., Eynaud, F., Lambert, C., Coste, P. O., Herlédan, M., et al., 2020.
1184 Oceanic *versus* continental influences over the last 7 kyrs from a mid-shelf record in
1185 the northern Bay of Biscay (NE Atlantic). *Quat. Sci. Rev.* 229, 106135.

1186 R Development Core Team., 2008. R: A language and environment for statistical computing.
1187 R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0.

1188 Radi, T., Bonnet, S., Cormier, M. A., de Vernal, A., Durantou, L., Faubert, É., et al., 2013.
1189 Operational taxonomy and (paleo-) autecology of round, brown, spiny dinoflagellate
1190 cysts from the Quaternary of high northern latitudes. *Mar. Micropaleontol.* 98, 41–57.

1191 Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B.,
1192 Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., et al., 2014. A stratigraphic framework
1193 for abrupt climatic changes during the Last Glacial period based on three synchronized
1194 Greenland ice-core records: Refining and extending the INTIMATE event
1195 stratigraphy, *Quat. Sci. Rev.* 106, 14–28.

1196 Reimnitz, E., Kempama, E.W., 1987. Field observations of slush-ice generated during freeze-
1197 up in Arctic coastal waters. *Mar. Geol.* 77, 219–231.

1198 Rochon, A., Vernal, A.d., Turon, J.-L., Matthießen, J., Head, M., 1999. Distribution of recent
1199 dinoflagellate cysts in surface sediments from the North Atlantic Ocean and adjacent
1200 seas in relation to sea-surface parameters. *American Association of Stratigraphic*
1201 *Palynologists Contribution Series* 35, 1–146.

1202 Rosell-Melé, A., Comes, P., 1999. Evidence for a Warm Last Glacial Maximum in the Nordic
1203 Seas or an example of shortcomings in UK 37' and UK 37 to estimate low sea surface
1204 temperature? *Paleoceanography* 14, 770–776.

1205 Salgueiro, E., Naughton, F., Voelker, A.H.L., de Abreu, L., Alberto, A., Rossignol, L.,
1206 Duprat, J., Magalhães, V.H., Vaqueiro, S., Turon, J.-L., Abrantes, F., 2014. Past
1207 circulation along the western Iberian margin: a time slice vision from the Last Glacial
1208 to the Holocene. *Quat. Sci. Rev.* 106, 316–329.

1209 Stanford, J.D., Rohling, E.J., Hunter, S.E., Roberts, A.P., Rasmussen, S.O., Bard, E.,
 1210 McManus, J., Fairbanks, R.G., 2006. Timing of mwp-1a and climate responses to
 1211 meltwater injections. *Paleoceanography* 21, PA4103.
 1212 Stanford, J. D., Rohling, E. J., Bacon, S., Roberts, A. P., Grousset, F. E., Bolshaw, M., 2011.
 1213 A new concept for the paleoceanographic evolution of Heinrich event 1 in the North
 1214 Atlantic. *Quat. Sci. Rev.* 30, 1047–1066.
 1215 Sutton, R., Allen, M., 1997. Decadal predictability of North Atlantic sea surface temperature
 1216 and climate. *Nature* 388, 563–567.
 1217 Svensson, A., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Davies, S.M.,
 1218 Johnsen, S.J., Muscheler, R., Rasmussen, S.O., Röthlisberger, R., Steffensen, J.P.,
 1219 Vinther, B.M., 2006. The Greenland ice core chronology 2005, 15–42 ka. Part 2:
 1220 comparison to other records. *Quat. Sci. Rev.* 25, 23–24.
 1221 Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl- Jensen, D., Davies, S. M., J
 1222 ohnson, S. J., Muscheler, R., Parrenin, F., Rasmussen, S. O., Röthlisberger, R.,
 1223 Seierstad, I., Steffensen, J. P., Vinther, B. M., 2008. A 60 000-year Greenland
 1224 stratigraphic ice core chronology. *Clim. Past.* 4, 47–57.
 1225 Toucanne, S., Zaragosi, S., Bourillet, J.F., Naughton, F., Cremer, M., Eynaud, F., Dennielou,
 1226 B., 2008. Activity of the turbidite levees of the Celtic-armoric margin (Bay of
 1227 Biscay) during the last 30,000 years: imprints of the last European deglaciation and
 1228 Heinrich events. *Mar. Geol.* 247, 84–103.
 1229 Toucanne, S., Zaragosi, S., Bourillet, J.F., Cremer, M., Eynaud, F., Turon, J.L., Fontanier, C.
 1230 Van Vliet Lanoë, B., Gibbard, P., 2009. Timing of massive ‘Fleuve Manche’
 1231 discharges over the last 350 kyr: insights into the European Ice Sheet oscillations and
 1232 the European drainage network from MIS 10 to 2. *Quat. Sci. Rev.* 28, 1238–1256.
 1233 Toucanne, S., Zaragosi, S., Bourillet, J. F., Marieu, V., Cremer, M., Kageyama, M., et al.,
 1234 2010. The first estimation of Fleuve Manche palaeoriver discharge during the last
 1235 deglaciation: Evidence for Fennoscandian ice sheet meltwater flow in the English
 1236 Channel ca 20–18 ka ago. *Earth Planet. Sci. Lett.* 290, 459–473.
 1237 Toucanne, S., Zaragosi, S., Bourillet, J. F., Dennielou, B., Jorry, S. J., Jouet, G., Cremer, M.
 1238 2012. External controls on turbidite sedimentation on the glacially-influenced
 1239 Armorican margin (Bay of Biscay, western European margin). *Mar. Geol.* 303, 137–
 1240 153.

1241 Toucanne, S., Soulet, G., Freslon, N., Jacinto, R. S., Dennielou, B., Zaragosi, S., et al., 2015.
1242 Millennial-scale fluctuations of the European Ice Sheet at the end of the last glacial,
1243 and their potential impact on global climate. *Quat. Sci. Rev.* 123, 113–133.

1244 Toucanne, S., Soulet, G., Vázquez Riveiros, N., Boswell, S. M., Dennielou, B., Waelbroeck,
1245 C., ... et al., 2021. The North Atlantic Glacial Eastern Boundary Current as a Key
1246 Driver for Ice-Sheet—AMOC Interactions and Climate Instability. *Paleoceanography*
1247 and *Paleoclimatology* 36, e2020PA004068.

1248 Turon, J. L., 1984. Le palynoplankton dans l'environnement actuel de l'Atlantique Nord-
1249 oriental. Evolution climatique et hydrologique depuis le dernier maximum glaciaire.
1250 Mémoires de l'Institut de Géologie du Bassin d'Aquitaine, 17, 313 pp.

1251 Turon, J.-L., Londeix, L., 1988. Dinoflagellate assemblages in the western Mediterranean,
1252 Alboran sea: evidence of the evolution of palaeoenvironments since the last glacial
1253 maximum [Les assemblages de kystes de Dinoflagelles en Méditerranée occidentale
1254 (Mer d'Alboran): mise en évidence de l'évolution des palaeoenvironnements depuis le
1255 dernier maximum glaciaire.]. *Bulletin – Centres de Recherche Exploration- Production*
1256 *Elf-Aquitaine* 12, 313–344.

1257 Turon, J.L., Lézine, A.M., Denèfle, M., 2003. Land-sea correlations for the last glaciation
1258 inferred from a pollen and dinocyst record from the Portuguese margin. *Quat. Res.* 59,
1259 88–96.

1260 Van Nieuwenhove, N., Pospelova, V., de Vernal, A., Rochon, A., 2020. A historical
1261 perspective on the development of the Northern Hemisphere modern dinoflagellate
1262 cyst database. 101824.

1263 Versteegh, G. J., Zonneveld, K. A., 1994. Determination of (palaeo-) ecological preferences of
1264 dinoflagellates by applying detrended and canonical correspondence analysis to Late
1265 Pliocene dinoflagellate cyst assemblages of the south Italian Singa section. *Rev.*
1266 *Palaeobot. Palynol.* 84, 181–199.

1267 Voelker, A. H., 2002. Global distribution of centennial-scale records for Marine Isotope Stage
1268 (MIS) 3: a database. *Quat. Sci. Rev.* 21, 1185–1212.

1269 Waelbroeck, C., Lougheed, B. C., Riveiros, N. V., Missiaen, L., Pedro, J., Dokken, T., et al.,
1270 2019. Consistently dated Atlantic sediment cores over the last 40 thousand years.
1271 *Scientific data* 6, 1–12.

1272 Wall, D., Dale, B., Lohmann, G.P., Smith, W.K., 1977. The environment and climatic
1273 distribution of dinoflagellate cysts in modern marine sediments from regions in the
1274 north and south Atlantic oceans and adjacent seas. *Mar. Micropaleontol.* 2, 121–200.

1275 Wary, M., Eynaud, F., Sabine, M., Zaragosi, S., Rossignol, L., Malaize, B., et al., 2015.
1276 Stratification of surface waters during the last glacial millennial climatic events: a key
1277 factor in subsurface and deep-water mass dynamics. *Clim. Past.* 11, 1507.

1278 Weaver, A.J., Saenko, O.A., Clark, P.U., Mitrovica, J.X., 2003. Meltwater pulse 1A from
1279 Antarctica as a trigger of the Bølling-Allerød warm interval. *Science* 299, 1709-1713.

1280 Weinelt, M., Sarnthein, M., Pflaumann, U., Schulz, H., Jung, S., Erlenkeuser, H., 1996. Ice-
1281 free Nordic Seas during the Last Glacial Maximum? Potential sites of deepwater
1282 formation. *Paleoclimates* 1, 283–309.

1283 Woerther, P., 2013. VT 133 /MERIADZEK cruise, RV Marion Dufresne.

1284 Zaragosi, S., Auffret, G. A., Faugères, J. C., Garlan, T., Pujol, C., Cortijo, E., 2000.
1285 Physiography and recent sediment distribution of the Celtic Deep-Sea Fan, Bay of
1286 Biscay. *Mar. Geol.* 169, 207–237.

1287 Zaragosi, S., Eynaud, F., Pujol, C., Auffret, G., Turon, J.-L., Garlan, T., 2001. Initiation of the
1288 European deglaciation as recorded in the northwestern Bay of Biscay slope
1289 environments (Meriadzek Terrace and Trevelyan Escarpment): a multi-proxy
1290 approach. *Earth Planet. Sci. Lett.* 188, 493–507.

1291 Zaragosi, S., Bourillet, J.F., Eynaud, F., Toucanne, S., Denhard, B., Van Toer, A., Lanfumeu,
1292 V., 2006. The impact of the last European deglaciation on the deepsea turbidite
1293 systems of the Celtic-Armorican margin (Bay of Biscay). *Geo-Marine Letters* 26,
1294 317–329.

1295 Zonneveld, K.A.F., 1997. New species of organic walled dinoflagellate cysts from modern
1296 sediments of the Arabian Sea (Indian Ocean). *Rev. Palaeobot. Palynol.* 97, 319–337

1297 Zonneveld, K.A.F., Hoek, R.P., Brinkhuis, H., Willems, H., 2001. Geographical distributions
1298 of organic-walled dinoflagellate cysts in surficial sediments of the Benguela upwelling
1299 region and their relationship to upper ocean conditions. *Progress in Oceanography* 48,
1300 25–72.

1301 Zonneveld, K. A., Versteegh, G., Kodrans-Nsiah, M., 2008. Preservation and organic
1302 chemistry of Late Cenozoic organic-walled dinoflagellate cysts: A review. *Marine*
1303 *Micropaleontology* 68, 179–197.

1304 Zonneveld, K.A.F., Marret, F., Versteegh, G.J.M., Bogus, K., Bonnet, S., Bouimetarhan, I., et
1305 al., 2013. Atlas of modern dinoflagellate cyst distribution based on 2405 data points
1306 *Rev. Palaeobot. Palynol.* 191, 1–197.

1307 Zumaque, J., Eynaud, F., de Vernal, A., 2017. Holocene paleoceanography of the Bay of
1308 Biscay: evidence for west-east linkages in the North Atlantic based on dinocyst data.
1309 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 468, 403–413.

Table 1

Core name	Geographic coordinates Depth (m)	Dinocyst assemblages	Ti/Ca-XRF ratio	% <i>N.pachyderma</i> s.	IRD conc (No g dry sed)	No Laminae/cm	BIT index + % COT
MD13-3438 Studied core	47° 27' N 8° 27' W 2180 m	This study	S. Toucanne in this study	L. Rossignol in this study	No data	No data	No data
MD95-2002 Reference core	47°27' N 8°32' W 2174 m	Eynaud, 1999 ; Zaragosi et al., 2001 ; Eynaud et al., 2012	Toucanne et al., 2015	Zaragosi et al., 2001 ; Eynaud et al., 2009	Zaragosi et al., 2001	Zaragosi et al., 2001	Ménot et al., 2006

Table 2

Tie points MD95-2002	Depth MD95-2002 (cm)	Equivalent depth MD13-3438 (cm)	Age cal BP (Toucanne et al.,2015)
Top core	0	0	1629
End YD	190	126	11488
Onset YD	262	170	12442
End HS1	390	242	14699
HS1-5	455	277	15908
HS1-4	515	314	16482
HS1-3(2)	548	324	16724
HS1-3	690	428	17388
HS1-2	745	475	17559
Onset HS1	865	567	18100
End LGM	940	619	18570
LGM-6	970	640	18739
LGM-5	1000	663	18904
LGM-4	1040	697	19127
LGM-3	1145	788	19807
LGM-2	1260	881	21077
LGM-1	1340	952	22788
Onset LGM	1355	960	23001
End HS2	1375	981	23185
HS2-4	1415	1011	23560
HS2-3	1460	1050	24249
HS2-2	1495	1084	25031
Onset HS2	1550	1123	26551
End GI-3	1573	1155	27186
Onset GI-3	1593	1179	27720
End HS3 / Onset GI-4	1640	1218	28953
mid HS3	1695	1287	30116
Onset HS3	1745	1339	31050
Onset GI-5	1793	1389	32419
Onset-GI-7	1858	1458	35694
End GI-8	1880	1482	36900
End HS4 / Onset GI-8	1906	1511	37908
mid-HS4	1968	1575	39300

Table 3

	Name of taxa	Abbreviation
Autotrophic taxa	<i>Bitectatodinium tepikiense</i>	Btep
	<i>Impagidinium aculeatum</i>	Iacu
	<i>Impagidinium pallidum</i>	Ipal
	<i>Impagidinium paradoxum</i>	Ipar
	<i>Impagidinium patulum</i>	Ipat
	<i>Impagidinium sphaericum</i>	Isph
	<i>Lingulodinium machaerophorum</i>	Lmac
	<i>Nematosphaeropsis labyrinthus</i>	Nlab
	<i>Operculodinium centrocarpum</i>	Ocen
	<i>Operculodinium janduchenei</i>	Ojan
	<i>Polysphaeridium zoharyi</i>	Pzoh
	<i>Spiniferites membranaceus</i>	Smem
	<i>Spiniferites delicatus</i>	Sdel
	<i>Spiniferites elongatus</i>	Selo
	<i>Spiniferites ramosus</i>	Sram
	<i>Spiniferites belerius</i>	Sbel
	<i>Spiniferites bentorii</i>	Sben
	<i>Spiniferites lazus</i>	Slaz
	<i>Spiniferites mirabilis</i>	Smir
	<i>Spiniferites</i> spp.	Sspp
	<i>Spiniferites septentrionalis</i>	Ssep
	Cyst of <i>Pentapharsodinium dalei</i>	Pdal
Heterotrophic taxa	<i>Islandinium minutum</i>	Imin
	<i>Brigantedinium</i> spp.	Bspp
	<i>Quinquecuspsis</i> spp.	Qspp
	<i>Lejeunecysta</i> spp.	Lspp
	<i>Dubridinium</i> spp.	Dspp
	<i>Votadinium</i> spp.	Vspp
	<i>Selenopemphix quanta</i>	Squa
	Cyst of <i>Protoperidinium nudum</i>	Pnud
	<i>Echinidinium</i> spp.	Espp

b)

MD95-2002 vs MD13-3438

MD95-2002

★ AMS ¹⁴C dates from core MD95-2002
 ★ AMS ¹⁴C dates from core MD03-2690
 ★ NGRIP tie-point

a)

b) Database n=1968 (de Vernal et al., 2020)

HS1-a: Laminated interval (18.2 - 16.7 ka BP)

Summer-prevailing mode (HS1-a2, HS1-a4)

Winter-prevailing mode (HS1-a1, HS1-a3, HS1-a5)

HS1-b: Early-HE1 (16.7- 15.6 ka BP)

HS1-c: Late-HE1 (15.6- 14.6 ka BP)