Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco: A latitudinal and biogeographic analysis

Soilam Boutoumit, Reda El Kamcha, Oussama Bououarour, Latifa Joulami, Loubna Boutahar, Abdelaziz Benhoussa, Mohamed Maanan, L. Godet, Abdellatif Bayed, Hocein Bazairi

► To cite this version:

Soilam Boutoumit, Reda El Kamcha, Oussama Bououarour, Latifa Joulami, Loubna Boutahar, et al.. Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco: A latitudinal and biogeographic analysis. *Regional Studies in Marine Science*, 2021, 44, pp.101689. 10.1016/j.rsma.2021.101689 . hal-03467914

HAL Id: hal-03467914

<https://hal.science/hal-03467914>

Submitted on 6 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco: A latitudinal and biogeographic analysis

Soilam Boutoumit, Reda El Kamcha, Oussama Bououraour,
Latifa Joulami, Loubna Boutahar, Abdelaziz Benhoussa,
Mohamed Maanan, Laurent Godet, Abdellatif Bayed, Hocein Bazairi

PII: S2352-4855(21)00081-5
DOI: <https://doi.org/10.1016/j.rsma.2021.101689>
Reference: RSMA 101689

To appear in: *Regional Studies in Marine Science*

Received date : 6 August 2020
Revised date : 13 November 2020
Accepted date : 20 February 2021

Please cite this article as: S. Boutoumit, R. El Kamcha, O. Bououraour et al., Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco: A latitudinal and biogeographic analysis. *Regional Studies in Marine Science* (2021), doi: <https://doi.org/10.1016/j.rsma.2021.101689>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Elsevier B.V. All rights reserved.

Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco: a latitudinal and biogeographic analysis

Soilam Boutoumit^{a, b}, Reda El Kamcha^a, Oussama Bououraour^a, Latifa Joulami^c, Loubna Boutahar^{a, d}, Abdelaziz Benhoussa^a, Mohamed Maanan^b, Laurent Godet^e, Abdellatif Bayed^f, Hocein Bazairi^{a*}

^a *BioBio Research Center, BioEcoGen Laboratory, Faculty of Sciences, Mohammed V University in Rabat, 4 Avenue Ibn Battouta, B.P. 1014 RP, 10106, Rabat, Morocco*

^b *Université de Nantes. UMR 6554 LETG-Nantes, BP 81227, F-44312 Nantes Cedex 03, France*

^c *Laboratoire d'Ecologie et Environnement, Faculté des Sciences Ben M'sik, B.P. 7955, Sidi Othman – Casablanca, Morocco*

^d *Laboratorio de Biología Marina, Departamento de Zoología, Universidad de Sevilla, Avda. Reina Mercedes 6, 41012, Sevilla, Spain*

^e *CNRS. UMR LETG-Nantes, Université de Nantes, BP 81227, F-44312 Nantes Cedex 03, France*

^f *Institut Scientifique, Unité de Recherche OCEMAR, Mohammed V University in Rabat, 4 Avenue Ibn Battouta, B.P. 703 RP, 10106, Rabat, Morocco*

***Corresponding author.** E-mail address: hocein.bazairi@um5.ac.ma (H.Bazairi)

ABSTRACT

Although soft-bottoms are the largest ecosystem on Earth in terms of area, only a small percentage of their macrobenthos has been studied and most of its species are not yet described. Herein, the most up-to-dated comprehensive inventory and the broad-scale baseline of the soft-sediment macrozoobenthos in semi-enclosed coastal systems (SECS) of the Moroccan Atlantic and Mediterranean coasts (3500 km) is presented. In total, 496 species (7 phyla, 21 classes, 65 orders and 201 families) were recorded among which 95 species were exclusively Mediterranean, 99 species were Atlantico-Mediterranean SECS and 302 species exclusively Atlantic.

The best multivariate model, explaining 33% of the total variation observed in benthic assemblages' composition, included the type of the SECS (estuaries vs lagoons vs bay), the marine ecoregion (Atlantic vs Mediterranean), the surface of the SECS and the environmental features (minimal temperature, minimal and maximal salinity) as predictors of benthic macrofauna composition in the Moroccan SECS. In contrast to the general latitudinal diversity gradient (LDG) pattern, our results showed that species richness and taxonomic diversity showed no relationship with latitude. Such differences in benthic macrofaunal composition across a large scale could result from the fact that each ecosystem has its own specific characteristics, which implies an individualistic approach to ecosystem ecology.

The sample of the 12 SECS considered in this study covers most of the range of variation along the coasts of Morocco. The current compilation is relevant in such poorly known area at the global scale and fulfills a knowledge gap on benthic macrofauna in SECS of the Southern part of the North-Eastern Atlantic and Mediterranean ecoregions. However, the knowledge gained here is insufficient to address perceived shortfalls in knowledge of biodiversity, its importance to ecosystem function, and the threats and consequences of disturbance by anthropogenic activities.

Key words: Benthic macrofauna; Checklist; Latitudinal diversity gradient; North-East Atlantic; Mediterranean.

1. Introduction

Ecologists have long been intrigued by global patterns in biodiversity (Piacenza et al., 2015) and the understanding of the distribution of life on earth is a main goal in ecology and biogeography (Gaston, 2000; Hillebrand, 2004). The latitudinal diversity gradient, hereafter LDG (Hawkins, 2001), with the highest numbers of species in the tropics and gradual decrease poleward, is the most famous large-scale biodiversity pattern (Hillebrand, 2004; Kinlock et al., 2018). Explanations for the LDG have been related to many potential mechanisms but no broad consensus on the causes of the LDG has emerged (Kinlock et al., 2018). The many explanations that have been proposed can be categorized broadly into ecological, evolutionary and historical processes (see Cruz-Motta et al., 2020 for recent summaries). Given the challenge of inferring process from pattern, disentangling these hypothesized drivers remains one of the great challenges in macroecology (Hurlbert & Stegen, 2014).

While the LDG is a well-recognized and long-established pattern in terrestrial ecology, the knowledge of global diversity patterns in marine ecosystems is limited to a small number of studies (Barboza & Defeo, 2015). For marine biota, the first suggestion of the latitudinal diversity cline was formulated in late 1950^{ies} for hard bottom epifauna (Thorson, 1957) with pronounced decrease in the species richness of hard substratum epifauna towards arctic areas, whereas the number of soft-sediment infauna species was roughly the same in tropical, temperate and arctic areas. Latitudinal clines in benthic diversity of shallow waters have been also reported for gastropods (Roy et al. 1998), bivalve molluscs (Crame 2000; Roy, Jablonski & Valentine 2000), in shallow and deep-seas benthos (Sanders, 1968; Poore & Wilson 1993; Rex et al. 1993), and for pelagic taxa (Angel 1997; Pierrot-Bults 1997). Nevertheless, many studies have shown a deviation from the general LDG for the shallow-water marine fauna (Kendall & Aschan 1993; Boucher & Lamshead 1995). Furthermore, in the southern hemisphere the evidence of a latitudinal gradient of decreasing richness from the tropics to Antarctica is less convincing than in the northern hemisphere (Clarke 1992; Poore & Wilson 1993; Crame 2000). Recently, Kinlock et al. (2018) revisited the challenge of synthesizing individual LDGs and indicated that the phenomenon is not ubiquitous among habitats of the marine realm. More precisely, they indicated that the phenomenon is non-significant in the benthic habitat. However, Menegotto et al. (2019), in their comment on Kinlock et al. (2018), suggest that the marine habitat categories used by them (i.e.,

benthic, coral reefs, coastal, open ocean) are not independent and that reclassifying the studies significantly alters one of their main results. By assigning the studies into benthic and pelagic categories, and additionally into coastal or oceanic zones, they show that non-ambiguous, evolutionarily meaningful marine habitats display a significant latitudinal decline in species richness. Thus, there is no convincing evidence of a latitudinal cline across all taxa in the sea comparable to that seen on land (Clarke 1992; Clarke & Crame 1997).

Meta-analysis represents a suitable technique to analyse latitudinal gradients across different biota and regions and thus to generalize or not findings on the latitudinal distribution of species richness (Hillbrand, 2004; Kinlock et al., 2019). Therefore, Open science and the accumulation of spatially explicit biodiversity data are crucial to document LDG patterns better and to evaluate hypotheses broadly or for particular groups of organisms (Kinlock et al., 2018; 2019).

Soft-bottom substrates cover most of the world's ocean bottom and maintain a substantial part of the world's biodiversity (Snelgrove, 1998; Labrune et al., 2008). Assessing their biodiversity and latitudinal patterns, even though it is complicated by the difficulty in sampling and sharply delineating habitats in these systems, is thus of special importance (Labrune et al., 2008). Macrobenthos constitute the dominant organism biomass of marine soft sediments (Snelgrove, 1998). All of the known non-symbiont phyla but one are found in the marine environment, with most being represented in marine sediments (Grassle et al., 1991). Soft bottom macrobenthos is a key biological component of marine ecosystems where it plays an important role in ecological processes such as nutrient cycling, pollutant metabolism or secondary production (Snelgrove, 1998; Pratt et al., 2014). Although soft-bottoms are the largest ecosystem on Earth in terms of area, only a small percentage of their macrobenthos has been studied and most of its species are not yet described (Snelgrove, 1998). In this way, it is useful to improve our knowledge of its biodiversity (Ellingsen, 2002; Veiga et al., 2016).

Semi-enclosed coastal systems (SECS), such as lagoons and estuaries, are particular ecosystems of the coastal zone where the macrobenthos is a dominant biological component. The SECS are largely distributed worldwide under all the latitudes and are therefore ideal systems to study LDGs and their potential drivers because they (a) are easily accessible and (b) have very diverse, abundant, macrobenthic organisms belonging to various taxonomic groups.

The Moroccan coast stretches over 3500 km and 15° of latitude along both the Mediterranean Sea and the Atlantic Ocean. Moroccan low coasts offer an important succession of particular geomorphological forms (Chafik et al., 2001) among which several lagoons, estuaries and bays. Nador and Smir lagoons lie on the Mediterranean coast, whereas Moulay Bousselham, Sidi Moussa, Oualidia and Khnifiss are located in the Atlantic coast. The largest estuaries are located in the Atlantic coast: Tahaddart, Loukkos, Sebou, Bouregreg and Oum Rbia. Finally, Dakhla bay is the most important bay of the Atlantic coast of Morocco, it is unique in North Africa as a migration, wintering and nesting area for thousands of waterbirds (Qninba et al., 2003). Those ecosystems are heterogeneous, mainly due to their typology (bay, estuaries, lagoons), geomorphology, catchment geology and the spatio-latitudinal variation in different environmental factors along the Moroccan coasts

Up-to-now, there is no synthesis of the benthic species of the Moroccan SECS, despite several studies conducted in the last decades (e.g. Elkaïm, 1974; Lacoste, 1984; Bekkali, 1987; Guelorget et al., 1987; Bayed et al., 1988; Cheggour, 1988; Zine, 1989; Chbicheb, 1996; Bazairi, 1999; Mergaoui, et al., 2003; Chaouti and Bayed, 2005; Zine, 2005; Azirar, 2006; Cherkaoui, 2006; Gauteur, 2006; Lefrere, 2012; Joulami, 2013; Bazairi et al., 2014; Boutahar, 2014; El Asri et al., 2015; El Asri et al., 2017; Touhami, 2018; El Asri, 2019). This was a stimulating fact to provide a first baseline meta-data on such ecosystems in such poorly known area at the global scale and to fulfill a knowledge gap on benthic macrofauna in SECS of the Southern North-East Atlantic and Mediterranean ecoregions.

The overall objectives of this study were (1) to provide the first national and comprehensive checklist of the soft bottom macrozoobenthic species in semi-enclosed coastal systems of Morocco, (2) to test for the presence of a latitudinal diversity gradients in soft bottom macrozoobenthic species of semi-enclosed coastal systems of Morocco and (3) to understand their drivers by comparing the benthic assemblages between the different sites according to their ecoregion, their latitudinal position, the type of ecosystem (lagoon, estuary or bay), the site surface area, temperature and salinity. Additionally, we identified the current knowledge and gaps and make recommendations on respective research in future years.

2. Materials and methods *Study sites*

12 SECS were considered in this study based of data availability (Fig. 1). There are distributed along a large latitudinal gradient and are situated both in the Mediterranean coast of Morocco (two lagoons) and the Atlantic coast of Morocco (four lagoons, five estuaries and one bay). These SECS differ in terms of configuration, surface area and environmental conditions (Table 1). These sites have a very important ecological interest; most of them have been listed to the RAMSAR Convention as wetlands of international importance. They represent the most important Moroccan wetlands for the migration and wintering of birds.

These SECS are contrasting in the climate: Mediterranean semi-arid to temperate variant at the level of Nador lagoon to a desert climate at Dakhla bay. On the other hand, anthropogenic activities differ from one site to another: traditional fishing (all sites), aquaculture (Oualidia and Dakhla), thermal power station (Tahaddart), intensive agriculture (Moulay Bousselham, Sidi Moussa and Oualidia), port's activities (Nador, Loukkos, Sebou, Bouregreg, and Dakhla), dredging activities (Sebou and Oum Rbia,), mining (Nador, Sidi Moussa, Oualidia and Khnifiss) and industrial effluents (Nador, Sidi Moussa and Dakhla).

2.2. *Data sources*

The checklist was based on published data from 1974 to 2020 as well as co-authors unpublished data (see Table S1 and Table 4 for references' details). The list of benthic macrofauna species and total species richness were compiled for each SECS. Species names were checked and updated to current nomenclature according to World Register of Marine Species (<http://www.marinespecies.org>) (consulted January 14, 2020). Only few species were not included in WORMS. Taxa identified at a higher taxonomic level than the species were removed from the checklist. Only the taxa sp. and cf. were retained if cited only once.

The heterogeneity and incompleteness of available information led to the selection of a subset of the SECS environment descriptors which fulfilled the criteria of data reliability (i.e. information coming from published sources), homogeneity (i.e. available for all the lagoons), and comparability (i.e. data which can be expressed with a shared measurement unit). Nine different SECS variables were included: type of aquatic system (1 = lagoon; 2 = estuary; 3 = bay), latitude

(LAT), longitude (LON), surface area, maximum annual water temperature (M Temp, °C), minimum annual water temperature (m Temp, °C), maximum annual water salinity (M Sal), minimum annual water salinity (m Sal).

2.3. Biological data

Biotic parameters including the species richness (S), average taxonomic distinctness with presence/absence data (Δ^+), total taxonomic distinctness ($S\Delta^+$), average phylogenetic diversity (Φ^+) and variation in taxonomic distinctness (Λ^+), total phylogenetic diversity ($S\Phi^+$) were calculated.

Sampling effort leading to the compilation of the species lists was also measured by the total number of samples collected in each SECS (calculated as number of sampling sites x sampling frequency).

2.4. Data analysis

The relation between SECS characteristics (independent variables) on species richness and taxonomic diversity indices (dependent variables) was tested using multiple linear regressions after testing the collinearity between independent variables. The possible effect of sampling effort on biotic parameters was previously tested by univariate regression. Then, residuals of the univariate regression with sampling effort were considered instead of the original data in multiple regression analysis. Moreover, the effect of the types of SECS (lagoon vs estuary) was tested using univariate PERMANOVA (Anderson et al., 2004).

Differences in the structure of the taxonomic assemblages between Moroccan SECS were explored using a cluster analysis based on a Bray-Curtis coefficient. Similarities (Bray-Curtis coefficient) among Moroccan SECS were calculated based on the taxonomical composition (Bray-Curtis coefficient calculated on (0, 1) species presence-absence data corresponding to Sorensen coefficient (Clarke and Warwick, 2001) (Table S2). Affinity groups differences were visualized through Principal Coordinates Ordination analysis (PCO) (Clarke and Warwick, 2001). The abiotic variables that were correlated (Spearman $\rho > 0.5$) to samples ordination were represented as superimposed vectors in the PCO graph.

The best subset of SECS characteristics explaining the observed variability in benthic macrofauna assemblages was selected by means of distance-based linear models (DISTLM) (Anderson et al., 2008), using appropriate permutation (9999 permutations) and with Adjusted R^2 criterion and stepwise procedure for the model selection. Distances among aquatic systems were visualized through a dbRDA plot. Predictors variables were partitioned to four sets of predictor variables: environmental variables (surface area, maximum annual water temperature (minimum annual water temperature, maximum annual water salinity, minimum annual water salinity), type of SECS (lagoon, estuary, bay), province (Mediterranean and Atlantic) and geographical variables (latitude and longitude).

All multivariate analyses were conducted in the Primer 7 space (Clarke and Gorley, 2006), while correlation and regressions tests were carried out in Statistica 12.0 (Statsoft, 2017).

3. Results

3.1. Checklist

36 sets of both published and unpublished data were compiled to obtain the checklist of benthic macrofauna of SECS of Morocco (Table S1). Most of them focused on a single SECS. There were 496 species recorded from the Moroccan SECS. They belong to 7 phyla, 21 classes, 65 orders and 201 families. Molluscs are the richest phylum with 179 species belonging to 5 classes, 31 orders and 71 families. Arthropods is the second richest phylum with 164 species belonging to 5 classes, 12 orders and 72 families. 120 species of Annelida were reported, with 2 classes, 7 orders and 33 families. Chordata presented 15 species, from 1 class, 5 orders and 9 families. The phylum of Echinodermata, Cnidaria, and Nemertea were represented by 8, 6 and 4 species respectively.

Among the 496 species reported in this study, 95 species showed an exclusive Mediterranean distribution (M), 99 species have an Atlantico-Mediterranean distribution (AM) and 302 have an Atlantic distribution (A) (Fig. 2).

3.2. Species composition and taxonomic diversity

Values estimated for the different indices are summarised in Table 2. The number of taxa compiled by site fluctuated between 32 (Loukkos estuary) and the 161 (Moulay Bousselham lagoon) (Table

3). Sampling effort showed a wide variability among the studied sites, with the maximal value in Moulay Bousselham lagoon and the minimum in Loukkos estuary (Table 4). With the exception of average phylogenetic diversity ($\Phi+$) and variation in taxonomic distinctness ($\Lambda+$), all the other indices showed significant dependence with the sampling effort ($p < 0.05$). Moreover, all the taxonomic indices were significantly ($p < 0.05$) related to species richness except the average taxonomic distinctness ($\Delta+$). In the majority of the SECS, both average taxonomic distinctness ($\Delta+$) and variation in taxonomic distinctness ($\Lambda+$) were within the 95% confidence funnel ($p \geq 0.05$). Only the Oum Rbia and the Sebou estuaries as well as the Khnifiss lagoon appeared out of the confidence funnels (Fig. 3).

Multiple regression analysis ($p > 0.05$) revealed that none of the SECS features considered here are structural abiotic features regarding the species richness and the taxonomic diversity indices. Moreover, permutational multivariate analysis of variance (PERMANOVA) showed no interactions between type of SECS ($p > 0.05$) (Table 3) in terms of species richness (S) and all taxonomic diversity indices (Table 4).

3.3. *Macrobenthic assemblages' affinity*

Cluster analysis, at a similarity distance of 40%, allowed to distinguish three affinity groups (Fig. 4): G1 (40% of similarity) composed by Bouregreg estuary and Moulay Bousselham lagoon, G2 (50% of similarity) composed by Tahaddart, Loukkos and Oum Rbia estuaries and G3 (60% of similarity) composed by Oualidia and Sidi Moussa lagoons; the others SECS (Nador and Smir lagoons on the Mediterranean coast; Khnifiss lagoon and Dakhla bay on the Atlantic coast) remain individually separated from the previous groups with similarity fluctuating between 20% and 30%.

The obtained affinity groups as well as the other sites were represented on the PCO ordination graph (Fig. 5). The first two ordination axes explained 36% (all sites considered) of the total variance in benthic macrofauna assemblages. The two-dimensional plots show a clear separation between lagoons and estuaries systems.

Through the DistLM analysis, only the SECS-type set of predictor variables had a significant relationship (type-estuary and type-lagoon) with species-derived multivariate cloud ($p < 0.01$), explaining 27% of the total variation. However, the best model obtained through the DistLM

procedure included height variables (Type-estuary, Type-lagoon, Type-bay, Province-Atlantic Surface, m-Temp, m-Sal and M-Sal) as predictors of benthic macrofauna composition, explaining 33% of the total variation (Adjusted $R^2=0.33$). When transposed to the dbRDA plot, the first two axes captured nearly 62% of the variability in the fitted model and 47% of the total variation in the data cloud (Fig. 6). Axis 1 (representing 20% of total variation) was negatively correlated to Type-estuary ($r = -0.58$) and to M-Sal ($r = -0.52$). Axis 2 (representing 14% of total variation) was negatively correlated to Surface ($r = -0.53$). The axis 3 (representing 13% of total variation) was negatively correlated to Province-Atl ($r = -0.52$) and Type-bay ($r = -0.55$).

3.4. Gaps and knowledge gained

Table 4 showed that there is a scarcity of studies and that there is no regular spatio-temporal monitoring. Sampling techniques, effort and objectives differ between sites and studies. Although all the existing studies are quantitative, most of them were single spot studies, had limited geographic scope (Mergaoui et al., 2003; Ait Mlik, 2009; Bououarour, 2013; Joulami, 2013), focused only on single taxonomic groups (e.g. Annelida: El asri et al., 2017; Arthropoda: Aksissou, 1997; Boussalwa et al., 2000), or on habitats (intertidal, subtidal, meadows). On the other hand, few studies have been carried out on benthos-predator interactions (Joulami, 2013; Touhami et al., 2019). Moreover, while few studies have considered biomass, there is no evaluation of secondary production and productivity in all the Moroccan SECS. As a result, the studies that can be considered as references for soft sediments benthic assemblages are those of Guelorget et al. (1987), Zine (1989) and El Kamcha (unpublished data) for the lagoon of Nador, Bazairi & Gam (2004) for the Loukkos estuary, Bazairi (1999) and Boutoumit (unpublished data) for the Moulay Bousselham lagoon, Elkaïm (1974) and Cherkaoui (2006) for the Bouregreg estuary, Gauteur (2006), Boutahar (2014) and El asri (2019) for the Oualidia lagoon, Bazairi & Zourarah (2001, 2007) and Bazairi et al. (2017) for the Oum Rbia estuary and finally El asri (2019) for Dakhla bay.

4. Discussion

Checklists of marine species at regional scale have multiple uses. In addition to offering comparative facts for biodiversity studies, they serve as a crucial device in spotting and delimiting regions in need of protection, inferring the capacity effect of anthropogenic interest, assessing the

complexity of organic communities, and estimating the provision of dwelling resources (Hendrickx and Harvey, 1999).

The sample of SECS considered in this study covers most of the range of variation along the coasts of Morocco. Therefore, the current compilation represents the first comprehensive annotated checklist that gives an overall view on soft-bottom benthic macrofauna of the semi-enclosed coastal systems within the Atlantic and Mediterranean oceanographic regions of Moroccan waters, which is relevant in such poorly known area at the global scale. Moreover, it fulfilled a knowledge gap on benthic macrofauna in SECS of the Southern part of the Northeastern Atlantic and Mediterranean ecoregions. While all the existing inventories on Moroccan marine fauna focused on single taxonomic group and have large scope, our ecosystem-based checklist is more than a simple list of species inhabiting comparable ecosystems and constitutes a synthetic illustration of the relationships that species have with each other and with their environment. Furthermore, the resulting metadata, available in open access, will serve in comparing the soft-bottom macrobenthos assemblages at a large scale.

This study revealed a diverse benthic macrofauna for the Moroccan SECS with overall 496 species dominated by Molluscs, Arthropods and Annelids. It represents almost 46% of the known marine fauna of marine waters of Morocco (1068 taxa, all groups combined) (ONEM, 1998). However, comparison of diversity results between Moroccan SECS may be done with caution since the sampling methods, units and scales are often different, and moreover the diversity of habitats in such ecosystems are high (Chardy and Clavier, 1988; Alongi, 1990).

Patterns in diversity are often related to latitude, a phenomenon known as the latitudinal diversity gradient (LDG), whereby a decrease with increasing latitude is found (Roy et al., 1998; Rex et al., 2000; Attrill et al., 2001; Willig et al., 2003; Hillebrand, 2004). Several hypotheses for the underlying causes for such pattern are suggested, yet none of them is solely sufficiently convincing (Willig et al., 2003; Hillebrand, 2004), although solar energy input (and for the marine territory, the sea surface temperature as its proxy) is most often mentioned as the main acting principle (Rohde, 1992; Roy et al., 1998). Species richness is the most elementary, easy to interpret and widely used measure of biodiversity (e.g. Dornelas et al., 2014). It has been shown to follow a – generally unimodal – large-scale ($>45^\circ$) latitudinal gradient for marine benthic invertebrates with

a peak in equatorial regions (Chaudhary et al., 2016). In contrast to the general LDG pattern, our results showed that species richness and taxonomic diversity indices showed no relationship with latitude. Deviations from the general LDG pattern have also been reported before for European marine benthos by Renaud et al. (2009) who found no or weakly positive relationships. The explanation for this kind of diverting trend is that the impact of local variation in environmental factors is stronger than that of latitude related factors (Gaston, 2000; Renaud et al., 2009). Marine diversity might not follow a strict latitudinal gradient as the drivers of marine diversity themselves are not usually correlated with latitude (Piacenza et al., 2015).

In terms of assemblages composition, the best multivariate model, explaining 33% of the total variation observed in benthic assemblages, included the type of the SECS (estuaries vs lagoons vs bay), the marine ecoregion (Atlantic vs Mediterranean), the surface of the SECS and the environmental features (minimal temperature, minimal and maximal salinity) as predictors of benthic macrofauna composition in the Moroccan SECS. Such differences in benthic macrofaunal composition across a scale could result from the fact that each ecosystem has its own specific characteristics, which implies an individualistic approach to ecosystem ecology. Indeed, all ecosystems are subject to climatic and environmental forces and it is assumed that their variations induce a response from communities (Möllmann and Diekmann, 2012). The diversity and distribution of organisms can be influenced by the stochastic, ecological and evolutionary processes at local and regional scales (Hubbell, 2001), the limits of dispersion and recruitment of macroinvertebrate taxa (Hurtt & Pacala, 1995), the structural heterogeneity of the transitional waters (Basset & Abbiati, 2004) and the consequent selection of macroinvertebrate taxa according to their functional traits and niche needs (MacArthur, 1970).

Variations in species richness and taxonomic composition of benthic macrofauna depends on local oceanographic processes (Aller et al., 2002; Coleman et al., 1997; McCallum et al., 2015), as well as on physiographic characteristics, such as surface area and outlet length (Basset et al., 2006). Variations might be related to ecosystem morphology, substrate type, organic residues (Galeron et al., 2001), salinity (Battaglia, 1959), degree of confinement (Guelorget and Perthuisot, 1983; Guelorget et al., 1983), changes in nutrient concentrations and changes in primary productivity (Galeron et al., 2001).

According to Spalding et al. (2007), the vast marine region of Morocco can be subdivided into two provinces, the Mediterranean Sea (Alboran Sea Ecoregion) and the Lusitanian (Saharan Upwelling Ecoregion). In corroboration to this subdivision, the type of province (Atlantic vs Mediterranean) was shown to be a significant predictor factor of the composition of the soft-bottom benthic fauna of the Moroccan SECS. Therefore, the currently described marine biogeographic boundaries in Morocco seem to apply to soft-bottom macrofauna of SECS and which environmental drivers were most associated with species differences among these two provinces. Significant differences have been found on the structure of the communities between the marine ecoregions, a fact which may be correlated with the specificities of the ecoregion's physiographical characteristics, as shown by Kong et al. (2013), which emphasizes that the distribution of the macroinvertebrate is correlated with ecoregional characteristics. Moreover, these marine ecoregions are affected by environmental factors (Lara-Lara et al., 2008).

5. Conclusion

Our results provide the first broad-scale baseline of composition and diversity patterns of soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco and elucidate the main environmental factors that shape their latitudinal and biogeographic patterns. To our knowledge, the presently reviewed meta-data would be considered as most up-to-dated checklist of the soft-sediment macrozoobenthos in Moroccan SECS. This checklist is relevant in such poorly known area and fulfills a knowledge gap on SECS in the Northeastern Atlantic and Mediterranean ecoregions. However, the knowledge gained here is insufficient to address perceived shortfalls in knowledge of biodiversity, its importance to ecosystem function, and the threats and consequences of disturbance by anthropogenic activities.

Acknowledgements

Data for the Tahaddart estuary were obtained in the framework of EU funded WADI Project. Those of the Dakhla bay and the Khnifiss lagoon were partially obtained in the framework of the

Saharian wetlands of Morocco. The authors are indebted to two anonymous reviewers for their kind and helpful contributions.

Authors' Contributions

Soilam Boutoumit: Investigation, Data curation, Writing – Original Draft, Visualisation. **Reda El Kamcha:** Investigation, Data curation, Writing – Original Draft, Visualisation. **Oussama Bououarour:** Investigation, Data curation, Writing – Original Draft, Visualisation. **Latifa Joulami:** Investigation, Data curation, Review. **Boutahar Loubna:** Investigation, Data curation, Review. **Abdelaziz Benhoussa:** Resources, Funding acquisition. **Mohamed Maanan:** Review & Editing, Supervision. **Laurent Godet:** Writing-Review & Editing, Validation. **Abdellatif Bayed:** Resources, Validation. **Hocein Bazairi:** Conceptualization, Methodology, Writing – Original Draft, Review & Editing, Supervision.

References

- Achab, M., 2011. Les plages et les vasières des environs des embouchures des oueds Tahaddart et Gharifa (NW du Maroc) : dynamique morphosédimentaire et impact des aménagements sur leur évolution récente. Travaux de l'Institut Scientifique, Rabat. 6, 1-12.
- Ait Mlik, K., 2009. Contribution à l'étude de l'évolution à long terme des structures benthiques de la lagune de Merja Zerga (Mémoire Master). Université Hassan II. Casablanca, 40pp.
- Aksissou, M., 1997. Dynamique des populations d'*Orchestia gammarellus* (Pallas, 1766) - Crustacea, Amphipoda, Talitridae- du littoral méditerranéen du Maroc occidental et impact des aménagements (Thèse Doctorat d'Etat). Université Abdelmalek Essaâdi. Tétouan, 145pp.
- Aller, J.Y., Aller, R.C., Green, M.A., 2002. Benthic faunal assemblages and carbon supply along the continental shelf/shelf break-slope off Cape Hatteras, North Carolina. Deep-Sea Res. Pt. II 49 (20), 4599-4625. [http://dx.doi.org/10.1016/S0967-0645\(02\)00131-5](http://dx.doi.org/10.1016/S0967-0645(02)00131-5)
- Alongi, D.M., 1990. The ecology of tropical soft-bottom benthic ecosystems. Oceanography and Marine Biology: An Annual Review. 28, 381-496.

- 387 Anderson, M.J., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA + for PRIMER: guide to
388 software and statistical methods. University of Auckland and PRIMER-E, Plymouth.
- 389 Anderson, M.J., Millar, R.B., 2004. Spatial variation and effects of habitat on temperate reef fish
390 assemblages in northeastern New Zealand. *J. Exp. Mar. Biol. Ecol.* 305, 191–221.
391 <https://doi.org/10.1016/j.jembe.2003.12.011>
- 392 Angel, M.V., 1997. Pelagic diversity in “Marine Biodiversity. Patterns and Processes” (eds R.F.G.
393 Ormond, J.D. Gage & M.V. Angel). Cambridge University Press, Cambridge, UK, 35-68. .
394 <https://doi.org/10.1017/CBO9780511752360>
- 395 Attrill, M.J., Stafford, R., Rowden, A.A., 2001. Latitudinal diversity patterns in estuarine tidal
396 flats: indications of a global cline. *Ecography*. 24, 318-324. [https://doi.org/10.1034/j.1600-](https://doi.org/10.1034/j.1600-0587.2001.240309.x)
397 [0587.2001.240309.x](https://doi.org/10.1034/j.1600-0587.2001.240309.x)
- 398 Azirar, A., 2006. Macrofaune benthique de l'estuaire d'Oum Rbia (Mémoire DESA). Université
399 Mohammed V. Rabat, 64pp.
- 400 Barboza, F.R., Defeo, O., 2015. Global diversity patterns in sandy beach macrofauna: a
401 biogeographic analysis. *Sci. Rep.* 5, 14515. <https://doi.org/10.1038/srep14515>
- 402 Basset, A., Abbiati, M., 2004. Challenges to transitional water monitoring: ecological descriptors
403 and scales. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 14, S1-S3.
404 <https://doi.org/10.1002/aqc.669>
- 405 Basset, A., Sabetta, L., Fonnesu, A., Mouillot, D., Do Chi, T., Viaroli, P., Giordani, G.,
406 Reizopoulou, S., Abbiati, M., Carrada, G.C., 2006. Typology in Mediterranean transitional waters:
407 new challenges and perspectives. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 16
408 (5), 441-455. <https://doi.org/10.1002/aqc.767>
- 409 Battaglia, B., 1959. Final resolution of the symposium on the classification of brackish waters.
410 *Archivio Di Oceanografia E Limnologia*. 11(suppl), 243-248.

- 411 Bayed, A., EL Agbani, M.A., Fekhaoui, M., Schouten, J.R., 1988. Benthos of soft substrates in the
412 intertidal zone of the Khnifiss lagoon; in DAKKI M. & W. LIGNY (eds.): The Khnifiss lagoon
413 and its surrounding environment (Province of Laayoune, Morocco). Trav. Inst. Sci., Mém. Hors-
414 série. 71-80.
- 415 Bazairi, H., 1999. La faune macrobenthique de la lagune de Moulay Bousselham : Structure et
416 successions spatio-temporelles (Thèse Doctorat). Université Mohammed V. Rabat, 199pp.
- 417 Bazairi, H., Bayed, A., 2006. L'écosystème benthique de l'estuaire de Tahaddart : biodiversité,
418 fonctionnement et qualité écologique. Unpublished report, WADI Project, 75pp.
- 419 Bazairi, H., Gam, M., 2004. Diagnostic pour l'élaboration du plan de gestion: macroinvertébrés
420 estuariens. Projet Gestion Intégrée du Complexe de Zones Humides du Bas Loukkos (Larache,
421 Maroc), Deuxième phase (2004-2005): Actions opérationnels, 40pp.
- 422 Bazairi, H., Zourarah, B., 2001. Faune macrobenthique de l'estuaire d'Oum Rbia: biodiversité,
423 distribution spatiale et état de la qualité écologique. Unpublished report, GREPEN, 25pp.
- 424 Bazairi, H., Zourarah, B., 2007. Faune macrobenthique de l'estuaire d'Oum Rbia: biodiversité,
425 distribution spatiale et état de la qualité écologique. Unpublished report, GREPEN, 30pp.
- 426 Bazairi, H., Zourarah, B., El Kamcha, R., Boutoumit, S., 2017. Faune macrobenthique de l'estuaire
427 d'Oum Rbia: biodiversité, distribution spatiale et état de la qualité écologique. Unpublished report,
428 OMAZINE, 50pp.
- 429 Bekkali, R., 1987. Les ostracodes du lac Smir (Maroc nord-occidental) (Thèse Doctorat 3ème
430 cycle). Université Mohammed V de Rabat, 165pp.
- 431 Boucher, G., Lamshead, P.J.D., 1995. Ecological biodiversity of marine nematodes in samples
432 from temperate, tropical, and deep sea regions. Conservation Biology, 9, 1594–1604.
433 <https://doi.org/10.1046/j.1523-1739.1995.09061594.x>

- 434 Bououarour, O., 2013. La Macrofaune benthiques des herbiers de *Zostera noltii* de la lagune de
435 Moulay Bousselham (Merja Zerga) : Structure des communautés et état de la qualité écologique
436 (Mémoire Master). Université Mohammed V. Rabat, 45pp.
- 437 Boussalwa, E., Douira, A., Mokhtar, N., 2000. Contribution à l'étude typologique d'une lagune
438 exposée à la pollution, la lagune de Nador (Maroc): distribution des Crustacés. *Cah. Biol. Mar.* 41,
439 255-263.
- 440 Boutahar, L., 2014. Evaluation à long terme de la macrofaune benthique dans la lagune de Oualidia
441 (Atlantique, Maroc) entre 2006 et 2014 (Mémoire Master). Université Mohammed V. Rabat,
442 105pp.
- 443 Chafik, A., Cheggour, M., Cossa, D., Benbrahim, S., Sifeddine, M., 2001. Quality of Moroccan
444 Atlantic coastal waters: water monitoring and mussel watching. *Aquat. Living Resour.* 14, 239-
445 249. [https://doi.org/10.1016/S0990-7440\(01\)01123-8](https://doi.org/10.1016/S0990-7440(01)01123-8)
- 446 Chaouti, A., Bayed, A., 2005. Diversité taxonomique et structure de la macrofaune benthique des
447 substrats meubles de la lagune de Smir. In: Bayed A. & Felicita F. (Eds) – Ecosystèmes côtiers
448 sensibles de la Méditerranée : cas du littoral de Smir. *Trav. Inst. Sci. Série générale.* 4, 33-42.
- 449 Chardy, P., Clavier, J., 1988. Biomass and trophic structure of the macrobenthos in the south-west
450 lagoon of New Caledonia. *Marine Biology.* 99, 195-202. <https://doi.org/10.1007/BF00391981>
- 451 Chaudhary, C., Saeedi, H., Costello, M.J., 2016. Bimodality of Latitudinal Gradients in Marine
452 Species Richness. *Trends in Ecology & Evolution.* 31, 670-676.
453 <https://doi.org/10.1016/j.tree.2016.06.001>
- 454 Chbicheb, A., 1996. Organisation biogéologique d'un bassin paralytique : la lagune de Oualidia
455 (côte atlantique marocaine) (Thèse Doctorat 3ème cycle). Université de Nantes, 167pp.
- 456 Cheggour, M., 1988. Contribution à l'étude d'un milieu paralytique : l'estuaire du Bou Regreg (Côte
457 atlantique marocaine). Condition écologique globale. Etude de la contamination métallique (Thèse
458 de 3ème cycle). ENS. Rabat, 337pp.

- 459 Cherkaoui, S., 2006. Structure et organisation des peuplements macrozoobenthiques de l'estuaire
 460 du Bou Regreg après la construction du barrage (Thèse Doctorat). Université Mohammed V.
 461 Rabat, 305 pp.
- 462 Clarke, A., 1992. Is there a latitudinal diversity cline in the sea? *Trends in Ecology and Evolution*.
 463 7, 286-287. [https://doi.org/10.1016/0169-5347\(92\)90222-W](https://doi.org/10.1016/0169-5347(92)90222-W).
- 464 Clarke, A., Crame, J.A., 1997. Diversity, latitude and time: patterns in the shallow sea. In "Marine
 465 Diversity: Patterns and Processes" (eds R.F.G. Ormond, J.D. Gage & M.V. Angel). Cambridge
 466 University Press, Cambridge, UK, 122-147. <https://doi.org/10.1017/CBO9780511752360>
- 467 Clarke, K.R., Gorley, R.N., 2006. PRIMER v.6: User Manual/Tutorial (Plymouth Routines in
 468 Multivariate Ecological Research). PRIMER-E, Plymouth.
- 469 Clarke, K.R., Warwick, R., 2001. Change in Marine Communities: An approach to statistical
 470 analysis and interpretation. PRIMER-E, Plymouth.
- 471 Coleman, N., Gason, A.S.H., Poore, C.B., 1997. High species richness in the shallow marine
 472 waters of south-east Australia. *Mar. Ecol. Prog. Ser.* 154, 17-26.
 473 <http://dx.doi.org/10.3354/meps154017>
- 474 Crame, J.A., 2000. Evolution of taxonomic diversity gradients in the marine realm: evidence from
 475 the composition of recent bivalve faunas. *Paleobiology*. 26, 188-241.
 476 [https://doi.org/10.1666/0094-8373\(2000\)026<0188:EOTDGI>2.0.CO;2](https://doi.org/10.1666/0094-8373(2000)026<0188:EOTDGI>2.0.CO;2)
- 477 Cruz-Motta, J.J., Miloslayich, P., Guerra-Castro, E., Hernández-Agreda, A., Herrera, C., Klein,
 478 E., Barros, F., Navarrete, S.A., Sepúlveda, R., Glasby, T.M., Bigatti, G., Cárdenas-Calle, M.,
 479 Carneiro, P.B.M., Carranza, A., Flores, A., Gil-Kodaka, P., Gobin, J., Gutiérrez, J., Klein, E.,
 480 Krull, M., Lazarus, J.F., Londoño-Cruz, E., Lotufo, T., Macaya, E.C., Mora, E., Palomo, G.,
 481 Parragué, M., Pellizzari, F., Retamales, R., Rocha, R., Romero., 2020. Latitudinal patterns of
 482 species diversity on South American rocky shores: Local processes lead to contrasting trends in
 483 regional and local species diversity. *Journal of Biogeography*. 47, 1966-1979.
 484 <https://doi.org/10.1111/jbi.13869>

- 485 Cuvelier, J., Riviere, V., Bazairi, H., Benhoussa, A., Cerniaut, C., Charrier, M., Moulis, D., Pupier,
 486 P., Qninba, A., 2014. Plan de gestion ; Lagune de Nador – Cordon de la Bocana – partie 1 :
 487 Diagnostic socio-économique et écologique. Conservatoire du Littoral. O2TERRE ; AGIR
 488 écologique, 177p.
- 489 Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., Magurran, A.E.,
 490 2014. Assemblage time series reveal biodiversity change but not systematic loss. *Science*. 344,
 491 296-299. <https://doi.org/10.1126/science.1248484>
- 492 El Amraoui, M., Salama, Y., El Amraoui, B., Lazrak, N., Mounkad, M., 2015. Evaluation of the
 493 physicochemical and some trace elements of the surface waters of the Moroccan Atlantic estuary:
 494 case of the estuary of the river Bouregreg. *Carpathian Journal of Earth and Environmental*
 495 *Sciences*. 10 (2), 189-198.
- 496 El Asri, F., 2019. Composition, organisation et dynamique des communautés zoobenthiques de
 497 deux écosystèmes côtiers : la lagune de Oualidia et la Baie de Dakhla (Atlantique Marocain)
 498 (Thèse Doctorat). Université Hassan II. Casablanca, 297 pp.
- 499 El Asri, F., Zidane, H., Errhif, A., Tamsouri, M.N., Maanan, M., Malouli Idrissi, M., Martin, D.,
 500 2017. Polychaete diversity and assemblage structure in the Oualidia Lagoon, Moroccan Atlantic
 501 coast. *J. Mar. Biol. Assoc. United Kingdom*. 98, 1337-1346.
 502 <https://doi.org/10.1017/S0025315417000388>
- 503 El Asri, F., Zidane, H., Maanan, M., Tamsouri, M., Errhif, A., 2015. Taxonomic diversity and
 504 structure of the molluscan fauna in Oualidia lagoon (Moroccan Atlantic coast). *Environmental*
 505 *Monitoring and Assessment*. 187, 545. <http://dx.doi.org/10.1007/s10661-015-4752-7>
- 506 El Houssaini, F., 2005. Composition et structure du macrozoobenthos associé à *Zostera noltii* de
 507 la lagune de Smir (Mémoire DESA). Université Hassan II Aïn Chock. Casablanca, 68 pp.
- 508 El Kamcha, R., Bououarour, O., Boutoumit, S., Bazairi, H., 2020. Occurrence of the invasive
 509 *Caprella scaura* Templeton, 1836 (Amphipoda: Caprellidae) in the Marchica coastal lagoon
 510 (Alboran Sea, Morocco). *BioInvasions Records*. 9. Article in press.

- 511 Elkaïm, B., 1974. Contribution à l'étude écologique d'un estuaire atlantique marocain : l'estuaire
512 du Bou Regreg (Thèse Doctorat d'Etat). Université de Bordeaux I- 2ème partie, 250 pp.
- 513 Ellingsen, K.E., 2002. Soft-sediment benthic biodiversity on the continental shelf in relation to
514 environmental variability. Marine Ecology Progress Series. 232, 15-27.
515 <https://doi.org/10.3354/meps232015>
- 516 Galeron, J., Sibuet, M., Vanreusel, A., Mackenzie, K., Gooday, A., Diné, A., Wolff, G., 2001.
517 Temporal patterns among meiofauna and macrofauna taxa related to changes in sediment
518 geochemistry at an abyssal NE Atlantic site. Progress in Oceanography. 50 (1-4), 303-324 Sp. Iss.
519 SI). [https://doi.org/10.1016/S0079-6611\(01\)00059-3](https://doi.org/10.1016/S0079-6611(01)00059-3)
- 520 Gam, M., de Montaudouin, X., Bazairi, H., 2010. Population dynamics and secondary production
521 of the cockle *Cerastoderma edule*: A comparison between Merja Zerga (Moroccan Atlantic Coast)
522 and Arcachon Bay (French Atlantic Coast). Journal of Sea Research. 63 (3-4), 191-201.
523 <https://doi.org/10.1016/j.seares.2010.01.003>
- 524 Gaston, K.J., 2000. Global patterns in biodiversity. Nature. 405, 220-227.
525 <https://doi.org/10.1038/35012228>.
- 526 Gauteur, M., 2006. Etablissement du diagnostic écologique d'un environnement lagunaire : la
527 lagune de Oualidia (Côte atlantique, Maroc) (Mémoire Master). Université de Bordeaux 1, 60 pp.
- 528 Geawhari, M.A., Huff, L., Mhammedi, N., Trakadas, A., Ammar, A., 2014. Spatial-temporal
529 distribution of salinity and temperature in the Oued Loukkos estuary, Morocco: using vertical
530 salinity gradient for estuary classification. Computational Statistics. 3 (1), 1-9.
531 <https://doi.org/10.1186/2193-1801-3-643>
- 532 Giangrande, A., Licciano, M., 2004. Factors influencing latitudinal pattern of biodiversity: An
533 example using Sabellidae (Annelida, Polychaeta). Biodiversity and Conservation 13, 1633-1646.
534 <https://doi.org/10.1023/B:BIOC.0000029327.63397.6b>

- 535 Grassle, J.F., Lasserre, P., McIntyre, A.D., Ray, G.C., 1991. Marine Biodiversity and Ecosystem
536 Function: A proposal for an international program of research. *Biology International Special Issue*.
537 23, 19p.
- 538 Guelorget, O., Frisoni, G.F., Perthuisot, J.P., 1983. Zonation biologique des milieux lagunaires:
539 définition d'une échelle de confinement dans le domaine paralique méditerranéen. *Journal de*
540 *Recherche océanographique*. 8, 15-35.
- 541 Guelorget, O., Perthuisot, J.P., 1983. Le domaine paralique. Expressions géologiques, biologiques
542 et économiques du confinement. *Travaux du laboratoire de géologie*. 16, 1-136.
- 543 Guelorget, O., Perthuisot, J.P., Frisoni, G.F., Monti, D., 1987. Le rôle du confinement dans
544 l'organisation biogéographique de la lagune de Nador (Maroc). *OceanolActa*. 10 (4), 435-444.
- 545 Haddout, S., Maslouhi, A., Magrane, B., Igouzal, M., 2015. Study of salinity variation in the Sebou
546 River Estuary (Morocco). *Desalination and Water Treatment*. 57 (36), 17075-17086.
547 <https://doi.org/10.1080/19443994.2015.1091993>
- 548 Hawkins, B.A., 2001. Ecology's oldest pattern? *Trends Ecol. Evol.*, 16, 470.
549 [https://doi.org/10.1016/S0160-9327\(00\)01369-7](https://doi.org/10.1016/S0160-9327(00)01369-7)
- 550 Hendrickx, M., Harvey, A.W., 1999. Checklist of anomuran crabs (Crustacea: Decapoda) from the
551 Eastern Tropical Pacific. *Belgian Journal of Zoology*. 129 (2), 363-389.
- 552 Hillebrand, H., 2004. On the generality of the latitudinal diversity gradient. *The American*
553 *Naturalist*. 163, 192-211. <https://doi.org/10.1086/381004>
- 554 Hilmi, K., Koutitonsky, V.G., Orbi, A., Lakhdar, J.I., Chagdali, M., 2005. Oualidia lagoon,
555 Morocco: An estuary without a river. *African Journal of Aquatic Science*. 30 (1), 1-10.
556 <https://doi.org/10.2989/16085910509503828>
- 557 Hubbell, S.P., 2001. The unified neutral theory of biodiversity and biogeography. Princeton
558 University Press, Princeton, 448 pp.

- 559 Hurlbert, A.H., Stegen, J.C., 2014. On the processes generating latitudinal richness gradients :
 560 identifying diagnostic patterns and predictions. *Evol Popul Genet.* 5: 420.
 561 <https://doi.org/10.3389/fgene.2014.00420>
- 562 Hurtt, G.C., Pacala, S.W., 1995. The consequences of recruitment limitation. Reconciling chance,
 563 history and competitive differences between plants. *Journal of Theoretical Biology.* 176, 1-12.
 564 <https://doi.org/10.1006/jtbi.1995.0170>
- 565 Joulami, L., 2013. Interactions entre les conditions écologiques, le benthos et les communautés
 566 des limicoles dans un écosystème côtier du Nord-Ouest de L'Afrique (Lagune de Sidi Moussa,
 567 Maroc) (Thèse Doctorat.) Université Hassan II. Casablanca, 129 pp.
- 568 Kendall, M.A., Aschan, M., 1993. Latitudinal gradients in the structure of macrobenthic
 569 communities: a comparison of Arctic, temperate and tropical sites. *Journal of Experimental Marine*
 570 *Biology and Ecology.* 172, 157–169. [https://doi.org/10.1016/0022-0981\(93\)90095-6](https://doi.org/10.1016/0022-0981(93)90095-6)
- 571 Khalki, A.E., Moncef, M., 2007. Etude du peuplement de copépodes de l'estuaire de l'Oum Er
 572 Rbia (côte Atlantique du Maroc): Effets des marées et des lachers de barrages. *Lebanese Science*
 573 *Journal.* 8 (1), 3-18.
- 574 Kinlock, N. L., Prowant, L., Herstoff, E. M., Foley, C. M., Akin-Fajiye, M., Bender, N., Umarani,
 575 M., Ryu, H.Y., Şen, B., Gurevitch, J., 2018. Explaining global variation in the latitudinal diversity
 576 gradient: Meta-analysis confirms known patterns and uncovers new ones. *Global Ecology and*
 577 *Biogeography.* 27, 125-141. <https://doi.org/10.1111/geb.12665>.
- 578 Kinlock, N. L., Prowant, L., Herstoff, E. M., Foley, C. M., Akin-Fajiye, M., Bender, N., Umarani,
 579 M., Ryu, H.Y., Şen, B., Gurevitch, J., 2019. Open science and meta-analysis allow for rapid
 580 advances in ecology: A response to Menegotto et al. (2019). *Global Ecology and Biogeography.*
 581 28, 1533-1534. <https://doi.org/10.1111/geb.12964>
- 582 Kong, W.J., Meng, W., Zhang, Y., Christopher, G., Qu, X.D., 2013. A freshwater ecoregion
 583 delineation approach based on freshwater macroinvertebrate community features and spatial

- 584 environmental data in Taizi River Basin, northeastern China. *Ecol. Res.* 28, 581-592.
585 <http://dx.doi.org/10.1007/s11284-013-1048-7>
- 586 Labrune, C., Grémare, A., Amouroux, J.M., Sardá, R., Gil, J., Taboada, S., 2008. Structure and
587 diversity of shallow soft-bottom benthic macrofauna in the Gulf of Lions (NW Mediterranean).
588 *Helgol Mar Res.* 62, 201-214. <https://doi.org/10.1007/s10152-008-0108-9>
- 589 Lacoste, M., 1984. Contribution à l'étude écologique de la lagune de Moulay Boussalham (Maroc)
590 (Thèse Doctorat 3ème cycle). Université Paul Sabatier. Toulouse, 207 pp.
- 591 Lara-Lara, J., Arreola, J., Calderon, L., Camacho, V., De la Lanza, G., Escofet, A., Espejel, M.,
592 Guzman, M., Ladah, L., Lopez, M., Meling, E., Moreno, P., Reyes, H., Rios, E., Zertuche, J., 2008.
593 Los ecosistemas costeros, insulares y epicontinentales. In: Capital natural de México Vol. I :
594 Conocimiento actual de la biodiversidad. Mexico City : CONABIO, 109-134.
- 595 Lefrere, L., 2012. Contribution à l'étude de deux lagunes atlantiques marocaines, Khnifiss et
596 Oualidia: faune malacologique, biologie de trois mollusques bivalves et contamination métallique
597 (Thèse Doctorat). Université Ibn Zohr. Agadir, 180 pp.
- 598 Maanan, M., Zourarah, B., Carruesco, C., Aajjane, A., Naud, J., 2004. The distribution of heavy
599 metals in the Sidi Moussa lagoon sediments (Atlantic Moroccan Coast). *Journal of African Earth*
600 *Sciences.* 39 (3-5), 473-483. <https://doi.org/10.1016/j.jafrearsci.2004.07.017>
- 601 MacArthur, R.H., 1970. Species packing and competitive equilibrium for many species.
602 *Theoretical Population Biology.* 1, 1-11.
- 603 McCallum, A.W., Woolley, S., Błażewicz-Paszkowycz, M., Browne, J., Gerken, S., Kloser, R.,
604 Poore, G.C.B., Staples, D., Syme, A., Taylor, J., Walker-Smith, G., Williams, A., Wilson, R.S.,
605 2015. Productivity enhances benthic species richness along an oligotrophic Indian Ocean
606 continental margin. *Global Ecol. Biogeogr.* 24 (4), 462-471. <http://dx.doi.org/10.1111/geb.12255>

- 607 Menegotto, A., Kurtz, M.N., Lana, P., 2019. Benthic habitats do show a significant latitudinal
608 diversity gradient: a comment on Kinlock et al. (2018). *Global Ecology and Biogeography*. 28,
609 1712–1717. <http://doi.org/10.5281/zenodo.3245484>
- 610 Mergaoui, L., Fekhaoui, M., Bouya, D., Gheit, A., Stambouli, A., 2003. Qualité des eaux et
611 macrofaune benthique d'un milieu estuarien du Maroc ; Cas de l'estuaire de Sebou. *Bulletin de*
612 *l'Institut Scientifique*. 25, 67-75.
- 613 Möllmann, C., Diekmann, R., 2012. Marine ecosystem regime shifts induced by climate and
614 overfishing a review for the northern hemisphere. *Advances in Ecological Research*. 47, 303-347.
615 <https://doi.org/10.1016/B978-0-12-398315-2.00004-1>
- 616 Observatoire National de l'Environnement au Maroc (ONEM)., 1998. Rapport National sur la
617 biodiversité : faune marine, 105pp.
- 618 Piacenza, S. E., Barner, A. K., Benkwitt, C. E., Boersma, K. S., Cerny-Chipman, E. B., Ingeman,
619 K. E., Kindinger, T.L., Lee, J.D., Lindsley, A.J., Reimer, J.N., Rowe, J.C., Shen, C., Thompson,
620 K.A., Thurman, L.L., Heppell, S.S., 2015. Patterns and variation in benthic biodiversity in a large
621 marine ecosystem. *PLOS One*. 10(8): e0135135. <https://doi.org/10.1371/journal.pone.0135135>
- 622 Pierrot-Bults, A.C., 1997. Biological diversity in oceanic macrozooplankton: more than counting
623 species in "Marine Biodiversity: Patterns and Processes" (eds R.F.G. Ormond, J.D. Gage & M.V.
624 Angel), pp. 69–93. Cambridge University Press, Cambridge, UK.
625 <https://doi.org/10.1017/CBO9780511752360>
- 626 Poore, G.C.B., Wilson, G.D.F., 1993. Marine species richness. *Nature*. 361, 597-598.
627 <https://doi.org/10.1038/361597a0>
- 628 Pratt, D.R., Pilditch, C.A., Lohrer, A.M., Thrush, S.F., 2014. The effects of short-term increases
629 in turbidity on sandflat microphytobenthic productivity and nutrient fluxes. *J. Sea Res.* 92, 170-
630 177. <https://doi.org/10.1016/j.seares.2013.07.009>

- 631 Qninba, A., Radi, M., Benhoussa, A., Bazairi, H., Menioui, M., 2003. Fiche descriptive sur les
632 zones humides Ramsar (FDR). Catégories approuvées dans la Recommandation 4.7 modifiée par
633 la résolution VIII.13 de la Conférence des Parties contractantes 2005, 9p.
- 634 Renaud, P.E., Webb, T.J., Bjørgesæter, A., Karakassis, I., Kędra, M., Kendall, M.A., Labrune, C.,
635 Lampadariou, N., Somerfield, P.J., Włodarska-Kowalczyk, M., Vanden Berghe, E., Claus, S.,
636 Aleffi, I.F., Amouroux, J.M., Bryne, K.H., Cochrane, S.J., Dahle, S., Degraer, S., Denisenko, G.,
637 Deprez, T., Dounas, C., Fleischer, D., Gil J., Grémare, A., Janas, U., Mackie, A.S.Y., Palerud, R.,
638 Rumohr, H., Sardá, R., Speybroeck, J., Taboada, S., Van Hoey, G., Węśławski, J.M., Whomersley,
639 P., Zettler, M.L., 2009. Continental-scale patterns in benthic invertebrate diversity: insights from
640 the MacroBen database. *Marine Ecology Progress Series*. 382, 239-252.
641 <https://doi.org/10.3354/meps07963>
- 642 Rex, M.A., Stuart, C.T., Coyne, G., 2000. Latitudinal gradients of species richness in the deep-sea
643 benthos of the North Atlantic. *Proceedings of the National Academy of Sciences of the USA*. 97,
644 4082-4085. <https://doi.org/10.1073/pnas.050589497>
- 645 Rex, M.A., Stuart, C.T., Hessler, R.R., Allen, J.A., Sanders, H.L., Wilson, G.D.F., 1993. Global-
646 scale latitudinal patterns of species diversity in the deep-sea benthos. *Nature*. 365, 636-639.
647 <https://doi.org/10.1038/365636a0>
- 648 Rohde, K., 1992. Latitudinal gradients in species diversity: the search for the primary cause. *Oikos*.
649 65, 514-527.
- 650 Roy, K., Jablonski, D., Valentine, J.W., 2000. Dissecting latitudinal diversity gradients: functional
651 groups and clades of marine bivalves. *Proceedings of the Royal Society of London B. Biological*
652 *Sciences*. 267, 293–299. <https://doi.org/10.1098/rspb.2000.0999>
- 653 Roy, K., Jablonski, D., Valentine, J.W., Rosenberg, G., 1998. Marine latitudinal diversity
654 gradients: tests of causal hypotheses. *Proceedings of the National Academy of Sciences USA*. 95,
655 3699-3702. <https://doi.org/10.1073/pnas.95.7.3699>

- 656 Sanders, H.L., 1968. Marine benthic diversity: a comparative study. *Am Nat.* 102, 243-282.
657 <https://doi.org/10.1086/282541>
- 658 Semlali, A., Chafik, A., Talbi, M., Budzinski, H., 2012. Origin and Distribution of Polycyclic
659 Aromatic Hydrocarbons in Lagoon Ecosystems of Morocco. *The Open Environmental Pollution*
660 *& Toxicology Journal.* 3, 37-46. <https://doi.org/10.2174/1876397901203010037>
- 661 Snelgrove, P.V., 1998. The biodiversity of macrofaunal organisms in marine sediments.
662 *Biodiversity and Conservation.* 7, 1123-1132. <https://doi.org/10.1023/A:1008867313340>
- 663 Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern,
664 B.S., Jorge, M.A., Lombana, A., Lourie, S.A., Martin, K.D., McManus, E., Molnar, J., Recchia,
665 C.A., Robertson, J., 2007. Marine ecoregions of the world: A bioregionalization of coastal and
666 shelf areas. *Bioscience.* 57, 573-583. <https://doi.org/10.1641/B570707>
- 667 Thorson, G., 1957. Bottom communities. In *Treatise on Marine Ecology and Paleoecology.* Vol.
668 1 Ecology (ed. G W Hedgepeth). *Memoir of the Geological Society of America.* 67, 461-534.
669 <https://doi.org/10.1130/MEM67V1>
- 670 Touhami, F., 2018. Caractérisation des peuplements benthiques des habitats intertidaux de Merja
671 Zerga et leur utilisation par les limicoles hivernants (Thèse Doctorat). Université Mohammed V
672 de Rabat. 219pp.
- 673 Touhami, F., Bazairi, H., Badaoui, B., Morabbi, A., Benhoussa, A., 2019. Structure and spatial
674 organization of macrobenthic fauna of intertidal habitats frequented by wintering shorebirds at
675 Merja Zerga lagoon (Moroccan Ramsar Site). *Cah. Biol. Mar.* 60, 41-50. <https://doi.org/10.21411/CBM.A.26046E89>
- 677 UN General Assembly, Convention of the Law of the Sea, 10 December., 1982. Available at:
678 <https://www.refworld.org/docid/3dd8fd1b4.html> (accessed 9 April 2019).
- 679 Veiga, P., Torres, A.C., Aneiros, F., Sousa-Pinto, I., Troncoso, J.S., Rubal M., 2016. Consistent
680 patterns of variation in macrobenthic assemblages and environmental variables over multiple

spatial scales using taxonomic and functional approaches. Mar. Environ. Res. 120, 191-201.
<http://dx.doi.org/10.1016/j.marenvres.2016.08.011>

Willig, M.R., Kaufman, D.M., Stevens, R.D., 2003. Latitudinal gradients of biodiversity: pattern, process, scale, and synthesis. Annual Review of Ecology, Evolution and Systematics. 34, 273-309.
<https://doi.org/10.1146/annurev.ecolsys.34.012103.144032>

WoRMS Editorial Board., 2020. World Register of Marine Species. Available from
<http://marinespecies.org/> (consulted January 14, 2020).

Zidane, H., Mannan, M., Mouradi, A., Maanan, M., El Barjy, M., Zourarah, B., Blais, J.F., 2017. Environmental and ecological risk of heavy metals in the marine sediment from Dakhla Bay, Morocco. Environmental Science and Pollution Research. 24, 7970-7981.
<https://doi.org/10.1007/s11356-017-8367-0>

Zine, N.E., 1989. Etude de la malacofaune de la lagune de Nador et dynamique de population de *Venerupis decussata* (Linné 1767) (Thèse Doctorat 3ème cycle). Université Mohammed V. Rabat, 97 pp.

Zine, N.E., 2005. Etude éco-biologique et dynamique des populations de la palourde (*Ruditapes decussatus*) et analyse typologique de la faune benthique de milieux paraliques atlantico-méditerranéens du Maroc (Thèse Doctorat d'Etat). Université Moulay Ismail. Meknès, 200 pp.

Table 1

Environment descriptors of the semi-enclosed coastal systems of Morocco in terms of aquatic system type (1 = lagoon; 2 = estuary; 3 = bay), latitude (LAT, N), longitude (LON, W), surface area (km²), maximum annual water temperature (M Temp, °C), minimum annual water temperature (m Temp, °C), maximum annual water salinity (M Sal), minimum annual water salinity (m Sal). NA: Nador, SM: Smir, TA: Tahaddart, LO: Loukkos, MB: Moulay Bousselham, SE: Sebou, BR: Bouregreg, OR: Oum Rbia, SI: Sidi Moussa, OU: Oualidia, KH: Khnifiss, DA: Dakhla.

System	Type	LAT	LON	Surface Area	M Temp	m Temp	M Sal	m Sal	Reference
Nador (NA)	1	35°10'	02°51'	115	28	14	38	32	El Kamch et al. (2020)
Smir (SM)	1	35°42'	05°20'	0.3	32	12	41	7.8	Chaouti & Bayed 2005
Tahaddart (TA)	2	35°46'	05°42'	10	26	13	41	21	Achab 2011
Loukkos (LO)	2	35°07'	06°00'	72	27	15	34	22	Geawhari et al., 2014
Moulay Bousselham (MB)	1	34°51'	06°16'	27	28	11	35	27	Gam et al., 2010
Sebou (SE)	2	34°16'	06°39'	17.5	30	16	35	12	Haddout et al., 2015
Bouregreg (BR)	2	34°	06° 50 '	4000	45	14	30	10	Cherkaoui 2006; El Amraoui et al., 2015
Oum Rbia (OR)	2	33°28'	08°34'	1.5	25	15	35	30	Khalki & Moncef 2007
Sidi Moussa (SI)	1	32°54'	08°49'	4.2	27	15	33	22	Maanan et al., 2004
Oualidia (OU)	1	32°45'	08°30'	3.0	21	16	36	28	Hilmi et al., 2005
Khnifiss (KH)	1	28°03'	12°15'	65	22	16	38	34	Semlali et al., 2012
Dakhla (DA)	3	23°45'	15°50'	400	26	14	40	37	Zidane et al., 2018

Table 2

Species richness (S) and taxonomic distinctness indices values in the semi-enclosed coastal ecosystems of Morocco. Δ^+ : taxonomic distinctness, $S\Delta^+$: total taxonomic distinctness, Φ^+ : average phylogenetic diversity, Λ^+ : variation in taxonomic distinctness, $S\Phi^+$: total phylogenetic diversity. NA: Nador, SM: Smir, TA: Tahaddart, LO: Loukkos, MB: Moulay Bousselham, SE: Sebou, BR: Bouregreg, OR: Oum Rbia, SI: Sidi Moussa, OU: Oualidia, KH: Khnifiss, DA: Dakhla.

Sites	Sampling effort	S	Δ^+	$S\Delta^+$	Λ^+	Φ^+	$S\Phi^+$
NA	366	158	90	14232	279.9	47.7	7533.3
SM	123	53	89.6	4748.1	293.3	56.3	2983.3
TA	99	40	88.7	3549.6	311.6	55.0	2200.0
LO	134	32	89.1	2852.7	300.5	60.4	1933.3
MB	599	161	90.5	14569.2	268.7	47.8	7700.0
SE	7	94	87.8	8162.0	275.5	52.9	4916.7
BR	120	101	89.9	9075.7	278.3	51.7	5216.7
OR	39	48	88.5	4247.5	328.8	51.4	2466.7
SI	117	57	90.2	5141.7	271.3	56.4	3216.7
OU	43	105	89.8	9428.2	268.2	48.6	5100.0
KH	29	62	88.6	5491.8	333.7	51.9	3216.7
DA	100	105	89.7	9414.4	255.7	50.6	5316.7

Table 3. Results of PERMANOVAs testing for the effect of the type of ecosystem (Lagoon - Estuary) on species richness and taxonomic.

Source	df	MS	Pseudo-F	P (perm)	df	MS	Pseudo-F	P (perm)
<i>Species richness (S)</i>					<i>Taxonomic distinctness (A⁺)</i>			
Type	1	312,26	1,9692	0,1845	1	0,20702	5,188	0,0528
Residual	9	158,57			9	3,99E-02		
Total	10				10			
<i>Total taxonomic distinctness (SA⁺)</i>					<i>Variation in taxonomic distinctness (A⁺)</i>			
Type	1	330,52	2,076	0,1825	1	3,5385	0,87669	0,3505
Residual	9	159,21			9	4,0363		
Total	10				10			
<i>Average phylogenetic diversity (Φ⁺)</i>					<i>Total phylogenetic diversity (SΦ⁺)</i>			
Type	1	5,0236	1,4911	0,2565	1	251,47	2,0287	0,1876
Residual	9	3,3689			9	123,95		
Total	10				10			

Table 4

Gaps and knowledge gained from studies performed on soft-bottom zoomacroenthos of the Moroccan SECS. NA: Nador. SM: Smir. TA: Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR: Bouregreg. OR: Oum Rbia. SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla. **I** = Intertidal **S**= Subtidal. **B** = Biomass. References (Numbers 1-36): **(1)** Elkaïm (1974); **(2)** Lacoste (1984); **(3)** Bekkali (1987); **(4)** Guelorget et al. (1987); **(5)** Bayed et al. (1988); **(6)** Cheggour (1988); **(7)** Zine (1989); **(8)** Chbicheb (1996); **(9)** Aksissou (1997); **(10)** Bazairi (1999); **(11)** Boussalwa et al. (2000); **(12)** Bazairi & Zourarah (2001); **(13)** Mergaoui et al. (2003); **(14)** Bazairi & Gam (2004); **(15)** Chaouti & Bayed (2005); **(16)** El Houssaini (2005); **(17)** Zine (2005); **(18)** Azirar (2006); **(19)** Bazairi & Bayed (2006); **(20)** Cherkaoui (2006); **(21)** Gauteur (2006); **(22)** Bazairi & Zourarah (2007); **(23)** Ait Mlik (2009); **(24)** Lefrere (2012); **(25)** Bououarour (2013); **(26)** Joulami (2013); **(27)** Boutahar (2014); **(28)** Cuvelier et al. (2014); **(29)** El Asri et al. (2015); **(30)** Bazairi et al. (2017); **(31)** El Asri et al. (2017); **(32)** Touhami (2018); **(33)** El Asri (2019); **(34)** Bououarour (unpublished data); **(35)** Boutoumit (unpublished data); **(36)** El Kamcha (unpublished data). Grey colour indicates studies that can be considered as references for respective sites.

	Mediterranean		Atlantic									
	NA	SM	TA	LO	MB	SE	BO	OR	SI	OU	KH	DA
1974							1, IS					
1984					2, I							
1987	4, S, B	3, S										
1988							6, I				5, I	
1989	7, S											
1995												
1996										8, I		
1997		9, I										
1999					10, IS							
2000	11, S											
2001								12, IS, B				
2003						13, S						
2004				14, IS, B								
2005		15, I	16, I			17, IS						
2006			19, IS				20, IS			21, IS		
2007								22, IS, B				
2009					23, IS							
2012											24, I	
2013					25, I				26, I, B			
2014	28, S									27, IS		
2015										29, IS		
2016												
2017								30, IS, B		31, IS		
2018					32, I, B							
2019										33, IS		33, IS
2020	36, S, B			34, I, B	34, I, B	35, IS, B			34, I, B	34, I, B	34, I, B	34, I, B

742

743 **Figure Captions:**

744 **Figure 1.** Map showing the geographical position of the semi-enclosed coastal systems of
 745 Morocco considered in this study. ★: lagoon; ▲: estuary; ■: bay.

746 **Figure 2.** Species richness by biogeographical repartition in the semi-enclosed coastal systems of
 747 Morocco. A: Atlantic. AM: Atlantico-Mediterranean. and M: Mediterranean. NA: Nador. SM:
 748 Smir. TA: Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR: Bouregreg. OR:
 749 Oum Rbia. SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla.

750 **Figure 3.** Confidence funnel (mean and 95% confidence interval) of the variation in taxonomic
 751 distinctness (A) and taxonomic distinctness (B) in the Moroccan Semi-enclosed coastal systems.
 752 NA: Nador. SM: Smir. TA: Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR:
 753 Bouregreg. OR: Oum Rbia. SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla.

754 **Figure 4.** Dendrogram of cluster analysis using group-average linkage of Bray-Curtis similarities
 755 based on benthic macrofauna composition in the semi-enclosed coastal systems of Morocco. NA:
 756 Nador. SM: Smir. TA: Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR:
 757 Bouregreg. OR: Oum Rbia. SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla.

758 **Figure 5.** Ordination of the semi-enclosed coastal systems using the Principal coordinates
 759 ordination (PCO) with vectors (longer than 0.7) and clusters overlay. NA: Nador. SM: Smir. TA:
 760 Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR: Bouregreg. OR: Oum Rbia.
 761 SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla.

762 **Figure 6.** Distance-based redundancy analysis plot and the correlated variables that explained
 763 the semi-enclosed coastal systems distribution based on benthic macrofauna composition. NA:
 764 Nador. SM: Smir. TA: Tahaddart. LO: Loukkos. MB: Moulay Bousselham. SE: Sebou. BR:
 765 Bouregreg. OR: Oum Rbia. SI: Sidi Moussa. OU: Oualidia. KH: Khnifiss. DA: Dakhla.

766

768 **Figure 1**

773 **Figure 2**

785 **Figure 3**

Figure 4

Figure 5

Figure 6

A

B

- The first broad-scale baseline of soft-bottom macrozoobenthos in semi-enclosed coastal systems (SECS) of Morocco is presented
- 496 species were recorded from the Moroccan SECS and belong to 7 phyla, 21 classes, 65 orders and 201 families
- Species richness and taxonomic diversity indices showed no relationship with SECS features including latitude
- The type of the SECS (estuaries vs lagoons vs bay), the marine ecoregion (Atlantic vs Mediterranean), the surface of the SECS and the environmental features (minimal temperature, minimal and maximal salinity) are the most predictors of benthic macrofauna composition in the Moroccan SECS

Author statement file:

Soilam Boutoumit: Investigation, Data curation, Writing – Original Draft, Visualisation. **Reda El Kamcha:** Investigation, Data curation, Writing – Original Draft, Visualisation. **Oussama Bououarour:** Investigation, Data curation, Writing – Original Draft, Visualisation. **Latifa Joulami:** Investigation, Data curation, Review. **Boutahar Loubna:** Investigation, Data curation, Review. **Abdelaziz Benhoussa:** Resources, Funding acquisition. **Mohamed Maanan:** Review & Editing, Supervision. **Laurent Godet:** Writing-Review & Editing, Validation. **Abdellatif Bayed:** Resources, Validation. **Hocein Bazairi:** Conceptualization, Methodology, Writing – Original Draft, Review & Editing, Supervision.

Declaration of interests

The authors (Soilam Boutoumit, Reda El Kamcha, Oussama Bououraour, Latifa Joulami, Loubna Boutahar, Abdelaziz Benhoussa, Mohamed Maanan, Laurent Godet, Abdellatif Bayed and Hocein Bazairi) declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in the paper entitled **"Soft-bottom macrozoobenthos in semi-enclosed coastal systems of Morocco"**

In behalf of all authors: Hocein Bazairi

Laboratoire de Biodiversité,
Ecologie et Génome

A handwritten signature in blue ink, appearing to read "Hocein", enclosed within a blue oval shape.

Pr Hocein Bazairi