Unexpected abundance and long-term relative stability of the brown alga *Cystoseira amentacea*, hitherto regarded as a threatened species, in the north-western Mediterranean Sea

Thierry Thibaut^{a,*}, Aurélie Blanfuné^a, Laurent Markovic^b, Marc Verlaque^a, Charles F. Boudouresque^a, Michèle Perret-Boudouresque^a, Vesna Mačić^c, Lorraine Bottin^d

^a Aix-Marseille University, and Toulon University, Mediterranean Institute of Oceanography (MIO), CNRS/INSU, IRD, UM 110, Campus of Luminy, 13288 Marseille cedex 9, France

^b European Commission, Directorate-General for Maritime Affairs and Fisheries, Avenue Joseph-II, 79 office 02/67, 1049 Brussels, Belgium

^c Institut za Biologiju Mora, Rukovodilac laboratorije, Naučni saradnik, P.Fah 69, 85330 Kotor, Montenegro

^d Université Nice-Sophia-Antipolis, EA 4228 ECOMERS, Faculté des Sciences, Parc Valrose, 06108 Nice cedex 2, France

ABSTRACT

Cystoseira amentacea is a Mediterranean endemic alga thriving on very shallow rocky substrates. It has been considered as a threatened species, having experienced a steady decline and is therefore protected by international conventions. The historical distribution of the species has been assessed along the French Mediterranean coast, on the basis of 467 articles and herbarium vouchers. We have produced an accurate map of its current distribution and abundance along 1832 km of coastline, through *in situ* surveys. *C. amentacea* was observed along 1125 km of shoreline, including 33% of almost continuous or continuous belt. In most of its range, there is no evidence of loss, except in 4 areas of Provence, French Riviera and Corsica. A significant relation was found between the absence or low abundance of *C. amentacea* and the vicinity of ports and large sewage outfalls. The status of conservation of the species should therefore be reassessed.

Keywords:

Conservation status
Cystoseira amentacea
Human pressure
Mapping
Mediterranean Sea
Long-term study

1. Introduction

The coastal ecosystems are highly impacted all over the world, due to the cumulative impact of increasing human pressure (e.g. destruction of habitats, pollution, species introduction, overfishing, coastal aquaculture and global warming). None of the forms of impact affecting the coastal ecosystems exists in isolation. Different pressures act over time and in unison, with a possible synergy effect, to affect the species, the ecosystems and their ability to deliver ecosystem services (e.g. Worm et al., 2006; Halpern et al., 2008; Waycott et al., 2009). The Mediterranean Sea is a hotspot of marine biodiversity that is under siege due to high demographic pressure, a high percentage of worldwide shipping and tourism and the highest rate of biological invasions (Bianchi and Morri, 2000; Galil, 2000; Boudouresque and Verlaque, 2002; Lotze et al., 2006; Coll et al., 2010; Lejeune et al., 2010; Zenetos et al., 2010; UNEP/MAP, 2012; Bianchi et al., 2014).

Along the temperate rocky coasts, the large canopy-forming kelps (Laminariales, Phaeophyceae, Ochrophyta) and fucoids

(Fucales, Phaeophyceae, Ochrophyta) represent the dominant species in pristine environment (Dayton, 1985; Steneck et al., 2002; Schiel and Foster, 2006). They provide shelter, food, habitat and nurseries to a multitude of species; they provide high primary production involved in the maintaining of diversified trophic levels; the largest species attenuate wave action (Steneck et al., 2002). These seaweeds can be controlled by a top-down mechanism mainly in the case of the sublittoral species, while the subsurface species are controlled by a bottom-up mechanism (Hereu et al., 2008; Cardona et al., 2013). The decline of kelps and fucoids is a worldwide phenomenon due, directly or indirectly, to human activities (Steneck et al., 2002; Díez et al., 2003; Helmuth et al., 2006; Worm and Lotze, 2006; Airoldi and Beck, 2007; Hawkins et al., 2008; Wernberg et al., 2010; Schiel, 2011; Lamela-Silvarrey et al., 2012; Raybaud et al., 2013; Filbee-Dexter and Scheibling, 2014). Some taxa, have been driven to regional extinction (Thibaut et al., 2005). These changes are leading to shifts in habitat structure from a state with canopy forming species to alternative states, in the worst cases to barren grounds composed of filamentous and encrusting species (Micheli et al., 2005; Perkol-Finkel and Airoldi, 2010; Sala et al., 2011, 2012; Parravicini et al., 2013; Filbee-Dexter and Scheibling, 2014).

In the Mediterranean Sea, species of the genus *Cystoseira* C. Agardh are habitat-forming species dominating several benthic assemblages from the littoral fringe down to the lower sublittoral zone (Feldmann, 1937; Molinier, 1960; Pignatti, 1962; Verlaque, 1987; Ballesteros, 1988, 1990a,b; Giaccone et al., 1994). Their zonation is dependent on different environmental conditions (light, temperature, hydrodynamism and grazing) (Sauvageau, 1912; Ollivier, 1929; Vergès et al., 2009). Losses of fucoids have been reported all around the Mediterranean Sea caused by habitat destruction, eutrophication and overgrazing by herbivores leading to a shift to less structural complexity, such as turf-forming, filamentous or other ephemeral seaweeds or barrens grounds where urchin density is a driver of habitat homogenisation (Munda, 1974, 1982, 1993; Thibaut et al., 2005; Devescovi and Ivesa, 2007; Airoidi et al., 2008; Falace et al., 2010; Frascchetti et al., 2011; Giakoumi et al., 2012; Sala et al., 2012; Bianchi et al., 2014; Templado, 2014).

Seaweeds living in the rocky intertidal zone or in the sublittoral fringe can endure hard ecological conditions (desiccation, high variation of temperature, salinity and insolation, strong wave actions, etc.).

Among the 51 taxonomically accepted taxa of the genus *Cystoseira* (Guiry and Guiry, 2014), 36 are present in the Mediterranean Sea, 30 are endemic to the Mediterranean, and only 3 live exclusively in the Mediterranean littoral fringe (i.e. the upper level of the sublittoral, from the mean sea level down to less than 1 m depth): *Cystoseira mediterranea* Sauvageau in the western basin, *Cystoseira tamarascifolia* (Hudson) Papenfuss in the Alboran Sea and *Cystoseira amentacea* (C. Agardh) Bory de Saint-Vincent in all the Mediterranean; two poorly characterised varieties var. *amentacea* and var. *stricta* Montagne, are sometimes considered (Ribera et al., 1992; Cormaci et al., 2012; Guiry and Guiry, 2014) but they will not be distinguished here. Because of their shallow habitat, these species escape overgrazing by most of the large herbivores, e.g. the sea urchins *Paracentrotus lividus* (Lamarck 1816) and *Arbacia lixula* (Linnaeus 1758) and the teleost *Sarpa salpa* (Linnaeus 1758) unlike the others subtidal fucoids (Cormaci and Furnari, 1999; Thibaut et al., 2005; Serio et al., 2006; Vergès et al., 2009). All these sub-surface species are considered as vulnerable to sea surface pollution, habitat destruction, trampling and a number of regression events have been reported all around the Mediterranean Sea (Bellan-Santini, 1968b; Desrosiers et al., 1982, 1986; Cecere et al., 1996; Rodríguez-Prieto and Polo, 1996; Cormaci et al., 2001; Soltan et al., 2001; Milazzo et al., 2002; Thibaut et al., 2005; Díez et al., 2009; Arévalo et al., 2007; Falace et al., 2010; Sales et al., 2011; Tsiamis et al., 2013). These taxa are also facing the low frequency natural disturbances called in Spanish 'minva' (very low tide for a long period of time, due to meteorological events), leading to mass mortality of the shallowest *Cystoseira* species (Huvé, 1970; Rodríguez-Prieto, 1992). In disturbed areas, *Cystoseira* species can be replaced by dense stands of fast growing *Corallina* spp. (Corallinales, Rhodophytes) and/or the mussel *Mytilus galloprovincialis* Lamarck, 1819 (Mytiloidea, molluscs), which become dominant, forming extensive and wide belts along the shore (Berner, 1931; Bellan-Santini, 1964, 1965, 1968a,b; Huvé, 1970; Arnoux and Bellan-Santini, 1972; Bellan and Bellan-Santini, 1972; Astier, 1975; Augier, 1977; Fernandez and Niell, 1982; Desrosiers et al., 1982, 1986; Thomas, 1983; Janssens et al., 1993; Soltan et al., 2001; Arévalo et al., 2007; Maggi et al., 2009; Falace et al., 2010). The interactions between *Cystoseira* species, turf-forming algae, herbivores and associated species have been extensively studied at different spatial and temporal scales and levels of protection (Benedetti-Cecchi and Cinelli, 1992a,b, 1995; Menconi et al., 1999; Ruitton et al., 2000; Benedetti-Cecchi, 2001; Benedetti-Cecchi et al., 2001; Bulleri et al., 2002; Maggi et al., 2009, 2012; Frascchetti et al., 2012; Tamburello et al.,

2013). *C. amentacea* can also be replaced by the tolerant *Cystoseira compressa* (Esper) Gerloff and Nizamuddin (Mangialajo et al., 2012).

Along the whole of the French Mediterranean coasts, two species are restricted to the upper sublittoral zone, *C. mediterranea* in the western part along the French Catalan coast (Sauvageau, 1912; Feldmann, 1937), and *C. amentacea* in Provence, French Riviera and Corsica. The latter taxon is widely distributed in the Mediterranean (Ribera et al., 1992). A non-Mediterranean record, from Sudan (East Africa) is regarded as a misidentification (Papenfuss, 1968). *C. amentacea* possesses a 40 cm high caespitose thallus and a basal holdfast a few centimetres wide. The species forms extensive stands on shallow rocky wave-exposed zones (Agardh, 1842; Molinier, 1960; Boudouresque, 1971a,b). As thalli are negatively buoyant and the maximum distance for egg dissemination is low (~few dozen cm, Mangialajo et al., 2012), the species is suspected of having low dispersal ability. *C. amentacea* is one of the most productive Mediterranean seaweeds (Bellan-Santini, 1963; Bellan and Bellan-Santini, 1972; Susini, 2006), because of steady exposure to turbulence (frequent emersion/immersion), which maintains a high nutrient uptake (Epiard-Lahaye et al., 1987).

Because of its habitat preferences, *C. amentacea* has been considered as one of the species most severely impacted by coastal development and sewage outfalls (Bellan-Santini, 1964, 1965, 1966, 1968b; Bellan and Pérès, 1970; Arnoux and Bellan-Santini, 1972; Bellan and Bellan-Santini, 1972; Astier, 1975; Belsher, 1977; Desrosiers et al., 1982, 1986; Thomas, 1983; Janssens et al., 1993; Sinnassamy et al., 1993; Soltan et al., 2001; Bulleri et al., 2002; Falace et al., 2010). *C. amentacea* is also harvested for the cosmetics industry (Max Pellegrini, pers. comm.). Furthermore, the species is thought to be vulnerable to oil spills and chemical releases at the sea surface (Bellan-Santini, 1968b). The species is considered to have a high ecological status within the framework of the EU Water-Framework Directive (2000/60/EC) (Orfanidis et al., 2001; Panayotidis et al., 2004; Ballesteros et al., 2007; Mangialajo et al., 2008; Cecchi et al., 2009; Sfriso and Facca, 2011; Nikolic et al., 2013).

C. amentacea is currently listed as a species strictly protected under the Berne Convention (Annex I 1979), together with 4 other taxa (*C. mediterranea*, *C. sedoides*, *C. spinosa*, *C. zosteroides*); all the Mediterranean species of the genus *Cystoseira*, except *C. compressa* have been listed under Annex II of the Barcelona convention (2010). All the Mediterranean *Cystoseira* species are under surveillance by international organizations such as the IUCN, the RAC/ASP and MedPan. Like all the species of the genus *Cystoseira*, *C. amentacea*'s stands are an habitat and listed as Habitat of Community Interest by the UE (Micheli et al., 2013).

The use of historical data in marine ecology is growing, as they provide a baseline of animal and plant populations in very lightly disturbed areas so changes can be analysed over a long time period (Jackson et al., 2001; Thibaut et al., 2005; Sáenz-Arroyo et al., 2005; Barsanti et al., 2007; Babalis, 2011; Lotze et al., 2010; Montefalcone et al., 2013; Husa et al., 2014). Most studies of habitat loss or coastal urbanisation, however, focus only on local scales and short periods over contemporary times. Whilst this focus is also necessary, the preoccupation with these local and present-day drivers means that we are often not aware of the past extension of the species and of the regional context of local studies.

In order to analyse the long-term changes in *C. amentacea* populations along the French Mediterranean coasts, we collected all available data (herbarium vouchers, published and grey literature) and we have compared them to our field surveys. Because most of the data collected over the last centuries are qualitative, the aim of this study was: (i) to give a present exhaustive quantitative map of the distribution of *C. amentacea* along the French Mediterranean

coasts, including Corsica; (ii) to compare the present distribution with historical data; (iii) and to highlight the eventual sites of regression and identify the possible causes.

2. Materials and methods

2.1. Study area

In France, *Cystoseira amentacea* was reported from four regions: the western and eastern Provence, the French Riviera (including the Principality of Monaco) and Corsica (Fig. 1). The species is absent from French Catalonia and Languedoc where it can be replaced on suitable hard substrates by *C. mediterranea*.

2.2. Data collection

2.2.1. Literature and herbaria

Historical data available concerning *C. amentacea* along the French Mediterranean coasts are numerous (Supplementary material Tables S1–S4). The species is easy to identify and to collect and, from the 18–19th to the early 21st century, many naturalists and phycologists surveyed the French Mediterranean coasts. They received the support of local natural history museums and they built an efficient web of collectors providing a large quantities of samples.

In addition to its present taxonomical status (Guiry and Guiry, 2014) *C. amentacea* has been mentioned in the literature and herbariums under several nomenclatural synonyms, with correct or

incorrect authorities: *C. amentacea* (C. Agardh) Bory de St-Vincent, *C. amentacea* var. *stricta* Montagne, *Cystoseira stricta* Sauvageau, *Cystoseira ericoides* (Linnaeus) C. Agardh var. *amentacea* C. Agardh. Its name was also misspelled as *C. amantacea*, *C. amentacea* Agardh and *C. amentacea* Bory. Finally, it was sometimes misidentified as *C. mediterranea* or as *Fucus ericoides* Linnaeus, *Fucus selaginoides* Linnaeus and *Fucus tamarascifolius* Hudson.

Vouchers held in herbaria are a valuable source of data providing a basis for checking the identification of the specimens. We surveyed several thousand of vouchers held in the following herbaria named after *Index Herbariorum*. **AIX**: The Muséum d'Histoire Naturelle d'Aix-en-Provence: Herbarium Berner. **ARAGO**: The herbaria held at Pierre and Marie Curie University – Laboratoire Arago: Herbarium Feldmann. **AV**: The Muséum d'Histoire Naturelle d'Avignon: Herbarium Requier. **HCOM**: The herbaria held at Aix-Marseille University: Herbarium C.F. Boudouresque, Herbarium Berner, Herbarium P. and H. Huvé, Herbarium Saint-Charles, Herbarium T. Thibaut, Herbarium M. Verlaque, **LAU**: The online herbarium of the Musée Botanique de Lausanne: Herbarium Ducommun. **MPU**: The herbaria held at the University of Montpellier: Herbarium Flahaut, Herbarium Général, Herbarium Lenormand, Herbarium Raphélis, Herbarium Salzmänn, Herbarium Schousboe. **Musée Océanographique of Monaco** (not indexed): Herbarium Mouret, Herbarium Prince Albert I. **NICE**: The Muséum d'Histoire Naturelle de Nice: Herbarium Algues Vertes 1890–1910, Herbarium Camous. **Nice-Sophia Antipolis University** (not indexed): Herbarium Meinesz. **PC**: The Muséum National d'Histoire naturelle (MNHN) in Paris: Carnet de Récolte de Feldmann,

Fig. 1. Studied areas.

Herbarium Algues de France non trié, Herbarium J. Arènes, Herbarium Bornet, Herbarium Bory de Saint-Vincent, Herbarium A. Bunge, Herbarium O. Debeaux, Herbarium Fabre, Herbarium Feldmann, Herbarium Général, Herbarium R. Lami, Herbarium Lebel, Herbarium Le Sourd, Herbarium Lamouroux, Herbarium Lenormand, Herbarium Magne, Herbarium Roger Melin, Herbarium Montagne, Herbarium Moquin-Tandon, Herbarium Roussel, Herbarium Sauvageau, Herbarium Schousboe, Herbarium Soleirol, Herbarium Solier, Herbarium Thuret, Herbarium Sébastien Vaillant, Herbarium A. Vickers, Excicata O. Debeaux, Excicata G. Hamel. **TLON**: The Muséum d'Histoire Naturelle de Toulon; Herbarium Mouret, Algues de Provence. **VTA**: The Botanical Garden of Villa Thuret in Antibes: a recently rediscovered collection of vouchers gathered by Gustave Thuret and Edouard Bornet in the 19th century.

2.2.2. Fieldwork

The fieldwork was carried out during four campaigns from April to June in 2008 and 2009 for the continental Mediterranean French coasts and from April to June 2010 and 2011 for Corsica. Late spring is the best season for fieldwork as primary branches of *C. amentacea* are fully developed.

C. amentacea populations were drawn on black and white A3 format aerial photographs from the IGN (French National Institute of Geographical and Forest Information): BD Ortho; Google Earth was also used). The scale was 1:2500. Three people were on board a small boat (length 5 m) moving at low speed (3–6 km h⁻¹) at few metres offshore. Identification of the species is easy and it cannot be confused with any other species, with the exception of *C. mediterranea*. For this reason, every 1-km, the presence of a caespitose base, the easiest to observe distinctive character of *C. amentacea*, absent in *C. mediterranea*, was checked via snorkelling. *C. amentacea* populations were recorded within 6 classes as defined by Ballesteros et al. (2007): C0 = absence, C1 = rare scattered individuals, C2 = abundant scattered individuals, C3 = abundant patches of dense stands, C4 = almost continuous belt and C5 = continuous belt.

For each 50-m sector, corresponding to a class of *C. amentacea* (C0–C5), we also recorded the slope of the substrate in three classes (vertical, sub-vertical, sub-horizontal to horizontal), the height of coast (High coast: vertical cliff >15 m; Low coast: all the other cases, including decimetric blocks), the nature of the substrate (natural or artificial including stone blocks used for jetties, ports, etc.) and when present the dominant taxa and functional groups other than *C. amentacea* (e.g. *Corallina* spp., *C. compressa*, turf algae and the mussel *M. galloprovincialis*).

2.3. GIS analysis

The changes over time of *C. amentacea* (past and present distribution) were analysed on a GIS (Geographical Information System) database (ArcGis10[®]) with Spatial Analyst tools. Each past or present location has been geo-localised. All historical maps have been digitized to fit the same coastal line used for the present distribution (scale 1:2500). Because no ready-to-use coastal line was available, the entire shoreline of the French Mediterranean coasts was drawn at a scale 1:2500.

The current distribution of *C. amentacea* was analysed regarding possible perturbations (i.e. sewage outfalls, fish farms, vicinity of ports). Although the general pattern of currents in the study area is well known, with the northern Mediterranean current flowing westward along the continental French coast (Millot, 1987; Albérola and Millot, 2003; Taupier-Letage et al., 2013), the near shore local currents are more variable in direction and intensity,

depending upon wind, swell, shore topography, bathymetry and season. For this reason, we drew a 500 m radius circular buffer from the centre of each source of possible perturbation and the *C. amentacea* populations and accompanying species were analysed within this area. In the studied area, are located 275 ports, 43 sewage outfalls and 13 fish farms (Supplementary material Table S5). Ports localised on a rocky shore (193 ports out of 275) were taken into consideration (Supplementary material Table S6), and classified according to their size, i.e. the surface area of the enclosed water body. As regards fish farms, the parameter considered was their distance to the nearest shore, since reared fish volume, a proxy of the pollution generated, was not available. As regards the sewage outfalls, only those where the pipe end is located at less than 500 m from a rocky shore (20 outfalls out of 43) were considered, as their effluents are prone to reach the shore and to impact coastal shallow assemblages (Supplementary material Table S7). The study parameter is the effluent volume (Agence de l'Eau Rhône Méditerranée Corse, 2014). It is worth noting that all these outfalls come from sewage treatment plants.

2.4. Statistical analysis

We tested the influence of (i) the size of ports (surface area, ha), (ii) the mean effluent flow (m³ d⁻¹) of sewage outfalls and (iii) the distance from the coast of fish cages on the abundance of *C. amentacea* and associated species along the natural coast and suitable substrate. Effects on assemblages (*C. amentacea*, and dominant taxon or functional group) composition were first studied through a multivariate exploratory approach using a non-metric multi-dimensional scaling (nMDS) (Clarke and Gorley, 2006). Similarity measure matrixes were calculated from the initial data matrix containing for each sample the percentage of coast occupied by each taxon or functional group, after square root transformation. The chosen similarity measure was the S17 Bray Curtis similarity, (Clarke and Warwick, 1994; Clarke and Gorley, 2006); nMDS represents samples as points in low-dimensional space such as the relative distances apart of all points are in the same rank order as the relative similarities of the samples. Finally, for each taxon or functional group, correlations of taxa-specific abundance with the 2-D ordination plot of samples were plotted by displaying correlation vectors. Spearman correlation was used given its non-parametric properties.

In order to compare assemblage composition as a function of the various factors of our design, we performed a PERMutational multivariate ANalysis Of VAriance (PERMANOVA) (Anderson, 2001) based on the S17 Bray Curtis similarity matrixes, as for the nMDS. *P*-values were obtained by 999 permutations of residuals under a reduced model. Monte Carlo *P*-values were considered when there were not enough possible permutations (<200). In order to compare assemblage abundances between levels of factors we performed the PERMANOVA on the model including terms and all interactions (Underwood, 1981).

Three separate models were successively applied, in order to test the effect of each factor was fit to assemblage abundances in order to test its response to 'size of port', 'effluent flow' and 'cages distance'. Factor 'size of port' has 4 levels (surface area: <2 ha, between 2.1 and 6 ha, between 6.1 and 100 ha, >101 ha) and is fixed; Factor 'effluent flow' has 5 levels (<100 m³ d⁻¹, between 100 and 1000 m³ d⁻¹, between 1000 and 10000 m³ d⁻¹, between 10000 m³ d⁻¹ and 100000 m³ d⁻¹, >100000 m³ d⁻¹ and is fixed; Factor 'cages distances' has 3 levels (<100 m, between 100 and 200 m, between 200 and 500 m) and is fixed. Since ecological data are by nature highly variable, terms were considered significant for

P-values < 0.01. Post-hoc pair-wise analysis were performed when necessary.

3. Results

3.1. Historical data

The first putative mention of *C. amentacea* in the literature dates back to the Roman naturalist Pliny the Elder (23–70 CE; CE means Common Era) in his *Naturalis Historiae* (liber XIII, 48–168 and 170) where [Fée \(1833\)](#) interpreted the description of a Mediterranean marine plant as corresponding to *C. ericoides*. This name is currently regarded as a taxonomic synonym of *Cystoseira tamariscifolia* (Hudson) Papenfuss ([Guiry and Guiry, 2014](#)), but we can hypothesise that [Fée \(1833\)](#) was actually thinking of the current *C. amentacea*.

467 Historical records of *C. amentacea* have been found from the 18th to the 21st centuries ([Table 1](#), [Figs. 2A–6A](#)).

3.1.1. 18th century

We found the oldest specimen of *C. amentacea* in the Sébastien Vaillant herbarium held in PC in Paris, identified as *F. selaginoides*, collected in Marseilles in 1700, this specimen was most probably collected in March of that year by Joseph Pitton de Tournefort while waiting for a boat in Marseilles for his expedition to the Levantine coasts ([Tournefort, 1717](#)). We also found three non-dated samples of *C. amentacea* collected in Toulon (as *Fucus ericoides*) and one in Marseilles (as *F. selaginoides*) in two bundles of vouchers from the 18th century (respectively herbarium Roussel and herbarium Lamouroux, both also held in PC).

3.1.2. 19th century

The 19th century was the golden age for the collection of *Cystoseira* along the French Mediterranean coasts, with famous botanists such as Jacob Georg Agardh (1813–1901), Edouard Bornet (1828–1911), Jean-Baptiste Bory de Saint-Vincent (1778–1846), Jean-Louis Martin Castagne (1785–1858); Jean Odon Debeaux (1826–1910), Camille Montagne (1784–1866), Jean Vincent Felix Lamouroux (1779–1825), Antoine Risso (1777–1845), Camille Sauvageau (1861–1936), Peter Schousboe (1766–1832), Gustave Thuret (1817–1875); they all collected specimens of *C. amentacea* in France. We found 106 specimens of *C. amentacea* collected in the 19th century. Some descriptions give information on its abundance. According to [Agardh \(1842\)](#), the species lives on the most sea-exposed rocks, very frequent on the French Mediterranean shore ('*Hab. ad rupes in ipso limite vehementiori mari expositas, ad littora Galloprovinciae frequentissima*'); he also reported the species precisely '*ad Nizzam ipse* (in Nice), *prope Frejus* (near Fréjus)'. [Montagne \(1846\)](#) studied specimens collected by De Notaris '*Ad Nicaeam*' (in Nice) and by J. Agardh '*Ad Massilliam*' (in Marseilles). Peter Schousboe, Danish consul in Tangier, while waiting for his boat in Marseilles, collected *C. amentacea* in 1818–1820 ([Bornet, 1892](#)): '*Sat frequens circa Massilliam per totum fere annum cum*

fructificatione reperitur' (stand near Marseilles throughout the year with fructifications observed). [Castagne \(1845\)](#) reported the species everywhere (in French '*partout*') around Marseilles. [Debeaux \(1873\)](#) recorded the species is Corsica. [Marion \(1883\)](#) reported the species as abundant throughout the Gulf of Marseilles including the islands and the Côte Bleue (west part of the gulf), but he first noticed that the species is replaced by the articulate Rhodophytes *Corallina officinalis* or the mussel *M. galloprovincialis* when the waters are polluted. He described massive developments of Ulvales all around Marseilles in small bays; at that time, domestic and industrial non-treated outflows to the sea were numerous to the north and the south of the city.

Despite these few records, very localised in space (the vicinity of large cities: Marseilles, Nice, Toulon), accurate maps are non-existent and large portions of the coast have not been surveyed. As a result, the absence of records of *C. amentacea* along most of the coastline only means that we did not find any historical observation. We can hypothesise that rest of the coast was pristine (witnessing all the paintings from this period held in art Museums) and poorly urbanised. Thus, when the favourable ecological conditions were met, it is highly probable that *C. amentacea* was abundant outside the large cities in the 19th century.

3.1.3. 20th century

During the first half of 20th century, the records and collections remained numerous along the whole of the French continental coasts ([Anonymous, 1911](#); [Mouret, 1911](#); [Camous, 1912](#); [Decrock, 1914](#); [Raphélis, 1924a,b,c](#); [Jahandiez, 1929](#); [Ollivier, 1929](#); [Berner, 1931](#); [Hamel, 1931–1939](#)). For Corsica the records are rarer ([Leblond, 1924](#); [Hamel, 1931–1939](#)). The species was reported as abundant and no evidence of regression was noted along most of the coast except at Marseilles, where [Berner \(1931\)](#) reported heavy pollution of the coast and the development of large stands of *Corallina* spp., *M. galloprovincialis* and proliferations of Ulvales interfering with navigation in some small bays. At that time, Marseilles was still an active industrial city.

From 1950 to 1989, the number of works concerning *C. amentacea* considerably increased with algal inventories, phytosociological and chemical studies. The species was collected along the whole coast, in western Provence ([Magne, 1956](#); [Huvé, 1960, 1970](#); [Bellan-Santini, 1961, 1962, 1964, 1965, 1966, 1968a,b](#); [Bellan and Pérès, 1970](#); [Pellegrini and Pellegrini, 1970, 1972](#); [Boudouresque, 1971a,b, 1972](#); [Arnoux and Bellan-Santini, 1972](#); [Bellan and Bellan-Santini, 1972](#); [Belsher, 1975, 1977](#); [Picard and Bourcier, 1975](#); [Belsher and Boudouresque, 1976](#); [Bitar, 1980, 1982](#); [Desrosiers et al., 1982, 1986](#); [Epiard-Lahaye, 1988](#); [Gravez et al., 1988](#)), Eastern Provence ([Molinier, 1956](#); [Bellan-Santini, 1961, 1962](#); [Augarde and Molinier, 1964](#); [Vignes, 1965](#); [Augier et al., 1966](#); [Bensimon et al., 1966](#); [Bensimon, 1967](#); [Augier and Boudouresque, 1967, 1969, 1970a,b, 1976](#); [Augier, 1974, 1978, 1981, 1985](#); [Gadea, 1967](#); [Maggi, 1967](#); [Canamella, 1968](#); [Carvou, 1967](#); [Carvou et al., 1968](#); [Gasquet, 1968](#); [Boudouresque, 1969, 1971a,b](#); [Pellegrini and Pellegrini, 1970](#); [Coppejans, 1972, 1977](#); [Astier, 1975](#); [Belsher et al., 1976](#); [Verlaque and Tiné, 1979](#); [Artaud et al., 1980](#); [Thomas, 1983](#); [Azzolina, 1985](#); [Augier and Nieri, 1987](#)), French Riviera ([Gilet, 1954](#); [Guglielmi, 1969](#)) and in Corsica ([Picard, 1954](#); [Gilet et al., 1954](#); [Molinier and Molinier, 1955](#); [Molinier, 1956, 1960](#); [Santini, 1961](#); [Deleuil and Molinier, 1964](#); [Boudouresque, 1969, 1971b, 1972](#); [Verlaque, 1975](#); [Boudouresque and Perret, 1977](#); [Coppejans, 1979](#); [Clarisse, 1980–1981, 1984](#)).

The first reports of a severe regression around a sewage outfall were documented near Marseilles in the vicinity of the untreated sewage outfall of Cortiou where the species was precisely mapped ([Bellan-Santini, 1964, 1965, 1966, 1968b](#); [Bellan and Bellan-Santini, 1972](#); [Bellan and Pérès, 1970](#); [Bitar, 1982](#)). A link

Table 1

Number of historical records of *Cystoseira amentacea* from the 18th to the 21st century along the French Mediterranean coasts. HV: Herbarium vouchers, P: publications.

Century	18th		19th		20th		21st	
	HV	P	HV	P	HV	P	HV	P
Western Provence	2	0	31	4	31	37	0	1
Eastern Provence	3	0	9	1	26	45	6	9
French Riviera	0	0	41	2	41	11	60	0
Corsica	0	0	25	3	16	25	36	2
Total	5	0	106	10	114	118	102	12

Fig. 2. Distribution of *Cystoseira amentacea* along the coastline of western Provence. Grey line: border of western Provence. (A) Historical records of *C. amentacea* (data from vouchers and literature). Only the oldest and the most recent dates have been reported. (B) Current distribution of *C. amentacea* and locations of sewage outfalls and fish farms.

between the anionic detergents and the regression of the species was hypothesised (Arnoux and Bellan-Santini, 1972). Studies and mapping of *C. amentacea* in the Gulf of Fos, impacted by a warm water outflow from a thermal power plant, the construction of one of the largest European commercial harbours and outfalls from petrochemical industries, evidenced their cumulative negative effects on the species (Desrosiers et al., 1982, 1986). A decline of *C. amentacea* on 2.4 km of shoreline was reported in the vicinity of the untreated sub-surface sewage outfall of Cap Sicié, near Toulon (Eastern Provence) (Astier, 1975; Thomas, 1983). Local depletion was reported in the vicinity of a small sewage outfall at La Madrague (Saint-Cyr-sur-Mer, Eastern Provence) (Thomas, 1983). A shift from *C. amentacea* to turf-forming algae or mussels was observed in the Gulf of Marseilles (Belsher, 1977). In contrast, a recovery of *C. amentacea* has been reported after the construction of sewage treatment plant in La Ciotat (western Provence) (Picard and Bourcier, 1975).

Mapping was undertaken in some places around the pristine Islands of Port-Cros (National Park since 1963) and Porquerolles

(Eastern Provence): *C. amentacea* was abundant and well developed, forming a more or less continuous belt (Augier et al., 1966; Augier and Boudouresque, 1967, 1969, 1970a,b, 1973, 1976; Augier, 1974, 1978, 1981, 1985). A survey (a few hundred metres) over time in the Bay of Port-Man (Port-Cros Island) showed the stability of the population over 20 years (Augier and Nieri, 1987).

Several studies of the vegetation of ports in eastern Provence first mention that small patches or isolated individuals of *C. amentacea* were able to grow on the jetties (Vignes, 1965; Bensimon et al., 1966; Carvou, 1967; Gadea, 1967; Maggi, 1967; Canamella, 1968; Carvou et al., 1968).

From 1990 to 2000, the number of scientific studies concerning *C. amentacea* decreased but was counterbalanced by the increase in the number of maps because of the demand for ecological impact studies and species inventories, as *C. amentacea* is reported as a priority habitat for the EU Natura 2000 programme (code Natura 2000: 1170-13). The species was reported as abundant in western Provence (Nieri et al., 1991; Charbonnel et al., 1993; Sinnassamy et al., 1993; Bonhomme et al., 1997, 1999; Augier et al., 1998), in

Fig. 3. Distribution of *Cystoseira amentacea* along the coastline of eastern Provence. Grey lines: borders of eastern Provence. (A) Historical records of *C. amentacea* (data from vouchers and literature). Only the oldest and the most recent dates have been reported. (B) Current distribution of *C. amentacea* and locations of sewage outfalls and fish farms.

eastern Provence (Valls et al., 1993; Augier, 1995; Francour et al., 1997; Bernard et al., 1999; Gravez et al., 1999; Meinesz et al., 2000a), on the French Riviera (Meinesz et al., 1994; Jaffrenou et al., 1996; Verlaque and Bernard, 1998) and in Corsica (Janssens, 1991; Janssens et al., 1993; Frick et al., 1996; Blachier et al., 1998; Mari et al., 1998; de Vaugelas et al., 1998; Verlaque et al., 1998; Meinesz et al., 1999).

3.1.4. 21st century

In the early 21st century, *C. amentacea* was observed as abundant in western Provence (Bonhomme et al., 2001, 2005; Cadiou et al., 2006), in eastern Provence (Bernard et al., 2001, 2002, 2004; Meinesz et al., 2001a,b,c, 2004a; Soltan, 2001; Bonhomme et al., 2004; Cottalorda et al., 2004; Ruitton et al., 2005; Gratiot et al., 2006, 2007), on the French Riviera (Meinesz et al., 2000b; Francour et al., 2003; Susini, 2006) and in Corsica (Ganteaume

et al., 2004; Meinesz et al., 2001d, 2004b; Javel et al., 2005; Ballesteros et al., 2007). Isolated individuals of *C. amentacea* have been reported on port jetties (Bernard et al., 2004; Bonhomme et al., 2005; Susini, 2006).

The survey of *C. amentacea* in the vicinity of a sewage outfall near Calvi (Corsica) and in Cortiou (Marseilles, western Provence) highlighted its decline over time (Janssens, 1991; Janssens et al., 1993; Soltan et al., 2001). A local disappearance was reported after a port enlargement at Corbières, near Marseilles (Sinnassamy et al., 1993).

3.2. Field data

C. amentacea is very common along the whole of the rocky coasts of Provence, French Riviera and Corsica. The species is present along 377 km (scale 1:2500) from Menton (Italian border) to

Fig. 4. Distribution of *Cystoseira amentacea* along the coastline of the French Riviera. Grey lines: borders of French Riviera. (A) Historical records of *C. amentacea* (data from vouchers and literature). Only the oldest and the most recent dates have been reported. (B) Current distribution of *C. amentacea* and locations of sewage outfalls and fish farms.

Bonnieu Cove, in the Gulf of Fos. In Corsica, the species is spread along 748 km (scale 1:2500). It is absent from the sandy eastern coast of Corsica, from Bastia to Sulenzara, due to the lack of suitable hard substrates.

Most of the populations constitute patchy to continuous belts, with high variation according to the area. Populations are more extensive and more abundant in eastern Provence and Corsica than in other areas (Table 2).

In western Provence, *C. amentacea* occupies only 45 km, i.e. 23% of the hard substrates (Table 2, Fig. 2B), because part of the coast is unsuitable for its development, with high cliffs shading the shore, vertical substrate and overhanging rocks. Furthermore, the northern part of the Marseilles region is directly under the cumulative impact of different human activities. Thus, even when substrates are favourable for the development of *C. amentacea*, the populations are sparsely distributed, outcompeted by large stands of

M. galloprovincialis and *Corallina* spp. The southern part of the city is under the influence of its sewage outlet and *C. amentacea* is lacking over 6.5 km in the vicinity of the outfall (Cortiou), being replaced by mussel beds and *Corallina* spp. assemblages (Fig. 2B).

In eastern Provence, where the coast is more favourable to the development of the species, cliffs are rarer and the region is less urbanised than western Provence, *C. amentacea* is very abundant and occupies 288 km: 72% of the available hard substrate (Table 2, Fig. 3B). The possible influence of the only large city (Toulon) is counterbalanced by its geographical position within a semi-enclosed bay. Near the outfall of the Cape Sicié sewage outlet, *C. amentacea* is reduced to isolated individuals or small patches over 500 m, while *Corallina* spp. assemblages are dominant.

On the French Riviera (including the Principality of Monaco), the species occupies 44 km: 42% of the available hard substrates (Table 2, Fig. 4B). Despite the profound changes that have occurred

Fig. 5. Distribution of *Cystoseira amentacea* along the coastline of northern Corsica. Grey line: border of northern Corsica. (A) Historical records of *C. amentacea* (data from vouchers and literature). Only the oldest and the most recent dates have been reported. (B) Current distribution of *C. amentacea* and locations of sewage outfalls.

along this coast (coastal development, land reclamations, building of a number of ports and marinas) *C. amentacea* is still abundant on all the suitable substrates; the populations are mainly constituted by a continuous belt. Populations are isolated from each other by wide beaches and mostly located around the capes and islands. In the vicinity of the ports, and off the main cities (Cannes, Antibes, Nice, Villefranche-sur-Mer, Saint-Jean-Cap-Ferrat), *Corallina* spp. assemblages dominate. At Monaco, where 87% of the coast has been completely artificialized, a few patches remain; they are located off the Oceanographic Museum, on the last natural rocky substrates of the Principality.

C. amentacea is very abundant in Corsica, especially on the west coast where the populations are the most highly developed. On the east coast, *C. amentacea* is found along the entire rocky coast but populations are more fragmented (Table 2, Fig. 5B and Fig. 6B). Close to Bastia, Ajaccio and Calvi, *C. amentacea* is less abundant and is replaced to a greater or lesser extent by *Corallina* spp. assemblages.

3.3. Impact of sewage outfalls

We analysed the abundance of each dominant species or functional group around 20 sewage outfalls. Each sewage outfall is a particular case, depending on location on the coast, depth, effluent flow, date of setting up, type of treatment and frequency of untreated release events during heavy rains. Field observations clearly showed a relationship between the benthic assemblages

and the existence of an outfall at Cortiou (Fig. 2B) and Cape Sicié (Fig. 3B), the two studied sewage treatment plants with the highest outflow. In addition, the treated effluent is released at the sea surface and 6 m depth, respectively. (Fig. 7, Table 3). Overall, this pattern is confirmed by the two-dimensional nMDS ordination (Fig. 7), although not statistically significant (PERMANOVA; Table 3). However, a significant difference is evidenced if only two classes of sewage volume (below and above 10000 m³ d⁻¹, respectively) are considered (results not shown).

3.4. Impact of fish farms

Like for the sewage outfalls, each fish farm is a particular case, according to the production, the food used (data not available) and the exposure to hydrodynamism. There is no clear relationship between the presence of a fish farm and the abundance of species and functional groups (*Corallina* spp., *C. amentacea*, *M. galloprovincialis*, turf algae), either from the field evidence and from statistical analyses (Fig. 8, Table 4).

3.5. Impact of ports

We analysed the impact of 193 ports, according to their surface area. During field surveys, an effect of the vicinity of the port on the abundance of each dominant species or functional group on the hard substrates was obvious. In the vicinity of ports, especially medium and large ones, *C. amentacea* is absent or only present as isolated individuals and patches, to the advantage of

Fig. 6. Distribution of *Cystoseira amentacea* along the coastline of southern Corsica. Grey line: border of southern Corsica. (A) Historical records of *C. amentacea* (data from vouchers and literature). Only the oldest and the most recent dates have been reported. (B) Current distribution of *C. amentacea* and locations of sewage outfalls and fish farms.

Table 2
Structure of *Cystoseira amentacea* population divided into 6 categories along the French Mediterranean coasts (% of hard substrates). C0: absence, C1: Rare scattered plants, C2: Abundant scattered plants, C3: Abundant patches of dense stands, C4: Almost continuous belt, C5: Continuous belt. Shore length: cumulative length (measured on a 1:2500 map) of the coast colonised by C1–C5 *C. amentacea*.

	C0	C1	C2	C3	C4	C5	Shore length (km)
Western Provence	77.05	2.16	4.45	5.38	5.40	5.56	45
Eastern Provence	27.59	0.77	3.40	11.66	13.10	43.48	288
French Riviera	57.51	0.91	2.12	5.67	6.54	27.24	44
Corsica	34.14	1.06	6.70	29.99	15.87	12.23	748
Total (France)	38.60	1.11	5.48	22.02	13.63	19.17	1125

Corallina spp. This effect is confirmed by statistical analyses (Fig. 9, Tables 5 and 6).

3.6. Development of *Cystoseira amentacea* on man-made structures

We found 41 sites with *C. amentacea* growing on man-made structures: 25 on harbour jetties and the others on breakwaters, groynes, pontoons and artificial platforms (Supplementary material Table S8). Its presence was limited to isolated individuals or small patches. The largest population was observed along 293 m of the jetty at Frioul Island port (Marseilles). In the immediate vicinity (<100 m) of these populations, natural populations were found, constituting putative source populations. However, on the breakwater enclosing the Bay of Toulon, we found isolated individuals up to 3320 m distant from the nearest natural population. Similar observations were made at Marseilles (Corbières, Digue

des Catalans). Most of jetties have been rebuilt at least once since their construction; this is not the case of the segment of the Toulon breakwater that harbours *C. amentacea*, which dates from 1880. Overall, the populations established on man-made structures represent a diminutive cumulative length (2.74 km), compared to the total length of artificial habitats (116.53 km), *Corallina* spp. assemblages being widely dominant.

3.7. Comparison between historical and current maps

The current maps exhibit extensive and numerous areas occupied by *C. amentacea* that were not mentioned in the historical maps. This does not mean a progression of the species distribution, but rather emphasises the vast lack of knowledge and the lacunae in the previous mappings. Here, we only compare reliable historical maps with current distribution (Table 7).

Fig. 7. (A) Two-dimensional nMDS ordination plot on abundance of each dominant species or functional group near each outfall according to their mean effluent flow. The grouping of the major sewage treatment plants on the right seems to show an effect of the effluent volume. (B) Correlation vectors on taxa of interest (Spearman). The opposition of long vectors (*Cystoseira amentacea* 4 and *Corallina* spp.) also seems to reflect an effect of sewage volume. 2D stress: 0.15.

Table 3

Permutation multivariate analysis of variance (PERMANOVA) on square root transformation of abundance of each dominant species or functional group of each sewage outfall, according to the effluent flow, using S17 Bray Curtis similarity.

Source	df	SS	MS	Pseudo-F	P (perm)	Perms	P (MC)
Effluent flow	4	11 760	2940.1	1.2559	0.194	999	0.254
Residual	15	35 114	2340.9				
Total	19	46 874					

Table 4

Permutation multivariate analysis of variance (PERMANOVA) on square root transformation of abundance of each dominant species or functional group of each fish farm, according to the distance from the coast, using S17 Bray Curtis similarity.

Source	df	SS	MS	Pseudo-F	P (perm)	Perms	P (MC)
Distance	2	6324.8	3162.4	0.79519	0.748	78	0.632
Residual	10	39 769	3976.9				
Total	12	46 094

Fig. 8. (A) Two-dimensional nMDS ordination plot on abundance of each dominant species or functional group near each fish farm, according to the distance from the coast. (B) correlation vectors on taxa and functional groups of interest (Spearman); 2D stress: 0.06.

Fig. 9. (A) Two-dimensional nMDS ordinations plot on abundance of each dominant species or functional group near each port, according to its surface area. The dispersion of the smallest ports reflects their low impact, in contrast with larger ones. (B) Correlation vectors on taxa of interest (Spearman); the opposition of long vectors (*Cystoseira amentacea* 3, 4 and 5 and *Corallina* spp.) reflects the port effect. 2D stress: 0.18.

Table 5
Permutation multivariate analysis of variance (PERMANOVA) on square root transformation of abundance of each dominant species or functional group, regarding each surface area of port, using S17 Bray Curtis similarities: small port ≤ 2 ha; middle port = 2.1–6 ha; large port = 6.1–100 ha; very large port ≥ 101 ha.

Source	df	SS	MS	Pseudo-F	P (perm)	Perms	P(MC)
Surface area	3	55927	18642	7.6304	0.001	997	0.001
Residual	189	4.6176E5	2443.2				
Total	192	5.1769E5					

Table 6
Pair-wise comparison on square root transformation of abundance of each dominant species or functional group, regarding each surface area of port, using S17 Bray Curtis similarities: small port ≤ 2 ha; medium sized port = 2.1–6 ha; large port = 6.1–100 ha; very large port ≥ 101 ha.

Pair-wise tests	t	P (perm)	Number of perms	P (Monte Carlo)
Large port/middle port	1.8478	0.01	999	0.02
Large port/small port	4.1901	0.001	999	0.001
Large port/very large port	1.6623	0.039	613	0.027
Medium sized port/small port	2.185	0.002	999	0.004
Medium sized port/very large port	1.8518	0.01	477	0.012
Small port/very large port	1.738	0.008	876	0.015

The comparison of the historical maps with the current maps, within exactly the same areas, shows that, in most cases, only small changes occurred, -3.8% to 4.4% (overall: -0.2%) (Table 7: non-bold lines). These slight differences (positive or negative) between historical and current maps probably result from differences in mapping techniques and accuracy of positioning of the historical maps. Considering these biases, these results reflect a remarkable stability over 1–5 decades. In contrast, a conspicuous decline occurred in 4 areas (Table 7: bold lines). The possible causes and biases for this decline are discussed below.

4. Discussion

For the first time, *C. amentacea* has been exhaustively mapped at a very large scale (1:2500), over 1832 km of its distribution area where hard substrates, putatively favourable to its dwelling, are present. It occurs along 1125 km of coast, from discontinuous

populations (isolated individuals) to continuous belts; the continuous and almost continuous belts are found along 369 km of shoreline. Thus, a general view of its current distribution in the French Mediterranean Sea is now available. This distribution has been compared with a comprehensive analysis of all available historical data from literature, grey literature and herbarium vouchers since 1700.

Most authors have emphasised the dramatic decline of species of the genus *Cystoseira* throughout the Mediterranean Sea (e.g. Munda, 1974, 1982, 1993; Boudouresque, 2003; Thibaut et al., 2005; Devescovi and Ivesa, 2007; Airoldi et al., 2008; Falace et al., 2010; Mačić et al., 2010; Fraschetti et al., 2011; Giakoumi et al., 2012; Sala et al., 2012; Bianchi et al., 2014; Templado, 2014). It is the case, in particular, for *C. amentacea* (e.g. Bellan-Santini, 1963, 1964, 1966, 1968b; Arnoux and Bellan-Santini, 1972; Bellan and Bellan-Santini, 1972; Belsher and Boudouresque, 1976; Clarisse, 1980–1981, 1984; Boudouresque

Table 7

Comparison of the length (in m) of shoreline occupied by *Cystoseira amentacea* in the past (various periods) and today, where reliable historical maps are available. Bold: conspicuous decline.

Source	Location	Historical	Current	Difference (%)
Ollivier (1929)	Between Eze-sur-Mer and Cap Martin	13269	9977	–24.8
Guglielmi (1969)	Beaulieu-sur-Mer and Saint-Jean-Cap-Ferrat	10425	7311	–29.9
Meinesz et al. (2000b)	Coastline of Nice	2369	2279	–3.8
Bonhomme et al. (2004)	Cap Dramont corniche d'Antheor, Saint-Raphaël	7333	7356	0.3
Augier and Boudouresque (1967)	Bay of La Palud, Port-Cros Island	689	709	2.9
Augier and Boudouresque (1970b)	Bay of Port-Man, Port-Cros Island	1571	1577	0.4
Augier and Boudouresque (1976)	Port-Cros Island (part)	10453	10537	0.8
Meinesz et al. (2000a, 2001a,b, 2004a) and Cottalorda et al. (2004)	All mapped areas of Port-Cros Island	19107	19296	1.0
Meinesz et al. (2001c)	Bagaud Island	6211	6120	–1.5
Augier (1978)	Bon Renaud Cove, Porquerolles Island	113	118	4.4
Augier (1981)	Port-Fay Cove, Porquerolles Island	554	555	0.2
Augier (1995)	Gorges du Loup Cove, Porquerolles Island	271	279	3.0
Gratiot et al. (2006)	Porquerolles Island (part)	12625	12614	–0.1
Picard and Bourcier (1975)	La Ciotat	5689	5610	–1.4
Bellan-Santini (1966)	Calanques of Marseilles	17465	8051	–53.9
Soltan (2001)	Calanques of Marseilles	13154	8051	–38.8
Cadiou et al. (2006)	Cassis, between Canaille Cape and Grand Caunet Cape	7528	7479	–0.7
Clarisse (1980–1981, 1984)	Between punta Revellata and the harbour of Calvi	7056	5801	–17.8
Blachier et al. (1998)	Between punta Cappicciolu and punta Sperone	6011	5826	–3.1

et al., 1990; Soltan et al., 2001; Boudouresque, 2003; Susini, 2006; Susini et al., 2007). This worrying trend resulted in the listing of *C. amentacea* under the Berne Convention, the Barcelona Convention and Habitat of Community Interest in the UE (Micheli et al., 2013).

We were therefore expecting a severe decline of *C. amentacea*, as reported for *C. mediterranea* along the French Catalan coast (Thibaut et al., 2005), at least in Provence and French Riviera, in relation with the increase in human pressure (population, summer touristic frequentation, coastal development and leisure boating). Since the last decades of the 20th century, the increase in marine habitat losses (275 ports, 144 land reclamations, 458 groynes and jetties) has been dramatic (Meinesz et al., 2013). As regard the pollution, it has conspicuously increased over the investigated period, especially during the 20th century (Serrano et al., 2011) and its decline in the study area, following the construction of sewage treatment plants, is too recent (1980–1990s) to have been likely to reverse the trend (Soltan et al., 2001; Thibaut et al., 2005).

In fact, among most of the precise locations where historical specimens were reported, we found the species present at the same sites; in some cases, when the abundance was indicated, it proved to be unchanged. In the Gulf of Fos (western Provence), which harbours one of the largest French port and industrial facilities (petrochemical and chemical industries, oil refineries, steelworks, power plant, etc.), the distribution of *C. amentacea* has remained unchanged since 1975 (Fig. 2B), but populations are now more fragmented and are in competition with mussel beds (Desrosiers et al., 1982, 1986) whereas industrial activities and perturbations have clearly increased over the past 40 years. Only 4 locations presented a conspicuous decline of *C. amentacea* populations between historical and current data (Table 7): the Calanques of Marseilles (Western Provence), between Eze-sur-Mer and Cap Martin, between Beaulieu-sur-Mer and Saint-Jean-Cap-Ferrat (French Riviera) and between Punta Revellata and the harbour of Calvi (Corsica). Ollivier (1929) mapped the species between Eze-sur-Mer and Cap Martin, including the Principality of Monaco. The 24.8% loss can be related to well-documented events, such as the near-total artificialisation of the coastline, especially at Monaco (between 1963 and 2002), where artificialisation encompasses 87% of the coastline (Meinesz et al., 2013). This loss, and its rate of development, is therefore convincing. The 29.9% loss observed between Beaulieu-sur-Mer and Saint-Jean-Cap-Ferrat is based upon a map of Guglielmi (1969). This map is highly reliable and the causes of the loss well identified: enlargement of the

coastal road at Beaulieu-sur-Mer (~1975), building of the port of Beaulieu-sur-Mer (1969), enlargement of the Saint-Jean-Cap-Ferrat port (1972), land reclamation of Le Lido (Saint-Jean-Cap-Ferrat, ~1998) (Meinesz et al., 2013). In Corsica, a similar decline trend has been documented between Punta A Rivellata (La Revellata) and the harbour of Calvi (–17.8%, Clarisse, 1980–1981, 1984), where the impact of a sewage outfall has been suspected (Janssens, 1991).

The greatest loss has been documented in the vicinity of the Cortiou sewage outfall at Marseilles (western Provence), the largest French Mediterranean city and one of the largest cities of the whole Mediterranean Sea, where a steady decrease of *C. amentacea* has apparently occurred since the 1960s (Bellan-Santini, 1966; Soltan, 2001; Soltan et al., 2001). The loss, with reference to the 1960s hypothetical baseline, would be 39% in the early 2000s (Soltan, 2001) and 54% (present work). However, the validity of the baseline can be questioned; Bellan-Santini (1966, 1968b) mapped the species in some places where its presence is unlikely, due to unsuitable habitats such as beaches, protected coves, under overhanging rocks, on cliffs and on *Lithophyllum byssoides* (Lamarck) Foslé rims. We therefore suspect an overestimation of the rate of loss since the 1960s. In any case, the 1960s do not constitute a reliable baseline since pollution in the Marseilles area was already worrying in the late 19th century, when the Cortiou sewage outfall was constructed (Marion, 1883), and in the first half of the 20th century (Berner, 1931). Nevertheless, the loss continued after 2000 and the unbiased map of Soltan (2001), which is unexpected due to the construction of a sewage treatment plant in 1987 and the improvement of the treatment process in 2007, for the largest discharge of sewage along the French coasts (1 630 000 population equivalent). The reason could lie in the fact that the outfall flow is extremely high (192 000–233 000 m³ d^{–1}), and located at sea level, it gives rise to a large plume of freshwater drifting at the sea surface (low salinity is known to have a negative effect on some species of Fucales, Zavodnik, 1975), and it is dumped without any treatment during the occasional heavy rains.

Our results confirm that *C. amentacea* is able to colonise man-made structures (we found the species at 41 sites). These populations cannot be perennial since the structures (rocks, concrete blocks, etc.) can be replaced from time to time; this is the probably the case in many ports (French Riviera and eastern Provence) where all the previously reported sightings were not observed again after the enlargement of the port (Ollivier, 1929;

Bensimon et al., 1966; Carvou, 1967; Gadea, 1967; Maggi, 1967; Canamella, 1968; Carvou et al., 1968; Vignes, 1968; Susini, 2006). In the Mediterranean, among the Fucales, only *C. compressa* was commonly observed on dykes (Susini, 2006) and, in the Adriatic *Fucus virsoides* J. Agardh is also able to grow on man-made substrates (Mačić, 2006), whereas in the Atlantic *Fucus* spp. are commonly observed on man-made substrates. The observation of *C. amentacea* on relatively recent man-made substrates (less than 40 years) and relatively distant from the nearest natural population invalidates the hypothesis of Mangialajo et al. (2012) suggesting very slow colonisation, step by step, by eggs disseminated over less than few dozen cm from the source population. Possible dispersion mechanisms, able to account for our results, involve the sea bird *Larus cachinnans* Pallas 1811 (Bartoli et al., 1997) and drifting fragments (e.g. Raphaélis specimens in Herbarium Général, MU; drift specimens on French Riviera beaches).

When disturbed, *C. amentacea* is replaced by *Corallina* spp, the mussel *M. galloprovincialis* and/or *C. compressa*. This phenomenon has been extensively documented; the fast growing *Corallina* spp. take the place of *C. amentacea* while mussels prevent the recruitment of *C. amentacea* (Berner, 1931; Desrosiers et al., 1982, 1986; Bulleri et al., 2002). A number of authors (Bellan and Bellan-Santini, 1972; Kautsky et al., 1992) have reported increments of mussels in polluted waters, particularly in the vicinity of domestic outfalls. Similarly, an increase in mussel populations has been related to algal stand declines in Great Britain and the Baltic Sea (Kautsky et al., 1992). Lastly, *C. compressa* is a tolerant opportunistic fast growing species, replacing *C. amentacea* when disturbed (Mangialajo et al., 2012). Loss of *C. amentacea* reduces the species diversity (point diversity, as defined by Boudouresque (2011) from 133 species in *C. amentacea* stands in pristine waters, 108 species in *C. elongata* stands and 76 species in mussels beds in polluted sites (Bellan and Bellan-Santini, 1972). Competition between rhodophytes and fucoids is a common feature worldwide (Worm and Chapman, 1996). In Asturias (Spain), coastal development and pollution on rocky coast result in long term changes, with the decline of *Fucus* and *Laminaria* in favour of *Gelidium* sp. (Fernandez and Niell, 1982). In the Atlantic, *C. tamarascifolia*, *C. baccata* (S.G. Gmelin) P.C. Silva and *Bifurcaria bifurcata* R. Ross forests have been replaced by turf algae in Abra Bilbao Bay (Biscayne Gulf) under a pollution gradient (Díez et al., 1999, 2009). Overall, the replacement of *C. amentacea* by *Corallina* spp., mussel beds and/or *C. compressa* can be interpreted as a phase shift from a canopy stage to a less structured stage (turf), inducing simplification of the zonation pattern, as reported when fucoids have been lost (Munda, 1980, 1993; Tewari and Joshi, 1988; Fairweather, 1990; Rueness and Fredriksen, 1991).

To date, *C. amentacea* has been regarded as a species very sensitive to human impact, e.g. pollution, coastal development and land reclamations. It was thought to have become relatively rare in the highly man-impacted north-western Mediterranean and therefore benefits from a protection status. Undoubtedly, our results confirm a positive correlation between the decline of the species and the vicinity of medium-sized and large ports; as regards the sewage outfalls, such a correlation only concerns very large outfalls flowing at the sea surface, such as that at Marseilles; finally, no relation may be observed between *C. amentacea* and fish farms. The species has a low nitrogen uptake rate (Orlandi et al., 2014) that could protect the species from non-permanent nitrogen pollution. However, at the scale of the French Mediterranean coasts (Provence, French Riviera and Corsica), this impact seems limited. The comparison of 14 accurate maps produced in the past (1967 through 2006) with current maps, under a variety of human pressure, along 81 km of coastline, failed to evidence any general trend (only 0.02% of difference). Loss is concentrated in 4 areas near Marseilles (Provence), Nice-Monaco (French Riviera) and Calvi (Corsica): from 75 km (1929–1984; may

be a slight overestimation) to 47 km (currently) of coastline. Overall, at the scale of the study area, *C. amentacea* extends along 1125 km, with long stretches of near-continuous to continuous belts. Although the sensitivity of the species to human impact is confirmed, its resilience may be higher than previously thought.

A large amount of historical data, in the form of rarely available old publications, grey literature and herbarium vouchers, do exist, although frequently neglected by authors. Taking into account these data makes it possible to reduce the sliding baseline syndrome, i.e., in the absence of any benchmark, the acceptance of an already degraded environment or of a hypothetically exaggerated degraded one, as the reference, i.e. the baseline. The reconstruction of historical data performed here demonstrates its feasibility, usefulness and efficiency. In addition, the accurate mapping of the current distribution of *C. amentacea* (2008–2011) will allow future comparisons.

Acknowledgments

This work was funded by the Agence de l'Eau Rhône Méditerranée-Corse, P. Boissery. We thank Eric Pironneau, Pauline Robvieux, J.M. Cottalorda for technical assistance at sea, Adrien Cheminée for his statistical advices, the rangers of the French MPAs (H. Bergère, N. Nègre, J.M. Culioli, J.M. Dominici, M.C. Santoni), the curators of all the following herbaria we visited at the Museum National d'Histoire Naturelle (B. Denetière and B. de Reviers), Villa Thuret (C. Ducatillon), Museum d'Histoire Naturelle d'Aix-en-Provence (C. Delnatte, M. Durand), Museum d'Histoire Naturelle de Toulon (L. Charrier), Museum d'Histoire Naturelle d'Avignon (P. Moulet), Herbar de l'Université de Montpellier 2 (V. Bourgade), Museum d'Histoire Naturelle de Nice (B. Rollier), Musée Océanographique de Monaco (M. Bruni). Thanks are due to M. Paul for proof-reading the English.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.marpolbul.2014.09.043>.

References

- Agardh, J.C., 1842. *Algae maris Mediterranei et Adriatici, observationes in diagnosis specierum et dispositionem generum*. Parisiis: Apud Fortin, Masson et Cie., Paris, 164 p.
- Agence de l'Eau Rhône Méditerranée Corse, 2014. <www.eaurmc.fr> (accessed 24.07.14).
- Airolidi, L., Beck, M.W., 2007. Loss, status and trends for coastal marine habitats of Europe. *Oceanogr. Mar. Biol. Annu. Rev.* 45, 345–405.
- Airolidi, L., Balata, D., Beck, M.W., 2008. The gray zone: relationships between habitat loss and marine diversity and their applications in conservation. *J. Exp. Mar. Biol. Ecol.* 366 (1), 8–15.
- Albérola, C., Millot, C., 2003. Circulation in the French Mediterranean coastal zone near Marseilles: the influence of wind and the Northern Current. *Cont. Shelf Res.* 23, 587–610.
- Anderson, M.J., 2001. Permutation tests for univariate or multivariate analysis of variance and regression. *Can. J. Fish. Aquat. Sci.* 58 (3), 626–639.
- Anonymous, 1911. Excursion du 11 juin 1911 au Cap Bénat. *Ann. Soc. Hist. Nat. Toulon* 2, 73–77.
- Arévalo, R., Pinedo, S., Ballesteros, E., 2007. Changes in the composition and structure of Mediterranean rocky-shore communities following a gradient of nutrient enrichment: descriptive study and test of proposed methods to assess water quality regarding macroalgae. *Mar. Pollut. Bull.* 55, 104–113.
- Arnoux, A., Bellan-Santini, D., 1972. Relations entre la pollution du secteur de Cortiou par les détergents anioniques et les modifications des peuplements de *Cystoseira stricta*. *Téthys* 4, 583–586.
- Artaud, J., Iatrides, M.C., Tisse, C., Estienne, J., 1980. Etudes chromatographiques des stéroïdes: application aux stéroïdes d'algues. *Analisis* 8 (7), 227–286.
- Astier, J.M., 1975. Cartographie des fonds marins de la région de Toulon par le groupe "ECOMAIR". *Ann. Soc. Sci. Nat. Arch. Toulon Var* 27, 15.
- Augarde, J., Molinier, R., 1964. Le Clydonomètre appareil de mesure et d'enregistrement des perturbations hydrodynamiques superficielles. *Bull. Inst. Oceanogr. Monaco* 61 (1288), 18 p.

- Augier, H., 1974. Les phytohormones des algues. II Etudes biochimiques. Ann. Sci. Nat. Bot. Biol. Vég. 15 (1), 1–64.
- Augier, H., 1977. Excursion au Cap Croisette. In: Sélection du Rider Digest. Guide Ecologique de la France, Paris, pp. 460–462.
- Augier, H., 1978. Peuplements benthiques de l'île de Porquerolles (Méditerranée, France). 1. Carte de l'anse du Bon Renaud et de la plage d'Argent. Trav. Sci. Parc Nat. Port-Cros 4, 89–100.
- Augier, H., 1981. Etude et cartographie des peuplements benthiques de l'île de Porquerolles (Méditerranée, France). II L'anse de Port-Fay et la Calanque de la Grotte. Trav. Sci. Parc Nat. Port-Cros 7, 103–117.
- Augier, H., 1985. Etude biocénotique et cartographique de l'anse de la Plage du Sud (Parc National de Port-Cros, Méditerranée, France). Trav. Sci. Parc Nat. Port-Cros 11, 23–38.
- Augier, H., 1995. Etude et cartographie des peuplements benthiques de l'île de Porquerolles (Méditerranée, France). III Transect permanent et cartographie biocénotique des gorges du Loup, Ile de Porquerolles (Var, S.E. France). Trav. Sci. Parc Nat. Port-Cros 16, 115–122.
- Augier, H., Boudouresque, C.F., 1967. Végétation marine de l'île de Port-Cros (Parc National). I. La baie de la Palud. Bull. Mus. Hist. Nat. Marseille 27, 93–124.
- Augier, H., Boudouresque, C.F., 1969. Enjeu et bilan d'un Parc National sous-marin. Ann. Soc. Sci. Nat. Arch. Toulon Var 21, 24.
- Augier, H., Boudouresque, C.F., 1970a. Végétation marine de l'île de Port-Cros (Parc National). VI. Le récif barrière de Posidonies. Bull. Mus. Hist. Nat. Marseille 30, 221–228.
- Augier, H., Boudouresque, C.F., 1970b. Végétation marine de l'île de Port-Cros (Parc National). V. La baie de Port-Man et le problème de la régression de l'herbier de Posidonies. Bull. Mus. Hist. Nat. Marseille 30, 145–164.
- Augier, H., Boudouresque, C.F., 1973. Dix ans de recherche dans la zone marine du Parc National de Port-Cros (France). Deuxième partie. Ann. Soc. Sci. Nat. Arch. Toulon Var 25, 34.
- Augier, H., Boudouresque, C.F., 1976. Végétation marine de l'île de Port-Cros (Parc National). XIII. Documents pour la carte des peuplements benthiques. Trav. Sci. Parc Nat. Port-Cros 2, 9–22.
- Augier, H., Nieri M., 1987. Cartographie des peuplements benthiques de la baie de Port-Man. Etude de son évolution depuis 1967–1969. Degré de contamination par le cuivre. Groupement de recherche de Biologie, Ecologie, Nuisances et Gestion Océanique (BENGO), Faculté des Sciences de Marseille-Luminy. Fr, 67p.
- Augier, H., Boudouresque, C.F., Molinier, R., 1966. Etudes des biocénoses marines de la baie de la Palud. Ile de Bagaud. Excursion des étudiants en Agrégation à l'île de Port-Cros. Mai 1966, 5 p.
- Augier, H., Baudin, J.C., Lion, R., 1998. Etude biocénotique et cartographique des fonds marins de la Calanque d'En Vau (Bouches-Rhône, Méditerranée, France. Cerimer Information, Fr., 7, 32 p.
- Azzolina, J.F., 1985. Etude écologique des secteurs acquis ou susceptibles d'être acquis par le Conservatoire du Littoral dans le département du Var: le domaine marin. Contrat Parc National de Port-Cros – Conservatoire du Littoral, 150 p.
- Babalís, T., 2011. Restoring the Past: Environmental history and oysters at Point Reyes national seashore. The George Wright Forum 28 (2), 199–215.
- Ballesteros, E., 1988. Estructura y dinámica de la comunidad de *Cystoseira mediterranea* Sauvageau en el Mediterráneo Noroccidental. Inv. Pesq. 52, 313–334.
- Ballesteros, E., 1990a. Structure and dynamics of the community of *Cystoseira zosteroides* (Turner) C. Agardh (Fuciales, Phaeophyceae) in the Northwestern Mediterranean. Sci. Mar. 54, 217–299.
- Ballesteros, E., 1990b. Structure and dynamics of the *Cystoseira caespitosa* Sauvageau (Fuciales, Phaeophyceae) community in the North-Western Mediterranean. Sci. Mar. 54, 155–168.
- Ballesteros, E., Torras, X., Pinedo, S., García, M., Mangialajo, L., de Torres, M., 2007. A new methodology based on littoral community cartography for the implementation of the European Water Framework Directive. Mar. Poll. Bull. 55, 172–180.
- Barsanti, M., Delbono, I., Ferretti, O., Peirano, A., Bianchi, C.N., Morri, C., 2007. Measuring change of Mediterranean coastal biodiversity: diachronic mapping of the meadow of the seagrass *Cymodocea nodosa* (Ucria) Ascherson in the Gulf of Tigullio (Ligurian Sea, NW Mediterranean). Hydrobiologia 580, 35–41.
- Bartoli, P., Bourgeau-Cause, M., Combes, C., 1997. Parasite transmission via a vitamin supplement. BioScience 47b (4), 251–253.
- Bellan, G., Bellan-Santini, D., 1972. Influence de la pollution sur les peuplements marins de la région de Marseille. In: Ruivo, M. (Ed.), Marine Pollution and Sea Life. Fishing News (Books) Ltd., Surrey, pp. 396–401.
- Bellan, G., Pères, J.M., 1970. Etat général des pollutions sur les côtes méditerranéennes de France. Quad. Civ. Sta Idrobiol. Milano 1, 36–65.
- Bellan-Santini, D., 1961. Note préliminaire sur la faune et la flore du peuplement à *Petroglossum nicaense* (Duby) Schotter et sur ses rapports avec le peuplement à *Cystoseira stricta* (Mont.) Sauvageau. Rec. Trav. St. Mar. End. Bull. 23 (37), 19–30.
- Bellan-Santini, D., 1962. Etude faunistique et floristique de peuplements infralittoraux de substrats rocheux. Rec. Trav. St. Mar. End. Bull. 26 (41), 237–298.
- Bellan-Santini, D., 1963. Etude quantitative du peuplement à *Cystoseira stricta* (Mont.) Sauv. Rapp. P.V. Réunion CIESM 17 (2), 133–138.
- Bellan-Santini, D., 1964. Influence de la pollution sur quelques peuplements superficiels de substrats rocheux. In: CIESM. Symp. Poll. Mar. Microorganismes Prod. pétrol., Monaco avril 1964, pp. 127–131.
- Bellan-Santini, D., 1965. Etude quantitative du peuplement à *Mytilus galloprovincialis* Lamarck en eau moyennement polluée. Rapp. P.V. Réunion CIESM 18 (2), 85–89.
- Bellan-Santini, D., 1966. Influence des eaux polluées sur la flore et la faune marines benthiques dans la région marseillaise. Techn. San. Mun. 61 (7), 285–292.
- Bellan-Santini, D., 1968a. Conclusions d'une étude quantitative dans la biocénose des algues photophiles en Méditerranée sur les côtes de Provence. Mar. Biol. 1 (3), 250–256.
- Bellan-Santini, D., 1968b. Influence de la pollution sur les peuplements benthiques. Rev. Intern. Oceanogr. Med. 10, 27–53.
- Belsher, T., 1975. Inventaire et analyse du milieu naturel en Rade d'Hyères. Contrat Université Aix-Marseille II et Parc National de Port-Cros, I–III, 246 p.
- Belsher, T., 1977. Analyse des répercussions de pollution urbaine sur le macrophytobenthos de Méditerranée (Marseille, Port-Vendres, Port-Cros). Thèse Université Aix-Marseille II, 287 p.
- Belsher, T., Boudouresque, C.F., 1976. L'impact de la pollution sur la fraction algale des peuplements benthiques de Méditerranée. Atti Tavola rotonda internazionale. "La biologia marina per la difesa e per la produttività del mar", Livorno, 215–260.
- Belsher, T., Augier, H., Boudouresque, C.F., Coppejans, E., 1976. Inventaire des algues marines benthiques de la rade et des îles d'Hyères (Méditerranée, France). Trav. Sci. Parc Nat. Port-Cros 2, 39–89.
- Benedetti-Cecchi, L., 2001. Variability in abundance of algae and invertebrates at different spatial scales on rocky sea shores. Mar. Ecol. Prog. Ser. 215, 79–92.
- Benedetti-Cecchi, L., Cinelli, F., 1992a. Canopy removal experiments in *Cystoseira*-dominated rockpools from the Western coast of the Mediterranean (Ligurian Sea). J. Exp. Mar. Biol. Ecol. 155 (1), 69–83.
- Benedetti-Cecchi, L., Cinelli, F., 1992b. Effects of canopy cover, herbivores and substratum type on patterns of *Cystoseira* spp. settlement and recruitment in littoral rockpools. Mar. Ecol. Prog. Ser. 90, 183–191.
- Benedetti-Cecchi, L., Cinelli, F., 1995. Habitat heterogeneity, sea urchin grazing and the distribution of algae in littoral rock pools on the west coast of Italy (western Mediterranean). Mar. Ecol. Prog. Ser. 126, 203–212.
- Benedetti-Cecchi, L., Pannacchiulli, F., Bulleri, F., Moschella, P.S., Airoldi, L., Relini, G., Cinelli, F., 2001. Predicting the consequences of anthropogenic disturbance: large-scale effects of loss of canopy algae on rocky shores. Mar. Ecol. Prog. Ser. 214, 137–150.
- Bensimon, E., 1967. Etude de la végétation des horizons superficiels de l'étage infralittoral dans le port de Bandol (Var). DES Faculté des Sciences, Marseille, 67 p.
- Bensimon, E., Meze, A., Pastore, M., Vignes, P., 1966. Rapports entre la forme et la distribution de quelques algues du littoral varois. Ann. Soc. Sci. Nat. Arch. Toulon Var 18, 3–22.
- Bernard, G., Bonhomme, P., Charbonnel, E., Cadiou, G., 1999. Etat de référence des biocénoses, peuplements, espèces et paysages remarquables – commune de St-Cyr-sur-Mer. Analyses des atouts pour valorisation du milieu marin. Contrat Ville de St-Cyr-Mer and GIS Posidonie. GIS Posidonie publ., Marseille Fr. 75 p. + annexes non num.
- Bernard, G., Denis, J., Deneux, F., Belsher, T., Sauzade, D., Boudouresque, C.F., Charbonnel, E., Emery, E., Hervé, G., Bonhomme, P., 2001. Etude et cartographie des biocénoses de la rade de Toulon. Rapport de synthèse final. Contrat d'étude pour le syndicat intercommunal de l'Aire Toulonnaise, Ifremer et GIS Posidonie. Ifremer Publ. La Seyne, Fr. 152 p.
- Bernard, G., Boudouresque, C.F., Charbonnel, E., Bonhomme, P., Cadiou, G., 2002. Cartographie et état de vitalité des peuplements marins du Brusuc (Commune de Six-Fours, Var) – conseil et recommandations pour l'aménagement du port du Brusuc. Contrat Conseil Général des ports and GIS Posidonie. GIS Publ. Fr. 87p.
- Bernard, G., Bonhomme, P., Ganteaume, A., Boudouresque, C.F., 2004. Relevé des herbiers de Magnolophytes marines au droit du port de St-Elme (commune de la Seyne-sur-mer, Var). Rapport Cartographique. GIS Posidonie and Conseil Général du Var Service des Ports. Gis Posidonie publ., Fr. 28 p. + annexes non num.
- Berner, L., 1931. Contributions à l'étude sociologique des algues marines dans le golfe de Marseille. Ann. Mus. Hist. Nat. Marseille 2, 3–84.
- Bianchi, C.N., Morri, C., 2000. Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. Mar. Pollut. Bull. 40, 367–376.
- Bianchi, C.N., Corsini-Foka, M., Morri, C., Zenetos, A., 2014. Thirty years after: dramatic changes in the coastal marine ecosystems of Kos Island (Greece), 1981–2013. Medit. Mar. Sci. 15, 482–497.
- Bitar, G., 1980. Etude de l'impact de la pollution par un émissaire urbain (collecteur Cortiou) sur les peuplements infralittoraux de substrats durs de la côte sud de Marseilleveyre (Marseille). PhD Thesis Marseille, 137 p.
- Bitar, G., 1982. Influence d'un grand émissaire urbain sur la distribution du zoobenthos de substrat dur dans la région de Marseille (Méditerranée nord-occidentale). Téthys 10 (3), 200–210.
- Blachier, J., Meinesz, A., de Vaugelas, J., 1998. Répartition de Lithophyllum lichenoides (Rhodophyta), de Cystoseira amentacea (Chromophyta), de Patella ferruginea (Mollusca) dans la réserve naturelle des îles Lavezzi: îlots et littoral de la pointe di U Cappicciolu à la pointe de Spérone. Trav. Sci. Parc Nat. Reg. Corse 17, 103–140.
- Bonhomme, P., Bernard, G., Gravez, V., 1997. Le milieu marin du site de Corbières (Bouches-du-Rhône). Deuxième suivi (1996). DGST Ville de Marseille and GIS Posidonie publ. Fr. 43 p. + annexes I–II.
- Bonhomme, P., Bernard, G., Daniel, B., Boudouresque, C.F., 1999. Archipel de Riou: étude socio-économique sur la plaisance, la pêche amateur, la plongée et la chasse sous-marine. Ville de Marseille and GIS Posidonie publ. Fr. 83 p.
- Bonhomme, P., Boudouresque, C.F., Bernard, G., Verlaque, M., Charbonnel, E., Cadiou, G., 2001. Espèce, peuplements et paysages marins remarquables de La Ciotat, de

- l'île Verte à la Calanque des Capucins (Bouches-du-Rhône). Contrat Ramoge and GIS Posidonie. GIS Posidonie publ, Fr, 133 p.
- Bonhomme, P., Cadiou, G., Bernard, G., Loques, F., Cottalorda, J.M., Talin, F., Meinesz, A., Chambard, V., Boudouresque, C.F., 2004. Espèces, peuplements et paysages marins remarquables du Cap Dramont et de la Corniche d'Anthéor (Var, France). Contrat Ramoge and GIS Posidonie, GIS Posidonie publ., Fr 78 p. + annexes 1-2.
- Bonhomme, P., Ganteaume, A., Bellan, G., Cadiou, G., Emery, E., Clabaut, P., Bernard, G., Hervé, G., Bourcier, M., Boudouresque, C. F., 2005. Etude et cartographie des biocénoses marines des calanques de Marseille à Cassis, y compris l'archipel de Riou. Phase 3. GIS Posidonie publ., Fr. 156 p.
- Bornet, E. 1892. Les algues de P.-K.-A. Schousboe. Mém. Soc. Sci. Nat. Math. Cherbourg, pls I-III, 28, 165-376.
- Boudouresque, C.F., 1969. Etude qualitative et quantitative d'un peuplement algal à *Cystoseira mediterranea* dans la région de Banyuls-sur-Mer. Vie et Milieu 20 (28), 437-452.
- Boudouresque, C.F., 1971a. Contribution à l'étude phytosociologique des peuplements algaux des côtes varoises. Vegetatio 22 (1-3), 83-184.
- Boudouresque, C.F., 1971b. Recherches de bionomie analytique, structurale et expérimentale sur les peuplements benthiques sciaphiles de Méditerranée occidentale (fraction algale). La sous-strate sciaphile des peuplements de grandes *Cystoseira* de mode battu. Bull. Mus. Hist. Nat. Marseille 31, 141-151, 1 Tabl.
- Boudouresque, C.F., 1972. Recherche de bionomie analytique structurale et expérimentale sur les peuplements benthiques sciaphiles de Méditerranée occidentale (Fraction algale): la sous-strate sciaphile d'un peuplement photophile de mode calme, le peuplement à *Cystoseira crinita*. Bull. Mus. Hist. Nat. Marseille 32, 253-263.
- Boudouresque, C.F., 2003. The erosion of Mediterranean biodiversity. In: Rodríguez-Prieto, C., Pardini, G. (Eds.), The Mediterranean Sea: An Overview of its Present State and Plans for Future Protection. Servei de Publicacions de la Universitat de Girona, pp. 53-112.
- Boudouresque, C.F., 2011. Insights into the diversity of the biodiversity concept. Mésogée 67, 13-26.
- Boudouresque, C.F., Perret, M., 1977. Inventaire de la flore marine de Corse (Méditerranée): Rhodophyceae, Phaeophyceae, Chlorophyceae et Bryopsidophyceae. Bibliotheca Phycol. 25, 171.
- Boudouresque, C.F., Verlaque, M., 2002. Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. Mar. Pollut. Bull. 44, 32-38.
- Boudouresque, C.F., Ballesteros, E., Ben Maiz, N., Boisset, F., Bouladier, E., Cinelli, F., Cirik, S., Cormaci, M., Jeudy De Grissac, A., Laborel, J., Lan franco, E., Lundberg, B., Mayhoub, H., Meinesz, A., Panayotidis, P., Sem roud, R., Sinnassamy, J.M., Span, A., Vuignier, G., 1990. Livre rouge "Gérard VUI GNIER" des végétaux, peuplements et paysages marins menacés de Méditerranée. Pro gramme des Nations Unies pour l'Environnement (UNEP), IUCN publ. 250 p.
- Bulleri, F., Benedetti-Cecchi, L., Acunoto, S., Cinelli, F., Hawkins, S.J., 2002. The influence of canopy algae on vertical patterns of distribution of low-shore assemblages on rocky coasts in the northwest Mediterranean. J. Exp. Mar. Biol. 267, 89-106.
- Cadiou, G., Bonhomme, P., Boudouresque, C.F., Bellan, G., Bourcier, M., Emery, E., 2006. Etude et cartographie des biocénoses de la partie maritime du sous-site Natura 2000 "Cap Canaille - Grand Caunet". Phase III. Rapport final. Contrat Office National des Forêts 13/84, GIS Posidonie/IFREMER/COM, GIS Posidonie publ., Marseille Fr. 82 p. + annexes.
- Camous, A., 1912. Listes des Algues marines de Nice. Bull. Assoc. Nat. Nice Alpes-Maritimes 33 p.
- Canamella, A., 1968. Etude phytosociologique des horizons superficiels de l'étage infralittoral du port de la Coudouillère. DES Sci. Nat. Univ. Aix-Marseille 43 p. + 38 planches.
- Cardona, L., Moranta, J., Reñones, O., Hereu, B., 2013. Pulses of phytoplanktonic productivity may enhance sea urchin abundance and induce state shifts in Mediterranean rocky reefs. Estuar. Coast. Shelf Sci. 133, 88-96.
- Carvou, A., 1967. Etude de la végétation des horizons superficiels de l'étage infralittoral dans le port des Lecques (Var). DES, Faculté des Sciences de Marseille, 61 p.
- Carvou, A., Gadea, E., Vignes, P., 1968. Distribution comparée de 20 dominantes du peuplement algal dans le port des Lecques et le port Saint-Elme (Var.) Ann. Soc. Sci. Nat. Arch. Toulon Var 20, 1-16.
- Castagne, L., 1845. Catalogue des plantes qui croissent naturellement aux environs de Marseille. Imprimerie Nicot et Pardigon, Pont Moreau, 263 p.
- Cecchi, E., Piazzini, L., Serena, F., 2009. Applicazione del metodo CARLIT lungo la costa di Calafuria (Livorno). Atti Soc. Tosc. Sci. Nat. Mem. Ser. B 116, 123-125.
- Cecere, E., Cormaci, M., Furnari, G., Petrocelli, A., Saracino, O., Serio, D., 1996. Benthic algal flora of Cheradi Islands (Gulf of Taranto, Mediterranean Sea). Nova Hedwigia 62 (1), 191-214.
- Charbonnel, E., Bourcier, M., Francour, P., Boudouresque, C.F., 1993. Evolution des peuplements marins après la mise en service de la station d'épuration de la ville de Cassis (Bouches-du-Rhône). Comparaison 1881-1993 et mise en place d'un système de surveillance permanent. GIS Posidonie publ., Marseille, Fr., pp. 1-90.
- Clarisse, S., 1980-1981. Contribution à la connaissance de la phénologie de l'autoécologie et de l'épiphytisme de *Cystoseira balearica* Sauvageau en Baie de la Revellata (Calvi, Corse). Mémoire de Licence. Université de Liège, 77 p.
- Clarisse, S., 1984. Apport de différentes techniques cartographiques à la connaissance de l'autoécologie de *Cystoseira balearica* Sauvageau, macroalgue marine dominante dans la région de Calvi (Corse). Lejeunia 113, 1-24.
- Clarke, K.R., Gorley, R.N., 2006. Primer v6: User Manual/Tutorial. Primer-E Ltd., 190 p.
- Clarke, K.R., Warwick, R.M., 1994. Changes in Marine Communities: An Approach to Statistical Analysis and Interpretation. Plymouth Marine Laboratory, Plymouth, 144 p.
- Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Lasram, F.B.R., Aguzzi, J., Ballesteros, E., Bianchi, C.N., Guilhaumon, F., Kesner-Reyes, K., Militiadis-Spyridon, K., Koukouras, A., Lampadariou, N., Laxamana, E., López-Fé de la Cuadra, C.M., Lotze, H.K., Martin, D., Mouillot, D., Oro, D., Raicevich, S., Rius-Barile, J., Saiz-Salinas, J.L., San Vicente, C., Somot, S., Templado, J., Turon, X., Vafidis, D., Villanueva, R., Voultsiadou, E., 2010. The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. PLoS one 5 (8), e11842.
- Coppejans, E., 1972. Résultats d'une étude systématique et écologique de la population algale des côtes rocheuses du Dramont, St-Raphaël (France). Biol. Jb. Dodonaea 40, 153-180.
- Coppejans, E., 1977. Bijdrage tot de studie van de wierpopulaties (Chlorophyceae, Phaeophyceae, Rhodophyceae) van het fotofiel infralittoraal in het noordwestelijk Midderraan bekken. PhD Thesis Rijksuniversiteit Gent - Fakulteit des Wetenschappen. I-III. 339 p.
- Coppejans, E., 1979. Végétation marine de la Corse (Méditerranée). III. Documents pour la flore des algues. Marine Vegetation of Corsica (Mediterranean Sea). III. Documents for the Algal Flora. Bot. Acta 22 (4), 257-266.
- Cormaci, M., Furnari, G., 1999. Changes of the benthic algal flora of the Tremiti Islands (Southern Adriatic) Italy. Hydrobiologia 398, 75-79.
- Cormaci, M., Furnari, G., Alongi, G., Catra, M., Pizzuto, F., Serio, D., 2001. Spring marine vegetation on rocky substrata of the Tremiti Islands (Adriatic Sea, Italy). In: Faranda, F.M., Gigliemo, L., Spezie, G. (Eds.), Mediterranean Ecosystems. Springer Milan, pp. 245-254.
- Cormaci, M., Furnari, G., Catra, M., Alongi, G., Giaccone, G., 2012. Flora marina biontica del Mediterraneo: Phaeophyceae. Bollet. dell'Accad. Gioenia di Scienze Nat. Catania 45, 1-508.
- Cottalorda, J.M., Meinesz, A., Thibaut, T., Chiaverini, D., 2004. Représentation cartographique de l'abondance de quelques algues et invertébrés sur le littoral des îlots du Rascass et de la Gabinière (Parc National de Port-Cros, Var; France). Trav. Sci. Parc Nat. Port-Cros, Fr. 20, 195-209.
- Dayton, P.K., 1985. Ecology of kelp communities. Ann. Rev. Ecol. Syst. 16, 215-245.
- de Vaugelas, J., Meinesz, A., Culioli, J.M., 1998. Premiers éléments sur les peuplements sous-marins des îles Cerbicales (Corse-du-Sud). Trav. Sci. Parc Nat. Rég. Rés. Nat. Corse 57, 11-41.
- Debeaux, O., 1873. Énumération des algues marines du littoral de Bastia (Corse). Rev. Sci. Nat. 2 (2), 193-205.
- Decrock, E., 1914. Les Bouches-du-Rhône. Troisième partie. Le sol et ses habitants. Tome XII. le Sol. Chapitre IX à XIV. Photogéographie. Masson, Marseille, 12 p.
- Deleuil, G., Molinier, R. 1964. Excursion en Corse du 13 au 21 mai 1964. Institut de Botanique de Montpellier, 56 p.
- Desrosiers, G., Bellan-Santini, D., Brêthes, J.C., 1982. Evolution spatio-temporelle des peuplements de substrats rocheux superficiels dans un golfe soumis à de multiples pollutions (Golfe de Fos, France). Téthys 10 (3), 245-253.
- Desrosiers, G., Bellan-Santini, D., Brêthes, J.-C., 1986. Organisation trophique de quatre peuplements de substrats rocheux selon un gradient de pollution industrielle (Golfe de Fos, France). Mar. Biol. 91, 107-120.
- Devescovi, M., Ivesa, L., 2007. Short term impact of planktonic mucilage aggregates on macrobenthos along the Istrian rocky coast (Northern Adriatic, Croatia). Mar. Pollut. Bull. 54, 887-893.
- Díez, I., Secilla, A., Santolaria, A., Gorostiaga, J.M., 1999. Phytobenthic intertidal community structure along an environmental pollution gradient. Mar. Pollut. Bull. 39 (6), 462-472.
- Díez, I., Santolaria, A., Gorostiaga, J.M., 2003. Relationships of environmental factors with the structure and distribution of subtidal seaweed vegetation of the western Basque coast (N. Spain). Estuar. Coast. Shelf Sci. 56, 1041-1054.
- Díez, I., Santolaria, A., Secilla, A., Gorostiaga, J.M., 2009. Recovery stages over long-term monitoring of the intertidal vegetation in the 'Abra de Bilbao' area and on the adjacent coast (N. Spain). Eur. J. Phycol. 44 (1), 1-14.
- Epiard-Lahaye, M., 1988. Effects of ammonium, nitrate and phosphate on the growth of *Cystoseira stricta* (Phaeophyta, Fucales) cuttings in culture. Crypt. Alg. 9 (3), 211-229.
- Epiard-Lahaye, M., Pellegrini, M., Weiss, H., 1987. Influence des rythmes émigration-immersion sur le développement des boutures de *Cystoseira stricta* Sauvageau (Phéophycées, Fucales) en culture. Bot. Mar. 30, 259-266.
- Fairweather, P.G., 1990. Sewage and the biota on seashores: assessment of impact in relation to natural variability. Environ. Monit. Assess. 14 (2-3), 197-210.
- Falace, A., Alongi, G., Cormaci, M., Furnari, G., Curiel, D., Cecere, E., Petrocelli, A., 2010. Changes in the benthic algae along the Adriatic Sea in the last three decades. Chem. Ecol. 26 (S1), 77-90.
- Fée, A.L.A., 1833. Commentaires sur la botanique et la matière médicale de Pline. Tome 3, Imprimerie de C.L.F. Panckoucke, Paris, 468 p.
- Feldmann, J., 1937. Recherches sur la végétation marine de la Méditerranée. La Côte des Albères. Rev. Algol. 10, 339.
- Fernandez, C., Niell, F., 1982. Patterns of zonation in rocky inter-tidal shores at Cape-Penas-region (Asturies, north of Spain). Investigacion Pesquera 46 (1), 121-141.
- Filbee-Dexter, K., Scheibling, R.E., 2014. Sea urchin barrens as alternative stable states of collapsed kelp ecosystems. Mar. Ecol. Prog. Ser. 495, 1-25.
- Francour, P., Roberts, N., Bernard, G., Bonhomme, P., 1997. Cartographie des fonds marins, et en particulier de l'herbier à *Posidonia oceanica*, à proximité du port des Ourisnières (Le Pradet, Var). Contrat GIS Posidonie/Ville du Pradet. GIS Posidonie Publ., Fr. 52p.

- Francour, P., Ganteaume, A., Bodilis, P., Cottalorda, S., 2003. Etat actuel des fonds du littoral de la ville d'Antibes. Rapport final. Contrat Ville d'Antibes et Laboratoire Environnement Marin Littoral. LEML Publ., Nice 74 p.
- Fraschetti, S., Terlizzi, A., Guarnieri, G., Pizzolante, F., D'Ambrosio, P., Maiorano, P., Beqiraj, S., Boero, F., 2011. Effects of unplanned development on marine biodiversity: a lesson from Albania (central Mediterranean Sea). *J. Coast. Res.* 58, 106–115.
- Fraschetti, S., Bevilacqua, S., Guarnieri, G., Terlizzi, A., 2012. Idiosyncratic effects of protection in a remote marine reserve. *Mar. Ecol. Progr. Ser.* 466, 21–34.
- Frick, H.G., Boudouresque, C.F., Verlaque, M., 1996. A checklist of marine algae of the Lavezzi Archipelago; with special attention to species rare or new to Corsica (Mediterranean). *Nova Hedwigia* 62 (1–2), 119–135.
- Gadea, E., 1967. Contribution à l'étude écologique et phytosociologique des peuplements marins des horizons superficiels de l'étage infralittoral dans le port de Saint-Elme (Var). DES, Faculté des Sciences de Marseille, 42 p.
- Galil, B.S., 2000. A sea under siege – alien species in the Mediterranean. *Biol. Inv.* 2, 177–186.
- Ganteaume, A., Bonhomme, P., Ruitton, S., Verlaque, M., Boudouresque, C.F., 2004. Etat des lieux et des potentialités du milieu marin autour des Îles Sanguinaires (Corse du Sud). Contrat GIS Posidonie et Association des Amis du Parc de Corse. GIS Posidonie Publ., 60 p.
- Gasquet, J., 1968. Contribution à l'étude écologique et phytosociologique des peuplements marins du port de Carqueiranne [Var]. DES, Faculté des Sciences de Marseille, 110 p.
- Giaccone, G., Alongi, G., Pizzuto, F., Cossu, A., 1994. La Vegetazione marina bentonica fotofila del Mediterraneo: 2. Infralitorale e Circalitorale: proposte di aggiornamento. *Boll. Accad. Gioenia Sci. Nat. Catania* 27 (346), 111–157.
- Giakoumi, S., Cebrian, E., Kokkoris, G.D., Ballesteros, E., Sala, E., 2012. Relationships between fish, sea urchins and macroalgae: the structure of shallow rocky sublittoral communities in the Cyclades, Eastern Mediterranean. *Est. Coast. Shelf Sci.* 109, 1–10.
- Gilet, R., 1954. Particularités de la zonation marine des côtes rocheuses s'étendant entre Nice et la frontière italienne. *Rec. Trav. Stat. Mar. Endoume* 12, 41–49.
- Gilet, R., Molinier, R., Picard, J., 1954. Etudes bionomiques littorales sur les côtes de Corse. *Rec. Trav. Stat. Mar. Endoume* 13, 25–53.
- Griati, J., Mannoni, P.A., Meinesz, A., 2006. Cartographie des espèces méditerranéennes et infralittorales supérieures du pourtour de l'île de Porquerolles (Var). LEML report, Nice 24 p.
- Griati, J., Mannoni, P.A., Meinesz, A., 2007. Cartographie des espèces méditerranéennes et infralittorales supérieures du pourtour de l'île de Porquerolles (Var). LEML report, Nice 30 p.
- Gravez, V., Bouladier, F., Boudouresque, C.F., 1988. Aménagement du site de Corbière: le milieu marin. In: Aménagement balnéaire sur le site de Corbières. Etude d'impact. GIS Posidonie publ., Marseille, 37 p. + annexes.
- Gravez, V., Charbonnel, E., Bonhomme, P., Cadiou, G., 1999. Inventaire biologique des fonds de l'anse des Roseaux, presqu'île de Saint-Mandrier (Var). Octobre 1999. BCEOM, Direction régionale de Fréjus et GIS Posidonie. GIS Posidonie publ., Marseille, 51 p.
- Guglielmi, G., 1969. Contribution à l'étude des algues du Cap Ferrat. DES, Faculté des Sciences de Nice, 100 p.
- Guiry, M.D., Guiry, G.M., 2014. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <<http://www.algaebase.org>>; searched on 24.07.14.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agora, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global map of human impact on marine ecosystems. *Science* 319 (5865), 948–952.
- Hamel, G., 1931–1939. *Phéophycées de France*, Imprimerie Wolf, Rouen, XLVI, 431 p.
- Hawkins, S.J., Moore, P.J., Burrows, M.C., Poloczanska, E., Mieszkowska, N., Herbert, R.J., Jenkins, S.R., Thompson, R.T., Genner, M.J., Southward, A.J., 2008. Complex interactions in a rapidly changing world: responses of rocky shore communities to recent climate change. *Clim. Res.* 37, 123–133.
- Helmuth, B., Mieszkowska, N., Moore, P., Hawkins, S.J., 2006. Living on the edge of two changing worlds: forecasting the responses of rocky intertidal ecosystems to climate change. *Annu. Rev. Ecol. Syst.*, 373–404.
- Hereu, B., Zabala, M., Sala, E., 2008. Multiple controls of community structure and dynamics in a sublittoral marine environment. *Ecology* 89 (12), 3423–3435.
- Husa, V., Steen, H., Sjøtun, K., 2014. Historical changes in macroalgal communities in Hardangerfjord (Norway). *Mar. Biol.* 10 (3), 226–240.
- Huvé, P., 1960. Résultats sommaires de l'étude expérimentale de la réinstallation d'un peuplement de *Cystoseira stricta* (Montagne) Sauvageau. *Rapp. P.V. Réunion CIESM* 15 (2), 121–125.
- Huvé, P., 1970. Recherche sur la genèse de quelques peuplements algaux marins de la roche littorale de la région de Marseille. Thèse Doctorat d'Etat. Faculté des Sciences de Paris 479 p.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlanson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R., 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293, 629–638.
- Jaffrenou, F., Bonnin, A., Charrier, S., 1996. Répartition d'algues remarquables de la frange littorale du Cap Ferrat (Alpes-Maritimes). *Riviera Scientifique*, 41–46.
- Jahandiez, E., 1929. Les Îles d'Hyères. Monographie des îles d'Or. Presqu'île de Giens, Porquerolles, Port-Cros, Ile du Levant. Histoire, description Géologie, Flore, Faune. Troisième édition, Rébua et Rouard publ., Toulon, 447 p.
- Janssens, M., 1991. Etude de la répartition de différentes macroalgues et algues bleues filamenteuses dans la région de Calvi. Mémoire de Licence. Université de Liège, 90 p.
- Janssens, M., Hoffmann, L., Demoulin, V., 1993. Cartographie des macroalgues dans la région de Calvi (Corse): comparaison après 12 ans. (1978–1979, 1990–1991). *Lejeunia* 141, 62 p.
- Javel, F., Meinesz, A., Chiaverini, D., Thibaut, T., Cottalorda, J.M., 2005. Cartographie des espèces méditerranéennes et infralittorales supérieures des falaises de Bonifacio au Cap di Feno (Corse du Sud). Contrat Office de l'Environnement de la Corse – GIS Posidonie. GIS Posidonie – LEML-UNSA publ., Nice 23 p.
- Kautsky, H., Kautsky, L., Kautsky, N., Kautsky, U., Lindblad, C., 1992. Aspects on *Fucus vesiculosus* in the Baltic Sea. *Acta Phytogeogr. Suec.* 78, 33–48.
- Lamela-Silvarrey, C., Fernández, C., Anadón, R., Arrontes, J., 2012. Fucoid assemblages on the north coast of Spain: past and present (1977–2007). *Bot. Mar.* 55 (3), 199–207.
- Leblond, E., 1924. Algues du littoral septentrional du Golfe d'Ajaccio. *Rev. Algol.* 1 (2), 156–161.
- Lejeune, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C.F., Pérez, T., 2010. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends Ecol. Evol.* 25 (4), 250–260.
- Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B., 2006. Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* 312 (5781), 1806–1809.
- Lotze, H.K., Coll, M., Dunne, J.A., 2010. Historical changes in marine resources, food-web structure and ecosystem functioning in the Adriatic Sea, Mediterranean. *Ecosystems* 14 (2), 198–222.
- Mačić, V., 2006. Distribution of seaweed *Fucus virsoides* J. Agardh in Boka Kotorska bay (South Adriatic Sea). *Ann. Ser. Hist. Nat.* 16 (1), 1–4.
- Mačić, V., Antolich, B., Thibaut, T., Svircev, Z., 2010. Distribution of the most common *Cystoseira* C. Agardh species (Heterokontophyta, Fucales) on the coast of Montenegro (south-east Adriatic sea). *Fres. Env. Bull.* 19 (6), 1191–1198.
- Maggi, P., 1967. Contribution à l'étude des peuplements des horizons superficiels de l'étage infralittoral du port de Saint-Mandrier (Var). DES Sciences Naturelles, Marseille, 59p.
- Maggi, E., Bertocci, I., Vaselli, S., Benetti-Cecchi, L., 2009. Effects of changes in number, identity and abundance of habitat-forming species on assemblages of rocky seashores. *Mar. Ecol. Progr. Ser.* 381, 39–49.
- Maggi, E., Bulleri, F., Bertocci, I., Benedetti-Cecchi, L., 2012. Competitive ability of macroalgal canopies overwhelms the effects of variable regimes of disturbance. *Mar. Ecol. Progr. Ser.* 465, 99–109.
- Magne, F., 1956. La végétation marine du Grand Congloué. Résultats scientifiques des campagnes de la "Calypto". Fascicule II. Masson et Cie Publ. Paris, pp. 163–184.
- Mangialajo, L., Chiantore, M., Cattaneo-Vietti, R., 2008. Loss of fucoid algae along a gradient of urbanisation, and structure of benthic assemblages. *Mar. Ecol. Progr. Ser.* 358, 63–74.
- Mangialajo, L., Chiantore, M., Susini, M.L., Meinesz, A., Thibaut, T., 2012. Zonation patterns and interspecific relationships of fucoids in microtidal environments. *J. Exp. Mar. Biol. Ecol.* 421, 72–80.
- Mari, X., Meinesz, A., de Vaugelas, J., 1998. Répartition de *Lithophyllum lichenoides* de *Cystoseira amantacea*, de *Patella ferruginea* et des zones polluées par les hydrocarbures de l'île Lavezzi (Réserve naturelle des Lavezzi, Corse). *Trav. Sci. Parc Nat. Rég. Rés. Nat. Corse* 57, 145–162.
- Marion, A.F., 1883. Esquisse d'une topographie zoologique du Golfe de Marseille. *Ann. Mus. Hist. Nat. Marseille* 1, 108p.
- Meinesz, A., Cottalorda, J.M., de Vaugelas, J., 1994. Valorisation du domaine maritime de la commune de Vallauris-Golf-Juan. Rapport final. LEML-UNSA publ., Nice, 127 p.
- Meinesz, A., de Vaugelas, J., Chiaverini, D., Bialecki, K., Cottalorda, J.M., Molenaar, H., 1999. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de la Réserve naturelle de Scandola (Corse). LEML-UNSA Publ., Nice, 8 p.
- Meinesz, A., Cottalorda, J.M., Chiaverini, D., de Vaugelas, J., 2000a. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de la face nord du Parc National de Port-Cros. LEML-UNSA Publ., 22p.
- Meinesz, A., de Vaugelas, J., Cottalorda, J.M., Chiaverini, D., Francour, P., Thibaut, T., 2000b. Flore faune et écosystèmes sous marins du littoral de Nice. LEML-UNSA Publ., 167 p.
- Meinesz, A., Chiaverini, D., Cottalorda, J.M., Gilletta, L., Javel, F., Molenaar, H., 2001a. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de la face sud du Parc National de Port-Cros. LEML-UNSA Publ., 27 p.
- Meinesz, A., Cottalorda, J.M., Thibaut, T., Chiaverini, D., de Vaugelas, J., 2001b. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de la face est de l'île de Port-Cros (Parc National). LEML-UNSA Publ., 9 p.
- Meinesz, A., Cottalorda, J.M., Chiaverini, D., de Vaugelas, J., 2001c. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de l'Îlot Bagaud (Parc National de Port-Cros). *Trav. Sci. Parc Nat. Port-Cros* 18, 123–141.
- Meinesz, A., Cottalorda, J.M., Chiaverini, D., de Vaugelas, J., 2001d. Représentation cartographique de l'abondance de quelques algues et invertébrés du littoral de l'île de Cavallo (archipel des Lavezzi). LEML-UNSA Publ., 114 p.
- Meinesz, A., Javel, F., Chiaverini, D., Cottalorda, J.M., Gilletta, L., 2004a. Représentation cartographique de l'abondance de quelques algues et

- invertébrés du littoral de la face ouest du Parc National de Port-Cros. LEML-UNSA Publ., 26 p.
- Meinesz, A., Javel, F., Cottalorda, J.M., Susini, M.L., Capiomont, A., Levi, F., Robert, A., 2004b. Cartographie des espèces méditerranéennes et infralittorales supérieures des falaises de Bonifacio (Corse-du-Sud) – Mission 2003 – Rapport intermédiaire. Contrat Office de l'Environnement de la Corse – GIS Posidonie. LEML-UNSA Publ., 14 p.
- Meinesz, A., Blanfuné, A., Chancollon, O., Javel, F., Longepierre, S., Markovic, L., Vaugelas, de J. de, Garcia, D., 2013. Côtes méditerranéennes françaises: inventaire et impacts des aménagements gagnés sur la mer. Lab. ECOMERS (ed.), Université Nice Sophia Antipolis, 156 p. and e-publication: <<http://www.medam.org>> (accessed 22.07.14).
- Menconi, M., Benedetti-Cecchi, L., Cinelli, F., 1999. Spatial and temporal variability in the distribution of algae and invertebrates on rocky shores in the northwest Mediterranean. J. Exp. Mar. Biol. Ecol. 233, 1–23.
- Micheli, F., Benedetti-Cecchi, L., Gambaccini, S., Bertocci, I., Borsini, C., Osio, G.C., Romano, F., 2005. Cascading human impacts, marine protected areas, and the structure of Mediterranean reef assemblages. Ecol. Mon. 75 (1), 81–102.
- Micheli, F., Levin, N., Giakoumi, S., Katsanevakis, S., Abdulla, A., Coll, M., Fraschetti, S., Kark, S., Koutsoubas, D., Mackelworth, P., Maiorano, L., Possingham, H.P., 2013. Setting priorities for regional conservation planning in the Mediterranean Sea. PLoS One 8 (4), e59038.
- Milazzo, M., Chemello, R., Badalamenti, F., Riggio, S., 2002. Short-term effect of human trampling on the upper infralittoral macroalgae of Ustica Island MPA (western Mediterranean, Italy). J. Mar. Biol. Ass. UK 82, 745–748.
- Millot, C., 1987. Circulation in the Western Mediterranean Sea. Ocean. Acta 10 (2), 143–149.
- Molinier, R., 1956. Note explicative de la carte des groupements végétaux terrestres et des peuplements marins superficiels de l'île du Grand Ribaud (Var). CRNS Publ., Fr. 8 p. + carte.
- Molinier, R., 1960. Etude des biocénoses marines du Cap Corse. Vegetatio 9 (4–5), 217–312.
- Molinier, R., Molinier, R., 1955. Eléments de bionomie marine et de phytosociologie aux îles Sanguinaires (Corse). Rév. Gén. Bot. 62, 1–8.
- Montagne, C., 1846. Phycée. In: Gaudichaud, C. (Ed.), Voyage autour du monde exécuté pendant les années 1836 et 1837 sur la corvette 'La Bonite' commandée par M. Vaillant. Botanique. Vol. 1: Cryptogames cellulaires et vasculaires, Paris 112p.
- Montefalcone, M., Rovere, A., Parravicini, V., Albertelli, G., Morri, C., Bianchi, C.N., 2013. Evaluating change in seagrass meadows: a time-framed comparison of side-scan sonar maps. Aquatic Botany 104, 204–212.
- Mouret, M., 1911. Liste des algues marines du Var. Ann. Soc. Sci. Nat. Arch. Toulon Var 2, 78–107.
- Munda, I.M., 1974. Changes and succession in the benthic algal associations of slightly polluted habitats. Rev. Int. Oceanogr. Med. 34, 37–52.
- Munda, I.M., 1980. Changes in the benthic algal associations of the vicinity of Rovinj (Istrian coast north Adriatic) caused by organic wastes. Acta Adriat. 21 (2), 299–332.
- Munda, I.M., 1982. The effects of organic pollution on the distribution of fucoid algae from the Istrian coast (vicinity of Rovinj). Acta Adriat. 23, 329–337.
- Munda, I.M., 1993. Changes and degradation of seaweed stands in the Northern Adriatic. Hydrobiologia 260 (261), 239–253.
- Nieri, M., Francour, P., Sinnasamy, J.M., Urscheller, F., Fleury, M.C., 1991. Les fonds marins de l'anse du Rouet (Carry-le-Rouet, Bouches-du-Rhône) Étude de site. Gis Posidonie Publ., Marseille, Fr. 107 p. + 1 carte.
- Nikolic, V., Zuljevic, A., Mangialajo, L., Antolic, B., Kuspilic, G., Ballesteros, E., 2013. Cartography of littoral rocky-shore communities (CARLIT) as a tool for ecological quality assessment of coastal waters in the Eastern Adriatic Sea. Ecol. Indic. 34, 87–93.
- Ollivier, G., 1929. Etude de la flore marine de la Côte d'Azur. Ann. Inst. Oceanogr. Paris 7 (3), 53–173.
- Orfanidis, S., Panayotidis, P., Stamatis, N., 2001. Ecological evaluation of transitional and coastal waters: a marine benthic macrophyte-based model. Medit. Mar. Sci. 2, 45–65.
- Orlandi, L., Bentivoglio, F., Carlino, P., Calizza, E., Rossi, D., Costantini, M.L., Rossi, L., 2014. $\Delta^{15}\text{N}$ variation in *Ulva lactuca* as a proxy for anthropogenic nitrogen inputs in coastal areas of Gulf of Gaeta (Mediterranean Sea). Mar. Pollut. Bull. 84 (1–2), 76–82.
- Panayotidis, P., Montesanto, B., Orfanidis, S., 2004. Use of low-budget monitoring of macroalgae to implement the European Water Framework Directive. J. Appl. Phyc. 16, 49–59.
- Papenfuss, G.F., 1968. A history, catalogue, and bibliography of Red Sea benthic algae. Israël J. Bot. 17, 118p.
- Parravicini, V., Micheli, F., Montefalcone, M., Morri, C., Villa, E., Castellano, M., Povero, P., Bianchi, C.N., 2013. Conserving biodiversity in a human-dominated world: degradation of marine sessile communities within a protected area with conflicting human uses. PLoS One 8 (10), e75767.
- Pellegrini, L., Pellegrini, M., 1970. Sur d'éventuelles possibilités d'exploitation de quelques Cystoseires méditerranéennes. Ann. Soc. Sci. Nat. Arch. Toulon Var 22, 133–138.
- Pellegrini, L., Pellegrini, M., 1972. Contribution à l'étude biochimique des Cystoseiracées méditerranéennes. Bull. Soc. Museum His. Nat. Marseille 32, 101–107.
- Perkol-Finkel, S., Airolidi, L., 2010. Loss and recovery potential of marine habitats: an experimental study of factors maintaining resilience in subtidal algal forests at the Adriatic Sea. PLoS One 5 (5), e10791.
- Picard, J., 1954. Modifications saisonnières des peuplements de l'horizon inférieur de la roche littorale. C. R. Hebd. Séances. Acad. Sci. 238 (12), 1358–1359.
- Picard, J., Bourcier, M., 1975. Evolution sous influence humaine des peuplements benthiques des parages de La Ciotat entre 1954 et 1972. Téthys 2 (2–3), 213–222.
- Pignatti, S., 1962. Associazioni di alghe marine sulla costa veneziana. Memorie del Reale Istituto Veneto di Scienze, Lettere ed Arti, Italy, 32, 134p.
- Raphélis, A., 1924a. Additions à la flore des algues de Cannes. Rev. Algol. 1 (2), 162–167.
- Raphélis, A., 1924b. Inventaire des algues du musée de Nice. I. Album Bonfils. Riviera Sci. 11 (2), 32–36.
- Raphélis, A., 1924c. Inventaire des algues du musée de Nice. II. Herbarier Sarato. Riviera Sci. 11 (3), 49–52.
- Raybaud, V., Beaugrand, G., Goberville, E., Delebecq, G., Destombe, C., Valero, M., Davoult, D., Morin, P., Gevaert, F., 2013. Decline in kelp in west Europe and climate. PLoS One 8 (6), e66044.
- Ribera, M.A., Gomez-Garreta, A., Gallardo, T., Cormaci, M., Furnari, G., Giaccone, G., 1992. Check-list of Mediterranean seaweeds. I. Fucophyceae (Warming, 1884). Bot. Mar. 35, 109–130.
- Rodriguez-Prieto, C., 1992. Estudi de l'estructura, la dinàmica i la fenologia de la comunitat de *Cystoseira mediterranea* Sauvageau: la importància de les minves d'hivern. Mem. Doct. Ciènc. Inst. Ecol. aquat. Univ. autonoma Barcelona, 187 p.
- Rodriguez-Prieto, C., Polo, L., 1996. Effects of sewage pollution in the structure and dynamics of the community of *Cystoseira mediterranea* (Fuciales, Phaeophyceae). Sci. Mar. 60, 253–263.
- Rueness, J., Fredriksen, S., 1991. An assessment of possible pollution effects on the benthic algae of the outer Oslofjord. Ophelia 17, 223–235.
- Ruitton, S., Francour, P., Boudouresque, C.F., 2000. Relationship between algae, benthic herbivorous invertebrates and fishes in rocky sublittoral communities of a temperate sea (Mediterranean). Estuar. Coast. Shelf Sci. 50, 217–230.
- Ruitton, S., Bonhomme, P., Cadiou, G., Harmelin, J.G., Perez, T., 2005. Inventaire du patrimoine naturel sous-marin des faces Est et Sud de Porquerolles – Substrats durs et Herbière à Posidonie oceanica. Rapport final. Contrat Parc national de Port-Cros et GIS Posidonie, GIS Posidonie Publ., France 63 p. + annexes.
- Sáenz-Arroyo, A., Roberts, C.M., Torre, J., Cariño-Olvera, M., 2005. Using fishers' anecdotes, naturalists' observations and grey literature to reassess marine species at risk: the case of the Gulf grouper in the Gulf of California. Mexico. Fish. Fish. 6 (2), 121–133.
- Sala, E., Kizilkaya, Z., Yildirim, D., Ballesteros, E., 2011. Alien marine fishes deplete algal biomass in the eastern Mediterranean. PLoS One 6 (2), e17356.
- Sala, E., Ballesteros, E., Dendrinis, P., Di Franco, A., Ferretti, F., Foley, D., Fraschetti, S., Foley, D., Fraschetti, S., Friedlander, A., Garrabou, J., Güçlüsoy, H., Guidetti, P., Halpern, B.S., Hereu, B., Karamanlidis, A., Kizilkaya, Z., Macpherson, E., Mangialajo, L., Mariani, S., Micheli, F., Pais, A., Riser, K., Rosenberg, A.A., Sales, M., Selkoe, K.A., Starr, R., Tomas, F., Zabala, M., 2012. The structure of Mediterranean rocky reef ecosystems across environmental and human gradients, and conservation implications. PLoS One 7 (2), e32742.
- Sales, M., Cébrian, E., Tomas, F., Ballesteros, E., 2011. Pollution impacts and recovery potential in three species of the genus *Cystoseira* (Fuciales, Heterokontophyta). Estuar. Coast. Shelf Sci. 92 (3), 347–357.
- Santini, D., 1961. Note sur les peuplements sciaphiles de l'étagé infralittoral rocheux de la région de Bonifacio. Rec. Trav. St. Mar. End. Bull. 23 (37), 61–70.
- Sauvageau, C., 1912. A propos des *Cystoseira* de Banyuls et de Guéthary. Bull. Stat. Biol. Arcachon 14, 424p.
- Schiel, D.R., 2011. Biogeographic patterns and long-term changes on New Zealand coastal reefs: Non-trophic cascades from diffuse and local impacts. J. Exp. Mar. Biol. Ecol. 400 (1), 33–51.
- Schiel, D.R., Foster, M.S., 2006. The population biology of large brown seaweeds: ecological consequences of multiphase life histories in dynamic coastal environments. Ann. Rev. Ecol. Evol. Syst. 37, 343–372.
- Serio, D., Alongi, G., Catra, M., Cormaci, M., Furnari, G., 2006. Changes in the benthic algal flora of Lìnos Island (Strait of Sicily, Mediterranean Sea). Bot. Mar. 49, 135–144.
- Serrano, O., Mateo, M.A., Dueñas-Bohóquez, A., Renom, P., López-Sáez, J.A., Martínez Cortizas, A., 2011. The *Posidonia oceanica* marine sedimentary record: a Holocene archive of heavy metal pollution. Sci. Tot. Environ. 409, 4831–4840.
- Sfriso, A., Facca, C., 2011. Macrophytes in the anthropic constructions of the Venice littorals and their ecological assessment by an integration of the "CARLIT" index. Ecol. Indic. 11 (3), 772–781.
- Sinnassamy, J.M., Gravez, V., Boudouresque, C.F., Conejero, S., Ripert, S., 1993. Le milieu marin du site de Corbières (Bouches-du-Rhône). Premier suivi (1993). DGST Ville de Marseille et GIS Posidonie Publ., France 62p.
- Soltan, D., 2001. Étude de l'incidence de rejets urbains sur les peuplements superficiels de macroalgues en Méditerranée nord-occidentale. PhD Thesis. Université de la Méditerranée-Centre d'Océanologie de Marseille. France, 157 p.
- Soltan, D., Verlaque, M., Bourouesque, C.F., Francour, P., 2001. Changes in macroalgal communities in the vicinity of a Mediterranean sewage outfall after the setting up of a treatment plant. Mar. Pollut. Bull. 42, 59–71.
- Steneck, R.S., Graham, M.H., Bourque, B.J., Corbett, D., Erlandson, J.M., Estes, J.A., Tegner, M.J., 2002. Kelp forest ecosystems: biodiversity, stability, resilience and future. Environ. Cons. 29 (4), 436–459.
- Susini, M.L., 2006. Statut et biologie de *Cystoseira amentacea* var. *stricta*. PhD Thesis. Université Nice-Sophia Antipolis, France, 236 p.

- Susini, M.L., Mangialajo, L., Thibaut, T., Meinesz, A., 2007. Development of a transplantation technique of *Cystoseira amentacea* var. *stricta* and *Cystoseira compressa*. *Hydrobiologia* 580, 241–244.
- Tamburello, L., Bulleri, F., Bertocci, I., Maggi, E., Benedetti-Cecchi, L., 2013. Reddened seascapes: experimentally induced shifts in 1/f spectra of spatial variability in rocky intertidal assemblages. *Ecology* 94 (5), 1102–1111.
- Taupier-Letage, I., Piazzola, J., Zakardjian, B., 2013. Les îles d'Hyères dans le système de circulation marine et atmosphérique de la Méditerranée. *Sci. Rep. Port-Cros Natl. Park* 27, 29–52.
- Templado, J., 2014. Future trends of Mediterranean biodiversity. In: Goffredo, S., Dubinsky, Z. (Eds.), *The Mediterranean Sea: its History and Present Challenges*. Springer, Netherlands, pp. 479–498.
- Tewari, A., Joshi, H.V., 1988. Effect of domestic sewage and industrial effluents on biomass and species diversity seaweeds. *Bot. Mar.* 31 (5), 389–397.
- Thibaut, T., Pinedo, S., Torras, X., Ballesteros, E., 2005. Long-term decline of the populations of Fucales (*Cystoseira*, *Sargassum*) in the Albères coast (northwestern Mediterranean). *Mar. Pollut. Bull.* 50, 1472–1489.
- Thomas, A., 1983. Contribution à une étude d'impact des activités humaines du littoral du Cap de l'Aigle (Bouches-du-Rhône), au Cap Garonne. La dynamique sédimentaire et l'herbier à Posidonies. Tome 2. PhD Thesis Université Aix-Marseille II. France 199 p.
- Tournefort, J.P., 1717. Relation d'un voyage du Levant fait par ordre du Roy. Tome 1. Paris, Imprimerie Royale, 653 p.
- Tsiamis, K., Panayotidis, P., Salomidi, M., Pavlidou, A., Kleinteich, J., Balanika, K., Küpper, F.C., 2013. Macroalgal community response to re-oligotrophication in Saronikos Gulf. *Mar. Ecol. Prog. Ser.* 472, 73–85.
- Underwood, A., 1981. Techniques of analysis of variance in experimental marine biology and ecology. *Oceanogr. Mar. Biol. Annu. Rev.* 19, 513–605.
- UNEP/MAP, 2012. State of the Mediterranean Marine and Coastal Environment, UNEP/MAP – Barcelona Convention, Athens, 96 p.
- Valls, R., Piovetti, L., Praud, A., 1993. The use of diterpenoids as chemotaxonomic markers in the genus *Cystoseira*. *Hydrobiologia* 260 (1), 549–556.
- Vergés, A., Alcoverro, T., Ballesteros, E., 2009. Role of fish herbivory in structuring the vertical distribution of canopy algae *Cystoseira* spp. in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 375, 1–11.
- Verlaque, M., 1975. Compte rendu de la première mission effectuée, sur la façade maritime du Parc Naturel régional de Corse, par l'équipe de phytosociologie benthique marine de Luminy. Université Aix-Marseille II, Marseille, 64 p.
- Verlaque, M., 1987. Contribution à l'étude du phytobenthos d'un écosystème photophile thermophile marin en Méditerranée occidentale. Etude structurale et dynamique du phytobenthos et analyses des relations Faune-Flore. PhD Thesis. Université d'Aix-Marseille II, Marseille 389p.
- Verlaque, M., Bernard, G., 1998. Inventaire de la flore marine de la principauté de Monaco. Avril 1998. Contrat Ministère d'état de la principauté de Monaco-Service environnement-GIS Posidonie. Gis Posidonie publ. France 39 p. + Annexes.
- Verlaque, M., Tiné, J., 1979. Végétation marine de Toulon (Var-France): Grande Rade et Rade-abri. Université Aix-Marseille II, Marseille, 83p.
- Verlaque, M., Francour, P., Sartoretto, S., 1998. Evaluation de la valeur patrimoniale des biocénoses de la face ouest de l'îlot de Gargalu (Réserve naturelle de Scandola). Contrat Direction Régionale de l'Environnement and GIS Posidonie. UMR DIMAR and GIS Posidonie publ. Marseille, 66p.
- Vignes, P., 1965. Contribution à l'étude écologique et phytosociologique des peuplements marins du port des Salins d'Hyères (Var). *L'information Sci.* 3, 21.
- Vignes, P., 1968. Etude écologique des peuplements marins superficiels sur substrat rocheux (baie de Cavalas, Var). *Annuaire du centre régional de documentation pédagogique*, 40p.
- Waycott, M., Duarte, C.M., Carruthers, T.J., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck Jr, K.L., Hugues, A.R., Kendrick, G.A., Kenworthy, W.J., Short, F.T., Williams, S.L., 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. Nat. Acad. Sci.* 106 (30), 12377–12381.
- Wernberg, T., Thomsen, M.S., Tuya, F., Kendrick, G.A., Staehr, P.A., Toohey, B.D., 2010. Decreasing resilience of kelp beds along a latitudinal temperature gradient: potential implications for a warmer future. *Ecol. Lett.* 13 (6), 685–694.
- Worm, B., Chapman, A.R.O., 1996. Interference competition among two seaweeds: *Chondrus crispus* strongly affects survival of *Fucus evanescens* recruits. *Mar. Ecol. Prog. Ser.* 145 (2–3), 297–301.
- Worm, B., Lotze, H.K., 2006. Effects of eutrophication, grazing, and algal blooms on rocky shores. *Limnol. Oceanogr.* 51 (1), 569–579.
- Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Watson, R., et al., 2006. Impacts of biodiversity loss on ocean ecosystem services. *Science* 314 (5800), 787–790.
- Zavodnik, N., 1975. Effects of temperature and salinity variations on photosynthesis of some littoral seaweeds of the north Adriatic Sea. *Bot. Mar.* 18, 245–250.
- Zenetos, A., Gofas, S., Verlaque, M., Cinar, M.E., García Raso, J.E., Bianchi, C.N., Morri, C., Azzurro, E., Bilecenoglu, M., Frogia, C., Siokou, S., Violanti, D., Sfriso, A., San Martín, G., Giangrande, A., Katağan, T., Ramos-Esplá, A., Mastrotoato, F., Ocaña, O., Zingone, A., Gambi, M.C., Streftaris, N., 2010. Alien species in the Mediterranean Sea by 2010 A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Medit. Mar. Sci.* 11 (2), 381–493.