Tunable construction of transition metal-coordinated helicene cages

Yongle Ding^{a, b}, Chengshuo Shen^{a, *}, Fuwei Gan^a, Jinghao Wang^a, Guoli Zhang^a, Lingling Li^b, Mouhai Shu^a, Bangshang Zhu^{b, *}, Jeanne Crassous^c, Huibin Qiu^{a, *}^a School of Chemistry and Chemical Engineering, Frontiers Science Center for Transformative Molecules, State Key Laboratory of Metal Matrix Composites, Shanghai Jiao Tong University, Shanghai 200240, China^b Instrumental Analysis Center, Shanghai Jiao Tong University, Shanghai 200240, China^c Univ Rennes, Institut des Sciences Chimiques de Rennes, UMR CNRS 6226, Campus de Beaulieu, Rennes 35042, France

ABSTRACT

We report a facile and tailored method to prepare globally twisted chiral molecular cages through tunable coordination of bis-bipyridine-terminated helicene ligands to a series of transition metals including Fe(II), Co(II), Ni(II) and Zn(II). This system shows an efficient remote transfer of stereogenicity from the helicene core to the bipyridine-metal coordination sites and subsequently the entire cages. While the Fe(II), Co(II) and Ni(II)-derived M_2L_3 (M for metal and L for ligand) cages exhibit quasi-reversible redox features, the Zn(II) analogues reveal prominent yellow circularly polarized luminescence. Interestingly, with the addition of Na_2SO_4 , the Zn_2L_3 cages reassemble into sextuple-stranded $Zn_6L_6(SO_4)_4$ cages in which three Zn_2L_2 units are bound together by four sulfates and further coalesced by offset inter-ligand π - π interactions.

Keywords:

Helicene

Coordination cage

Chiral

Global helicity

Circularly polarized luminescence

Molecular cages with well-defined structures and internal cavities have raised abundant interest in recent years [1,2]. The synthesis of these intrinsic hollow species normally involved tailored connections of molecular building blocks either through covalent bonding or coordination to metals [3–9]. The flexible linkage chemistry along with the diversity in building blocks has favored an unprecedentedly wide access to cages with tunable geometries and functions [10–12]. Amongst, chiral molecular cages have drawn significant attentions due to their potential applications in chiral sensing, chiral separation and asymmetric catalysis [13–17]. So far, a series of chiral molecular building blocks have been adopted to enantioselectively create these chiral entities [18,19]. However, the facile construction of chiral cages with highly extended, globally distorted unsymmetrical structures remains a great challenge, majorly as a consequence that the conventional chiral molecular building blocks normally fail to efficiently transfer the local chirality to the whole cage skeletons in spite of a few advances (mainly in helicates) achieved by axially chiral building blocks [20–22].

Helicenes are a type of chiral molecules constituted by *ortho*-fused aromatic rings with unique helical shapes [23–25]. Helicene-cored molecule generally exhibits an apparent three-dimensionally twisted structure with chirality effectively expressed across the whole molecule along with the remote substituents [26,27]. Normally, carbo[*n*]helicenes with the number of rings (*n*) larger than 5 possess stable configuration against racemization [28]. We have previously demonstrated that helicene derivatives can serve as building blocks to construct covalent organic cages, which presented a globally distorted, triple-stranded helical structure with high recognition ability towards chiral guest molecules [29]. Clever *et al.* further synthesized bis-pyridine-terminated helicenes and used them as

ligands to prepare a series of Pd-coordinated chiral cages, which also showed promising guest recognition features [30]. However, the structure and function of these helicene cages is so far limited by the diversity of helicene precursor and linkage chemistry. Herein, we introduce two 2,2'-bipyridine moieties onto the 4,13-positions of [6]helicene through two alkyl bridges and develop a globally bent ligand, which favors a tunable coordination to a variety of transition metals with efficient transfer of stereogenicity to the coordination sites, and hence allow the formation of coordination cages with unique triple-stranded binuclear or sextuple-stranded hexanuclear structures and distinctive properties (Scheme 1).

Scheme 1. Schematic representation for the synthesis of helicene-derived chiral coordination cages.

We first synthesized the helicene ligand through a Sonogashira coupling reaction between 5-bromo-2,2'-bipyridine and 4,13-diethynyl[6]helicene in both racemic and enantiopure forms (Scheme S1 in Supporting information). The resulting ligand showed a bent, non-planar structure with two extended

* Corresponding authors.

E-mail addresses: shenchengshuo@sjtu.edu.cn (C. Shen), bshzhu@sjtu.edu.cn (B. Zhu), hbqiu@sjtu.edu.cn (H. Qiu)

bipyridine arms organized in a $\bar{\Delta}$ (for bis-bipyridine-terminated (*P*)-helicene, *i.e.*, (*P*)-ligand) or Δ (for bis-bipyridine-terminated (*M*)-helicene, *i.e.*, (*M*)-ligand) chiral fashion (the dihedral angle between two arms was *ca.* 165°, Figs. S4-1 and S4-15 in Supporting information) [31]. We then mixed the ligand with $\text{Fe}(\text{CF}_3\text{SO}_3)_2$ in a ratio of 3:2, and subsequently observed an immediate change of color from pale-yellow to deep-red, corresponding to a fast chelating process between the bipyridine groups and Fe(II) [32]. The collected product showed a clean and simple ^1H NMR pattern with the same number of proton environments as the ligand (Fig. 1a, Figs. S2-1, S3-1 and S3-2 in

Supporting information). Notably, protons *H*-22 and *H*-28 adjacent to the nitrogen atoms drastically shifted upfield from 8.98 and 8.73 ppm to 7.92 and 7.30 ppm, respectively, demonstrating the complexation between the bipyridine modules and Fe(II) [33]. Electrospray ionization high resolution mass spectrometry (ESI-HRMS) revealed three main peaks at $m/z = 541.3949$ ($z = 4$), 771.5036 ($z = 3$) and 1231.7511 ($z = 2$), which can be indexed as $[\text{Fe}_2\text{L}_3]^{4+}$, $[\text{Fe}_2\text{L}_3(\text{CF}_3\text{SO}_3)]^{3+}$ and $[\text{Fe}_2\text{L}_3(\text{CF}_3\text{SO}_3)_2]^{2+}$, respectively (L for the helicene ligand, Fig. 1b).

Fig. 1. Fe(II)-, Co(II)- and Ni(II)-based M_2L_3 cages. (a) ^1H NMR spectrum of (*rac*)- Fe_2L_3 cage (500 MHz, CD_3CN , 298 K) with comparison to (*rac*)-bis-bipyridine-terminated helicene ligand (L) (500 MHz, CDCl_3 , 298 K). (b) ESI-HRMS spectrum of (*rac*)- Fe_2L_3 cage. (c) Chemical structure of Fe_2L_3 cage. (d) Crystal structures of (*M*)- and (*P*)- Fe_2L_3 cages with the largest possible spheres inside the cavity. (e) UV-vis (bottom) and ECD (top) spectra of enantiopure helicene ligands (in CH_2Cl_2 , $c = 2 \times 10^{-5}$ mol/L) and Fe_2L_3 , Co_2L_3 and Ni_2L_3 cages (in CH_3CN , $c = 1 \times 10^{-5}$ mol/L). (f) Cyclic voltammograms of racemic Fe_2L_3 , Co_2L_3 and Ni_2L_3 cages (1×10^{-3} mol/L in CH_3CN , with 1×10^{-1} mol/L of *n*- Bu_4NPF_6 as the electrolyte, scan rate = 50 mV/s).

Single-crystal X-ray diffraction further confirmed the yield of Fe_2L_3 coordination cages (Fig. 1c), in which three helicene ligands were bridged by two Fe atoms (distance of *ca.* 18.3 Å) and formed a triple-stranded helical structure (Fig. 1d). The three surrounding helicene ligands created a sizeable internal cavity, which could accommodate a sphere with a diameter of *ca.* 9 Å (with its surface just attaching the van der Waals surface of the cage framework). For the cage derived from the (*P*)-ligand (*i.e.*, (*P*)- Fe_2L_3), both two Fe-bipyridine coordination sites were in Δ stereogenicity, and the whole cage framework adopted a *P* helical conformation [34]. Similarly, in the enantiomeric mirror series, the cage formed by the (*M*)-ligand (*i.e.*, (*M*)- Fe_2L_3) displayed an *M* helical shape with two Fe-bipyridine sites solely in Δ stereogenicity. It seemed that the helicene core substantially dominates the stereogenicity of the coordination domains as well as the helicity of the entire cage, which renders the cage

formation reaction highly diastereoselective and prevents the formation of other diastereomers. This was actually also demonstrated by the simple ^1H NMR spectrum, which was indicative of the absence of other isomers. Such efficient remote transfer of stereogenic information probably can be interpreted by the minimal-energy stacking of the relatively rigid, globally bent helicene ligands.

The Fe_2L_3 cage showed a group of intensive UV-vis absorption bands from 230 to 450 nm (Fig. 1e), roughly reprinting the nature of the helicene ligand. The ECD signals in the region of 360 ~ 450 nm were dramatically enhanced compared to that of the helicene ligands, which was probably due to the coordination to Fe(II). The Fe_2L_3 cage also revealed a distinctive absorption band with moderate intensity in the region of 450 ~ 650 nm, which could be attributed to the metal to ligand charge transfer (MLCT) from the Fe(II) centers to the bipyridine

modules [35,36]. The corresponding ECD curves were resolved in a bisignate fashion. The (*P*)-Fe₂L₃ cage displayed a positive Cotton effect at 570 nm and a negative one at 490 nm, which is in accordance with the *A* configuration of the Fe-bipyridine coordination sites [35]. Meanwhile, the (*M*)-Fe₂L₃ cage presented a mirror-imaged ECD curve, indicating a *A* configuration.

We further switched the transition metals to Co(II) and Ni(II) and obtained yellow-colored products. ESI-HRMS analysis (Figs. S3-4 and S3-5) and/or single-crystal X-ray diffraction (Fig. S4-4) confirmed the formation of Co₂L₃ and Ni₂L₃ cages in a triple-stranded helical shape with the selective formation of stereogenic coordination sites analogous to the Fe₂L₃ cage. The Co₂L₃ and Ni₂L₃ cages revealed similar UV-vis and ECD spectra to the Fe₂L₃ cage in the shorter wavelength region (< 450 nm), but in the longer wavelength region (> 450 nm) no obvious absorption band was detected (Fig. 1e), indicating a distinct metal to ligand electron transition mechanism [35]. Besides, due to the introduction of electrochemically active metal coordination sites [37,38], all these three helicene cages exhibited a single redox event at +0.68 (Fe₂L₃ cage), +0.78 (Co₂L₃ cage) and +0.69 V (Ni₂L₃ cage) vs. Fc⁺/Fc⁰, respectively, which appeared to be quasi-reversible (Fig. 1f).

Similarly, with the addition of Zn(II) salts into the solution of the bis-bipyridine-helicene ligand, we obtained a yellow-colored compound, which showed a simple ¹H NMR curve (Fig. S3-6) and well-resolved ESI-HRMS peaks at *m/z* = 546.1373 (*z* = 4, Fig. 2a). Single-crystal X-ray diffraction again confirmed the formation of Zn₂L₃ cages with homologous triple-stranded helical topology (Fig. 2b). ECD spectra further reflected the chiral fashion of the resulting Zn₂L₃ cage in solution (Fig. 2c), which was similar to the Co₂L₃ and Ni₂L₃ cage. Like other Zn-bipyridine coordination complexes [39], the Zn₂L₃ cages showed yellow luminescence ($\lambda_{\max}$ = 580 nm, Fig. 2d) under the irradiation of UV light, which was considerably bathochromically shifted compared with the blue luminescence of the helicene ligand ($\lambda_{\max}$ = 430 nm). As a result of the global transfer of stereogenic information throughout the whole cage, these Zn₂L₃ cages presented prominent circularly polarized luminescence (CPL) with dissymmetric factor g_{lum} of ± 0.002 at 580 nm (negative for (*P*)-Zn₂L₃ cage and positive for (*M*)-Zn₂L₃ cage, *i.e.*, with the signs in agreement with the lowest ECD-active band).

Fig. 2. Zn₂L₃ cages. (a) ESI-HRMS and calculated mass spectra of (*rac*)-Zn₂L₃ cages. (b) Crystal structures of (*M*)- and (*P*)-Zn₂L₃ cages with the largest possible spheres inside the cavity. (c) UV-vis (bottom) and ECD (top) spectra of enantiopure Zn₂L₃ cages (in CH₃OH, *c* = 4×10^{-5} mol/L, compared with the helicene ligands). (d) Fluorescence (bottom) and CPL (top) spectra of enantiopure helicene ligands and Zn₂L₃ cages. The inset photographs show their luminescence under irradiation at 365 nm.

Interestingly, when treating the Zn₂L₃ cage solution with Na₂SO₄ powder, we obtained a green-luminescent compound with a splitting of all proton environments into two sets in the ¹H NMR spectrum (Fig. 3b). For example, the signals of protons *H*-22 and *H*-28 at 8.56 and 8.01 ppm were split, with one set drastically shifted upfield to 7.52 and 7.03 ppm and the other set moved downfield to 9.98 and 9.39 ppm, respectively. ESI-HRMS showed that this new composite possesses a relatively large molecular weight (*m* = 4884.7608, Figs. S3-15 in Supporting information), which can be assigned as Zn₆L₆(SO₄)₄. Single-crystal X-ray diffraction of the racemic product further illustrated a six-stranded helical structure with the integration of three tilted homochiral Zn₂L₂ macrocyclic modules by two pairs of sulfates aligned on the vertical axis (Figs. 3a and c). The intricate trigonal-bipyramid-like Zn₃(SO₄)₂ cluster also revealed a helical conformation with the two SO₄ vertexes slightly twisted with each other. The neighboring helicene ligands belonging to two Zn₂L₂ modules were closely packed in a short distance of *ca.* 3.6 Å and revealed unsymmetrical offset π - π interactions (Figs. S4-16 in Supporting information), which rendered a more sophisticated environment and hence the split of the ¹H NMR signals. This giant cage showed an expanded cavity compared with the original Zn₂L₃ cage and could host a sphere with a diameter of *ca.* 10.0 Å.

By gradual addition of Na₂SO₄ into the Zn₂L₃ cage solution in a mixture of CD₃OD/CDCl₃, the ¹H NMR spectra revealed an apparent evolution with the rising of the Zn₆L₆(SO₄)₄ cage and the free ligand in a molar ratio of 1:3, and the simultaneous

consumption of the Zn_2L_3 cage precursor (Figs. S3-16 in Supporting information). This strongly indicated that each $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cage was assembled from three Zn_2L_3 cages *via* ligand exchange to sulfates, with the corresponding release of three free helicene ligands. Notably, no obvious peaks for other intermediates were observed, demonstrating that this assembly process might undergo in a cooperative pathway [40].

The $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cages showed similar absorption bands as the Zn_2L_3 cages, suggesting that the major electron transition modes were generally preserved in this new coordination complex. However, the $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cages presented intense ECD bands in the region of 370 ~ 450 nm with $\Delta\epsilon_{\text{max}} = \pm 1040 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$, which was *ca.* 1.6 folders of the Zn_2L_3 cages (with consideration of the number of helicene ligands, Fig. 3d). Such drastic enhancement might be resulted from the unique chiral packing of the helicene ligands as well as the twisted structure of the whole $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cage. On the other hand, the changes on the coordination sites apparently affected the luminescence

property and hence the maximum emission hypsochromically shifted to 520 nm. Notably, the $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cages revealed remarkable CPL with g_{lum} of ± 0.004 (doubled compare to the Zn_2L_3 cages, Fig. 3e vs. Fig. 2d).

In conclusion, we have prepared a series of chiral coordination cages *via* facile coordination of bis-dipyridine-terminated helicene ligands with a variety of transition metals. The flexible selection of metal centers substantially enriches the chemistry and functions (potentially for electrochemical or luminescence-sensitive recognition and separation of chiral guest molecules) of the resultant cages. Besides, the relatively dynamic association between the bipyridine modules and Zn(II) may also favor the fabrication of more complex chiral coordination cages *via* ligand exchanges using other suitable inorganic and organic species. We are currently exploring the construction of other distinctive chiral molecular cages utilizing the diverse helicene coordination chemistry [41], aiming at innovative structures and functions.

Fig. 3. Sextuple-stranded $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cages. (a) Schematic representation for the synthesis of $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cage. (b) ^1H NMR spectrum of (*rac*)- $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cage (500 MHz, CD_3OD , 298 K) with comparison to (*rac*)- Zn_2L_3 cage (500 MHz, CD_3OD , 298 K). (c) Crystal structure of (*rac*)- $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cage with the largest possible sphere inside the cavity, and partially enlarged views of the $\text{Zn}_3(\text{SO}_4)_2$ cluster. (d) UV-vis (bottom) and ECD (top) spectra of enantiopure $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ (in CH_3OH , $c = 1 \times 10^{-5} \text{ mol/L}$) and Zn_2L_3 cages (the intensity for Zn_2L_3 cages were doubled for clearer comparison). (e) Fluorescence (bottom) and CPL (top) spectra of enantiopure $\text{Zn}_6\text{L}_6(\text{SO}_4)_4$ cages. The inset photograph shows the luminescence under irradiation at 365 nm.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by the National Key R&D Program of China (No. 2020YFA0908100), the National Natural Science Foundation of China (Nos. 92056110 and 22075180), the Science and Technology Commission of Shanghai Municipality (Nos.

18JC1415500, 195271040, 20JC1415000). The authors thank Prof. Shunai Che's group in Tongji University for assistance with CPL measurement.

References

- [1] R. Chakrabarty, P.S. Mukherjee, P.J. Stang, Chem. Rev. 111 (2011) 6810–6918.
- [2] T. Hasell, A.I. Cooper, Nat. Rev. Mater. 1 (2016) 16053.
- [3] M.M.J. Smulders, I.A. Riddell, C. Browne, et al., Chem. Soc. Rev. 42 (2013) 1728–1754.

- [4] A.M. Castilla, W.J. Ramsay, J.R. Nitschke, *Acc. Chem. Res.* 47 (2014) 2063–2073.
- [5] R. Custelcean, *Chem. Soc. Rev.* 43 (2014) 1813–1824.
- [6] G. Zhang, M. Mastalerz, *Chem. Soc. Rev.* 43 (2014) 1934–1947.
- [7] W. Wang, Y.X. Wang, H.B. Yang, *Chem. Soc. Rev.* 45 (2016) 2656–2693.
- [8] G.H. Clever, P. Punt, *Acc. Chem. Res.* 50 (2017) 2233–2243.
- [9] T.R. Cook, P.J. Stang, *Chem. Rev.* 115 (2015) 7001–7045.
- [10] H. Amouri, C. Desmarets, J. Moussa, *Chem. Rev.* 112 (2012) 2015–2041.
- [11] S. Zarra, D.M. Wood, D.A. Roberts, et al., *Chem. Soc. Rev.* 44 (2015) 419–432.
- [12] D. Zhang, T.K. Ronson, J.R. Nitschke, *Acc. Chem. Res.* 51 (2018) 2423–2436.
- [13] M. Yoshizawa, J.K. Klosterman, M. Fujita, *Angew. Chem. Int. Ed.* 48 (2009) 3418–3438.
- [14] C.J. Brown, F.D. Toste, R.G. Bergman, et al., *Chem. Rev.* 115 (2015) 3012–3035.
- [15] S.H.A.M. Leenders, R. Gramage-Doria, B. de Bruin, et al., *Chem. Soc. Rev.* 44 (2015) 433–448.
- [16] T. Hasell, M.A. Little, S.Y. Chong, et al., *Nanoscale* 9 (2017) 6783–6790.
- [17] P. Howlader, E. Zangrando, P.S. Mukherjee, *J. Am. Chem. Soc.* 142 (2020) 9070–9078.
- [18] L.J. Chen, H.B. Yang, M. Shionoya, *Chem. Soc. Rev.* 46 (2017) 2555–2576.
- [19] M. Pan, K. Wu, J.H. Zhang, et al., *Coord. Chem. Rev.* 378 (2019) 333–349.
- [20] A. Lützen, M. Hapke, J. Griep-Raming, et al., *Angew. Chem. Int. Ed.* 41 (2002) 2086–2089.
- [21] J. Bunzen, T. Bruhn, G. Bringmann, et al., *J. Am. Chem. Soc.* 131 (2009) 3621–3630.
- [22] C. Gütz, R. Hovorka, N. Struch, et al., *J. Am. Chem. Soc.* 136 (2014) 11830–11838.
- [23] Y. Shen, C.F. Chen, *Chem. Rev.* 112 (2012) 1463–1535.
- [24] M. Gingras, *Chem. Soc. Rev.* 42 (2013) 968–1006.
- [25] K. Dhbaibi, L. Favereau, J. Crassous, *Chem. Rev.* 119 (2019) 8846–8953.
- [26] C. Shen, G. Loas, M. Srebro-Hooper, et al., *Angew. Chem. Int. Ed.* 55 (2016) 8062–8066.
- [27] K. Dhbaibi, L. Favereau, M. Srebro-Hooper, et al., *Chem. Sci.* 9 (2018) 735–742.
- [28] J. Barroso, J.L. Cabellos, S. Pan, et al., *Chem. Commun.* 54 (2018) 188–191.
- [29] A.U. Malik, F. Gan, C. Shen, et al., *J. Am. Chem. Soc.* 140 (2018) 2769–2772.
- [30] T.R. Schulte, J.J. Holstein, G.H. Clever, *Angew. Chem. Int. Ed.* 58 (2019) 5562–5566.
- [31] T. Damhus, C.E. Schäffer, *Inorg. Chem.* 22 (1983) 2406–2412.
- [32] E.M. Kober, T.J. Meyer, *Inorg. Chem.* 21 (1982) 3967–3977.
- [33] P. Bonakdarzadeh, F. Pan, E. Kalenius, et al., *Angew. Chem. Int. Ed.* 54 (2015) 14890–14893.
- [34] N. Struch, C. Frömbgen, G. Schnakenburg, et al., *Eur. J. Org. Chem.* (2017) 4984–4989.
- [35] M. Ziegler, A. von Zelewsky, *Coord. Chem. Rev.* 177 (1998) 257–300.
- [36] G. Auböck, M. Chergui, *Nat. Chem.* 7 (2015) 629–633.
- [37] Q.T. He, X.P. Li, Y. Liu, et al., *Angew. Chem. Int. Ed.* 48 (2009) 6156–6159.
- [38] S.H.R.W. Singh, H.L.C. Feltham, S. Brooker, *Dalt. Trans.* 48 (2019) 15435–15444.
- [39] K.D. Oyler, F.J. Coughlin, S. Bernhard, *J. Am. Chem. Soc.* 129 (2007) 210–217.
- [40] C.T. McTernan, T.K. Ronson, J.R. Nitschke, *J. Am. Chem. Soc.* 143 (2021) 664–670.
- [41] E.S. Gauthier, R. Rodríguez, J. Crassous, *Angew. Chem. Int. Ed.* 59 (2020) 22840–22856.

Graphical abstract

Coordination of bis-dipyridine-terminated helicene ligands with a variety of transition metals leads to the formation of various triple-stranded binuclear chiral cages and the addition of sulfate further promotes the transformation to sextuple-stranded hexanuclear chiral cages.