Role of Proteasome Inhibitors in Relapsed and/or Refractory Multiple Myeloma

Xavier Leleu, Thomas G. Martin, Hermann Einsele, Roger M. Lyons, Brian G.M. Durie, Karim S. Iskander, Sikander Ailawadhi

► To cite this version:

Xavier Leleu, Thomas G. Martin, Hermann Einsele, Roger M. Lyons, Brian G.M. Durie, et al.. Role of Proteasome Inhibitors in Relapsed and/or Refractory Multiple Myeloma. Clinical Lymphoma, Myeloma & Leukemia, 2019, 19, pp.9 - 22. 10.1016/j.clml.2018.08.016 . hal-03484591

HAL Id: hal-03484591

<https://hal.science/hal-03484591>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Role of Proteasome Inhibitors in Relapsed and/or Refractory Multiple Myeloma

Xavier Leleu¹, Thomas G Martin², Hermann Einsele³, Roger M Lyons⁴, Brian GM Durie⁵, Karim S Iskander⁶
and Sikander Ailawadhi⁷

¹CHU la Milettrie and Inserm CIC 1402, Poitiers, France, Xavier.LELEU@chu-poitiers.fr; ²University of California, San Francisco, CA, USA, Tom.Martin@ucsf.edu; ³University Hospital Würzburg, Würzburg, Germany, Einsele_H@ukw.de; ⁴Texas Oncology and US Oncology Research, San Antonio, TX, USA, Roger.Lyons@USONCOLOGY.COM; ⁵Cedars-Sinai Comprehensive Cancer Center, Los Angeles, CA, USA, bdurie@myeloma.org; ⁶Amgen Inc., Thousand Oaks, CA, USA, iskander@amgen.com; ⁷Mayo Clinic, Jacksonville, FL, USA, Ailawadhi.Sikander@mayo.edu.

Correspondence: Xavier Leleu, CHU la Milettrie and Inserm CIC 1402, Poitiers, France. E-mail: Xavier.LELEU@chu-poitiers.fr

Running title: Proteasome inhibitors in relapsed multiple myeloma

Competing Interests

XL reports consulting, honoraria, and membership on an entity's Board of Directors or advisory committees for Janssen, Novartis, BMS, Celgene, Merck, Amgen, Sanofi, Pierre Fabre, and Abbvie; and honoraria and membership and membership on an entity's Board of Directors or advisory committees for Takeda, Incyte, and Gilead. HE reports honoraria, research funding, and advisory participation for Amgen, Celgene, and Janssen; and honoraria and advisory role participation for BMS, Novartis, and Takeda. TGM reports consultancy for Sanofi and receipt of research funding through USCF from Amgen. RML reports no conflicts of interest to disclose. BGMD reports consulting fees from Takeda and Janssen. KSI reports employment and stock ownership with Amgen. SA reports research funding from Pharmacyclics; consultancy and honoraria from Takeda, Amgen, and Novartis; and consultancy for Celgene.

ABSTRACT

Proteasome inhibition is an established treatment strategy for patients with multiple myeloma as proteasome inhibitors selectively target and disrupt the protein metabolism of aberrant plasma cells. Since the introduction of the first-in-class proteasome inhibitor bortezomib, the therapeutic landscape for multiple myeloma has shifted with the development of next-generation proteasome inhibitors (carfilzomib and ixazomib) and new classes of agents. Treatment with modern combination therapies has been shown to result in deep responses and improved outcomes, and these potent regimens are increasingly used as frontline therapy. As patients continue to live longer with modern frontline therapy, there will be an increased need for effective regimens after initial treatment failure. Several recent studies have shown that treatment with combination therapy incorporating proteasome inhibitors induces deep and durable responses in patients with relapsed and/or refractory multiple myeloma. In this review, we review pivotal data and discuss the role of proteasome inhibitor-based doublet and triplet regimens for the management of relapsed/refractory multiple myeloma in the era of modern combination therapy.

Keywords: carfilzomib, bortezomib, ixazomib, efficacy, safety

INTRODUCTION

Multiple myeloma is a plasma cell dyscrasia characterized by uncontrolled proliferation and bone marrow infiltration of aberrant plasma cells and the presence of abnormal monoclonal protein secreted by these cells in the serum and/or urine.¹ It is the second most common hematologic malignancy in the United States (30 280 estimated new cases in 2017)^{1,2} and constitutes 0.8% of all new cancer cases worldwide.³

Multiple myeloma is an aggressive and incurable disease for most patients, characterized by periods of remission and relapse. Patients with relapsed multiple myeloma have been defined by Dimopoulos et al as those with active disease who have received at least one prior therapy and are not refractory to the last therapy; patients with relapsed/refractory multiple myeloma are defined as those with disease relapse who have previously achieved at least a minimal response or those with disease progression on salvage therapy or within 60 days of discontinuation of last therapy.⁴ With each relapse, patients face increasingly worse outcomes and each subsequent line of therapy tends to result in a shorter duration of response than previous lines. Each line of therapy is associated with an increased risk of treatment- and disease-related complications.⁵ The poor prognosis also reflects the genomic complexity of relapsed multiple myeloma, with tumors acquiring multiple genetic and epigenetic alterations that promote treatment resistance and refractory disease.^{1,4} The introduction of new classes of drugs, such as proteasome inhibitors, has improved survival outcomes in these difficult-to-treat populations.⁶⁻⁸

The ubiquitin-proteasome pathway regulates cellular homeostasis through protein catabolism (proteolysis).⁹⁻¹² Inhibition of proteasome function disrupts several important cellular pathways and affects malignant plasma cells to a greater extent than normal cells. Proteasome inhibition as a treatment strategy for multiple myeloma was clinically validated with the accelerated approval of the first-in-class proteasome inhibitor bortezomib in 2003 for relapsed/refractory multiple myeloma. Since

63 this approval, 2 next-generation proteasome inhibitors, carfilzomib and ixazomib, have also been
64 approved for patients with relapsed/refractory multiple myeloma.¹³

65 With the availability of several distinct proteasome inhibitors and other new drug classes (eg, histone
66 deacetylase inhibitors, monoclonal antibodies) with established efficacy, the potential options for
67 multidrug regimens based on proteasome inhibitors have increased. In addition, the treatment
68 paradigm for multiple myeloma has shifted to a more aggressive approach emphasizing the early use of
69 potent combination therapies to achieve deep responses and prolonged survival.¹⁴ More than half of
70 patients who are treated with modern multidrug regimens along with transplantation and maintenance
71 achieve a complete response,¹⁵ which has been associated with improved survival outcomes.¹⁶ Based on
72 these observations, frontline therapy with 3-drug regimens is currently considered a standard of care.¹⁷
73 Importantly, recent data from the SWOG S0777 study showed that treatment with the 3-drug
74 combination of bortezomib, lenalidomide, and dexamethasone significantly improved progression-free
75 survival (PFS) and overall survival (OS) compared with lenalidomide and dexamethasone alone in
76 patients with newly diagnosed multiple myeloma.¹⁸ Thus, as frontline triplet therapy becomes more
77 established and patients continue to live longer, there will be greater need for potent regimens after
78 initial treatment failure. In a study of Medicare patients with multiple myeloma diagnosed between
79 2008 and 2011, it was found that among patients who received frontline doublets, 58% continued
80 doublets as second-line therapy.¹⁹ Among patients receiving frontline triplets, 26% continued triplets,
81 47% switched to doublets, and 27% switched to monotherapy as second-line therapy.¹⁹ These treatment
82 patterns underscore the ongoing need for effective doublet and triplet therapy at first relapse. Modern
83 combination regimens, including those based on proteasome inhibitors, have been found to result in
84 deep responses, even in patients with relapsed/refractory disease.^{14,20}

85 Although not every modern combination regimen includes a proteasome inhibitor, proteasome
86 inhibition is an important treatment strategy for patients with multiple myeloma. Given the essential

role of proteasome inhibition in multiple myeloma and the availability of several potential treatment options for patients with relapsed/refractory disease, it is important to assess the role of proteasome inhibitors in this setting to aid in the selection of optimal treatment strategies. Here, we review pivotal data and discuss the role of proteasome inhibitor-based doublet and triplet regimens for the management of relapsed/refractory multiple myeloma in the era of modern combination therapy.

DISCUSSION

The Proteasome and Proteasome Inhibitors

The proteasome, first observed in 1968 and characterized in the late 1980's,⁹⁻¹² is a multicatalytic protease complex located in the nucleus and cytoplasm of eukaryotic cells. Each catalytic subunit ($\beta 1$, $\beta 2$, $\beta 5$) of the proteasome is named after an enzyme with similar proteolytic activity as the corresponding subunit: caspase-like, trypsin-like, and chymotrypsin-like, respectively.²¹ The proteasome is a key component of the ubiquitin-proteasome pathway, which is a central platform for recycling proteins involved in important cellular processes including cell cycle progression, DNA repair, and transcription.²² The scientists responsible for the discovery of ubiquitin-mediated protein degradation were awarded the Nobel Prize in Chemistry in 2004.²³ Timely proteolysis through the ubiquitin-proteasome pathway is an important regulator of homeostasis in normal cells.²² Malignant plasma cells almost always secrete a greater amount of immunoglobulins (ie, proteins) than normal plasma cells and are consequently more dependent on the ubiquitin-proteasome pathway for survival.²¹ Therefore, the proteasome is a rational target for antimyeloma therapies. Preclinical studies have demonstrated that proteasome inhibition results in the build-up of proteins within malignant plasma cells leading to activation of the unfolded protein stress response and decreased NF- κ B activity, both of which induce apoptosis.²¹

Three proteasome inhibitors have been approved for the treatment of patients with multiple myeloma: bortezomib, carfilzomib, and ixazomib. Structurally, bortezomib is a dipeptide boranate.²¹ Functionally, it is a slowly reversible inhibitor targeting primarily the $\beta 5$ subunit of the proteasome, with less potent inhibition effects on the $\beta 1$ and $\beta 2$ subunits.²¹ Since the introduction of bortezomib, 2 new proteasome inhibitors have entered the clinic for patients with relapsed/refractory multiple myeloma, carfilzomib (first approved in 2012) and ixazomib (first approved in 2015). Carfilzomib, bortezomib, and ixazomib have unique structural and mechanistic properties, resulting in differential proteasome inhibition profiles. Carfilzomib, a tetrapeptide epoxyketone, selectively targets and irreversibly inhibits the proteasome. Compared with bortezomib, carfilzomib has greater selectivity for the $\beta 5$ subunit²¹ of the proteasome and less off-target activity against nonproteasomal proteases.²⁴ In patients with relapsed or refractory multiple myeloma, carfilzomib has been shown to induce deeper and more sustained proteasome inhibition compared with bortezomib.²⁵ Ixazomib is structurally similar to bortezomib and is a citrate ester of boronic acid.¹³ After aqueous exposure, ixazomib hydrolyzes to the free boric acid MLN2238, which is the pharmacologically active metabolite.¹³ Ixazomib reversibly inhibits the proteasome with selectivity for the $\beta 5$ subunit. Ixazomib is orally bioavailable and has a shorter 20S proteasome disassociation half-life than bortezomib.²⁶ In preclinical studies, ixazomib demonstrated more favorable pharmacokinetic, pharmacodynamic, and antitumor activity profiles compared with bortezomib.²⁶

Activity of Approved Proteasome Inhibitor-Based Combination Therapies in Relapsed/Refractory Multiple Myeloma

To date, a total of 7 proteasome inhibitor-based combination therapies have been approved for the treatment of patients with relapsed/refractory multiple myeloma (doublet, $n=3$; triplet, $n=4$). The approved doublets are bortezomib plus dexamethasone, bortezomib plus pegylated doxorubicin, and carfilzomib plus dexamethasone. The approved triplets are bortezomib plus panobinostat and

dexamethasone, bortezomib plus daratumumab and dexamethasone, carfilzomib plus lenalidomide and dexamethasone, and ixazomib plus lenalidomide and dexamethasone. Pivotal data supporting the approval of these therapies are summarized in Tables 1 and 2. Results from these pivotal studies and studies supporting other combination therapies recommended by the National Comprehensive Care Network are described below. The current National Comprehensive Care Network Guidelines evaluate treatments with categories of evidence and consensus (eg, category 1: based upon high-level evidence and there is uniform consensus the intervention is appropriate; category 2A: based upon lower-level evidence and there is uniform consensus the intervention is appropriate) and categories of preference (eg, preferred intervention: based on superior evidence; other recommended intervention: interventions may be somewhat less effective, more toxic, or based on less mature data).¹⁷ These categories of evidence and consensus from the National Comprehensive Care Network guidelines are included in the descriptions of the proteasome inhibitor-based therapies in this review. All proteasome inhibitor-based doublets recommended by the National Comprehensive Care Network guidelines were included in the review. Because there are a large number of recommended proteasome inhibitor-based triplets in these guidelines, only triplets based on a proteasome inhibitor plus an immunomodulatory agent or monoclonal antibody were included, as these were deemed to be the most clinically relevant based on the authors' judgment.

Approved Bortezomib-Based Regimens

Doublets

Bortezomib plus dexamethasone has been a standard of care in relapsed/refractory multiple myeloma and was approved in the United States in 2014 when used as re-treatment (see Re-treatment with Proteasome Inhibitors) and in the European Union in 2014 for patients with progressive multiple myeloma who have received at least one prior therapy. The broader European Union approval was

based on a retrospective matched-pairs analysis of the MMY-2045, APEX, and DOXIL-MMY-3001 clinical trials. The analysis demonstrated improved overall response rate (75% vs. 41%; odds ratio, 3.467; $P<0.001$), time to progression (median, 13.6 vs. 7.0 months; hazard ratio [HR], 0.394; $P=0.003$), and PFS (median, 11.9 vs. 6.4 months, HR, 0.595; $P=0.051$) after second-line treatment with bortezomib and dexamethasone versus bortezomib alone.³⁹ The National Comprehensive Care Network considers bortezomib plus dexamethasone a treatment option under “other recommended regimens” (category 1; not preferred) for patients with relapsed/refractory multiple myeloma.¹⁷

Bortezomib plus liposomal doxorubicin was approved in 2007 for the treatment of patients with relapsed or refractory multiple myeloma who have not previously received bortezomib and have received at least one prior therapy. This approval was based on the DOXIL-MMY-3001 study, which was a randomized, phase 3 trial comparing bortezomib plus liposomal doxorubicin with bortezomib monotherapy in patients with relapsed or refractory multiple myeloma. At the interim analysis, time to progression was longer for combination treatment compared with monotherapy (median, 9.3 vs. 6.5 months; HR, 1.82 [monotherapy vs. combination]; $P<0.0001$).²⁷ The results from the final OS analysis of this study showed that OS was similar between the treatment arms (median, 33.0 vs. 30.8 months; HR, 1.05; $P=0.61$).²⁸ The National Comprehensive Care Network considers bortezomib plus doxorubicin (with dexamethasone) a category 1 option (not preferred) for patients with relapsed/refractory multiple myeloma.¹⁷

Triplets

Bortezomib has been evaluated as part of triplets combining a proteasome inhibitor with new classes of drugs, as demonstrated in the CASTOR and PANORAMA1 studies. The randomized, phase 3 CASTOR study showed that treatment with daratumumab in combination with bortezomib and dexamethasone resulted in significantly longer PFS compared with bortezomib and dexamethasone alone (median, 16.7

vs. 7.1 months; HR, 0.31; $P<0.0001$).³⁸ Based on results from the CASTOR trial, this combination was approved in 2016 for the treatment of patients with multiple myeloma who have received at least one prior therapy. This combination is also considered a preferred regimen (category 1) for the treatment of patients with relapsed/refractory multiple myeloma.¹⁷

The panobinostat plus bortezomib and dexamethasone regimen was approved in 2015 for patients with multiple myeloma who have received at least 2 prior regimens, including bortezomib and an immunomodulatory drug. The pivotal trial supporting this approval was PANORAMA-1, which evaluated panobinostat or placebo plus bortezomib and dexamethasone in patients with relapsed or relapsed/refractory multiple myeloma. The primary analysis of this trial demonstrated that treatment with panobinostat plus bortezomib and dexamethasone resulted in significantly longer PFS compared with placebo plus bortezomib and dexamethasone (median, 11.99 vs. 8.08 months; HR, 0.63; $P<0.0001$).³⁰ The final OS analysis for this trial showed a trend toward OS benefit for panobinostat over placebo, but the difference was not found to be statistically significant (median, 40.3 vs. 35.8 months; HR, 0.94; $P=0.54$).³¹ The National Comprehensive Care Network has included this combination as a category 1 option for patients with at least 2 prior therapies, including an immunomodulatory drug and bortezomib.¹⁷

Approved Carfilzomib-Based Regimens

Doublets

In a head-to-head study of 2 proteasome inhibitors (carfilzomib vs. bortezomib), the randomized, phase 3 ENDEAVOR trial compared carfilzomib (56 mg/m² infused over 30 minutes) and dexamethasone with bortezomib and dexamethasone in patients with relapsed or refractory multiple myeloma (1 to 3 prior lines of therapy). This carfilzomib regimen was based on results from a phase 1 study, which established the maximum tolerated dose of carfilzomib to be 56 mg/m² when administered as a 30-minute

infusion.⁴⁰ The first interim analysis of ENDEAVOR showed that there was a statistically significant increase in PFS for the carfilzomib group compared with the bortezomib group (median, 18.7 vs. 9.4 months; HR, 0.53, $P<0.0001$).³² The second interim analysis of ENDEAVOR showed that carfilzomib also provided a significant and clinically meaningful OS improvement compared with bortezomib, prolonging median OS by 7.6 months (47.6 vs. 40.0 months; HR, 0.79; 1-sided $P=0.010$).³⁶ Use of bisphosphonates was balanced between treatment arms and did not confound the OS analysis.⁴¹ Notably, ENDEAVOR is the first and currently the only phase 3 study to demonstrate a significant OS advantage in the relapsed or refractory multiple myeloma setting in a head-to-head comparison of 2 different proteasome inhibitors.³⁶

The results of the ENDEAVOR study established the combination of carfilzomib with dexamethasone as a new treatment option for patients with relapsed or refractory multiple myeloma who are being considered for doublet therapy.³⁶ The carfilzomib (56 mg/m²) and dexamethasone combination was approved in 2016 for the treatment of patients with relapsed or refractory multiple myeloma (with 1 to 3 prior lines of therapy), and this combination is the only doublet that is recognized as a preferred treatment option (category 1) for previously treated multiple myeloma by the National Comprehensive Care Network.¹⁷

Triplets

Results from the interim analysis of the randomized, phase 3 ASPIRE study showed that treatment with carfilzomib (27 mg/m² infused over 10 minutes), lenalidomide, and dexamethasone significantly improved PFS compared with lenalidomide and dexamethasone alone (median, 26.3 vs. 17.6 months; HR, 0.69; $P=0.0001$) in patients with relapsed or refractory multiple myeloma (1 to 3 prior lines of therapy). The median PFS duration of 26 months in the carfilzomib arm was noted to be the longest observed to date in the relapsed setting.²⁰ In the ASPIRE regimen, carfilzomib was given for 18 cycles

and then patients continued on lenalidomide and dexamethasone alone, similar to the control arm. The current standard of care is to use antimyeloma therapies until disease progression, which is the case for the active arms of other phase 3 trials in relapsed/refractory multiple myeloma reviewed here. A post-hoc analysis of ASPIRE showed that the HR for PFS at 18 months was 0.58.⁴² Recently, results from a prespecified final analysis of OS in ASPIRE showed that carfilzomib plus lenalidomide and dexamethasone reduced the risk of death compared with lenalidomide and dexamethasone, prolonging median OS by 7.9 months (48.3 vs. 40.4 months; HR, 0.79; 95% CI, 0.67–0.95).³⁷ Notably, this is the first and currently the only study in patients with relapsed or refractory multiple myeloma to demonstrate an OS advantage for a triplet regimen against lenalidomide and dexamethasone.

Based on results from the ASPIRE study, the combination of carfilzomib with lenalidomide and dexamethasone was approved in 2015 for the treatment of patients with relapsed or refractory multiple myeloma (1 to 3 prior lines of therapy in the United States and 1 or more prior line in the European Union). This regimen is also considered a category 1 preferred treatment option for patients with previously treated multiple myeloma by the National Comprehensive Care Network.¹⁷

Approved Ixazomib-Based Regimens

Doublets

Ixazomib is currently not approved by the US Food and Drug Administration to be used as part of a doublet for the treatment of relapsed and/or refractory multiple myeloma.

Triplets

Interim results from the randomized, phase 3 TOURMALINE-MM1 study demonstrated that the addition of ixazomib (versus placebo) to lenalidomide and dexamethasone significantly prolonged PFS in patients with relapsed and/or refractory multiple myeloma (median 20.6 vs. 14.7 months; HR 0.74; $P=0.01$).³⁵ These data supported the approval (2015) of ixazomib plus lenalidomide and dexamethasone for the

treatment of patients with multiple myeloma who have received at least one prior therapy. In addition, this regimen is a category 1 preferred treatment option for patients with previously treated multiple myeloma.¹⁷ Notably, this regimen is the only approved all-oral triplet based on a proteasome inhibitor. Longer follow-up results from TOURMALINE-MM1 demonstrated a smaller improvement in PFS compared with the interim results (median, 20.0 vs. 15.9 months; $P=0.05$).⁴³

Activity of Other Proteasome Inhibitor-Based Combination Therapies in Relapsed/Refractory Multiple Myeloma

Several other proteasome-inhibitor based combination regimens are recommended by clinical guidelines, but have not yet been approved by regulatory agencies. This section summarizes the data supporting treatment recommendations from the National Comprehensive Care Network, with a focus on doublets and triplets containing a proteasome inhibitor and dexamethasone with either an immunomodulatory drug or a monoclonal antibody.

Doublets

The combination of once-weekly carfilzomib plus dexamethasone for patients with relapsed or refractory multiple myeloma was evaluated in the phase 1/2 CHAMPION-1 study. The phase 1 portion of this trial established the maximum tolerated dose of carfilzomib administered once weekly (30-minute infusion) as 70 mg/m².⁴⁴ Among patients treated at this dose in the phase 1 and 2 portions of the study ($n=104$), the overall response rate was 77% and median PFS was 12.6 months, demonstrating the activity of this regimen.⁴⁴ Once-weekly administration of carfilzomib may be more convenient for patients compared with twice-weekly administration. The carfilzomib regimen used in CHAMPION-1 is included in the National Comprehensive Care Network guidelines as a recommended regimen (category 2A).¹⁷ This regimen is being investigated in the phase 3 ARROW study (NCT02412878), which compares once-weekly carfilzomib (70 mg/m²) versus twice-weekly carfilzomib (27 mg/m²). At the interim analysis,

271 this trial met its primary endpoint. Patients who received the once-weekly carfilzomib had significantly
272 longer PFS compared with those who received the twice-weekly carfilzomib regimen (median, 11.2 vs.
273 7.6 months; HR, 0.69; 95% CI, 0.54–0.88).⁴⁵

274 Ixazomib is currently not approved in any doublet regimen, but the combination of ixazomib and
275 dexamethasone has shown promising activity in 2 phase 2 studies. In 1 study (*N*=33), patients with
276 relapsed multiple myeloma with limited prior exposure to bortezomib received weekly ixazomib with
277 dexamethasone added for suboptimal responses (lack of a minor response by cycle 2, partial response
278 by cycle 4, or disease progression at any time). Dexamethasone was added for 22 (67%) patients, 4 of
279 whom achieved a partial response or better after suboptimal response. With or without the addition of
280 dexamethasone, the overall response rate was 34%, and the median event-free survival was 11.5
281 months.⁴⁶ In the other study (*N*=70), patients with relapsed multiple myeloma (who were proteasome
282 inhibitor-naïve or received fewer than 6 cycles of therapy with bortezomib and had a partial response or
283 better with no progression at the time of discontinuation) were randomized to receive ixazomib 4 or 5.5
284 mg in combination with dexamethasone. Overall response rates were 31% in the 4 mg arm and 51% in
285 the 5.5 mg arm.⁴⁷ Median event-free survival was similar for both arms (8.4 and 8.2 months,
286 respectively). Patients receiving the 5.5-mg dose had higher rates of grade ≥3 adverse events considered
287 possibly related to treatment (54% vs. 21%) and adverse events requiring dose reductions (43% vs. 17%)
288 compared with patients receiving the 4-mg dose. Based on these data, the ixazomib plus
289 dexamethasone regimen is recognized as a category 2A treatment option for patients with
290 relapsed/refractory multiple myeloma (at least one prior therapy) by the National Comprehensive Care
291 Network.¹⁷

292 In addition to proteasome inhibitor-dexamethasone doublet regimens, the pairing of a proteasome
293 inhibitor with a histone deacetylase inhibitor has also been evaluated.⁴⁸ In a phase 1/2 study of
294 carfilzomib plus panobinostat (*N*=42), the overall response rate was 67% and median PFS was 7.7

months in patients with relapsed/refractory multiple myeloma.⁴⁹ Based on this promising phase 1/2 data, carfilzomib plus panobinostat is a recommended treatment option (category 2A) for patients with previously treated multiple myeloma.¹⁷

Triplets Containing a Proteasome Inhibitor and an Immunomodulatory Drug

Although not yet approved by regulatory agencies, the combination of bortezomib, lenalidomide, and dexamethasone is considered a preferred regimen (category 1) for the treatment of patients with relapsed/refractory multiple myeloma by the National Comprehensive Care Network.¹⁷ This recommendation is based on results from a phase 2 study of 64 patients with relapsed or relapsed/refractory multiple myeloma. In this study, the overall response rate was 64%, median PFS was 9.5 months, and median OS was 30 months.⁵⁰ Notably, the SWOG S0777 study showed that treatment with this triplet improved PFS and OS compared with lenalidomide and dexamethasone in patients with newly diagnosed multiple myeloma.¹⁸

Combination regimens consisting of pomalidomide and dexamethasone with bortezomib, carfilzomib, or ixazomib are considered recommended therapeutic options (category 2A) by the National Comprehensive Care Network for patients who have received at least 2 prior therapies, including an immunomodulatory drug and a proteasome inhibitor (specifically bortezomib for the bortezomib- and carfilzomib-based triplets), and have demonstrated disease progression on or within 60 days of completion of the last therapy.¹⁷ Phase 1 studies have demonstrated promising activity for bortezomib with pomalidomide and dexamethasone;¹⁷ the randomized phase 3 OPTIMISMM trial (NCT01734928) recently showed that bortezomib with pomalidomide and dexamethasone improved PFS compared with bortezomib and dexamethasone alone in patients with relapsed or refractory multiple myeloma.⁵¹ In phase 1 and 2 studies, treatment with carfilzomib plus pomalidomide and dexamethasone was found to be highly active in both heavily pretreated patients with relapsed or refractory multiple myeloma

(median of 6 prior regimens; $N=72$), with an overall response rate of 64%,⁵² and less pretreated, proteasome inhibitor naïve/sensitive patients ($n=64$), with an overall response rate of 84%.⁵³ In phase 1/2 studies, ixazomib plus pomalidomide and dexamethasone has shown early promising activity in patients with double-refractory disease (partial response or better in 8 of 13 patients)⁵⁴ and in patients with relapsed/refractory multiple myeloma who received at least one prior therapy ($n=25$, overall response rate of 48%).⁵⁵

Triplets Containing a Proteasome Inhibitor and a Monoclonal Antibody

A randomized, phase 2 study compared elotuzumab plus bortezomib and dexamethasone versus bortezomib and dexamethasone in patients with relapsed/refractory multiple myeloma ($N=152$). At the time of the primary analysis, the addition of elotuzumab to bortezomib and dexamethasone resulted in statistically significantly longer PFS compared with bortezomib and dexamethasone alone (median, 9.7 vs. 6.9 months; HR, 0.72; $P=0.09$ [statistical significance threshold of 2-sided $P\leq 0.05$ was used]).⁵⁶ In an updated analysis, the 2-year PFS rates were 18% for bortezomib plus elotuzumab and dexamethasone and 11% for bortezomib and dexamethasone.⁵⁶ This triplet combination is considered a recommended regimen (category 2A; not preferred) for patients with relapsed/refractory multiple myeloma who have received at least one therapy by the National Comprehensive Care Network.¹⁷

Activity of Proteasome Inhibitor-Based Regimens at First Relapse

The first relapse is a critical juncture on the myeloma disease continuum. Each subsequent relapse is associated with increased genomic complexity and drug resistance, resulting in increasingly shorter response durations.^{57,58} Furthermore, data suggest that achievement of deep responses at first relapse versus subsequent relapses is more strongly associated with improved survival outcomes.⁵⁹ Thus, early treatment with highly effective therapy at first relapse, when the disease is still at a sensitive stage, is critical for maximizing OS for patients with relapsed/refractory multiple myeloma.^{57,60} Consistent data

from pivotal trials show that proteasome inhibitor-based combination therapies are highly effective treatment options for first relapse (Table 3).

Doublets

The matched-pairs analysis, described previously, compared bortezomib and dexamethasone with single-agent bortezomib. This analysis showed that bortezomib and dexamethasone resulted in greater response and delayed disease progression compared with bortezomib monotherapy when used as second-line treatment.³⁹

A secondary analysis of the ENDEAVOR study established the benefit of carfilzomib and dexamethasone when used at first relapse, and demonstrated that it is more effective than bortezomib and dexamethasone in this setting. Results from this analysis showed that in the subgroup of patients who had received 1 prior line of therapy, there was a clinically meaningful improvement in PFS for patients treated with carfilzomib plus dexamethasone versus bortezomib and dexamethasone (median, 22.2 vs. 10.1 months; HR, 0.447; $P<0.0001$).⁶¹ Additionally, the final OS analysis of ENDEAVOR showed that patients at first relapse had a 17% lower risk of death if treated with carfilzomib plus dexamethasone versus bortezomib plus dexamethasone (median OS was not estimable in either arm).⁶²

Triplets

The ASPIRE study established the benefit of adding carfilzomib to the immunomodulatory backbone of lenalidomide and dexamethasone for the treatment of patients at first relapse. A secondary analysis of the ASPIRE study demonstrated that the addition of carfilzomib to lenalidomide and dexamethasone led to a 12-month improvement in median PFS compared with lenalidomide and dexamethasone alone (29.6 vs. 17.6 months; HR, 0.71; $P=0.0118$) in patients with 1 prior line of therapy.⁶⁴ Importantly, the final analysis of OS in ASPIRE showed that the OS benefit for the carfilzomib group compared with the

control group was most pronounced at first relapse, with an 11-month prolongation of median OS (47.3 vs. 35.9 months; HR, 0.81).³⁷

A subgroup analysis of the TOURMALINE-MM1 study demonstrated a trend for improvement in PFS with ixazomib, lenalidomide, and dexamethasone compared with lenalidomide and dexamethasone alone for patients at first relapse (median, 20.6 vs. 16.6 months; HR, 0.88; 95% CI, 0.65–1.20).⁶⁵ However, the magnitude of benefit (HR, 0.88) appeared to be lower than that in the overall population (HR, 0.74). It was noted that patients with 1 prior therapy without transplant had greater benefit with ixazomib, lenalidomide, and dexamethasone than patients with 1 prior therapy and transplant (HR 1.232 vs. 0.604) and further investigation into this effect is ongoing.⁶⁵

Another pivotal study demonstrated the benefit of adding a novel agent to the proteasome inhibitor backbone of bortezomib and dexamethasone for patients at first relapse. In a subgroup analysis of CASTOR, it was found that the addition of daratumumab to bortezomib and dexamethasone led to significantly improved PFS for patients with 1 prior line of therapy (median, not reached vs. 7.9 months; HR, 0.19; $P < 0.0001$).³⁸

Re-treatment with Proteasome Inhibitors in Proteasome Inhibitor-Sensitive Patients

The choice of salvage therapy at relapse is an important decision that depends on several factors, including the type of previous treatment and duration of response to previous treatment. As bortezomib-based regimens are among the most commonly used frontline therapies,⁶⁶ it is important that treatments used at relapse retain efficacy and tolerability in patients with previous bortezomib exposure. Several proteasome inhibitor-based doublets and triplets have demonstrated activity in patients with prior bortezomib exposure.

Doublets

Guidelines from the National Comprehensive Care Network state that if relapse occurs more than 6 months after treatment completion, re-treatment with the same therapy may be considered.¹⁷ In concordance with these guidelines, the phase 2 RETRIEVE trial ($n=126$) showed that re-treatment with bortezomib (with or without dexamethasone) was effective (overall response rate of 40%) in patients with multiple myeloma who had achieved a partial response or better upon completion of prior bortezomib therapy and relapsed more than 6 months after prior bortezomib therapy.²⁹ These data supported approval of bortezomib re-treatment (as a single agent or with dexamethasone) for patients with relapsed multiple myeloma.

The ENDEAVOR study included a large group of patients ($n=502$) who were previously exposed to bortezomib but were not bortezomib refractory (similar to the RETRIEVE population), which allowed for a robust comparison of carfilzomib and dexamethasone versus bortezomib and dexamethasone among patients with prior bortezomib exposure (Table 4). Median PFS was 7.5 months longer in the carfilzomib group compared with the bortezomib group among patients with previous bortezomib exposure (median, 15.6 vs. 8.1 months; HR, 0.56; $P<0.0001$).^{32,61} Treatment with carfilzomib also extended OS by 14.8 months versus re-treatment with bortezomib in this subgroup (median, 47.6 vs. 32.8 months; HR, 0.84; 95% CI, 0.65–1.08).⁶² Although bortezomib re-treatment has been a common therapeutic strategy,⁶⁶ these head-to-head results suggest that treatment with carfilzomib at relapse is more effective (complete response rates, PFS, and OS) than re-treatment with bortezomib for patients with prior bortezomib exposure.

Triplets

Triplet regimens that combine proteasome inhibitors with a lenalidomide-dexamethasone backbone have also improved outcomes compared with lenalidomide-dexamethasone alone in patients with previous bortezomib exposure. In the ASPIRE study, the addition of carfilzomib to lenalidomide and

dexamethasone resulted in an approximate 8-month improvement in median PFS in bortezomib-exposed patients (median, 24.4 vs. 16.6 months; HR, 0.699) and for patients treated at first relapse after bortezomib-containing therapy, carfilzomib plus lenalidomide and dexamethasone resulted in 13.4 months of improvement (median, 29.3 vs. 15.9 months; HR, 0.577).^{64,67} Further, the addition of carfilzomib to lenalidomide and dexamethasone prolonged median OS by 12 months (median, 45.9 vs. 33.9 months; HR, 0.82; 95% CI, 0.56–1.19) among patients treated at first relapse after prior bortezomib.³⁷

A subgroup analysis of the TOURMALINE-MM1 study showed a consistent PFS benefit for ixazomib, lenalidomide, and dexamethasone versus lenalidomide and dexamethasone, regardless of prior bortezomib exposure (bortezomib-exposed: median, 18.5 vs. 13.6 months; HR, 0.746; bortezomib-naïve: median, not estimable vs. 15.9 months; HR, 0.747).⁶⁵

Results from subgroup analyses of the CASTOR and PANORAMA-1 trials suggest that treatment with potent triplet combinations based on the bortezomib-dexamethasone backbone may be more effective than treatment with this backbone alone in patients with prior bortezomib exposure. In the CASTOR study, PFS was improved with daratumumab plus bortezomib and dexamethasone compared with bortezomib and dexamethasone alone, regardless of prior bortezomib exposure (bortezomib-exposed: median, 12.3 vs. 6.7 months; HR, 0.46).³³ Among prior treatment subgroups examined in the PANORAMA-1 study (group 1: prior immunomodulatory drug; group 2: prior bortezomib and immunomodulatory drug; group 3: 2 or more prior regimens including bortezomib and an immunomodulatory drug), the PFS benefit for panobinostat plus bortezomib and dexamethasone versus bortezomib and dexamethasone alone was most pronounced in patients who had received at least 2 prior therapies, including bortezomib and an immunomodulatory drug, with a 7.8-month prolongation in median PFS (median, 12.5 vs. 4.7 months; HR, 0.47).³⁴

Re-treatment with Proteasome Inhibitors in Proteasome Inhibitor-Refractory Patients

As bortezomib is a commonly used frontline antineoplastic agent, there is a need for therapies that can recapture responses in patients who become bortezomib-refractory. Proteasome inhibitor-based combinations have demonstrated activity in this setting (Table 5), and several general approaches have been used. The first is the addition of a novel agent with a unique mechanism of action to a bortezomib-based backbone. This was tested in the phase 2 PANORAMA trial where the histone deacetylase inhibitor panobinostat was added to bortezomib and dexamethasone. Treatment with this combination resulted in an overall response rate of 35% in bortezomib-refractory patients (n=55).⁵⁰ More recent studies have evaluated the addition of venetoclax (a BCL-2 inhibitor) or selinexor (a selective exportin 1 inhibitor) to bortezomib and dexamethasone with promising activity; however, these combinations are not yet endorsed by regulatory agencies or the National Comprehensive Care Network. Another approach for proteasome inhibitor-based treatment of bortezomib-refractory patients is the use of a different proteasome inhibitor, ie carfilzomib or ixazomib. Preclinically, carfilzomib and ixazomib are able to overcome bortezomib resistance in multiple myeloma cell lines and patient samples.^{71,72} Building on this preclinical rationale, several carfilzomib or ixazomib-based combinations have demonstrated activity in bortezomib-refractory patients (Table 5). Notably, ASPIRE is the only pivotal phase 3 trial of proteasome inhibitor-based combination therapy to have reported efficacy data for bortezomib-refractory patients. Among these patients in ASPIRE, median PFS was extended 3 months with the addition of carfilzomib to lenalidomide and dexamethasone (22.3 vs. 19.4 months; HR, 0.80), and the HR for OS (0.73) also favored the carfilzomib group.^{37,64}

The addition of novel agents (eg, panobinostat, venetoclax, or selinexor) to a carfilzomib-based regimen has also been investigated recently, and has generally shown promising activity in patients refractory to a proteasome inhibitor. With the exception of carfilzomib plus panobinostat, regimens based on this approach have not yet been endorsed by the National Comprehensive Care Network.

Safety of Proteasome Inhibitor-Based Combination Therapies in Relapsed/Refractory Multiple

Myeloma

Bortezomib, carfilzomib, and ixazomib each have their own unique safety profiles, which reflects their distinct structural and mechanistic properties. In the ENDEAVOR study, rates of grade ≥ 3 adverse events and serious adverse events were slightly more common with carfilzomib versus bortezomib. Grade ≥ 3 adverse events that were more common ($\geq 2\%$) with carfilzomib were anemia, hypertension, dyspnea, decreased lymphocyte count, pyrexia, and cardiac failure whereas those that were more common in the bortezomib group were diarrhea and peripheral neuropathy.³⁶

The safety profiles of bortezomib and carfilzomib are affected by the route of administration and/or dosing schedule used. Subcutaneous administration has been shown to result in a lower rate of peripheral neuropathy compared with intravenous bortezomib administration.⁷³ For carfilzomib, a longer infusion time (ie, 30 vs. 10 minutes) permits safe administration of higher doses of carfilzomib compared with the originally approved dose of 27 mg/m².^{32,44} The CHAMPION-1 study showed that once-weekly administration of carfilzomib (70 mg/m²; 30-minute infusion) in combination with dexamethasone was generally well tolerated in patients with relapsed or refractory multiple myeloma, with only 12% of patients discontinuing treatment due to adverse events.⁴⁴

In randomized studies comparing a triplet versus a doublet, adverse event rates tended to be slightly higher in the triplet arm than in the doublet arm, as expected (Table 6). In the PANORAMA-1 study, the addition of panobinostat to bortezomib and dexamethasone was associated with an increased incidence of grade 3 or 4 thrombocytopenia, lymphopenia, neutropenia, diarrhea, and asthenia or fatigue.³⁰ In the ASPIRE study, the addition of carfilzomib to lenalidomide and dexamethasone was associated with an increased incidence of diarrhea, cough, fever, and hypertension.²⁰ In the TOURMALINE-MM1 study, the addition of ixazomib to lenalidomide and dexamethasone was associated with an increased incidence of

thrombocytopenia, rash, gastrointestinal events, and peripheral neuropathy.³⁵ In the CASTOR study, the addition of daratumumab to bortezomib and dexamethasone led to an increased incidence of thrombocytopenia and infusion-related reactions.³³

Factors to Consider When Selecting Proteasome Inhibitor Therapy

Several disease-, treatment-, and patient-related factors should be considered when selecting therapy. These factors have been reviewed in detail elsewhere.⁷⁴ We briefly highlight some of these factors and how they apply to selection of proteasome inhibitor-based therapy.

The presence of high-risk cytogenetic features is associated with a poor prognosis and there is a need for therapies that can overcome (or partly overcome) this adverse prognosis. Subgroup analyses of the ASPIRE, ENDEAVOR, TOURMALINE-MM1, and CASTOR studies have shown that modern proteasome inhibitor-based combination therapy improves outcomes regardless of cytogenetic risk status, including in patients with high-risk cytogenetics.^{75–78} Safety and tolerability should also be considered when selecting therapy. As peripheral neuropathy is associated with bortezomib, regimens with other proteasome inhibitors (ie, carfilzomib or ixazomib) may be preferred in patients for whom this toxicity is a concern. Similarly, the potential for cardiac toxicity should be considered when evaluating carfilzomib-based therapy.⁷⁹ Safety and efficacy should be considered together and the overall benefit-risk profile assessed. Another factor to consider is convenience. For patients who cannot easily travel to the clinic, an oral proteasome inhibitor such as ixazomib may be preferred. Once-weekly administrations of carfilzomib and bortezomib have been evaluated,^{44,80} and these may be more convenient than the currently approved twice-weekly administrations.

As survival times have improved for patients with multiple myeloma, quality of life considerations and patient-reported outcomes have gained in importance when selecting therapy. However, until recently, there were limited data regarding effect of proteasome inhibitor-based therapies on quality of life in

patients with relapsed/refractory multiple myeloma.⁸¹ A prespecified exploratory analysis of the phase 3 APEX trial showed that patients with relapsed multiple myeloma who received bortezomib had better health-related quality of life (HRQoL) than those who received dexamethasone.⁸² Recent data from the ENDEAVOR, ASPIRE, and TOURMALINE-MM1 studies have provided important additional information regarding the effects of next-generation proteasome inhibitors on quality of life. A prespecified analysis of the ENDEAVOR study showed that treatment with carfilzomib resulted in statistically significant improvements in patient-reported health-related quality of life, fatigue, pain, side effects of treatment, and neurotoxicity compared with bortezomib treatment.⁸³ In ASPIRE, the addition of carfilzomib to lenalidomide resulted in a statistically significant and clinically meaningful improvement in health-related quality life with no detrimental effect (compared with the control group) on other patient-reported outcomes.⁸⁴ The addition of ixazomib to lenalidomide and dexamethasone in the TOURMALINE-MM1 study did not have a statistically significant impact on health-related quality of life.⁸⁵ Overall, these studies show that combination regimens based on next-generation proteasome inhibitors are able to improve (carfilzomib) or maintain (ixazomib) health-related quality of life compared with recent standards of care in relapsed/refractory multiple myeloma.

Conclusion

Proteasome inhibitor-based regimens are essential components of modern therapy for patients with relapsed/refractory myeloma, with demonstrated activity including in key subgroups such as patients in first relapse and patients with prior proteasome inhibitor exposure. Consistent data from randomized studies show that modern combination regimens incorporating proteasome inhibitors improve clinical outcomes (including OS in some cases) compared with standard of care in patients with relapsed/refractory multiple myeloma. Both doublet and triplet regimens are approved in this disease setting and both are demonstrated to be effective. The only head-to-head study directly comparing 2 proteasome inhibitors in relapsed/refractory multiple myeloma, ENDEAVOR, demonstrated

525 improvements in complete responses, PFS, and OS for the carfilzomib-treated patients compared with
526 the bortezomib-treated patients. ASPIRE is the only phase 3 study in relapsed/refractory multiple
527 myeloma to date to demonstrate an improvement in OS with the addition of a proteasome inhibitor.
528 The benefit was most prominent when the carfilzomib, lenalidomide, and dexamethasone combination
529 was used at first relapse, especially in proteasome inhibitor-sensitive patients (after bortezomib-based
530 frontline therapy). Ixazomib, lenalidomide, and dexamethasone is the only approved, all-oral
531 proteasome inhibitor-based regimen and it offers a convenient choice when infusion therapy is not an
532 option. Bortezomib re-treatment can be an option in certain markets with limited access to modern
533 therapies.

534 There are currently 3 approved proteasome inhibitors, each with their own unique efficacy and safety
535 profiles. Bortezomib is rapidly active and has a defined efficacy and safety profile that is well-understood
536 by clinicians. It will remain a standard of care option in the frontline setting for the foreseeable future.
537 The duration of bortezomib treatment is limited by peripheral neuropathy and development of
538 relapsed/refractory disease. Like bortezomib, carfilzomib is also rapidly active. In our experience,
539 carfilzomib can be administered for an extended period (2 to 3 years), especially when given once
540 weekly. It is the proteasome inhibitor of choice at relapse because of its PFS and OS advantage
541 compared with other therapies (including bortezomib retreatment) and its ability to be combined with
542 many other agents. Several different dose densities (once- vs. twice- weekly) and dose intensities (twice-
543 weekly carfilzomib with lenalidomide and dexamethasone: 27 mg/m², twice-weekly carfilzomib with
544 dexamethasone: 56 mg/m², once-weekly carfilzomib with dexamethasone: 70 mg/m²) of carfilzomib
545 have been evaluated in randomized phase 3 trials, and the various doses and schedules have the
546 potential to offer greater flexibility in the administration and utility of carfilzomib. In contrast to
547 bortezomib and carfilzomib, ixazomib is an oral drug and thus a convenient option for patients who can
548 come to treatment centers.

The role of proteasome inhibitors as a class and the 3 currently approved agents will continue to evolve as bortezomib becomes generic worldwide, the next-generation proteasome inhibitors move to the frontline, and additional classes of drugs are incorporated in the therapeutic landscape. The cost-effectiveness of generic bortezomib will need to be evaluated and it is likely that generic bortezomib will have a prominent role in markets with limited financial resources, although a recent analysis of ENDEAVOR predicted that twice-weekly carfilzomib 56 mg/m² with dexamethasone would be cost-effective versus bortezomib and dexamethasone, even at an 85% price reduction for bortezomib and dexamethasone.⁸⁶ As an increasing number of studies evaluate frontline carfilzomib- and ixazomib-based regimens, the comparative effectiveness of the various proteasome inhibitor-based, upfront combinations will need to be addressed, as well as the benefit-risk profile in specific patient populations. Real-world evidence from claims data and patient registries will be important in this regard. As novel agents with unique mechanisms of action (eg, venetoclax, selinexor, monoclonal antibodies) are introduced and become established, the scope and potential of proteasome inhibitor-based combinations will continue to expand, allowing effective therapy for heavily pretreated and refractory patients as well as tailored combination therapies for distinct patient subgroups.

There are several key knowledge gaps that remain regarding the role of proteasome inhibitors. Compared with the relapsed/refractory setting, phase 3 data in the upfront setting for carfilzomib and ixazomib are limited. In addition, the evaluation of the best partners for proteasome inhibitors is still an active area of study, and in the context of immunomodulation, there remains a question of whether proteasome inhibitors should be combined with immunomodulatory drugs, monoclonal antibodies, or both. To address an aspect of this question, a randomized phase 2 study (NCT02874742) is evaluating the addition of daratumumab to the triplet of bortezomib, lenalidomide, and dexamethasone in patients with newly diagnosed multiple myeloma. Several single-arm trials are also investigating carfilzomib and dexamethasone in combination with daratumumab or lenalidomide plus daratumumab and are already

demonstrating positive results.⁸⁷ Beyond optimal combinations, there is no uniform standard of care or data from randomized trials regarding how proteasome-based combination therapies should be sequenced. This will become increasingly important as carfilzomib and ixazomib move to the frontline and questions arise as to which proteasome inhibitor to use upfront, which to reserve for later lines, and the efficacy of bortezomib salvage therapy after frontline carfilzomib or ixazomib. Overall, several studies have found that switching to a different proteasome inhibitor or retreatment with the same proteasome inhibitor can be effective strategies for patients progressing after prior proteasome inhibitor exposure.^{29,32,64,65,88,89} Regarding specific sequences, a single-center, retrospective study found that treatment with frontline carfilzomib-based therapy followed by bortezomib-based therapy (n=15) resulted in higher response rates and deeper responses than the reverse sequence of bortezomib-based therapy followed by carfilzomib-based therapy (n=39); however, larger prospective studies are warranted.⁹⁰ Data from ENDEAVOR suggest it is more effective to switch to carfilzomib rather than retreat with bortezomib. Regarding retreatment with carfilzomib, 1 phase 2 study (n=13) found that retreatment with carfilzomib 56 mg/m² recaptured responses in patients who were refractory to carfilzomib at a lower dose (27 mg/m²).⁹¹ Other knowledge gaps include the value of minimal residual disease testing to assess response to proteasome inhibitor-based therapy, development of alternative oral proteasome inhibitors, and effects of proteasome inhibitors on immune fitness and immunomodulation.

The current treatment paradigm in multiple myeloma emphasizes use of potent combination therapy capable of producing deep and sustained responses. Modern proteasome inhibitor-based regimens are highly effective and are an important option for treating patients with relapsed/refractory disease.

Acknowledgements

595 Medical writing assistance was provided by BlueMomentum, an Ashfield company, part of UDG
596 Healthcare plc, and funded by Amgen Inc.

597

598 **Funding sources**

599 Funding: This work was supported by **Amgen Inc.**

600

601

References

1. Kumar SK, Rajkumar V, Kyle RA, et al. Multiple myeloma. *Nat Rev Dis Primers*. 2017;3:17046.
2. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin*. 2017;67:7–30.
3. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136:E359–E386.
4. Dimopoulos MA, Richardson PG, Moreau P, Anderson KC. Current treatment landscape for relapsed and/or refractory multiple myeloma. *Nat Rev Clin Oncol*. 2015;12:42–54.
5. Song X, Cong Z, Wilson K. Real-world treatment patterns, comorbidities, and disease-related complications in patients with multiple myeloma in the United States. *Curr Med Res Opin*. 2016;32:95–103.
6. Kumar SK, Dispenzieri A, Lacy MQ, et al. Continued improvement in survival in multiple myeloma: changes in early mortality and outcomes in older patients. *Leukemia*. 2014;28:1122–1128.
7. Kumar SK, Rajkumar SV, Dispenzieri A, et al. Improved survival in multiple myeloma and the impact of novel therapies. *Blood*. 2008;111:2516–2520.
8. Costa LJ, Brill IK, Omel J, Godby K, Kumar SK, Brown EE. Recent trends in multiple myeloma incidence and survival by age, race, and ethnicity in the United States. *Blood Advances*. 2017;1:282–287.
9. Arrigo AP, Tanaka K, Goldberg AL, Welch WJ. Identity of the 19S 'prosome' particle with the large multifunctional protease complex of mammalian cells (the proteasome). *Nature*. 1988;331:192–194.
10. Falkenburg PE, Haass C, Kloetzel PM, et al. Drosophila small cytoplasmic 19S ribonucleoprotein is homologous to the rat multicatalytic proteinase. *Nature*. 1988;331:190–192.
11. Harris JR. Release of a macromolecular protein component from human erythrocyte ghosts. *Biochim Biophys Acta*. 1968;150:534–537.

- 626 12. Voges D, Zwickl P, Baumeister W. The 26S proteasome: a molecular machine designed for
627 controlled proteolysis. *Annu Rev Biochem.* 1999;68:1015–1168.
- 628 13. Muz B, Ghazarian RN, Ou M, Luderer MJ, Kusdono HD, Azab AK. Spotlight on ixazomib: potential in
629 the treatment of multiple myeloma. *Drug Des Devel Ther.* 2016;10:217–226.
- 630 14. Landgren O, Iskander K. Modern multiple myeloma therapy: deep, sustained treatment response
631 and good clinical outcomes. *J Intern Med.* 2017;281:365–382.
- 632 15. Kumar S, Paiva B, Anderson KC, et al. International Myeloma Working Group consensus criteria for
633 response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol.*
634 2016;17:e328–e346.
- 635 16. van de Velde H, Londhe A, Ataman O, et al. Association between complete response and
636 outcomes in transplant-eligible myeloma patients in the era of novel agents. *Eur J Haematol.*
637 2017;98:269–279.
- 638 17. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology.
639 Multiple Myeloma Version 3.2018.
640 https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf. Accessed 4 June 2018.
- 641 18. Durie BG, Hoering A, Abidi MH, et al. Bortezomib with lenalidomide and dexamethasone versus
642 lenalidomide and dexamethasone alone in patients with newly diagnosed myeloma without
643 intent for immediate autologous stem-cell transplant (SWOG S0777): a randomised, open-label,
644 phase 3 trial. *Lancet.* 2017;389:519–527.
- 645 19. Li S, Natwick T, Yusuf A, Vidito IS, Mezzi K, Werther W. Treatment sequencing patterns in
646 Medicare-enrolled patients with multiple myeloma. *Blood.* 2016;128:2120.
- 647 20. Stewart AK, Rajkumar SV, Dimopoulos MA, et al. Carfilzomib, lenalidomide, and dexamethasone
648 for relapsed multiple myeloma. *N Engl J Med.* 2015;372:142–152.

- 649 21. Manasanch EE, Korde N, Zingone A, et al. The proteasome: mechanisms of biology and markers of
650 activity and response to treatment in multiple myeloma. *Leuk Lymphoma*. 2014;55:1707–1714.
- 651 22. Cavo M. Proteasome inhibitor bortezomib for the treatment of multiple myeloma. *Leukemia*.
652 2006;20:1341–1352.
- 653 23. Neefjes J, Groothuis TA, Dantuma NP. [The 2004 Nobel Prize in Chemistry for the discovery of
654 ubiquitin-mediated protein degradation]. *Ned Tijdschr Geneesk*. 2004;148:2579–2582.
- 655 24. Arastu-Kapur S, Anderl JL, Kraus M, et al. Nonproteasomal targets of the proteasome inhibitors
656 bortezomib and carfilzomib: a link to clinical adverse events. *Clin Cancer Res*. 2011;17:2734–2743.
- 657 25. Ludwig H, Spencer A, Kovacsics T, et al. Comparison of proteasome inhibition activity between
658 carfilzomib and bortezomib in the phase 3 Endeavor study. *Blood*. 2017;130:3125.
- 659 26. Kupperman E, Lee EC, Cao Y, et al. Evaluation of the proteasome inhibitor MLN9708 in preclinical
660 models of human cancer. *Cancer Res*. 2010;70:1970–1980.
- 661 27. Orlowski RZ, Nagler A, Sonneveld P, et al. Randomized phase III study of pegylated liposomal
662 doxorubicin plus bortezomib compared with bortezomib alone in relapsed or refractory multiple
663 myeloma: combination therapy improves time to progression. *J Clin Oncol*. 2007;25:3892–3901.
- 664 28. Orlowski RZ, Nagler A, Sonneveld P, et al. Final overall survival results of a randomized trial
665 comparing bortezomib plus pegylated liposomal doxorubicin with bortezomib alone in patients
666 with relapsed or refractory multiple myeloma. *Cancer*. 2016;122:2050–2056.
- 667 29. Petrucci MT, Giraldo P, Corradini P, et al. A prospective, international phase 2 study of bortezomib
668 retreatment in patients with relapsed multiple myeloma. *Br J Haematol*. 2013;160:649–659.
- 669 30. San-Miguel JF, Hungria VT, Yoon SS, et al. Panobinostat plus bortezomib and dexamethasone
670 versus placebo plus bortezomib and dexamethasone in patients with relapsed or relapsed and
671 refractory multiple myeloma: a multicentre, randomised, double-blind phase 3 trial. *Lancet Oncol*.
672 2014;15:1195–1206.

- 673 31. San-Miguel JF, Hungria VTM, Yoon S-S, et al. Overall survival of patients with relapsed multiple
674 myeloma treated with panobinostat or placebo plus bortezomib and dexamethasone (the
675 PANORAMA 1 trial): a randomised, placebo-controlled, phase 3 trial. *Lancet Haem.* 2016;3:e506–
676 e515.
- 677 32. Dimopoulos MA, Moreau P, Palumbo A, et al. Carfilzomib and dexamethasone versus bortezomib
678 and dexamethasone for patients with relapsed or refractory multiple myeloma (ENDEAVOR): a
679 randomised, phase 3, open-label, multicentre study. *Lancet Oncol.* 2016;17:27–38.
- 680 33. Palumbo A, Chanan-Khan A, Weisel K, et al. Daratumumab, bortezomib, and dexamethasone for
681 multiple myeloma. *N Engl J Med.* 2016;375:754–766.
- 682 34. Richardson PG, Hungria VT, Yoon SS, et al. Panobinostat plus bortezomib and dexamethasone in
683 previously treated multiple myeloma: outcomes by prior treatment. *Blood.* 2016;127:713–721.
- 684 35. Moreau P, Masszi T, Grzasko N, et al. Oral ixazomib, lenalidomide, and dexamethasone for
685 multiple myeloma. *N Engl J Med.* 2016;374:1621–1634.
- 686 36. Dimopoulos MA, Goldschmidt H, Niesvizky R, et al. Carfilzomib or bortezomib in relapsed or
687 refractory multiple myeloma (ENDEAVOR): an interim overall survival analysis of an open-label,
688 randomised, phase 3 trial. *Lancet Oncol.* 2017;18:1327–1337.
- 689 37. Siegel DS, Dimopoulos MA, Ludwig H, et al. Improvement in overall survival with carfilzomib,
690 lenalidomide, and dexamethasone in patients with relapsed or refractory multiple myeloma. *J Clin*
691 *Oncol.* 2018; e-pub ahead of print 17 January 2018; doi: 10.1200/JCO.2017.76.5032.
- 692 38. Spencer A, Hungria VTM, Mateos M-V, et al. Daratumumab, bortezomib, and dexamethasone
693 (DVd) versus bortezomib and dexamethasone (Vd) in relapsed or refractory multiple myeloma
694 (RRMM): updated efficacy and safety analysis of Castor. *Blood.* 2017;130:3145.

39. Dimopoulos MA, Orlowski RZ, Facon T, et al. Retrospective matched-pairs analysis of bortezomib plus dexamethasone versus bortezomib monotherapy in relapsed multiple myeloma. *Haematologica*. 2015;100:100–106.
40. Papadopoulos KP, Siegel DS, Vesole DH, et al. Phase I study of 30-minute infusion of carfilzomib as single agent or in combination with low-dose dexamethasone in patients with relapsed and/or refractory multiple myeloma. *J Clin Oncol*. 2015;33:732–739.
41. Dimopoulos MA, Kimball AS. Carfilzomib for relapsed or refractory multiple myeloma - Authors' reply. *Lancet Oncol*. 2018;19:e2.
42. Dimopoulos MA, Stewart AK, Wang M, et al. Carfilzomib, lenalidomide, and dexamethasone vs lenalidomide and dexamethasone in patients with relapsed multiple myeloma: analysis of response and progression-free survival hazard ratio over time. *Haematologica*. 2016;101:Abstract P275.
43. Center for Drug Evaluation and Research. Ninlaro (ixazomib) medical review. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2015/208462Orig1s000MedR.pdf. Accessed 6 March 2018.
44. Berenson JR, Cartmell A, Bessudo A, et al. CHAMPION-1: a phase 1/2 study of once-weekly carfilzomib and dexamethasone for relapsed or refractory multiple myeloma. *Blood*. 2016;127:3360–3368.
45. Moreau P, Mateos MV, Berenson JR, et al. Once weekly versus twice weekly carfilzomib dosing in patients with relapsed and refractory multiple myeloma (A.R.R.O.W.): interim analysis results of a randomised, phase 3 study. *Lancet Oncol*. 2018; e-pub ahead of print 31 May 2018; doi: 10.1016/S1470-2045(18)30354-1
46. Kumar SK, LaPlant B, Roy V, et al. Phase 2 trial of ixazomib in patients with relapsed multiple myeloma not refractory to bortezomib. *Blood Cancer J*. 2015;5:e338.

- 719 47. Kumar SK, Laplant BR, Reeder CB, et al. Randomized phase 2 trial of two different doses of
720 ixazomib in patients with relapsed multiple myeloma not refractory to bortezomib. *Blood*.
721 2015;126:3050.
- 722 48. Hideshima T, Richardson PG, and Anderson KC. Mechanism of action of proteasome inhibitors and
723 deacetylase inhibitors and the biological basis of synergy in multiple myeloma. *Mol Cancer Ther*.
724 2011;10:2034–2042.
- 725 49. Berdeja JG, Hart LL, Mace JR, et al. Phase I/II study of the combination of panobinostat and
726 carfilzomib in patients with relapsed/refractory multiple myeloma. *Haematologica*.
727 2015;100:670–676.
- 728 50. Richardson PG, Schlossman RL, Alsina M, et al. PANORAMA 2: panobinostat in combination with
729 bortezomib and dexamethasone in patients with relapsed and bortezomib-refractory myeloma.
730 *Blood*. 2013;122(14):2331–2337.
- 731 51. Richardson PG, Rocafiguera AO, Beksac M, et al. Pomalidomide (POM), bortezomib, and low-dose
732 dexamethasone (PVd) vs bortezomib and low-dose dexamethasone (Vd) in lenalidomide (LEN)-
733 exposed patients (pts) with relapsed or refractory multiple myeloma (RRMM): Phase 3
734 OPTIMISMM trial. *J Clin Oncol*. 2018;36(15_suppl):8001.
- 735 52. Stadtmauer EA, Abonour R, Cohen AD, et al. Phase I/II dose expansion of a multi-center trial of
736 Carfilzomib and pomalidomide with dexamethasone (Car-Pom-d) in patients with
737 relapsed/refractory multiple myeloma. *Blood*. 2013;122:690.
- 738 53. Jakubowiak AJ, Rosenbaum CA, Stephens LAL, et al. Final results of phase (Ph) 1/2 study of
739 carfilzomib, pomalidomide, and dexamethasone (KPd) in patients (Pts) with relapsed/refractory
740 multiple myeloma (RRMM): A Multi-center MMRC study. *Haematologica*. 2017;102:P680.
- 741 54. Voorhees PM, Mulkey F, Hassoun H, et al. Alliance A061202. A phase I/II study of pomalidomide,
742 dexamethasone and ixazomib versus pomalidomide and dexamethasone for patients with

multiple myeloma refractory to lenalidomide and proteasome inhibitor based therapy: phase I results. *Blood*. 2015;126:375.

55. Krishnan A, Kapoor P, Palmer JM, et al. Phase I/II trial of the oral regimen ixazomib, pomalidomide, and dexamethasone in relapsed/refractory multiple myeloma. *Leukemia*. 2017.

56. Jakubowiak A, Offidani M, Pegourie B, et al. Randomized phase 2 study: elotuzumab plus bortezomib/dexamethasone vs bortezomib/dexamethasone for relapsed/refractory MM. *Blood*. 2016;127:2833–2840.

57. Harousseau JL, Attal M. How I treat first relapse of myeloma. *Blood*. 2017;130:963–973.

58. Kumar SK, Therneau TM, Gertz MA, et al. Clinical course of patients with relapsed multiple myeloma. *Mayo Clin. Proc* 2004;79:867–874.

59. Sonneveld P, Broijl A. Treatment of relapsed and refractory multiple myeloma. *Haematologica*. 2016;101:396–406.

60. Dingli D, Ailawadhi S, Bergsagel PL, et al. Therapy for relapsed multiple myeloma: guidelines from the Mayo stratification for myeloma and risk-adapted therapy. *Mayo Clin Proc*. 2017;92:578–598.

61. Moreau P, Joshua D, Chng WJ, et al. Impact of prior treatment on patients with relapsed multiple myeloma treated with carfilzomib and dexamethasone vs bortezomib and dexamethasone in the phase 3 ENDEAVOR study. *Leukemia*. 2017;31:115–122.

62. Weisel KC, Siegel D, San Miguel JF, et al. Overall survival of patients with relapsed multiple myeloma treated with carfilzomib and dexamethasone versus bortezomib and dexamethasone according to prior line of therapy and previous exposure to bortezomib: secondary analysis of the phase 3 Endeavor study. *Blood*. 2017;130:1850.

63. Mateos M-V, Estell J, Barreto W, et al. Efficacy of daratumumab, bortezomib, and dexamethasone versus bortezomib and dexamethasone in relapsed or refractory myeloma based on prior lines of therapy: updated analysis of Castor. *Blood*. 2016;128:1150.

- 767 64. Dimopoulos MA, Stewart AK, Masszi T, et al. Carfilzomib-lenalidomide-dexamethasone vs
768 lenalidomide-dexamethasone in relapsed multiple myeloma by previous treatment. *Blood Cancer*
769 *J.* 2017;7:e554.
- 770 65. Mateos MV, Masszi T, Grzasko N, et al. Impact of prior therapy on the efficacy and safety of oral
771 ixazomib-lenalidomide-dexamethasone vs placebo-lenalidomide-dexamethasone in patients with
772 relapsed/refractory multiple myeloma in TOURMALINE-MM1. *Haematologica.* 2017;102:1767–
773 1775.
- 774 66. Jagannath S, Roy A, Kish J, et al. Real-world treatment patterns and associated progression-free
775 survival in relapsed/refractory multiple myeloma among US community oncology practices. *Expert*
776 *Rev Hematol.* 2016;9:707–717.
- 777 67. Mateos M-V, Goldschmidt H, San Miguel JF, et al. Relapsed or refractory multiple myeloma
778 patients treated with second-line carfilzomib and categorized by prior exposure to bortezomib: a
779 subgroup analysis of the randomized phase 3 Aspire and Endeavor trials. *Blood.* 2017;130:1840.
- 780 68. Siegel DS, Martin T, Wang M, et al. A phase 2 study of single-agent carfilzomib (PX-171-003-A1) in
781 patients with relapsed and refractory multiple myeloma. *Blood.* 2012;120(14):2817–2825.
- 782 69. Kaufman JL, Zimmerman T, Rosenbaum CA, et al. Phase I study of the combination of carfilzomib
783 and panobinostat for patients with relapsed and refractory myeloma: a Multiple Myeloma
784 Research Consortium (MMRC) clinical trial. *Blood.* 2014;124(21):32.
- 785 70. Shah JJ, Stadtmauer EA, Abonour R, et al. Carfilzomib, pomalidomide, and dexamethasone for
786 relapsed or refractory myeloma. *Blood.* 2015;126(20):2284–2290.
- 787 71. Kuhn DJ, Chen Q, Voorhees PM, et al. Potent activity of carfilzomib, a novel, irreversible inhibitor
788 of the ubiquitin-proteasome pathway, against preclinical models of multiple myeloma. *Blood.*
789 2007;110(9):3281–3290.

790 72. Chauhan D, Tian Z, Zhou B, et al. In vitro and in vivo selective antitumor activity of a novel orally
791 bioavailable proteasome inhibitor MLN9708 against multiple myeloma cells. *Clin Cancer Res.*
792 2011;17(16):5311–5321.

793 73. Moreau P, Pylypenko H, Grosicki S, et al. Subcutaneous versus intravenous administration of
794 bortezomib in patients with relapsed multiple myeloma: a randomised, phase 3, non-inferiority
795 study. *Lancet Oncol.* 2011;12:431–440.

796 74. Nooka AK, Kastiris E, Dimopoulos MA, Lonial S. Treatment options for relapsed and refractory
797 multiple myeloma. *Blood.* 2015;125:3085–3099.

798 75. Avet-Loiseau H, Bahlis NJ, Chng WJ, et al. Ixazomib significantly prolongs progression-free survival
799 in high-risk relapsed/refractory myeloma patients. *Blood.* 2017;130:2610–2618.

800 76. Avet-Loiseau H, Fonseca R, Siegel D, et al. Carfilzomib significantly improves the progression-free
801 survival of high-risk patients in multiple myeloma. *Blood.* 2016;128:1174–1180.

802 77. Chng WJ, Goldschmidt H, Dimopoulos MA, et al. Carfilzomib-dexamethasone vs bortezomib-
803 dexamethasone in relapsed or refractory multiple myeloma by cytogenetic risk in the phase 3
804 study ENDEAVOR. *Leukemia.* 2017;31:1368–1374.

805 78. Weisel KC, Miguel JS, Cook G, et al. Efficacy of daratumumab in combination with lenalidomide
806 plus dexamethasone (DRd) or bortezomib plus dexamethasone (DVd) in relapsed or refractory
807 multiple myeloma (RRMM) based on cytogenetic risk status. *J Clin Oncol.* 2017;35:8006.

808 79. Waxman AJ, Clasen S, Hwang WT, et al. Carfilzomib-associated cardiovascular adverse events: a
809 systematic review and meta-analysis. *JAMA Oncol.* 2017:e174519.

810 80. Bringhen S, Larocca A, Rossi D, et al. Efficacy and safety of once-weekly bortezomib in multiple
811 myeloma patients. *Blood.* 2010;116:4745–4753.

81. Sparano F, Cavo M, Niscola P, Caravita T, Efficace F. Patient-reported outcomes in relapsed/refractory multiple myeloma: a systematic review. *Support Care Cancer*. 2018;26(7):2075–2090.
82. Lee SJ, Richardson PG, Sonneveld P, et al. Bortezomib is associated with better health-related quality of life than high-dose dexamethasone in patients with relapsed multiple myeloma: results from the APEX study. *Br J Haematol*. 2008;143:511–519.
83. Ludwig H, Moreau P, Dimopoulos MA, et al. Health related quality of life results from the open-label, randomized, phase III Endeavor trial evaluating carfilzomib and dexamethasone versus bortezomib and dexamethasone in patients with relapsed or refractory multiple myeloma. *Blood*. 2016;128:3309.
84. Stewart AK, Dimopoulos MA, Masszi T, et al. Health-related quality-of-life results from the open-label, randomized, phase III ASPIRE trial evaluating carfilzomib, lenalidomide, and dexamethasone versus lenalidomide and dexamethasone in patients with relapsed multiple myeloma. *J Clin Oncol*. 2016;34:3921–3930.
85. Leleu X, Masszi T, Bahlis NJ, et al. Patient-reported health-related quality of life from the phase III TOURMALINE-MM1 study of ixazomib-lenalidomide-dexamethasone versus placebo-lenalidomide-dexamethasone in relapsed/refractory multiple myeloma. *Am J Hematol*. 2018; E-pub ahead of print 4 May 2018; doi: 10.1002/ajh.25134
86. Jakubowiak AJ, Housse I, Májer I, et al. Cost-effectiveness of carfilzomib plus dexamethasone compared with bortezomib plus dexamethasone for patients with relapsed or refractory multiple myeloma in the United States. *Expert Rev Hematol*. 2017;10(12):1107–1119.
87. Chari A, Martinez-Lopez J, Mateos MV, et al. Daratumumab (DARA) in combination with carfilzomib and dexamethasone (D-Kd) in lenalidomide (Len)-refractory patients (Pts) with

835 relapsed multiple myeloma (MM): Subgroup analysis of MMY1001. *J Clin Oncol*.
836 2018;36(suppl):8002.

837 88. Berenson JR, Hilger JD, Yellin O, et al. Replacement of bortezomib with carfilzomib for multiple
838 myeloma patients progressing from bortezomib combination therapy. *Leukemia*.
839 2014;28(7):1529–1536.

840 89. Berenson A, Vardanyan S, David M, et al. Outcomes of multiple myeloma patients receiving
841 bortezomib, lenalidomide, and carfilzomib. *Ann Hematol*. 2017;96(3):449–459.

842 90. Diamond B, Rossi AC, Pearse RN, et al. Sequencing of proteasome inhibitors in patients with
843 multiple myeloma. *Blood*. 2016;128(22):4522.

844 91. Chari A, Cho HJ, Parekh S, et al. Recapturing disease response: a phase II study of high dose
845 carfilzomib in patients with relapsed or refractory multiple myeloma who have progressed on
846 standard dose carfilzomib. *Blood*. 2015;126(23):3051.
847

Table 1. Baseline Characteristics From Pivotal Trials in RRMM

Study Name (reference)	Patient Population	Treatment Group (n)	Median (range) Age, years	ECOG PS, %	Cytogenetic Risk Status, ^a %	Median (range) Number of Prior Regimens, n
2-drug regimens						
DOXIL-MMY- 3001 ^{27,28}	RRMM, V-naive	V+PLD (324)	61 (28–85)	0: 43; 1: 57	Had cytogenetic abnormality: 43; Did not have cytogenetic abnormality: 57	NR (34% and 66% had 1 or ≥2 prior therapies, respectively)
RETRIEVE ²⁹	Patients with MM who had relapsed after achieving ≥PR with initial V-based therapy	V±DEX re- treatment (130)	67 (38–86)	NR	NR	2 (1–7)
PANORAMA1 ^{30,31}	Relapsed or relapsed and refractory MM, 1– 3 previous treatments	Vd (381)	63 (IQR, 56–68)	0: 43; 1: 49; 2: 8	NR	NR (52%, 28%, and 20% had 1, 2, or 3 prior lines, respectively)
ENDEAVOR ³²	RRMM, 1–3 previous treatments	Kd56 (464) Treatment	65 (35–89)	0: 48; 1: 45;	High: 25; Standard: 75	2 (IQR, 1–2)

		until PD		2: 7		
	RRMM, 1–3 previous treatments	Vd (465) Treatment until PD	65 (30–88)	0: 50; 1: 44; 2: 6	High: 28; Standard: 72	2 (IQR, 1–2)
CASTOR ³³	Relapsed or relapsed and refractory MM, ≥1 previous treatment	Vd (247) Up to 8 cycles of treatment [6 months]	64 (33–85)	NR	High: 21; Standard: 79	2 (1–10)
3-drug regimens						
PANORAMA1 ^{30,31,34}	Relapsed or relapsed and refractory MM, 1–3 previous treatments	PanVd (387)	63 (IQR, 56–69)	0: 45; 1: 49; 2: 5	NR	NR (51%, 32%, and 17% had 1, 2, or 3 prior lines, respectively)
ASPIRE ²⁰	RRMM, 1–3 previous treatments	KRd (396)	64 (38–87)	0 or 1: 90; 2: 10	High: 25; Standard: 75	2 (1–3)
TOURMALINE-MM1 ³⁵	Relapsed, refractory, or relapsed and refractory MM, 1–3 previous treatments	IRd (360)	66 (38–91)	0: 51; 1: 44; 2: 5	High: 27; Standard: 73	NR (62%, 27%, and 11% had 1, 2, or 3 prior therapies, respectively)
CASTOR ³³	Relapsed or relapsed and refractory MM, ≥1	DVd (251)	64 (30–88)	NR	High: 23; Standard: 77	2 (1–9)

previous treatment

^aPercentages are based on the number of patients with known cytogenetics information.

Abbreviations: DEX=dexamethasone; DVd=daratumumab, bortezomib, and dexamethasone; ECOG PS=Eastern Cooperative Oncology Group performance status; IQR=interquartile range; IRd=ixazomib, lenalidomide, and dexamethasone; Kd56=carfilzomib (56 mg/m²) and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; MM=multiple myeloma; NR=not reported; PanVd=panobinostat, bortezomib and dexamethasone; PD=progressive disease; PI=proteasome inhibitor; PLD=pegylated liposomal doxorubicin; PR=partial response; Rd=lenalidomide and dexamethasone; RRMM=relapsed or refractory multiple myeloma; V=bortezomib; Vd=bortezomib and dexamethasone.

849

Table 2. Activity of Approved PI-Based Combination Therapies in RRMM^a

Study Name (reference)	Patient Population	Treatment	ORR (≥CR), %	Median	PFS Rates	Median	OS Rates
		Group (n)		PFS, months		OS, months	
2-drug regimens							
DOXIL-MMY-3001 ^{27,28}	RRMM, V-naive	V+PLD (324)	44 (4 ^b)	9.0	1-year: 26% ^c ; 2-year: NR	33.0	1-year: 81% ^c ; 2-year: 64% ^c
PANORAMA1 ^{30,31,34}	Relapsed or relapsed and refractory MM, 1– 3 previous treatments	Vd (381)	55 (6 ^b)	8.1	1-year: 36% ^c ; 2-year: 8% ^c	35.8	1-year: 81% ^c ; 2-year: 62% ^c
	Relapsed or relapsed and refractory MM, ≥2 previous regimens including V and an immunomodulatory agent	Vd (74)	39 (≥CR not reported)	4.7	1-year: 16% ^c ; 2-year: 5% ^c	19.5	1-year: 67% ^c ; 2-year: 44% ^c
ENDEAVOR ^{32,36}	RRMM, 1–3 previous treatments	Kd56 (464) Treatment until PD	77 (13)	18.7	1-year: 63% ^c ; 2-year: 44% ^c	47.6	1-year: 83% ^c ; 2-year: 70% ^c

	RRMM, 1–3 previous treatments	Vd (465) Treatment until PD	63 (6)	9.4	1-year: 42% ^c ; 2-year: 23% ^c	40.0	1-year: 83% ^c ; 2-year: 63% ^c
CASTOR ³³	Relapsed or relapsed and refractory MM, ≥1 previous treatment	Vd (247) Up to 8 cycles of treatment [6 months]	63 (9)	7.1 ^c	1-year: 27%; 2-year: 5% ^d	NR	1-year: 82% ^c ; 2-year: NR
3-drug regimens							
ASPIRE ^{20,37}	RRMM, 1–3 previous treatments	KRd (396)	87 (32)	26.3	1-year: 77% ^c ; 2-year: 55% ^c	48.3	1-year: 89% ^c ; 2-year: 73% ^c
TOURMALINE-MM1 ³⁵	Relapsed, refractory, or relapsed and refractory MM, 1–3 previous treatments	IRd (360)	78 (14)	20.6	1-year: 67% ^c ; 2-year: NR	NR	NR
CASTOR ³³	Relapsed or relapsed and refractory MM, ≥1 previous treatment	DVd (251)	83 (19)	16.7 ^c	1-year: 61%; 2-year: 37% ^e	NR	1-year: 82% ^c ; 2-year: NR
PANORAMA1 ^{30,31,34}	Relapsed or relapsed and refractory MM, 1–3 previous treatments	PanVd (387)	61 (11 ^b)	12.0	1-year: 49% ^c ; 2-year: 21% ^c	40.3	1-year: 80% ^c ; 2-year: 67% ^c

Relapsed or relapsed and refractory MM, ≥2 previous regimens including V and an immunomodulatory agent	PanVd (73)	59 (≥CR not reported)	12.5	1-year: 51% ^c ; 2-year: 6% ^c	25.5	1-year: 74% ^c ; 2-year: 58% ^c
---	------------	--------------------------	------	---	------	--

^aOnly data from pivotal studies supporting regulatory approval are presented. All studies used IMWG response criteria except for DOXIL-IMMY-3001 and PANORAMA1, which used EBMT criteria. ^bRate of CR. ^cEstimated from Kaplan-Meier curves. ^dValue is from an updated analysis of CASTOR.³⁸ ^ePresented at the 2017 Annual Meeting of the American Society of Hematology.³⁸

Abbreviations: CR=complete response; DVd=daratumumab, bortezomib, and dexamethasone; EBMT=European Group for Blood and Marrow Transplant; IMWG=International Myeloma Working Group; IRd=ixazomib, lenalidomide, and dexamethasone; Kd56=carfilzomib (56 mg/m²) and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; MM=multiple myeloma; NR=not reported, not reached, or not estimable; ORR=overall response rate; OS=overall survival; PanVd=panobinostat, bortezomib and dexamethasone; PD=progressive disease; PFS=progression-free survival; PI=proteasome inhibitor; PLD=pegylated liposomal doxorubicin; Rd=lenalidomide and dexamethasone; RRM=relapsed or refractory multiple myeloma; V=bortezomib; Vd=bortezomib and dexamethasone.

Table 3. Activity of Approved PI-Based Combination Therapies at First Relapse (1 previous line of therapy)

Study Name (reference)	Patient Population	Treatment Group (n)	ORR (≥CR), %	Median PFS, months	PFS Rates	Median OS, months
2-drug regimens						
ENDEAVOR ^{61,62}	Relapsed or refractory MM	Kd56 (232)	82 (12)	22.2	1-year: 71% ^a ; 2-year: 43% ^a	NR
	Relapsed or refractory MM	Vd [until PD] (232)	66 (8)	10.1	1-year: 45% ^a ; 2-year: NR	NR
CASTOR ^{38,63}	Relapsed or relapsed and refractory MM	Vd [6 months] (113)	74 (15) ^b	7.9	1-year: 29%; 2-year: 8% ^b	NR
3-drug regimens						
ASPIRE ^{37,64}	RRMM	KRd (184)	87 (34)	29.6	1-year: 80% ^a ; 2-year: 58% ^a	47.3
TOURMALINE-MM1 ⁶⁵	Relapsed, refractory, or relapsed and refractory MM	IRd (212)	77 (9)	20.6	1-year: 65% ^a ; 2-year: NR	NR
CASTOR ^{38,63}	Relapsed or relapsed and refractory MM	DVd (122)	92 (43) ^b	26.2 ^b	1-year: 78%; 2-year: 55% ^b	NR

^aEstimated from Kaplan-Meier curves. ^bPresented at the 2017 Annual Meeting of the American Society of Hematology.

Abbreviations: BTZ=bortezomib; CR=complete response; DVd=daratumumab, bortezomib, and dexamethasone; IRd=ixazomib, lenalidomide, and dexamethasone;

Kd56=carfilzomib (56 mg/m²) and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; MM=multiple myeloma; NR=not reported, not reached, not

estimable; ORR=overall response rate; OS=overall survival; PFS=progression-free survival; PI=proteasome inhibitor; Rd=lenalidomide and dexamethasone;
RRMM=relapsed or refractory multiple myeloma; Vd=bortezomib and dexamethasone.

853

Table 4. Activity of Approved PI-Based Combination Therapies When Used After Prior PI Exposure (Re-Treatment With PIs)

Study Name (reference)	Patient	Treatment	ORR	Median	PFS Rates	Median OS, months
	Population	Group (n)	(≥CR), %	PFS, months		
2-drug regimens						
RETRIEVE ²⁹	Patients with MM who had relapsed after achieving ≥PR with initial V-based therapy	V±DEX re-treatment (126)	40 (1 ^a)	8.4 (TTP)	1-year TTP rate: 12% ^b ; 2-year TTP rate: NR	NR
PANORAMA1 ^{31,34}	Relapsed or relapsed and refractory MM, prior exposure to V and an immunomodulatory drug	Vd (99)	41 (≥CR not reported)	5.8	1-year: 21% ^b ; 2-year: 4% ^b	24.7
ENDEAVOR ^{61,62}	Relapsed or refractory MM, prior V exposure	Kd56 (250)	71 (≥CR not reported)	15.6	1-year: 57% ^b ; 2-year: 35% ^b	47.6

	Relapsed or refractory MM, prior V exposure	Vd [until PD] (252)	60 (≥CR not reported)	8.1	1-year: 36% ^b ; 2-year: NR	32.8
CASTOR ³³	Relapsed or relapsed and refractory MM, prior V exposure	Vd [6 months] (164)	NR	6.7	NR	NR
3-drug regimens						
PANORAMA1 ^{31,34}	Relapsed or relapsed and refractory MM, prior exposure to V and an immunomodulatory drug	PanVd (94)	58.5 (≥CR not reported)	10.6	1-year: 48% ^b ; 2-year: 9% ^b	27.2
ASPIRE ⁶⁴	RRMM, prior V exposure	KRd (261)	86 (30)	24.4	1-year: 75% ^b ; 2-year: 52%	NR
TOURMALINE-MM1 ⁶⁵	Relapsed, refractory, or relapsed and refractory MM, PI-	IRd (250)	77 (9)	18.4	1-year: 67% ^b ; 2-year: NR	NR

	exposed					
CASTOR ³³	Relapsed or relapsed and refractory MM	DVd (162)	NR	12.3	NR	NR

^aResults reported for CR. ^bEstimated from Kaplan-Meier curve.

Abbreviations: CR=complete response; DVd=daratumumab, bortezomib, and dexamethasone; IRd=ixazomib, lenalidomide, and dexamethasone; Kd56=carfilzomib (56 mg/m²) and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; MM=multiple myeloma; NR=not reported; ORR=overall response rate; OS=overall survival; PFS=progression-free survival; PI=proteasome inhibitor; PanVd=panobinostat, bortezomib and dexamethasone; PD=progressive disease; PLD=pegylated liposomal doxorubicin; Rd=lenalidomide and dexamethasone; RRMM=relapsed or refractory multiple myeloma; TTP=time to progression; V=bortezomib; Vd=bortezomib and dexamethasone.

855

856

Table 5. Activity of PI-Based Combination Therapies When Used in PI-Refractory Patients

Study Name or NCT#	Phase	Patient Population	Treatment Group (n)	ORR, %	Median PFS, months	Median OS, months
Approved regimens						
PX-171-003-A1 ⁶⁸	2	RRMM, refractory to V in the last prior line	Single-agent carfilzomib (45% of 266 enrolled patients)	19	NR	NR
ASPIRE ⁶⁴	3	RRMM, refractory to V in any previous regimen	KRd (60)	80	22.3	NR
NCCN-recommended regimens						
PANORAMA 2 ⁵⁰	2	RRMM, refractory to V	PanVd (55)	35	5.4	NR
CHAMPION-1 ¹⁷	1/2	RRMM, refractory to V	Once-weekly Kd (54)	63	NR	NR
NCT01496118 ⁴⁹	1/2	RRMM, refractory to V	Carfilzomib+panobinostat (15)	67	NR	NR
NCT01549431 ^{a,69}	1	RRMM, refractory to V	Carfilzomib+panobinostat (16)	44	NR	NR
NCT01464034 ⁷⁰	1	RRMM	KPd (32 ^b)	50	7.2	20.6

NCT02004275 ⁵⁴	1/2	MM, refractory to LEN and a PI	IPd (17)	62 ^c	NR	NR
NCT02119468 ⁵⁵	1/2	RRMM, refractory to LEN and V	IPd (14)	29	NR	NR

^aPresented at the 2014 ASH Annual Meeting.

^b30 of 32 patients were refractory to V.

^cORR was reported for 13 patients who received at least 1 cycle of therapy.

Abbreviations: IPd=ixazomib, pomalidomide, and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; LEN=lenalidomide; MM=multiple myeloma; NCCN, National Comprehensive Care Network; NR=not reported or not reached; ORR=overall response rate; OS=overall survival; PFS=progression-free survival; PI=proteasome inhibitor; PanVd=panobinostat, bortezomib and dexamethasone; RRMM=relapsed or refractory multiple myeloma; V=bortezomib; Vd=bortezomib and dexamethasone.

Table 6. Safety of Approved PI-Based Combination Therapies in RRMM

Study Name (reference)	Treatment Group, Safety Population (n)	Median	Any AE, %	Grade ≥3 AEs, %	SAEs, %	Discontinuations Due to AEs, (%)	Dose
		Treatment					Reductions
		Duration					Due to AEs, (%)
2-drug regimens							
DOXIL-MMY- 3001 ²⁷	V+PLD (318)	5 cycles; 105 days	98	80 ^a	36	V: 30; PLD: 36	NR
PANORAMA1 ³¹	Vd (377)	NR	NR	82 ^a	42	18	NR
ENDEAVOR ³⁶	Kd56 (463)	48.0 weeks	99	81	59	14	32
	Vd [until PD] (456)	27.0 weeks	99	71	40	16	50
CASTOR ³³	Vd [6 months] (237)	Median follow-up period was 7.4 months	95	62 ^a	NR	9	NR
3-drug regimens							
PANORAMA1 ³¹	PanVd (381)	NR	NR	96 ^a	60	34	NR
ASPIRE ³⁷	KRd (392)	72 weeks for	98	87	65	20	11 ^b

carfilzomib							
TOURMALINE-MM1 ³⁵	IRd (361)	17 cycles	98	74	47	17	56
CASTOR ³³	DVd (243)	Median follow-up period was 7.4 months	99	76 ^a	NR	7	NR

^aAny grade 3 or 4 AE. ^bReported in the primary analysis.²⁰

Abbreviations: AE=adverse event; DVd=daratumumab, bortezomib, and dexamethasone; IRd=ixazomib, lenalidomide, and dexamethasone; Kd56=carfilzomib (56 mg/m²) and dexamethasone; KRd=carfilzomib, lenalidomide, and dexamethasone; NR=not reported; PI=proteasome inhibitor; PanVd=panobinostat, bortezomib and dexamethasone; PD=progressive disease; PLD=pegylated liposomal doxorubicin; RRMM=relapsed or refractory multiple myeloma; SAE=serious AE; V=bortezomib; Vd=bortezomib and dexamethasone.