Learning condyle repositioning during orthognathic surgery with a surgical navigation system

R. Lartizien, I. Zaccaria, C. Savoldelli, L. Noyelles, E. Chamorey, J.L. Cracowski, G. Bettega

► To cite this version:

R. Lartizien, I. Zaccaria, C. Savoldelli, L. Noyelles, E. Chamorey, et al.. Learning condyle repositioning during orthognathic surgery with a surgical navigation system. International Journal of Oral and Maxillofacial Surgery, 2019, 48, pp.952 - 956. 10.1016/j.ijom.2019.01.018 . hal-03486119

HAL Id: hal-03486119

<https://hal.science/hal-03486119>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Financial disclosure statement:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Running heads:

Recto: Learning condyle repositioning with a surgical navigation system

Verso: Lartizien et al.

Correspondence:

R. Lartizien

Maxillofacial and Plastic Surgery Department

University Hospital of Grenoble

Avenue Maquis-du-Grésivaudan

38700 La Tronche

France

E-mail: rodolphe.lartizien@gmail.com

[ARTICLE TYPE – CLINICAL PAPER, ORTHOGNATHIC SURGERY]

Learning condyle repositioning during orthognathic surgery with a surgical navigation system

R. Lartizien^{1,2,3*}, I. Zaccaria⁴, C. Savoldelli⁵, L. Noyelles³, E. Chamorey⁶, J.L. Cracowski^{2,7}, G. Bettega³

¹Maxillofacial and Plastic Surgery Department, University Hospital of Grenoble, Avenue Maquis-du-Grésivaudan, 38700 La Tronche, France

²Université Grenoble Alpes, Medical Faculty, 23 Avenue Maquis du Grésivaudan, 38700 La Tronche, France

³Maxillofacial Surgery Department, Annecy Genevois Hospital, 1 Avenue de l'Hôpital, 74370 Epagny Metz-Tessy, France

⁴Clinical Research Department, Annecy Genevois Hospital, 1 Avenue de l'Hôpital, 74370 Epagny Metz-Tessy, France

⁵Oral and Maxillofacial Surgery, Head and Neck Institute, University Hospital of Nice, Nice, France

⁶Epidemiology and Biostatistics Unit, Antoine Lacassagne Center, Nice, France

⁷Clinical Pharmacology Unit, INSERM CIC1406, CHU Grenoble-Alpes, F-38000 Grenoble, France

Address:

R. Lartizien

Maxillofacial and Plastic Surgery Department

University Hospital of Grenoble

Avenue Maquis-du-Grésivaudan

38700 La Tronche

France

E-mail: rodolphe.lartizien@gmail.com

Abstract. Condyle repositioning during bilateral sagittal splint osteotomy (BSSO) is a challenging step for the inexperienced surgeon. We aimed to demonstrate the benefit of navigation for learning the condyle repositioning. We treated 100 patients who underwent a BSSO. A trainee performed the condyle repositioning of one side in two phases. In the first one, the trainee positioned without watching the screen of the Orthopilot Navigation system (ONS). In the second one, the trainee could use the ONS to replace the condyle. Heuristic, anatomical and functional scores of each phase were recorded. Heuristic (17% vs. 75%; $p<0.0001$), anatomical (35% vs. 86%; $p<0.0001$) and functional (14% vs. 56%; $p<0.0001$) scores were significantly greater with the ONS. The ONS is a promising and original intraoperative learning tool for the repositioning of the condyle during BSSO.

Key words: orthognathic surgery; computer-assisted surgery; condyle repositioning; surgical education; medical education.

The final result of bilateral sagittal split osteotomies (BSSOs) mainly depends on the correct repositioning of the condyle intraoperatively¹. For the inexperienced surgeon, condyle repositioning is a challenging step. Many techniques and devices have been described to improve this skill²⁻⁷ but manual seating remains the method used by the majority of surgeons. The accuracy and reproducibility of this empirical repositioning requires extensive experience and its learning is very complex.

We routinely use in our unit the Orthopilot™ navigation system (ONS) for orthognathic surgery and particularly for condyle repositioning^{2,3}. While this device improved our postoperative occlusal and functional results, its pedagogical interest has not been studied to date.

In this work, our aim was to evaluate the benefit of navigation for learning condyle repositioning during BSSO and to see how resident and inexperienced surgeons acquired this surgical skill.

Materials and methods

This monocentric, observational and retrospective study had included 100 patients needing a BSSO (isolated or associated to a Le Fort I osteotomy), between October 2012 and December 2014. Study ethics approval was obtained on 6 July 2018 (CECIC Rhône-Alpes-Auvergne, Clermont-Ferrand, IRB 5891).

The surgical indication was confirmed after clinical examination, sagittal and frontal cephalometry, and a plaster cast study. The condyle centric relation was recorded preoperatively and documented with a dental splint. The BSSO itself was made by the same experienced surgeon on both sides; on the left side the condyle was repositioned

and the bicortical screw osteosynthesis was performed by the trainee under supervision of the senior surgeon who positioned and fixed the contralateral condyle. Only the data from the trainee side were considered in this study.

Nineteen trainees were involved in 100 BSSOs. They performed between one and 14 surgical procedures.

Surgical procedure

The navigation procedure has been described elsewhere². Infrared reflectors (rigid bodies) were anchored via a small vice on each coronoid process. An intermediary fixation between the rigid body and the osseous anchorage allowed the removal and replacement of the rigid body without losing the landmark. Condyle centric relation position was recorded by the system using the specific splint prepared preoperatively and sterilized. BSSOs were performed according to the Obwegeser–Dal Pont technique. Final occlusion was stabilized by a maxillomandibular fixation. The rigid body was fixed on its vice for condyle navigation. Condyle position was controlled on the computer screen, the surgeon trying to minimize the three translations and the three rotations superimposing the current virtual image of the condyle on the planned one (Fig. 1). The goal was to be as close as possible to zero. The condyle repositioning procedure was conducted in two phases. During the first phase, the trainee manually repositioned the condyle; the screen of the computer was occluded so that he had no access to the navigation data. Once the surgeon believed that the condyle was correctly positioned, the condyle the data were collected by the navigation system and the screen was freed. These data were stocked in a file entitled “without ONS”. During the second phase, the trainee tracked the position on the computer screen and replaced the condyle minimizing the values of rotation and translation. When values of

discrepancies were as closed to 0 as possible, the osteosynthesis was performed with three bicortical screws. This second set of data was recorded in a second file entitled “with ONS”.

Statistical analysis

The collected data included: (1) seniority and number of intervention for each surgeons; (2) type of surgical procedure (BSSO isolated or associated to a Le Fort I osteotomy, asymmetry, vertical movement); (3) discrepancies from the reference position of the condyle without and with the ONS (translation top/bottom (Ttb), translation back/front (Tbf), translation left/right (Tlr), frontal rotation (Rf), axial rotation (Ra) and sagittal rotation (Rs)).

The primary outcome was the heuristic score, calculated as follows:

$$SH = \sqrt{Ttb^2 + Tlr^2 + Tbf^2 + Rs^2 + Ra^2 + Rf^2}$$

Heuristic score was considered good when ≤ 5 . The secondary outcomes were the anatomical score and the functional score. Anatomical score was defined by Ttb, Tbf and Rs. Functional score was defined by Ra, Rf and Tlr. Anatomical and functional scores were considered good when each value was ≤ 2 .

Scores could be both qualitative (good/not good) and quantitative variables (values).

Each score was calculated with and without the ONS (paired series). SA, SH and SF with and without the ONS were analysed by a McNemar test for paired series⁸, in order to know whether the condyle repositioning was significantly better with ONS.

Results

Fifty-seven patients were operated by isolated BSSO and 43 also had a Le Fort I osteotomy. Seventy-four patients had an asymmetry. A maxillary impaction was carried out on 31 patients.

The number of interventions performed by each surgeon is listed in Table 1.

The heuristic score was ≤ 5 and considered as good in 17% of the procedures without the ONS and in 75% with the ONS. The anatomical score was ≤ 2 in 35% of the procedures without the ONS and in 86% with the ONS. The functional score was ≤ 2 in 14% of the procedures without the ONS and in 56% with the ONS. The use of the ONS significantly improved the three scores of condyle repositioning ($p < 0.0001$). Table 2 (upper) shows the heuristic scores with and without the ONS: 38 patients had a bad score without the ONS and remained bad with the ONS, 55 patients were bad without the ONS and achieved a good score with the ONS, one patient had a good score without the ONS and became bad with the ONS, and six patients had a good score without the ONS and remained good with the navigation system. The remainder of Table 2 shows the same type of information for anatomical and functional scores.

The distribution of the discrepancy of each variable (Ra, Rf, Rs, Tbf, Trl and Ttb) with and without the ONS is listed in Fig. 2. Graphically, mean and the dispersion of the values were greater without the ONS, especially for the Ra and the Rs.

Fig. 3 shows the linear relation between the heuristic score with and without the ONS and the number of interventions performed by the surgeon. The ONS seemed to allow a better repositioning, but with increasing numbers of interventions (and experience) the regression lines tended to join. The intersection was around 15 interventions.

Discussion

The surgical education environment has changed in the past few decades with the tightening of the medical legislation (leading to a decreased autonomy of the practice of trainees), demands for patient safety, the development of new technologies and the limitation of working hours^{9,10}. This change imposed the development of new surgical educational tools: cadaveric dissection, simulations¹⁰, videos¹¹ and now virtual reality¹².

Our study demonstrated the educational value of the ONS for condyle repositioning during BSSO. The trainees obtained heuristic (17% vs. 75%; $p<0.0001$), anatomical (35% vs. 86%; $p<0.0001$) and functional (14% vs. 56%; $p<0.0001$) scores significantly greater with the ONS. The ONS gave a virtual image of the condyle and of the target (the final position), so for the trainee the good position is better understood, easier to reach and more accurate than with the empirical (blind) positioning.

The heuristic score was chosen to evaluate the improvement of the condyle repositioning because it took into account all of the parameters and considered the control of the three dimensions. The anatomical score represented the condyle position in the sagittal plane. We indicated in a previous study that it was the principal factor in the quality of postoperative occlusion and skeletal stability². The functional score reflected the axial and frontal control of the condyle position and was correlated in the same study with better functional results, especially considering postoperative mandibular motion.

In the present study, the functional score was the less improved by the ONS. It proved that perfect repositioning even with tool control is not always possible. Sometimes compromises are needed. The discrepancies are generally due to anatomical conflicts between the proximal and the distal bone segment¹³, especially in asymmetric cases. Rigid osteosynthesis may increase this problem. The improvement of the functional

score required a remodelling of the osseous valves¹⁴, but this not always possible without damaging the infra-alveolar nerve.

The analysis of the distribution of the values of each direction with and without the ONS showed that the dispersion of the variables was smaller with the ONS with fewer outliers. The use of the ONS could avoid large discrepancies in the repositioning of the condyle. Berger et al. reached the same conclusion on this point⁴. Without the ONS the dispersions were greater, especially for Rs and Ra. These two dimensions were the most difficult to control without the ONS. This could be explained by the lack of landmarks in the exiguous exposition of the bone intra-orally.

The benefit of a condyle positional tool is the subject of a controversial and long-lasting debate among experts. It is not the subject of this work. We only wanted to emphasize the relevance of navigation in learning this difficult task.

Surgical practice is an essential step for the learning of trainees^{15,16} and the use of the ONS can be beneficial for their educational in orthognathic surgery. Indeed, it allows an inexperienced surgeon to perform a critical step of the surgical procedure without compromising patient safety and postoperative results, his actions being watched continuously. Thus, the ONS allowed to inexperienced surgeons to directly acquire and improve skills to replace the condyle to its reference position (whatever the position chosen preoperatively). This point was demonstrated by the representation of the linear relation between the heuristic score and the number of interventions with and without the ONS: with an increasing number of procedures, the trainee improved his heuristic score without the use of the ONS. The lines seemed to cross at around 15 procedures. In other words, a trainee would be able to place the condyle in the right position without using the ONS after 15 procedures with the ONS. Of course, learning can not be

summarized by a linear relation but it would be interesting to compare this number with those obtained from other learning tools.

Training is more efficient if it is multimodal¹⁷. The ONS is a new way to teach condylar repositioning and is complementary to other existing tools. To our knowledge, it is the only tool that can be used in the operating room.

Furthermore, evaluating technical skills and competence of a trainee in the operating room is difficult¹⁸, especially because the methods are subjective and unreliable. The ONS could be a good way of objectively following the heuristic scores of trainees and thus their progression.

Our study presented several limitations. It was a monocentric study and the number of procedures performed by each trainee was variable. Our results were mainly dependent on the scores obtained by the trainee who performed 14 procedures. A multicentric study with a mixed model (taking into account the correlation of the data) is necessary to confirm our results. The side performed by the trainee was not randomized. This would have introduced a bias. The side should be randomized in future explorations. Another bias was the failure to take into account the trainees' dominant hand. Indeed, a right-handed surgeon would find it more difficult to operate on the left side, for example. This bias had little consequence because the trainee only performed the condyle repositioning and the osteosynthesis (and not the entire intervention) so he could use his dominant hand whatever the operated side. Consequently, a learning effect was observed regardless of the operated side or the dominant hand. However, we should take this point into account in the future.

In conclusion, the ONS is a promising and original intraoperative learning tool for the repositioning of the condyle during BSSO. Its use allowed trainees to improve their

241 accuracy and their surgical skills during this challenging step. However, further
242 explorations are needed to confirm these preliminary results.

243

244 **Funding**

245 None.

246 **Competing interests**

247 None.

248 **Ethical approval**

249 Study ethics approval was obtained on 06 July 2018 (CECIC Rhône-Alpes-Auvergne,
250 Clermont-Ferrand, IRB 5891).

251 **Patient consent**

252 Not required.

253

REFERENCES

1. Landes CA, Sterz M. Proximal segment positioning in bilateral sagittal split osteotomy: intraoperative controlled positioning by a positioning splint. *J Oral Maxillofac Surg* 2003;61:1423-1431.
2. Bettega G, Cinquin P, Lebeau J, Raphael B. Computer-assisted orthognathic surgery: clinical evaluation of a mandibular condyle repositioning system. *J Oral Maxillofac Surg* 2002;60:27-34.
3. Bettega G, Leitner F. [Computer assisted orthognathic surgery: Condyle repositioning.]. *Rev Stomatol Chir Maxillofac Chir Orale* 2013. Jul 17. doi: 10.1016/j.revsto.2013.06.001.
4. Berger M, Nova I, Kallus S, Ristow O, Eisenmann U, Dickhaus H, Engel M, Freudlsperger C, Hoffmann J, Seeberger R. Electromagnetic navigated condylar positioning after high oblique sagittal split osteotomy of the mandible: a guided method to attain pristine temporomandibular joint conditions. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2018;125(5):407-414.e1.
5. Luhr HG. The significance of condylar position using rigid fixation in orthognathic surgery. *Clin Plast Surg* 1989;16:147-156.
6. Beziat JL, Babic B, Ferreira S, Gleizal A. [Justification for the mandibular-maxillary order in bimaxillary osteotomy]. *Rev Stomatol Chir Maxillofac*. 2009;110:323-326.
7. Veyssiere A, Leprovost N, Ambroise B, Prevost R, Chatellier A, Benateau H. Study of the mechanical reliability of an S-shaped adjustable osteosynthesis plate for bilateral sagittal split osteotomies. Study on 15 consecutive cases. *J Stomatol Oral Maxillofac Surg*. 2018;119:19-24.
8. Mc NQ. Note on the sampling error of the difference between correlated proportions or percentages. *Psychometrika* 1947;12:153-157.

- 279 **9.** Haluck RS, Krummel TM. Computers and virtual reality for surgical education in
280 the 21st century. *Arch Surg*. 2000;135:786-792.
- 281 **10.** Reznick RK, MacRae H. Teaching surgical skills—changes in the wind. *N Engl J*
282 *Med*. 2006;355:2664-2669.
- 283 **11.** Mota P, Carvalho N, Carvalho-Dias E, Joao Costa M, Correia-Pinto J, Lima E. Video-
284 based surgical learning: improving trainee education and preparation for
285 surgery. *J Surg Educ* 2018;75:828-835.
- 286 **12.** Pulijala Y, Ma M, Pears M, Peebles D, Ayoub A. An innovative virtual reality
287 training tool for orthognathic surgery. *Int J Oral Maxillofac Surg* 2018;47:1199-
288 1205.
- 289 **13.** Suojanen J, Leikola J, Stoor P. The use of patient-specific implants in orthognathic
290 surgery: A series of 30 mandible sagittal split osteotomy patients. *J*
291 *Craniomaxillofac Surg* 2017;45:990-994.
- 292 **14.** Ellis E, 3rd. A method to passively align the sagittal ramus osteotomy segments. *J*
293 *Oral Maxillofac Surg* 2007;65:2125-2130.
- 294 **15.** Kotsis SV, Chung KC. Application of the "see one, do one, teach one" concept in
295 surgical training. *Plast Reconstr Surg* 2013;131:1194-1201.
- 296 **16.** Sutkin G, Littleton EB, Kanter SL. Intelligent cooperation: A framework of
297 pedagogic practice in the operating room. *Am J Surg* 2018;215:535-541.
- 298 **17.** Kim S, Dunkin BJ, Paige JT, Eggerstedt JM, Nicholas C, Vassiliou MC, Spight DH,
299 Pliego JF, Rush RM, Lau JN, Carpenter RO, Scott DJ. What is the future of training
300 in surgery? Needs assessment of national stakeholders. *Surgery* 2014;156:707-
301 717.
- 302 **18.** Winckel CP, Reznick RK, Cohen R, Taylor B. Reliability and construct validity of a
303 structured technical skills assessment form. *Am J Surg* 1994;167:423-427.

304 *Table 1.* Number of interventions by surgeon

Number of interventions	Number of surgeons	Percentage	Cumulated number of surgeons	Cumulated percentage
1	4	21.05	4	21.05
2	3	15.79	7	36.84
3	1	5.26	8	42.11
4	2	10.53	10	52.63
5	2	10.53	12	63.16
6	1	5.26	13	68.42
8	2	10.53	15	78.95
9	1	5.26	16	84.21
11	1	5.26	17	89.47
13	1	5.26	18	94.74
14	1	5.26	19	100.00

305

306

307 *Table 2. Agreement of each score with and without Orthopilot Navigation system (ONS)*
308

Heuristic score (HS)			
HS without ONS	HS with ONS		
%	Bad	Good	Total
Bad	38	55	93
Good	1	6	7
Total	39	61	100
Anatomical score (AS)			
AS without ONS	AS with ONS		
%	Bad	Good	Total
Bad	12	53	65
Good	2	33	35
Total	14	86	100
Functional score (FS)			
FS without ONS	FS with ONS		
%	Bad	Good	Total
Bad	41	44	85
Good	4	11	15
Total	45	55	100

309

310 **CAPTIONS TO ILLUSTRATIONS**

311 *Fig. 1.* Navigation system software interface.

312 *Fig. 2.* Distribution of the discrepancies of each dimension with and without Orthopilot
313 Navigation system (ONS).

314 *Fig. 3.* Linear representation between the heuristic score and the number of
315 interventions with and without Orthopilot Navigation system (ONS).

316

L

0%

1

Front

Score
3

0

Left

1°

Anterior

1

Top

0°

Right

2° Left

Navigation : Left Condyle

Box-Plot : Discrepancies with/without OSN