Can we refine the management of blunt liver trauma?

L. Barbier, M. Calmels, M. Lagadec, T. Gauss, P.-S. Abback, F. Cauchy, M. Ronot, O. Soubrane, C. Paugam-Burtz

► To cite this version:

L. Barbier, M. Calmels, M. Lagadec, T. Gauss, P.-S. Abback, et al.. Can we refine the management of blunt liver trauma?. Journal of Visceral Surgery, 2019, 156, pp.23 - 29. 10.1016/j.jviscsurg.2018.03.013 . hal-03486497

HAL Id: hal-03486497

<https://hal.science/hal-03486497>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Can we refine the management of blunt liver trauma?

Peut-on améliorer la prise en charge des traumatismes hépatiques fermés ?

Running title: Blunt liver trauma

Titre courant : Traumatismes hépatiques fermés

Louise Barbier^{1,4}, Mélanie Calmels¹, Matthieu Lagadec², Tobias Gauss³, Paer-Selim Abback³,
François Cauchy¹, Maxime Ronot², Olivier Soubrane¹, Catherine Paugam-Burtz³

Affiliations: 1: HPB surgery, 2: Radiology, 3: Departments of Anesthesiology and Critical
Care, Hôpital Beaujon, Université Paris 7 Diderot, DHU Unity-FRANCE

4: Department of Digestive Surgery, Hôpital Trousseau, Université Rabelais, Tours, FHU
SUPPORT-FRANCE

Corresponding author:

Louise Barbier, MD

Department of HPB surgery

Hôpital Beaujon, 100 avenue du Général Leclerc, 92110 Clichy-la-Garenne

Ph: +33-1-40-87-58-95; Fax: + 33-1-40-87-09-26

Email to: louisebarbier@hotmail.fr

Can we refine the management of blunt liver trauma?

Peut-on améliorer la prise en charge des traumatismes hépatiques fermés ?

Abstract

Aim: To describe the management of blunt liver injury and to study the potential relation between delayed complications, type of trauma mechanisms and liver lesions.

Patients and Methods: This is a retrospective single center study including 116 consecutive patients admitted with blunt liver injury between 2007 and 2015.

Results: Initial CT-scan identified an active bleeding in 33 (28%) patients. AAST (American Association for the Surgery of Trauma) grade was 1 to 3 in 82 (71%) patients and equal to 5 in 15 (13%) patients. Eighty (69%) patients had NOM, with a success rate of 96%. Other abdominal organ lesions were associated to invasive initial management. A follow-up CT-scan was useful to detect hepatic and extra-hepatic complications (46 complications in 80 patients), even without clinical or biological abnormalities. Subsequent hepatic complications such as bleeding, pseudo aneurysms, biloma and biliary peritonitis developed in 15 patients and were associated with the severity of blunt liver injury according to AAST classification (3.7 ± 1.0 vs. 3.0 ± 1.1 , $p=0.010$). Total biliary complications occurred in 13 patients and were significantly more frequently observed in patients with injury of central segments 1, 4 and 9 (69% vs. 36%, $p=0.033$).

Conclusions: Non-operative management is possible in most blunt liver injury with a success rate of 96%. A systematic CT-scan should be advocated during follow-up, especially when AAST grade is equal or superior to 3. Biliary complications should be suspected when lesions involve segments 1, 4 and 9.

Key words: Trauma centers; radiology, interventional; liver, trauma; non-operative management; CT-scan; biliary fistula; therapeutic embolization

Résumé

But: Décrire la prise en charge des traumatismes hépatiques fermés (THF) et étudier la relation entre les complications, le type de mécanisme et les lésions observées.

Patients et Méthodes: Etude rétrospective monocentrique de 116 patients consécutifs entre 2007 et 2015.

Résultats: Un saignement actif a été identifié au scanner chez 33 (28%) patients. Le grade AAST (American Association for the Surgery of Trauma) était inférieur ou égal à 3 chez 82 (71%) patients, et égal à 5 chez 15 (13%) des patients. Quatre-vingts (69%) patients ont eu une prise en charge non opératoire, avec un taux de succès de 96%. La prise en charge initiale chirurgicale était plus fréquente en cas de lésions associées d'autres organes. Un scanner de contrôle a permis d'identifier 46 complications hépatiques et extra-hépatiques chez 80 patients, même en l'absence d'anomalies cliniques ou biologiques. Quinze patients ont présenté des complications hépatiques ultérieures (saignement, pseudo anévrysme, biliome, péritonite biliaire), associées à la sévérité du traumatisme hépatique initial selon la classification AAST (3.7 ± 1.0 vs. 3.0 ± 1.1 , $p=0.010$). Les complications biliaires (13 patients) étaient plus fréquentes en cas de lésion des segments I, IV et IX (69% vs. 36%, $p=0.033$).

Conclusion: La prise en charge non opératoire des THF est possible, avec un taux succès de 96%. Un scanner de contrôle systématique est préconisé lors du suivi, plus particulièrement pour les lésions avec un grade AAST supérieur ou égal à 3. Des complications biliaires doivent être suspectées en cas d'atteinte des segments I, IV et IX.

Key words:

Trauma centers ; radiologie interventionnelle ; traumatisme hépatique; traitement non-opératoire; scanner; fistule biliaire; embolisation thérapeutique

Introduction

Non-operative management (NOM) is the reference treatment of blunt liver injuries (BLI) in hemodynamically stabilized patients [1–3]. NOM can be proposed in 65% to 85% of patients with BLI according to liver injury grading and associated lesions [2–4]. Computed Tomography-scan (CT-scan) is the corner stone in the assessment of BLI and the diagnostic modality of choice [5]. CT-scan allows the grading of the liver injury as established by the American Association for the Surgery of Trauma (AAST) [6] and the identification of associated lesions.

Failure of NOM is defined by the need for a delayed operation [3] and represents 5% [2] to 11% [7] of patients without early surgery. Systolic blood pressure on admission of 100 mmHg or less and the presence of other abdominal organ injury have been identified as independent factors of failure of NOM in AAST grades 4 and 5 BLI [3]. There is also no consensus on the monitoring of patients with NOM, specifically regarding the systematic imaging re-assessment.

Biomechanical and modelization studies of livers subjected to injury have focused on the mechanisms of the lesions and the relationship with both liver anatomy and its attachment system [8-10]. However, there is no radio-anatomical study focusing on the relationship between the type of mechanism, the type of liver injury and the potential vascular and biliary complications.

The aims of this study were to describe the initial management of BLI and to study the potential relation between delayed complications, type of trauma mechanisms and liver lesions.

Patients and Methods

Type of study

This retrospective study was conducted in the Beaujon Hospital a designated Trauma Center in the Paris region (France). Data from all consecutive patients with BLI who were admitted at the Trauma Center between 2007 and 2015 were collected from computerized clinical files. The trauma system and trauma management in the Paris region has been previously described [11].

Management of patients with trauma

Trauma patients were admitted in the resuscitation area after pre-alert from a physician-lead enhanced care team. The trauma leader, a senior intensivist, coordinated the trauma team including other specialists such as general surgeon, orthopedic surgeon, and radiologist. If the patient could be hemodynamically stabilized, a body CT-scan was performed. If not, the patient went directly to the dedicated operating room after minimum assessment included Chest X-Ray, pelvis X-Ray and FAST ultrasound. When there was no active hemorrhage or significant associated lesions, the NOM was the treatment of choice for BLI. In case of identified isolated active hemorrhage, interventional radiology with selective intra-arterial embolization was proposed, using either temporary agent such as Gel foam cubes or a permanent one such as micro coil depending on the type of vascular lesion and the selectivity of the catheterization. Indications for emergency laparotomy were uncontrolled hemorrhage or associated lesions such as perforation of hollow organs. Surgery was based on damage control laparotomy [12] with peri hepatic packing in case of massive venous hemorrhage involving hepatic veins, splenectomy, hollow organs sutures or stoma in case of perforation...

Failure of NOM of BLI was defined as delayed surgery (but not interventional radiology) or death related to BLI during in-hospital stay [3].

CT-scan

The initial CT-scan was performed in hemodynamically stabilized patients and with the presence of the trauma leader and surgeons. Radiologists perform a body CT-scan according to trauma protocol with intravenous contrast enhancement and different time points: arterial phase or mixed arterio-portal phase and delayed phase. Liver lesions were described according to the recommendations using the AAST liver injury scale [6,13]. However, this scale has limitations and additional imaging findings were searched for: extension of the injury to the hepatic vessels, active bleeding, hemoperitoneum or hemoretroperitoneum, presence of a pseudo aneurysm, anatomic localization of the injury, and imaging findings of non-liver injuries (thorax, spleen, pancreas, bowel). The locations of the liver injuries that were specifically studied were: central segments (1, 4 and 9) that include the main biliary convergence, right and main portal scissures, frequently involved in case of acceleration or deceleration trauma, that respectively include the posterior and right biliary ducts on their path, and the right liver. CT-scan during the follow-up was performed according to the attending practitioner and depending on the initial lesions.

Definitions

Hemodynamic instability was defined [14] by mean arterial pressure less than 65 mm Hg and/or systolic pressure less than 90 mmHg, and the need for crystalloids and/or red blood cells transfusions. Hypothermia was defined by body temperature less than 35 °C, and acidosis by pH less than 7.36 [15]. Coagulopathy was defined by the association of low

platelets count (less than 100 000/mm³), low fibrinogen (less than 1 g/L), prothrombin time above 3 s, and increase in soluble fibrin or fibrinogen degradation products [16].

Statistical analysis

Qualitative variables were described as absolute numbers and percentages. Two groups of qualitative variables were compared using the Chi-2 test, with Yates' correction as appropriate and the Fischer's exact test according to the number of data [and the distribution of the variable](#). Quantitative variables were described as mean and standard deviation of the mean. Two groups of quantitative variables were compared using the unpaired t-test or the Mann-Whitney test, according to the number of data.

Statistical analysis was performed using Microsoft® Excel® for Mac 2011 v 14.7.1 and Prism 7 for Mac OS X v 7.0b, GraphPad Software, Inc.

Ethics

Data for this study were extracted from a regional prospective trauma registry. The registry approval was obtained from the Institutional Review Board (Comité de Protection des Personnes, Paris VI) from the Advisory Committee for Information Processing in Health Research (CCTIRS, 11.305bis) and from the National Commission on Informatics and Liberties (CNIL, 911461). The structure of the database integrates algorithms for consistency and coherence. A central administrator assures the data monitoring. Missing data were collected anonymously from medical records.

The present study being an observational, non-interventional, retrospective study, neither informed consent nor approval of the ethics committee was required to use data from patients' records before 2017 according to French legislation.

Results

Description of the total cohort (Table 1 and Table 2)

During the study period, 116 patients were admitted at the Beaujon Hospital Trauma Center with BLI. Most of them were men (n=86, 74%) and mean age was 31.5 ± 12.1 years old. At their admission (see *Table 1*), 35 (30%) patients presented with hemodynamic instability and 30 (26%) patients required transfusions of red blood cell packs. Overall, 46 (40%), 29 (25%) and 25 (22%) patients presented with acidosis, hypothermia and coagulopathy, respectively. Injury Scale Score was 32.1 ± 16.8 and **SAPS** was 34.8 ± 20.4 . Motorcycle accidents were the most common type of trauma (n=38, 33%), followed by car accident and falls from heights in the context of suicide (n=25, 22% each). The mechanism of trauma was a direct shock in most cases (n=76, 66%). Acceleration and deceleration mechanisms accounted for 31 (27%) patients.

Fifty-one (44%) patients had a FAST ultrasound in the resuscitation area: FAST was considered normal in 17 (33%), a minimal effusion was found in 29 (57%), and a major effusion in 5 (10%) patients. Twenty-one (41%) of patients with FAST were hemodynamically unstable, and two of them had emergency surgery straight after FAST ultrasound.

CT-scan was performed after resuscitation in all but three patients. In these three patients emergency surgery was performed first. Findings of CT-scan are detailed in table 2. The most common liver injury (88%) was parenchymal lesions with lacerations and contusions. An active arterial bleeding was identified in 33 (28%) patients, and pseudo aneurysms in 6 (5%) patients. There was no hepatic vein injury. Most of the vascular injuries were located in the right liver (n=51, 77%). Most of the patients had 1 to 3 AAST grade (n=82, 71%), and 15

(13%) patients had grade 5 lesions. Associated lesions were present in 94 (81%) patients, predominantly thoracic injuries.

Initial management (Figure 1 and Table 1)

Figure 1 describes the initial management of the 116 patients: 80 (69%) patients were hemodynamically stabilized and had NOM, while 35 (30%) patients had an invasive management with surgery and/or interventional radiology. One patient deceased from associated cerebral lesions after initial assessment with FAST ultrasound and CT-scan.

Indications for emergency laparotomy (n=14, 12%) included: massive hepatic bleeding (n=7), associated spleen trauma (n=5), associated perforation of a hollow organ (n=9). Totally, emergency laparotomy was performed for an isolated BLI in only 2 patients. Interventional radiology was performed in 27 patients, and in combination with surgery in 6 patients.

As expected, patients with initial invasive management (*see Table 1*) were more frequently hemodynamically instable with acidosis, coagulopathy and hypothermia at admission. They required more often transfusions, and the [SAPS](#) was higher. There was more often active bleeding at the initial CT-scan and the AAST liver injury grade was higher.

Follow-up: subsequent management and complications (Table 1 and Table 3)

Overall, 20 (17%) patients required further interventional radiology and/or surgery (*see Table 1*), which was more frequent in the initial invasive management group vs. NOM group (36% vs. 9%, $p<0.001$).

A total of 55 (47%) patients presented complications (Table 3). The most frequent complications were pulmonary complications (pleural effusion, pneumopathy, pulmonary embolism). Iterative bleeding and pseudo aneurysms were found in 5 (4%) patients each, and biliary complications in 7 (6%) patients. Mean total in-hospital stay was 29 ± 52 days.

In-hospital mortality rate was 4% (n=3) in the NOM group and 17% (n=6) in the invasive management group. Causes of death were as follows: associated neurological trauma (n=4), respiratory distress in patients with thorax trauma (n=2), uncontrollable hemorrhage arising from thorax, abdomen and/or pelvis (n=3). Six of the deceased patients had management with early laparotomy and/or interventional radiology. The death occurred at a mean of 35 ± 109 days following the trauma. All patients but one who deceased during in-hospital stay had a SAPS (simplified acute physiology score) score superior to 60 at initial management.

Totally, NOM was a success in 77 (96%) patients, with three patients who required delayed surgery after NOM trial (*see Table 1*). Regarding the three patients with failure of NOM, none of them had hemorrhagic shock or hemodynamic instability at initial presentation, and none of them required initial red blood cells transfusions. All of them presented lesions of the right liver and the central segments (1,4 and 9). Only one of them did not have other organ associated lesions. A delayed surgical intervention was required in one patient for hemorrhage and in two patients for biloma/biliary peritonitis. Causes of death of the three patients in the NOM group were not related to BLI (and were not considered as a failure of NOM, according to the definition [3]): neurological trauma (n=1), and thoracic trauma (n=2).

CT-scan during the follow-up

A CT-scan was performed during the in-hospital stay for 80 (69%) patients. The CT-scan was performed systematically in 46 (58%) patients, and based on clinical or biological abnormalities in 34 (42%). Totally, the follow-up CT-scan detected 33 (41%) complications and 4 (5%) hepatic vein thrombi. Systematic CT-scan detected 7 (15% of patients with systematic CT-scan) hepatic complications (2 pseudo aneurysms, 2 biloma and 3 hepatic vein thrombi) and 13 (28% of patients with systematic CT-scan) thoracic complications (pleural effusion and pneumopathy).

Risk factors for hepatic and biliary complications

Subsequent hepatic complications (bleeding, pseudo aneurysms, biloma and biliary peritonitis) that developed during follow-up arose in a total of 15 patients. Initial AAST liver injury-grade was higher in patients who developed hepatic complications: 3.7 ± 1.0 vs. 3.0 ± 1.1 ($p=0.010$); all patients but one who developed hepatic complications had AAST liver injury grade equal or superior to 3.

Total biliary complications (at initial assessment and during follow-up) included biloma, biliary peritonitis and gallbladder injury and were present in 13 patients. The only factor associated with total biliary complications was injury of segments 1/4/9 (69% vs. 36%, $p=0.033$, see Table 4).

Discussion

Our findings confirm that NOM is feasible in most patients with BLI. The rate of 69% in the present study is consistent with the largest series of the literature [2-4], as is its success rate of 96% [3]. Patients most likely to undergo initial invasive management presented with more hemodynamic instability and higher SAPS. Associated lesions, such as spleen trauma and perforation of hollow organs were responsible for most indications of emergency surgery. Other abdominal organ injury has as well been found to be associated with failure of NOM [3]. Furthermore, neurological and thoracic lesions were considered as the cause of death in 6 of the 9 patients who died at hospital, with uncontrollable hemorrhage accounting for the three remaining deaths.

Classification of BLI according to the AAST classification [6] is a standardized tool for all radiologists and clinicians. In this study, AAST grade was associated with initial invasive management and development of biliary and vascular hepatic complications during follow-up. Systematic CT-scan during in-hospital stay after BLI and spleen trauma has been questioned in children [17] and in adults [18]. In the present study, although more complications were diagnosed on follow-up CT-scan when clinical or biological abnormalities were present, a systematic CT-scan detected hepatic and thoracic complications in respectively 15% and 28% of the patients. We confirm the results of Osterballe et al. [18], who showed that hepatic pseudo aneurysms develop in 4% of the BLI patients, and recommended systematic CT-scan during follow-up.

The only factor associated with biliary complications was the injury of central segments of the liver, i.e. segments 1, 4 and 9. Unlike Yuan et al. [19], we did not identify an association

between trans arterial embolization or high-grade liver injury and major bile leak. Segments 1, 4 and 9 are the segments in contact with the biliary confluence; if injured, a biloma or biliary peritonitis should be suspected and searched for systematically with imaging.

The main limit of this study is our failure to identify more associations between the mechanism of trauma, the detailed anatomical location of BLI and the subsequent complications. The retrospective nature of the study and the complexity of liver injuries mechanisms may be responsible for a certain lack of power and the incapacity to study complex mechanisms. Nonetheless, our results confirmed that NOM was possible in most BLI with a success rate of 96%, and that a systematic CT-scan should be advocated during follow-up, especially when AAST grade is equal or superior to 3. Biliary complications should always be suspected when lesions involve central liver segments.

Conclusion

This study demonstrated that NOM was feasible in 69% of BLI patients with a success rate of 96%. Associated organ lesions accounted for most of emergency surgery. We recommend a systematic follow-up CT-scan that allows detection of both hepatic and thoracic complications, in particular when AAST is equal or superior to 3. A specific attention must be paid to injuries to hepatic segments 1,4 and 9 were associated with biliary complications.

No funding was received for this work. No author has conflicts of interest to disclose related to this manuscript.

References

- [1] Letoublon C, Amariutei A, Taton N, et al. Management of blunt hepatic trauma. *J Visc Surg* 2016;153(4 Suppl):33-43.
- [2] Hommes M, Navsaria PH, Schipper IB, Krige JEJ, Kahn D, Nicol AJ. Management of blunt liver trauma in 134 severely injured patients. *Injury* 2015;46:837–42
- [3] van der Wilden GM, Velmahos GC, Emhoff T, et al. Successful nonoperative management of the most severe blunt liver injuries: a multicenter study of the research consortium of new England centers for trauma. *Arch Surg* 2012;147:423–8
- [4] Asfar S, Khoursheed M, Al-Saleh M, Alfawaz AA, Farghaly MM, Nur AM, Liver Trauma Registry Group. Management of liver trauma in Kuwait. *Med Princ Pract* 2014;23:160–6
- [5] Yoon W, Jeong Y, Kim J, Seo J. CT in Blunt Liver Trauma¹. *Radiographics* 2005;87–104
- [6] Tinkoff G, Esposito TJ, Reed J, et al. American Association for the Surgery of Trauma Organ Injury Scale I: Spleen, Liver, and Kidney, Validation Based on the National Trauma Data Bank. *J Am Coll Surg* 2008;207:646–55
- [7] Norrman G, Tingstedt B, Ekelund M, Andersson R. Non-operative management of blunt liver trauma: feasible and safe also in centres with a low trauma incidence. *HPB (Oxford)* 2009;11:50–6

- [8] Arnoux PJ, Serre T, Cheynel N, et al. Liver injuries in frontal crash situations a coupled numerical - experimental approach. *Comput Methods Biomech Biomed Engin* 2008;11:189–203
- [9] Cheynel N, Serre T, Arnoux P-J, Ortega-Deballon P, Benoit L, Brunet C. Comparison of the Biomechanical Behavior of the Liver During Frontal and Lateral Deceleration. *J Trauma Inj Infect Crit Care* 2009;67:40–4
- [10] Cheynel N, Serre T, Arnoux P-J, et al. Biomechanic Study of the Human Liver During a Frontal Deceleration. *J Trauma Inj Infect Crit Care* 2006;61:855–61
- [11] Hamada SR, Gauss T, Duchateau F-X, et al. Evaluation of the performance of French physician-staffed emergency medical service in the triage of major trauma patients. *J Trauma Acute Care Surg* 2014;76:1476–83
- [12] Voiglio EJ, Dubuisson V, Massalou D, et al. Abbreviated laparotomy or damage control laparotomy: Why, when and how to do it? *J Visc Surg* 2016;153:13–24
- [13] Soto JA, Anderson SW. Multidetector CT of Blunt Abdominal Trauma. *Radiology* 2012;265:678–93
- [14] Spahn DR, Bouillon B, Cerny V, et al. Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Crit Care* 2013;17:R76
- [15] Eddy VA, Morris JA, Cullinane DC. Hypothermia, coagulopathy , and acidosis. *Surg Clin North Am* 2000;80:845–54
- [16] Taylor FJ, Toh C, Hoots W, Wada H, Levi M. Scientific and Standardization Committee Communications Towards Definition , Clinical and Laboratory Criteria , and a Scoring System for Disseminated Intravascular Coagulation * On behalf of the

Scientific Subcommittee on Disseminated Intravascular Coagul. Thromb Haemost
2001;86 (5):1327–30

- [17] Safavi A, Beaudry P, Jamieson D, Murphy JJ. Traumatic pseudoaneurysms of the liver and spleen in children: is routine screening warranted? J Pediatr Surg 2011;46:938–41
- [18] Østerballe L, Helgstrand F, Axelsen T, Hillingsø J, Svendsen LB. Hepatic pseudoaneurysm after traumatic liver injury; is CT follow-up warranted? J Trauma Manag Outcomes 2014;8:18
- [19] Yuan K-C, Wong Y-C, Fu C-Y, Chang C-J, Kang S-C, Hsu Y-P. Screening and management of major bile leak after blunt liver trauma: a retrospective single center study. Scand J Trauma Resusc Emerg Med 2014;22:26

Figure legends

Figure 1. Initial management of the 116 with BLU admitted at the intensive care unit.

ICU: Intensive Care Unit

Table 1. Management of patients with BLI: comparison between NOM and invasive management. Invasive management group includes initial management with surgery and interventional radiology, and the patient who deceased at entrance.

		Total patients (n=116)	Invasive management (n=36)	NOM¹ (n=80)	P²
Type of trauma	- Motorbike road traffic accident	38 (33%)	14 (39%)	24 (30%)	0.343
	- Fall from heights	25 (22%)	7 (19%)	18 (23%)	0.711
	- Car accident	25 (22%)	9 (25%)	16 (20%)	0.545
	- Equestrian accident	3 (4%)	0	3 (4%)	0.586
	- Fall	13 (11%)	3 (8%)	10 (13%)	0.510
	- Other	12 (10%)			
Mechanism of trauma	- Direct shock	76 (66%)	28 (78%)	48 (60%)	0.062
	- Acceleration/Deceleration	31 (27%)	7 (19%)	24 (30%)	0.234
	- Crushing	6 (5%)	1 (3%)	5 (6%)	0.435
	- Other	3 (3%)			
State at initial presentation	- Hemodynamic instability	35 (30%)	20 (56%)	15 (19%)	<0.001
	- RBC ³ transfusions (Yes)	30 (26%)	21 (58%)	9(11%)	<0.001
	- Number of RBC packs	4.8 ± 4.1	5.5 ± 4.7	3.5 ± 1.7	0.448
	- Fluid expansion	48 (41%)	24 (67%)	24 (30%)	<0.001
	- Norepinephrine	41 (35%)	20 (56%)	21 (26%)	0.002
	- Acidosis	46 (40%)	22 (61%)	24 (30%)	0.002
	- Hypothermia	29 (25%)	14 (39%)	15 (19%)	0.021
	- Coagulopathy	25 (22%)	14 (39%)	11 (14%)	0.002

	- Hemorrhagic shock	24 (21%)	17 (47%)	7 (9%)	<0.001
	- ISS ⁴	32.1 ± 16.8	36.6 ± 20.0	31.4 ± 14.9	0.337
	- SAPS ⁵	34.8 ± 20.4	41.0 ± 23.0	31.6 ± 18.2	0.043
CT-scan lesions	- Active bleeding	33 (28%)	18 (50%)	15 (19%)	<0.001
	- AAST ⁶ classification	3.1 ± 1.1	3.5 ± 1.2	2.9 ± 1.0	0.001
Subsequent	- Surgery	11 (9%)	8 (22%)	3 (4%)	0.002
management	- Interventional radiology	9 (8%)	5 (14%)	4 (5%)	0.100
	- Surgery <i>OR</i> Interventional radiology	20 (17%)	13 (36%)	7 (9%)	<0.001
Intra hospital death		9 (8%)	6 (17%)	3 (4%)	0.020

1: NOM: Non Operative Management, 2: Comparison between invasive group and NOM group, 3: RBC: Red Blood Cells, 4: Injury Severity Score, 5: Simplified Acute Physiology Score, 6: American Association for the Surgery of Trauma, Quantitative data are expressed as mean ± standard deviation

Table 2. Features of liver trauma and associated lesions on initial CT-scan (n=116)

Type of lesions	Localization / Size	N	%
Hematoma		54	46
	- Sub-capsular	27	23
	- Intraparenchymal	27	23
	- < 1 cm	4	7
	- 1 to 3 cm	11	20
	- > 3 cm	39	72
Active arterial bleeding		33	28
Major hepatic venous injury		0	0
Parenchymal liver injury	- Lacerations	102	88
	- < 1 cm	0	0
	- 1 to 3 cm	18	17
	- > 3 cm	89	76
	- Contusions	5	4
	- Segments 1/4/9	46	43
	- Right liver	94	88
	- Left lobe	15	14
	- Along falciform ligament	2	2
Area nuda injury		44	38
Vascular injury		66	57
	- Right portal fissure/Right hepatic vein/Right portal vein/Right hepatic artery	51	77

	- Main portal fissure/Median hepatic vein/Segments 1/4	11	17
	- Left portal vein/Left hepatic vein/Left hepatic artery	9	14
Right adrenal gland injury		43	37
Hepatic hilum injury		26	22
Biliary injury		7	6
	- Including gallbladder	3	3
Pseudo aneurysm		6	5
Periportal infiltration		30	26
Flat inferior vena cava		6	5
Hemoperitoneum		33	28
AAST grade	- 1	10	9
	- 2	22	19
	- 3	50	43
	- 4	19	16
	- 5	15	13
Pneumoperitoneum		1	1
Other associated thoraco-		94	81
abdominal lesions	- Spleen	12	13
	- Hollow organ	3	3
	- Thorax (hemothorax/pneumothorax/rib fractures)	81	86

Table 3. Complications during hospitalization

Complications		N	%
Total		55	47
Hemorrhagic complications	- Bleeding	5	4
	- Pseudo aneurysm	5	4
	- Red blood cells transfusions after initial management	12	10
	- Requirement for interventional radiology	5	4
Biliary complications		7	6
	- Biloma	3	3
	- Biliary peritonitis	4	3
	- Hemobilia	0	0
Intrahepatic abscess/collection		2	2
Hepatic necrosis		1	1
Acute coronary syndrome		2	2
Pulmonary complications	- Pleural effusion requiring drainage	16	14
	- Pneumopathy	17	15
	- Pulmonary embolism	5	4

Table 4. Factors associated with total biliary complications (univariate analysis)

		Biliary complications (N=13)	No biliary complications (N=103)	P
Initial Assessment	Hemodynamic instability	5 (38%)	30 (29%)	0.528
	SAPS ¹	29.5 ± 14.6	35.5 ± 21.0	0.476
Mechanism of trauma	Direct shock	10 (77%)	66 (64%)	0.538
	Acceleration/Deceleration	3 (23%)	28 (27%)	>0.999
BLI ²	AAST score	3.4 ± 1.0	3.0 ± 1.1	0.295
	Laceration>3cm	11 (85%)	78 (76%)	0.730
Location of liver injury	Segments 1/4/9	9 (69%)	37 (36%)	0.033
	Right portal scissure	6 (46%)	45 (44%)	>0.999
	Main portal scissure	2 (15%)	9 (9%)	0.356
	Right liver	12 (92%)	82 (80%)	0.528
Percutaneous TAE ³		4 (31%)	28 (27%)	0.751

1: SAPS: Simplified Acute physiology Score; 2: BLI: Blunt Liver Injury; 3: TAE: Trans Arterial Embolization, during initial management or subsequent in-hospital stay