Incorporation and stability of carotenoids in a functional fermented maize yogurt-like product containing phytosterols

Magali Gies, Adriana Maria Descalzo, Adrien Servent, Claudie Dhuique-Mayer

► To cite this version:

Magali Gies, Adriana Maria Descalzo, Adrien Servent, Claudie Dhuique-Mayer. Incorporation and stability of carotenoids in a functional fermented maize yogurt-like product containing phytosterols. LWT - Food Science and Technology, 2019, 111, pp.105 - 110. 10.1016/j.lwt.2019.04.103 . hal-03486755

HAL Id: hal-03486755

<https://hal.science/hal-03486755>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Incorporation and stability of carotenoids in a functional fermented maize yogurt-like product containing phytosterols

Magali Gies^{a,b}, Adriana Maria Descalzo^c, Adrien Servent^{a,b}, Claudie Dhuique-Mayer^{a,b}

^a Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UMR Qualisud, 34398 Montpellier Cedex 5, France

^b Qualisud, Univ. Montpellier, CIRAD, Montpellier SupAgro, Université d'Avignon, Université de La Réunion, Montpellier, France

^c Instituto Nacional de Tecnología Agropecuaria INTA-LABINTEX, Center of Agroindustry, de la tradición & Aristizabal Str. s/n
1686 Hurlingham, Buenos Aires, Argentina

Corresponding author: claudie.dhuique-mayer@cirad.fr

Declarations of interest: none

Abstract

A functional yogurt-like product based on fermented maize enriched in carotenoids and phytosterols was developed to be an alternative to existing functional dairy products. Incorporation and stability of fat-soluble compounds were challenging in this complex matrix. Moreover, it is known that phytosterols decrease carotenoids absorption, but this decline can be offset by increasing carotenoids in foodstuffs. Thus, the aims of this work were to optimize the manufacturing process of a fermented yogurt-like product containing phytosterols by incorporating carotenoids taking into account two *Lactobacilli* strains as starters. The stability of β -cryptoxanthin, β -carotene and lycopene was studied during the whole process in order to confer a nutritional balance between them and phytosterols. After 15h of fermentation, a starter ratio 50-50 (*L. plantarum*-*L. casei*) gave the best final biomass with 10^9 CFU/g necessary to obtain a probiotic potential. Relative carotenoid losses (20 - 27 %) occurred during the pasteurization step while no negative impact on carotenoids was due to fermentation. Fat-soluble compounds remained relatively stable during the whole process with 75 % of retention. These results suggest that incorporation and stability of carotenoids in a fermented-maize yogurt-like product containing phytosterols are necessary steps to induce a cholesterol-lowering effect without detrimental effects on carotenoids.

Keywords

Fermented-maize; Probiotic cereal product; Fat-soluble phytonutrients; β -carotene; β -sitosterol

Introduction

The increasing demand of consumers for healthy food has reinforced the necessity to design new functional foods. A functional food containing bioactive compounds can enhance health or reduce the risk of diseases (Prado, Parada, Pandey, & Soccol, 2008; I. Salmerón, 2017; Serafini, Stanzione, & Foddai, 2012). Among functional foods, dairy products with incorporation of fat-soluble compounds (e.g. phytosterols) and fatty acid omega 3, or addition of probiotic starters represent 60 to 70 % of the functional foods market (Ivan Salmerón, Thomas, & Pandiella, 2015; Chen, McClements, & Decker, 2013). However, lactose intolerance or protein allergies and cholesterol content are factors that have recently increased the demand for non-dairy products (Prado et al., 2008). Cereal-based products are a general response to consumers for health benefits and as an alternative to dairy products (Marsh, Hill, Ross, & Cotter, 2014) by being both probiotic (living microorganisms 10^8 to 10^9 cells per gram of consumed product) and containing fibers, minerals and vitamins (Kandyliis, Pissaridi, Bekatorou, Kanellaki, & Koutinas, 2016).

However, there is a lack of available data in the literature about cereal-based fermented foods with incorporation of phytosterols and/or carotenoids. Chen et al. (2013) described the importance of added carotenoids in functional foods for their positive impact on health problems. The authors also reported the challenges associated with the incorporation of these fat-soluble compounds, because of their poor water solubility, their high sensitivity to oxidation and their structural and physico-chemical properties. It is important that these dietary molecules remain stable during the whole process, transport and storage, in order to maintain their potential bioactivity and a high bioavailability during digestion to be efficient on health.

Among carotenoids, β -carotene and β -cryptoxanthin are pro-vitamin A, known to help the organism to maintain normal eye health, epithelial function, embryonic development and immune system function (Stephensen, 2001). Beside these carotenoids, lycopene, which is a non-provitamin A carotenoid, has some other health benefits, such as decreasing the development of prostate, cervical, colon, rectal, stomach and other types of cancers (Giovannucci, 1999). This carotenoid is also known for its great antioxidant activity (Fiedor & Burda, 2014). Moreover, supplementing functional food containing phytosterols with carotenoids seems necessary because it is suggested that phytosterols reduce the incorporation of carotenoids into mixed micelles during the first steps of digestion (Baumgartner, Ras, Trautwein, Mensink, & Plat, 2017). Furthermore, the competition between phytosterols and β -carotene (which is the most important pro-vitamin A micronutrient) during intestinal absorption has been reported by Fardet, Morise, Kalonji, Margaritis, & Mariotti (2015).

From another side, carotenoid stability must be performed during food processing such as pasteurization and fermentation. The degradation of carotenoids and especially β -carotene during food processing is mainly due to oxidation and thermal treatments knowing that β -carotene oxidation generates losses of color and pro-vitamin A activity (Pénicaud, Achir, Dhuique-Mayer, Dornier, & Bohuon, 2011). Considering fermentation process, some strains such as *Lactobacillus plantarum* can enhance the nutritional value of fermented food by increasing the nutrient density (mostly due to their consumption of sugar), hydrolyzing polymers from the raw material, biosynthesizing bioactive peptides, degrading toxic or anti-nutritional factors and by synthesizing promoters for absorption (Septembre-Malaterre, Remize, & Poucheret, 2018).

Recently, Descalzo et al. (2018) have developed a traditional fermented maize yogurt-like product derived from an African preparation, with added phytosterols. The authors suggested

adding carotenoids from natural fruits to increase the bioactive potential of the yogurt-like product. They also proposed to turn the product probiotic with pertinent strains and to use dispersible phytosterols instead of esterified phytosterols emulsion because of the lipid oxidation due to the presence of unsaturated fatty acids in the esterified formulation. Finally in this work, the manufacturing process should have been standardized in fermenters by monitoring temperature and pH parameters during pasteurization and fermentation steps as it was advised by Marsh et al. (2014).

Consequently, this paper focuses on the standardization of a fermented functional product by incorporating carotenoids such as β -cryptoxanthin, β -carotene and lycopene in a fermented-maize yogurt-like product containing dispersible phytosterols. Specifically our aims were (1) to optimize the manufacturing process of this product containing phytosterols by incorporating natural carotenoids from papaya/melon fruit extracts before pasteurization step; (2) to improve the fermentation of the product with a selection of Lactobacilli starters in order to obtain a probiotic potential; (3) to study the behavior of these fat-soluble compounds during the whole process in order to determine which step has the main impact on their stability. In fine, the aim was obtaining a balanced product with carotenoids and phytosterols giving an interesting nutritional value and a cholesterol-lowering potential to this maize fermented yogurt-like product.

1. Materials and methods

1.1. Bacterial strains and inocula preparation

Two pure lyophilized strains *Lactobacillus plantarum* (Lp352-S1616 / CNCM I-3069) and *Lactobacillus casei* (Lc1 S42-2015 / CNCM I-4592) were provided as starters by Ennolys-

Lesaffre (Marq-en-Baroeul, France) and stored under vacuum pack at + 4°C. Individual strains and a 50-50 % ratio (*L. plantarum* - *L. casei*) were inoculated at $1 \cdot 10^6$ CFU/g in the maize-based matrix.

1.2. Manufacture of the maize-based matrix

The manufacturing process including maize soaking, crushing, sieving, formulation, pasteurization and lactic fermentation was described in Figure 1. 100 g of maize grains from a French local market were soak and then crushed at 5000 rpm for 1 min (Retsch Grindomix GM 200, Germany). Ingredients such as 3.6 % of sugar (Daddy, France) and 2.4 % of semi-skimmed powdered milk (Régilait, France) were added in the maize juice among others. Commercial dispersible phytosterols (Vitaesterol® S-80 WDP 90 % non GMO with 80 % of β -sitosterol, Vitae Naturals, Spain) were added. Freeze-dried papaya (*Carica papaya* var. Formosa, Brazil) and melon (*Cucumis melo* var. Cantalupensis, France) were incorporated as sources of carotenoids (β -cryptoxanthin - lycopene and β -carotene respectively). Pasteurization and fermentation were performed in a six-hundred-milliliter glass double wall fermenter (Legallais, France) as described by de J. C. Munanga, Loiseau, Grabulos, & Mestres (2016). The intern temperature and pH were determined per minute by food penetration probes connected with a central data acquisition (Almemo® 2690-8A, Ahlborn, Germany). After pasteurization, the lactic fermentation started with *Lactobacilli* strains added either pure or mixed (50-50 ratio) at a concentration of $1 \cdot 10^6$ CFU/g. Fermentations were conducted at 37 ± 0.2 °C for 15 h. The products were kept frozen and in darkness at -20 °C until analyses.

1.3. Microbiological analysis

1.3.1. Determination of growth kinetics

The number of *Lactobacilli* was estimated each hour during fermentation by plating on MRS agar medium (Biokar Diagnostics, France – Ref BK089HA) at 37 ± 0.2 °C for 48 h. Growth kinetics and pH values were then associated to each bacteria or ratio. Growth parameters were calculated: specific growth rate ($\mu_{\max}$) and generation time corresponding to $G = \ln(2)/\mu_{\max}$. Growth kinetics were determined and modelled in agreement with the Rosso equation:

$$\log N_{(t)} = \begin{cases} \log N_0 & \text{si } t < \lambda \\ \log N_{\max} - \log \left(\left[1 + \left(\frac{N_{\max}}{N_0} - 1 \right) \exp(-\mu_{\max}(t - \lambda)) \right] \right) & \text{si } t \geq \lambda \end{cases}$$

(1)

where $\mu_{\max}$ is the specific growth rate (h^{-1}), λ the latency period (h), N_0 the initial population (CFU/g), $N_{\max}$ the maximal population (CFU/g) and log the decimal logarithm.

1.3.2. Determination of the final bacterial ratio

In order to determine the final ratio of 50-50 ratio after 15h of fermentation the strains were differentiated by their sugar consumption with an API 50 CHL gallery (Analytical Profile Index - Biomérieux, France). Then, a 50 CHL agar medium commercialized (Biomérieux, France) was used with 16 g/L added agar-agar. A control was made without any sugar, although two others were made with 20 g/L added glucose (Glc medium) and 20 g/L added arabinose (Ara medium). Just at the end of the fermentation, 1 g of the product was sampled and diluted from 10^{-1} to 10^{-8} . 100 μL of the dilutions 10^{-6} , 10^{-7} and 10^{-8} were sowed on the control medium, the Glc and the Ara mediums and incubated during 48 h at 37 ± 0.2 °C.

169 **1.3.3. Determination of the microbiological efficiency of**
170 ***pasteurization: sterility of the product***

171 Just after pasteurization, 1 g of the product was sampled and diluted from 10^{-1} to 10^{-4} . 100 μ L
172 of each dilution were sowed on Plate Count Agar (PCA) medium in triplicates and incubated
173 during 72 h at 30 ± 0.2 °C according to the ISO 4833-2:2013 Afnor norm.

174 **1.4. Proximate analysis**

175 The food composition of the product was analyzed in terms of proteins, lipids and
176 carbohydrates contents. Proteins were measured following Kjeldahl method (AOAC- 2001),
177 (N X 6.25) using a Foss Techator Digester and an automatic Foss analytical AB Kjeltect™
178 8400 apparatus (Foss, Sweden). Lipid content was obtained with the Folch extraction
179 procedure (Schäfer, 1998). Starch was dosed according to Clegg (1956) with anthrone reagent
180 method. Analysis and separation of soluble sugars were determined using an UPLC – 1290
181 System Infinity II (Agilent, USA) equipped with a refractometer detector. A SHODEX
182 SH1011 column 300x8 mm (Tokyo, Japan) was used with an isocratic system of water with
183 H₂SO₄ (0.01 %) and a flow rate of 0.7 mL/min. Temperature was set at 30 °C, injection
184 volume at 10 μ L and spectrophotometric detection at 210 and 245 nm. External calibration of
185 glucose, fructose, lactose and sucrose were realized using standards from Sigma-Aldrich
186 (France). Dry matters were obtained in a vacuum oven at 70 °C during 24 h according AOAC
187 method 1991.

188 **1.5. Phytomicronutrients analysis**

189 Fat-soluble compounds such as carotenoids (β -cryptoxanthin, β -carotene and lycopene),
190 phytosterols (β -sitosterol) and tocopherols (α -tocopherol and γ -tocopherol) were analyzed by
191 UPLC-DAD. The fat-soluble extraction was adapted from Rossetti et al. (2010). Briefly, 1 g

of product was saponified with 1.5 ml of 12N KOH for 30 min at 70 °C and extracted twice with 5 mL of n-hexane. Hexanic phases were evaporated under nitrogen and dissolved in 1 mL of a MTBE/methanol solution (4:1, v:v) before injection in UPLC system. An UPLC – 1290 System Infinity II (Agilent, USA), with a diode array detector (DAD) and a fluorescence detector (FLD) was used. The column was a C30 YMC (150 x 4.6 mm; 3 µm) (YMC Europe GmbH, Germany). Mobile phases were methanol as eluant A, water as eluant B and MTBE as eluant C, set at 1.5 mL/min flow rate. The gradient used to separate carotenoids, phytosterols and tocopherols was the following: 0-1.5 min [60 % A, 40 % B]; 1.5-3 min [80 % A, 20 % B]; 3-12.5 min [80 % A, 5 % B, 15 % C]; 12.5-15 min [15 % A, 85 % C]; 15-17 min [100 % A] and back to the initial conditions for re-equilibration. The column temperature was 20 °C and the injection volume was 10 µL. Detection was set at 210 nm (DAD) for phytosterols, 450 and 470 nm (DAD) for carotenoids. Fluorescence detection (FLD) for tocopherols was set at 296 nm (excitation) and 330 nm (emission). Quantification was achieved using calibration curve with β-carotene, β-cryptoxanthin, lycopene (Extrasynthese, France), α/γ-tocopherols standards (Sigma S^t Louis, USA) and β-sitosterol standard (Supelco, Bellefonte, USA).

Statistical analyses

All statistical analyses were performed using XLSTAT software version 19.6 (Addinsoft, France). All data were reported as means ± standard deviation (SD) from three replicates of each experiment. Data were analyzed statistically using one-way analysis of variance (ANOVA) in order to determine significant differences ($p < 0.05$). Tukey's multiple comparison method was used to further examine any significant difference between results.

2. Results and discussion

2.1. Standardisation of the manufacturing process with incorporation of carotenoids

According to the manufacturing process (Figure 1), the standardization in fermenters was conducted to monitor pasteurization conditions and microbial parameters during fermentation. This optimization of batch fermentation allowed a better control of parameters such as temperature and pH, thus predicting kinetics bacterial growth and ensuring product safety. A model of fermentation of a traditional Beninese beverage called gowé using this batch fermentation was described by de J. C. Munanga et al. (2016). The incorporation of lyophilized fruits enhanced the homogeneity of the product just as dispersible phytosterols. Moreover, this form of fruit concentrate was recommended as functional food ingredients because its richness in antioxidant bioactive compounds and dietary fibers (Rocha, Fávaro-Trindade, & Grosso, 2012). The addition of bioactive compounds like carotenoids led to a fortification in fat-soluble pro-vitamin A as well as antioxidants. This carotenoids supplementation induced a functional equilibrium with a balanced ratio between carotenoids and phytosterols, which should have no detrimental impact on carotenoids absorption.

In order to obtain a probiotic functional product, the cooking step was set before fermentation, acting as pasteurization in our present study. In that case, after the pasteurization, the sterility test on PCA proved that the product was microbiologically stable and safe with only $3.33.10^1$ CFU/g which entered in the Afnor norm.

In regard to the nutritional composition of the functional product, the macronutrient contents were reported in Table 1. The product is balanced in proteins, lipids and sugars. It contained

twice more proteins and 30 % less sugars than a dairy yogurt containing fruits, regarding the table of the USDA (2018). Dry matter was 21.3 ± 0.01 % and the calorific value 121.64 ± 6.08 kcal/100 g.

2.2. Selection of starters and microbiological optimization of the functional product

In order to select the starters to initiate the fermentation, the kinetic growth, the latency period, the final pH and the final biomass were compared between different strains *L. casei* versus *L. plantarum* versus their ratio 50-50. Therefore, these strains were used either pure or mixed in a ratio 50-50.

Data were modelled in agreement with Rosso model (1) on Figure 2. Strains fermented into the maize-based matrix until reaching the stationary phase around 15 h, by increasing three times their biomass from $1 \cdot 10^5 - 1 \cdot 10^6$ CFU/g to $1 \cdot 10^9 - 1 \cdot 10^{9.5}$ CFU/g. Thus, the final product contained enough living *Lactobacilli* at the end of the process to be potentially probiotic, that is to say more than 10^9 UFC/g (Prado et al., 2008). All the final products had a pH between 4 and 4.8; it is the known pH for classic yogurt (FAO, 1995) and could guarantee its shelf life by insuring a low contamination rate as well as its microbiological stability. Moreover, the latency period of the 50-50 ratio is only 3 h (against 4 h and 5 h for *L. plantarum* and *L. casei* respectively) with a specific growth rate reaching 0.94 h^{-1} (Table 2). Thus, this ratio ensures a brief fermentation start with a swift exponential growth. Moreover, multistrain or multispecies probiotic beverages may provide greater beneficial effects than monostrain cultures (Marsh et al., 2014). *L. plantarum* was chosen for its ubiquity: it is found in the environment, especially on plants and therefore on maize. *L. casei* species are known to be used in a lot of probiotic dairy products. Consequently, the 50-50 ratio is the best compromise to ferment the maize-based product, regarding to the microbiological results.

In order to determine the real final ratio of strains, the two *Lactobacilli* strains have been counted individually at the end of the fermentation. The API 50 CHL gallery showed that the final ratio was around 45 % of *L. plantarum* and 55 % of *L. casei* which is highly correlated with the kinetic growth. Regarding those results, there is no competition for nutrients between these two strains: they kept their initial ratio and their own probiotic potential.

2.3. Carotenoids content and other fat-soluble phytonutrients in the functional yogurt-like product

Carotenoids, tocopherols and β -sitosterol were quantified in the functional yogurt-like product and reported in Table 1. Three main carotenoids were found in the final product: β -cryptoxanthin and lycopene related to papaya lyophilized extract and β -carotene from melon lyophilized extract. The major carotenoids were pro-vitamin A β -carotene and the well-known antioxidant carotenoid called lycopene. The total carotenoids content of the product was 16 mg/kg representing 2.01 ± 0.14 mg per serving portion of 125 g. This high content of carotenoids also produced the final orange color of this food. Using the retinol activity equivalent (RAE) as an estimation of vitamin A in the product, 125 g would bring 12 % of the Recommended Daily Allowance (RDA) for an adult. Moreover, approximately 2 mg of carotenoids per portion is enough to counteract the competition for micellization of carotenoids with phytosterols during digestion. Consequently, it is recommended to increase carotenoids intake during phytosterols consumption (Fardet et al., 2015; Noakes et al., 2002). Tocopherols originated from fruit extracts bring 4 % of RDA vitamin E. These fat-soluble compounds are antioxidants *in vivo* for metabolism functions and also efficient for the protection of other phytonutrients against oxidative damage. α -tocopherols represented 0.34 ± 0.04 mg/kg and γ -tocopherols 3.47 ± 0.30 mg/kg. In fine, one portion of 125 g of the

yogurt-like product contained 2.5 g of free phytosterols (mainly β -sitosterol). This content is enough to obtain a cholesterol-lowering effect. Indeed, according to the report of ANSES (2014), an ingestion of 2 g of phytosterols per day is the approximate effective dose essential to reduce the Low-Density Lipoprotein cholesterol (LDL-cholesterol) concentration in plasma.

2.4. Stability of fat-soluble phytonutrients during the manufacturing process

Carotenoids and tocopherols stability were studied during pasteurization and fermentation process. To evaluate the effect of lactic fermentation on carotenoid content, the yogurt-like product was first fermented with *L. plantarum* or *L. casei* during 10, 15 and 20 h. After 20 h of fermentation with *L. casei*, lycopene significantly ($p < 0.05$) increased from 5.1 to 7.7 mg/kg (+ 33 %), β -carotene from 6.9 to 7.4 mg/kg (+ 7 %) and tocopherols from 5.1 to 7.2 mg/kg (+ 30 %). No changes were observed for tocopherols when the product was fermented with *L. plantarum* and a smaller increase occurred for β -carotene (from 8 to 9.2 mg/kg; + 13 %) but no significant increase for lycopene. Together, these results supported that fermentation allowed the recovery of carotenoids and tocopherols to a significant extent particularly when the product was fermented with the starter *L. casei*. It was probably due to a production of enzymes like lipases or proteinases by strains, which allowed the liberation of carotenoids from complexes, then permitted a better extractability of these compounds. Moreover, it is also possible that these Lactobacilli strains demonstrated a carotenogenesis, when they are not in co-culture, as it is described by Kot, Błażej, Gientka, Kieliszek, & Bryś (2018) and Kot, Błażej, Kurcz, Gientka, & Kieliszek (2016). The increase of nutritional value and the changes in bioactive compound contents over lactic fermentation were reported by Katina et al. (2007) and recently by Septembre-Malaterre et al. (2018). The

310 proteolytic activity of lactic acid bacteria culture could result in a better recovery of
311 carotenoids. By disrupting the protein-carotenoid complexes in vegetables, carotenoid
312 extraction was improved (Bhaskar, Suresh, Sakhare, & Sachindra, 2007). All these results
313 represented another argument to support the choice of the fermentation with the 50-50 ratio of
314 strains.

315 Considering this ratio of starters, the fat-soluble phyto-micronutrient contents were then
316 analyzed at three different steps of the manufacturing process: raw product (neither
317 pasteurized nor fermented), pasteurized product and final product pasteurized and fermented
318 for 15 h. The Figure 3 describes the evolution of bioactive compounds in the fermented
319 product obtained with the 50-50 ratio. While carotenoids were slightly impacted during
320 pasteurization, tocopherols and phytosterols remained stable.

321 There were significant ($p < 0.05$) losses in carotenoids after pasteurization but no significant
322 difference between the pasteurized product before and after 15 h of fermentation. Indeed, the
323 averages of losses after pasteurization were $21.4 \pm 6.0 \%$, $26.9 \pm 6.0 \%$ and $20.2 \pm 7.0 \%$ for
324 β -cryptoxanthin, β -carotene and lycopene respectively; while the losses after 15 h of
325 fermentation was $23.2 \pm 1.0 \%$, $28.1 \pm 1.0 \%$ and $19.2 \pm 3.0 \%$. Therefore, in this case, β -
326 carotene was more sensitive to the pasteurization step than β -cryptoxanthin and lycopene. The
327 lower thermal degradation of lycopene, compared to other carotenoids, was generally
328 observed in vegetable matrix (tomato, citrus juice) because lycopene is bounded with
329 proteins, giving it a better structural protection (Achir, Hadjal, Madani, Dornier, & Dhuique-
330 Mayer, 2015). Moreover, the lycopene degradation was also reported to be lower in
331 lyophilized fruit form than in model system (Rocha et al., 2012).

332 It means that there is a “pasteurization effect” which represents the whole “process impact”
333 on the bioactive compounds, because there is no “fermentation impact”. Thus, the
334 fermentation of lactic strains did not impact the stability of the compounds, probably because

these bacteria released bioactive peptides improving the antioxidant capacity of the product (Septembre-Malaterre et al., 2018). Even with these relative losses of carotenoids after pasteurization (20 to 27 %), all fat-soluble compounds remained relatively stable during the whole process with a high level of retention (between 73 % and 100 %). Note that *cis*-isomerisation of β -carotene represents 6.5 % of total β -carotene and this low percentage of isomerization did not really affect the concentration of β -carotene during the process. In the final product, the *cis*-isomerization of the lycopene reaches 10 %, with the half formed during pasteurization. Similar results on carotenoids degradation in pumpkin puree was observed by Provesi, Dias, & Amante (2011) with a carotenoid retention > 75 %. According to Pinheiro Sant'Ana, Stringheta, Cardoso Brandão, & Cordeiro de Azeredo (1998), a water pasteurization without pressure, exactly what was performed in this new formulation, is the best way to keep the most of carotenoids, between 56 and 89 %.

Conclusion

The standardization of this yogurt-like product, presented here as a generic functional fermented food, demonstrated that incorporation and stabilization of fat-soluble phytonutrients during a whole process is possible in a probiotic cereal-based product. Moreover, the nutritional balance between phytosterols and carotenoids was respected, to provide potential health effects such as a reduction of the absorption of cholesterol during digestion and a non-negligible intake of pro-vitamin A. Indeed, only the pasteurization step had a relative impact on carotenoids. The fermentation with the 50-50 ratio of *L. plantarum* and *L. casei* had no significant impact on the content of fat-soluble compounds. In the contrary, it seems that pure strains could demonstrate carotenogenesis. Finally, the 50-50 ratio presented the best growth parameters into this matrix and it is known that a co-culture is always better to enhance a probiotic potential. Further researches are needed in order to

optimize this product on nutritional value. It is essential to know better this new cereal fermented yogurt-like product in terms of nutritional and sensory qualities. In that purpose, carotenoid bioaccessibility in this matrix has to be assessed in the future, just as sensory analyses.

Acknowledgments

We particularly would like to thank Ennolys-Lesaffre for providing the *L. plantarum* (CNCM I-3069) and the *L. casei* (CNCM I-4592). All our gratitude to Sylvie Lortal (Institut National de la Recherche Agronomique (INRA), UMR Science et Technologie du Lait et de l'Œuf (STLO), 35000 Rennes, France) for her priceless help for this work.

References

- Achir, N., Hadjal, T., Madani, K., Dornier, M., & Dhuique-Mayer, C. (2015). Carotene Reactivity in Pink Grapefruit Juice Elucidated from Model Systems and Multiresponse Modeling. *Journal of Agricultural and Food Chemistry*, 63(15), 3970–3979.
<https://doi.org/10.1021/acs.jafc.5b00509>
- ANSES. (2014). *OPINION of the French Agency for Food, Environmental and Occupational Health & Safety on Assessment of the risks and benefits of consuming food products fortified with phytosterols and phytostanols* (No. 2010-SA-0057; p. 15). Maisons-Alfort: ANSES.
- Baumgartner, S., Ras, R. T., Trautwein, E. A., Mensink, R. P., & Plat, J. (2017). Plasma fat-soluble vitamin and carotenoid concentrations after plant sterol and plant stanol

consumption: a meta-analysis of randomized controlled trials. *European Journal of Nutrition*, 56(3), 909–923. <https://doi.org/10.1007/s00394-016-1289-7>

Bhaskar, N., Suresh, P. V., Sakhare, P. Z., & Sachindra, N. M. (2007). Shrimp biowaste fermentation with *Pediococcus acidolactici* CFR2182: Optimization of fermentation conditions by response surface methodology and effect of optimized conditions on deproteinization/demineralization and carotenoid recovery. *Enzyme and Microbial Technology*, 40(5), 1427–1434. <https://doi.org/10.1016/j.enzmtec.2006.10.019>

Chen, B., McClements, D. J., & Decker, E. A. (2013). Design of Foods with Bioactive Lipids for Improved Health. *Annual Review of Food Science and Technology*, 4(1), 35–56. <https://doi.org/10.1146/annurev-food-032112-135808>

Clegg, K. M. (1956). The application of the anthrone reagent to the estimation of starch in cereals. *Journal of the Science of Food and Agriculture*, 7(1), 40–44. <https://doi.org/10.1002/jsfa.2740070108>

de J. C. Munanga, B., Loiseau, G., Grabulos, J., & Mestres, C. (2016). Modeling Lactic Fermentation of Gowé Using *Lactobacillus* Starter Culture. *Microorganisms*, 4(4), 44. <https://doi.org/10.3390/microorganisms4040044>

Descalzo, A. M., Rizzo, S. A., Servent, A., Rossetti, L., Lebrun, M., Pérez, C. D., ... Dhuique-Mayer, C. (2018). Oxidative status of a yogurt-like fermented maize product containing phytosterols. *Journal of Food Science and Technology*, 55(5), 1859–1869. <https://doi.org/10.1007/s13197-018-3102-5>

FAO. (1995). *Le lait et les produits laitiers dans la nutrition humaine*. In *Collection FAO: Alimentation et nutrition: Vol. 28*. Retrieved from <http://www.fao.org/docrep/t4280f/T4280F00.htm#Contents>

Fardet, A., Morise, A., Kalonji, E., Margaritis, I., & Mariotti, F. (2015). Influence of Phytosterol and Phytostanol Food Supplementation on Plasma Liposoluble Vitamins

and Provitamin A Carotenoid Levels in Humans: An Updated Review of the Evidence.
Critical Reviews in Food Science and Nutrition, 00–00.

<https://doi.org/10.1080/10408398.2015.1033611>

Fiedor, J., & Burda, K. (2014). Potential Role of Carotenoids as Antioxidants in Human
Health and Disease. *Nutrients*, 6(2), 466–488. <https://doi.org/10.3390/nu6020466>

Giovannucci, E. (1999). Tomatoes, Tomato-Based Products, Lycopene, and Cancer: Review
of the Epidemiologic Literature. *JNCI Journal of the National Cancer Institute*, 91(4),
317–331. <https://doi.org/10.1093/jnci/91.4.317>

Kandylis, P., Pissaridi, K., Bekatorou, A., Kanellaki, M., & Koutinas, A. A. (2016). Dairy and
non-dairy probiotic beverages. *Current Opinion in Food Science*, 7, 58–63.
<https://doi.org/10.1016/j.cofs.2015.11.012>

Katina, K., Liukkonen, K.-H., Kaukovirta-Norja, A., Adlercreutz, H., Heinonen, S.-M.,
Lampi, A.-M., ... Poutanen, K. (2007). Fermentation-induced changes in the
nutritional value of native or germinated rye. *Journal of Cereal Science*, 46(3), 348–
355. <https://doi.org/10.1016/j.jcs.2007.07.006>

Kot, A. M., Błażej, S., Gientka, I., Kieliszek, M., & Bryś, J. (2018). Torulene and
torularhodin: “new” fungal carotenoids for industry? *Microbial Cell Factories*, 17(1).
<https://doi.org/10.1186/s12934-018-0893-z>

Kot, A. M., Błażej, S., Kurcz, A., Gientka, I., & Kieliszek, M. (2016). Rhodotorula
glutinis—potential source of lipids, carotenoids, and enzymes for use in industries.
Applied Microbiology and Biotechnology, 100(14), 6103–6117.
<https://doi.org/10.1007/s00253-016-7611-8>

Marsh, A. J., Hill, C., Ross, R. P., & Cotter, P. D. (2014). Fermented beverages with health-
promoting potential: Past and future perspectives. *Trends in Food Science &
Technology*, 38(2), 113–124. <https://doi.org/10.1016/j.tifs.2014.05.002>

433 Noakes, M., Clifton, P., Ntanos, F., Shrapnel, W., Record, I., & McInerney, J. (2002). An
 434 increase in dietary carotenoids when consuming plant sterols or stanols is effective in
 435 maintaining plasma carotenoid concentrations. *The American Journal of Clinical*
 436 *Nutrition*, 75(1), 79–86. <https://doi.org/10.1093/ajcn/75.1.79>

437 Pénicaud, C., Achir, N., Dhuique-Mayer, C., Dornier, M., & Bohuon, P. (2011). Degradation
 438 of β -carotene during fruit and vegetable processing or storage: reaction mechanisms
 439 and kinetic aspects: a review. *Fruits*, 66(6), 417–440.
 440 <https://doi.org/10.1051/fruits/2011058>

441 Pinheiro Sant’Ana, H. M., Stringheta, P. C., Cardoso Brandão, S. C., & Cordeiro de Azeredo,
 442 R. M. (1998). Carotenoid retention and vitamin A value in carrot (*Daucus carota* L.)
 443 prepared by food service. *Food Chemistry*, 61(1–2), 145–151.
 444 [https://doi.org/10.1016/S0308-8146\(97\)00084-8](https://doi.org/10.1016/S0308-8146(97)00084-8)

445 Prado, F. C., Parada, J. L., Pandey, A., & Soccol, C. R. (2008). Trends in non-dairy probiotic
 446 beverages. *Food Research International*, 41(2), 111–123.
 447 <https://doi.org/10.1016/j.foodres.2007.10.010>

448 Provesi, J. G., Dias, C. O., & Amante, E. R. (2011). Changes in carotenoids during processing
 449 and storage of pumpkin puree. *Food Chemistry*, 128(1), 195–202.
 450 <https://doi.org/10.1016/j.foodchem.2011.03.027>

451 Rocha, G. A., Fávaro-Trindade, C. S., & Grosso, C. R. F. (2012). Microencapsulation of
 452 lycopene by spray drying: Characterization, stability and application of microcapsules.
 453 *Food and Bioproducts Processing*, 90(1), 37–42.
 454 <https://doi.org/10.1016/j.fbp.2011.01.001>

455 Rossetti L., Langman L., Grigioni G. M., Biolatto A., Sancho A. M., Comerón E., & Descalzo
 456 A. M. (2010). Antioxidant status and odour profile in milk from silage or lucerne-fed
 457 cows. *Australian Journal of Dairy Technology*, 65(1), 3–9.

- Salmerón, I. (2017). Fermented cereal beverages: from probiotic, prebiotic and synbiotic towards Nanoscience designed healthy drinks. *Letters in Applied Microbiology*, 65(2), 114–124. <https://doi.org/10.1111/lam.12740>
- Salmerón, Ivan, Thomas, K., & Pandiella, S. S. (2015). Effect of potentially probiotic lactic acid bacteria on the physicochemical composition and acceptance of fermented cereal beverages. *Journal of Functional Foods*, 15, 106–115. <https://doi.org/10.1016/j.jff.2015.03.012>
- Schäfer, K. (1998). Accelerated solvent extraction of lipids for determining the fatty acid composition of biological material. *Analytica Chimica Acta*, 358(1), 69–77. [https://doi.org/10.1016/S0003-2670\(97\)00587-4](https://doi.org/10.1016/S0003-2670(97)00587-4)
- Septembre-Malaterre, A., Remize, F., & Poucheret, P. (2018). Fruits and vegetables, as a source of nutritional compounds and phytochemicals: Changes in bioactive compounds during lactic fermentation. *Food Research International*, 104, 86–99. <https://doi.org/10.1016/j.foodres.2017.09.031>
- Serafini, M., Stanzione, A., & Foddai, S. (2012). Functional foods: traditional use and European legislation. *International Journal of Food Sciences and Nutrition*, 63(sup1), 7–9. <https://doi.org/10.3109/09637486.2011.637488>
- Stephensen, C. B. (2001). VITAMIN A, INFECTION, AND IMMUNE FUNCTION *. *Annual Review of Nutrition*, 21(1), 167–192. <https://doi.org/10.1146/annurev.nutr.21.1.167>
- USDA. (2018). *USDA - Basic Report: 43261, Yogurt, fruit variety, nonfat*. Retrieved from <https://ndb.nal.usda.gov/ndb/foods/show/43261?fgcd=&manu=&format=&count=&max=25&offset=&sort=default&order=asc&qlookup=soya+yogurt&ds=SR&qt=&qp=&qq=&qn=&q=&ing=>

Figure captions

- **Figure 1:** Schematic overview of the manufacturing process
- **Figure 2:** Growth kinetics with Rosso model and pH drop of fermented products with A) 100 % *L. plantarum*; B) 100 % *L. casei*; C) ratio 50-50 *L. plantarum* /*L. casei*
— Rosso model; — pH; ♦ ▲ ■ log(biomass)
- **Figure 3:** Contents of carotenoids, tocopherols and β -sitosterol at different steps of the process with a and b as different statistic groups ($p < 0.05$);
■ Not pasteurized; ■ Pasteurized, not fermented; ■ Fermented for 15h

Figure 1

Figure 2

Figure 3

Table 1. Fat-soluble phytomicronutrients and macronutrients content of the functional yogurt-like product

Phytomicronutrients	Content	Macronutrients	g per 100 g
Carotenoids	(mg/kg)		
β -cryptoxanthin	1.43 ± 0.09	Proteins	8.12 ± 0.39
β -carotene	7.65 ± 0.48	Lipids	3.48 ± 0.08
Lycopene	7.02 ± 0.55	Fruits sugars	13.50 ± 0.12
Tocopherols	(mg/kg)	<i>Glucose</i>	6.16 ± 0.05
α -tocopherol	0.34 ± 0.04	<i>Lactose</i>	1.08 ± 0.01
γ -tocopherol	3.47 ± 0.30	<i>Fructose</i>	6.26 ± 0.06
Phytosterols	(g/kg)	Added sugar (<i>Sucrose</i>)	3.6 ± 0.18
β -sitosterol	19.88 ± 1.10	Starch	2.34 ± 0.29

Table 2. Parameters of strains' growth kinetics

Lactobacillus strains	Specific growth rate (h ⁻¹)	Generation time (h)	Latency period (h)	Initial pH – Final pH
<i>L. plantarum</i>	0.86 ^b	0.81	4	1.7
<i>L. casei</i>	1.16 ^a	0.60	5	1.6
<i>L. plantarum</i> / <i>L. casei</i> 50-50	0.94 ^b	0.74	3	1.9

a and b letters significantly different (p < 0.05)