Microplastics pollution along the Lebanese coast (Eastern Mediterranean Basin):

Occurrence in surface water, sediments and biota samples

Maria Kazour^{a,b}, Sharif Jemaa^b, Christelle Issa^b, Gaby Khalaf^b, Rachid Amara^a

^a Univ. Littoral Côte d'Opale, CNRS, Univ. Lille, UMR 8187, LOG, Laboratoire d'Océanologie et de Géosciences, 32 Avenue Foch, Wimereux, France

^b National Center for Marine Sciences, CNRS-L, , PO Box 534, Batroun, Lebanon

The Mediterranean Sea is the largest semi-enclosed sea and one of the worst affected regional seas with sub-basin scale heterogeneity in plastics concentration. Few studies on microplastics (MPs) pollution have been conducted in the Eastern part of the Mediterranean basin. This study aims to evaluate, for the first time, the MPs pollution of the Lebanese coast (Levantine Basin) as well as the most common polymers found, and to assess the potential role of coastal landfills in this pollution. Two important seafood species that are wholly consumed by the Lebanese community: the European anchovy *Engraulis encrasicolus* and the spiny oysters *Spondylus spinosus* were sampled in three different sites englobing the littoral (Tripoli, Beirut and Sidon). Sea water and sediment samples were also collected from the same sites. Results showed different patterns of MPs concentration in the analyzed matrices: Sidon water sample was highly contaminated in MPs (6.7 MPs/m³) whilst Tripoli had the highest MPs in sediments (4.68 MPs/g). The occurrence of MPs in the biota was high (83.4% and 86.3% in anchovies and spiny oysters, respectively). Both anchovies and oysters from Beirut region had the highest ingested MPs/ individual (2.9 ± 1.9 and 8.3 ± 4.4 MPs/individual, respectively). This study is the first that investigated microplastics ingestion by *Spondylus spinosus* while indicating the most common polymers found in the three matrices (water, sediments and biota) in the Eastern Mediterranean. These results highlighted the high MPs pollution found in the Levantine Basin in comparison to other Western Mediterranean regions. In addition, the obtained results indicate a potential contribution of coastal landfills to this pollution.

Corresponding author: Maria Kazour; maria.kazour@univ-littoral.fr

Keywords: Microplastics, Levantine Basin, coastal landfill, filter-feeders, micro-Raman

1. Introduction

The world plastic production is continuously increasing, reaching 350 million tons in 2017 (PlasticsEurope, 2018). Among the produced plastics, only 6–26% are recycled. Landfills receive 21–42% of wastes, whereas the rest are wildly released into the environment due to mismanagement through a variety of pathways (Ellen MacArthur Foundation, 2016; O'Connor et al., 2016). Most of the manufactured plastics will eventually end up in the ocean. By the year 2025, about 250 million tons of wastes are expected to enter the ocean water (Jambeck et al., 2015) leading to their accumulation in the marine environment (Thompson et al., 2009; UNEP, 2014). Plastic pollution in the marine environment is becoming a problem of global concern. These plastics are capable of transforming into dangerous small particles known as microplastics (MPs) existing as particles of a size below 5 mm ubiquitously available in the aquatic environments (Thompson et al., 2004; Andrady, 2011). They are either intentionally produced in small sizes such as polystyrene beads for example, or they are the resultant of macroplastics degradation (whether by Photochemical reaction, or mechanical actions.).

The Mediterranean Sea is prone to several anthropic pressures and has recently been described as one of the most affected marine environments with regards to marine litter and microplastics (Lebreton et al., 2012; Cincinelli et al., 2019). Mainly owing to the semi-enclosed situation and limited outflow of surface waters, a densely populated coastline and intensive fishing, shipping, touristic and industrial activities, substantial amounts of marine litter are accumulating in the Mediterranean Basin. Also, most studies on microplastics have been conducted in the northwestern Mediterranean Sea (e.g. Collignon et al., 2014; Fossi et al., 2016; Ruiz-Orejón et al., 2016; Schmidt et al., 2018; Lefebvre et al., 2019). Results showed that their distribution and composition are not homogeneous and geographical differences exist between Mediterranean sub-basins (Suaria et al., 2016; Cincinelli et al., 2019). Such heterogenicity in MPs distribution

and of its concentration levels are likely reflecting a complex interplay between pollution sources, hydrodynamic features such as currents, up and down-welling, gyres, fronts and the shape of the coastline (Suaria et al., 2016; Schmidt et al., 2018; Cincinelli et al., 2019). In addition, the absence of a standardized sampling protocol for the identification and quantification levels of MPs could also be responsible for this heterogeneity in results.

Plastics in general, and microplastics specifically, are prone to be ingested by a wide range of organisms (Lusher et al., 2016) and about one hundred species in the Mediterranean have been affected either via ingestion or entanglement (Deudero and Alomar, 2015). Microplastics were already found in several Mediterranean fish species such as *Boops boops*, *Sardina pilchardus*, *Mullus barbatus* and *Merluccius merluccius* (Giani et al., 2019; Bellas et al., 2016; Rios-Fuster et al., 2019) and in wild mussels (*Mytilus galloprovincialis*) caught off the Adriatic Sea and central Tyrrhenian Sea (Gomiero et al., 2019; Renzi et al., 2018a).

Although it is suggested that it is the most polluted region, few studies have been conducted in the Eastern part of the Mediterranean Basin (Van der Hal et al., 2017; Güven et al., 2017; Gündoğdu and Çevik, 2017). In the Levantine basin, the Lebanese coast is highly affected by numerous anthropogenic pressures such as harbor activities, fishing, domestic and industrial wastewater discharges (generally without prior treatment). Waste production in Lebanon has increased significantly during the last decade and the country has experienced waste crises. The regulated dumping sites could no longer manage the tremendous quantity of municipal garbage generated and, therefore, numerous unregulated and open dumping sites were created throughout the country mainly along the coast (Khalil et al., 2018). The average composition of municipal solid waste generated shows an important percentage of plastics (15%) which is two to three times higher than that found in Sweden, France and the United States (El-Hoz, 2019). Though the processes may be slow, landfills are affected by UV radiation and mechanical abrasions leading to the fragmentation of plastics to microplastics, and small plastic debris would find their way into the aquatic environment via leachates discharges (Alimi et al., 2018).

Despite these concerns, there have been no studies specifically examining the occurrence and abundance of microplastics along the Lebanese coastal environment. The aim of this study was to investigate the quantity and types of microplastics found in three matrices (sea water,

sediments and marine biota) along the Lebanese coast. All suspected MPs were characterized by size, color and identified according to their polymer nature using micro-Raman spectroscopy. Fish and bivalves are known to ingest microplastics (Collard et al., 2015; Phuong et al., 2018; Kazour et al., 2019), making them a potential vector of toxic chemicals through food chains and into human diets. In the present study we analyzed two marine species that are wholly consumed by the Lebanese community: one pelagic fish species: the European anchovy, *engraulis encrasicolus* (Linnaeus, 1758) and a bivalve species, the spiny oyster, *Spondylus spinosus* (Schreibers, 1793).

2. Materials and Methods

2.1. Study and sampling area

Lebanon is located in the Eastern Mediterranean (Levantine Basin) with a coastal length of 220 km (Figure 1). Three sampling sites were chosen along the coast according to their degree of anthropogenic pressures making them pollution hotspots (Shaban, 2008). The first site, Tripoli, located in the north, is facing the town's public beaches, characterized by the presence of a fishing harbor, and the existence of a coastal landfill. This latter was constructed in the year 1999 and reached today 36 m height receiving about 400 tons of wastes/day. The second site Beirut, located in the middle, is the country's capital. This site is near the most important harbor in Lebanon and surrounded by three coastal landfills: One closed and two active landfills that reached their maximum capacities (55 meters of height) in July 2019. The third site Sidon, located in the south, is facing a river's mouth and is near a closed landfill operated between 1982 and 2013. With a height of 58 m, this landfill was the biggest in the country receiving 300 tons of wastes/day. Actually, its transformation to a public park led to the burying of all wastes underground; traces of these wastes can always be observed underwater. These landfills lack an appropriate leachates treatment or it is not operational, and leachates are directly dumped to the sea. The sampling for all three matrices was done in May 2018 (spring).

2.2. Contamination control

Several precautions were followed to minimize air-born contamination during sampling, samples preparation and analysis. Cotton lab coats were worn and different procedures (filtration, measurements and digestion) were done under a laminar flow hood. All surfaces and equipment were rinsed before usage with filtered ethanol 70% and distilled water. Laboratory supplies were made of glass (bottles, Petri dishes, Erlenmeyers, filtration system) and they were covered immediately after manipulation. All solutions (distilled water, Ethanol 70%, potassium hydroxide KOH 10% and zinc chloride ZnCl_2) were filtered 3 times on glass fibers filter (GF/A filters, 1.6 μm porosity, Whatman, France) to eliminate unwanted particles or fibers. Controls were made to follow up the contamination occurring with every step of preparation and observation.

2.3. Water and sediments sampling and analyses

Sea surface water and sediments samples were collected during spring 2018. Sea surface water was collected using a manta trawl net (52 μm mesh size) with an opening of 0.63 m and a length of 0.255 m equipped with a flowmeter. The net was trawled for 10 minutes parallel to the coast at about 500 m from the coast with a small artisanal vessel (3 m of length) at a speed of 2 knots. On average, 100 m^3 of sea water was sampled at each site (Supplementary Table 1). The net was washed with filtered distilled water and all collected residues were rinsed inside clean glass bottles. The collected water was filtered on a stainless-steel sieve of a mesh size of 850 μm and the recovered water was vacuum filtered on GF/A filters. The suspected MPs present on the sieve were sorted visually under the stereomicroscope, fixed on a microscopic slide and conserved inside a clean glass Petri dish.

The first two centimeters of sublittoral sediments were sampled using a steel ring with two stainless-steel plates, rinsed with filtered distilled water before usage. The samples were put inside clean glass jars and stored at -4°C . Samples were thawed at room temperature 12 hours prior to analysis. Then particles with a size range of 1 – 5 mm were visually sorted and the remaining sediments underwent a density separation using filtered ZnCl_2 ($d = 1.8 \text{ g/cm}^3$) (Imhof et al., 2013; Nuelle et al., 2014). Briefly, 50 mL of filtered ZnCl_2 was added to the sample; the solution was then shaken for 5 minutes twice (once every 4 hours) and let rest for 16 hours. The

supernatant was then recovered and filtered on GF/A filters. Microplastics found were reported per gram of sediments sample and per m³ in sea water samples.

2.4. Biota sampling and analyses

Anchovies and spiny oysters were also collected during spring 2018. Anchovies were caught using purse seine nets. Ten individuals of approximately the same size (9 cm ± 0.9) were chosen from each sampling site and were immediately individually conserved in aluminum foil and conserved at -4°C. In the laboratory, their morphometric characteristics (total length and weight, digestive tube weight) were measured and their digestive tracts were dissected, conserved in aluminum foil and frozen at -4°C. For the spiny oysters, ten individuals were collected by fishermen from each sampling site. In the laboratory, the whole individual was removed from its shell and was conserved in aluminum foil. The net weight of the animal ranged between 4.8 g to 38.2 g with an average of 20.2 g ± 9.4.

All samples were defrosted 12 hours prior to digestion. Fish digestive tracts and oyster soft tissues were digested using potassium hydroxide 10% KOH (Dehaut et al., 2016). In short, the organic material of each individual was put inside separate Erlenmeyer containing solution of KOH 10% and then they were incubated at 60°C for 24 hours. Then, the digestat was recovered and vacuum filtered on GF/A filters and put inside clean glass Petri dishes.

2.5. Stereomicroscope and micro-Raman spectroscopy

All samples were first observed under a Leica M165 C Stereomicroscope with a 120x magnification. Photos of items were captured using a Leica M170 HD camera connected to LAS (Leica Application Suite) software. They were listed, measured, and their morphology, color and shape were noted. Several criteria were taken into consideration while counting the suspected items: (1) absence of cellular or organic structures; (2) a homogenous thickness across the particles; and, (3) homogenous colors (Hidalgo-Ruz et al. 2012). Suspected microplastics were divided into 6 size classes: <200 µm, 200-400 µm, 400-600 µm, 600-800 µm, 800-1000µm, and > 1000 µm.

Suspected microplastics were analyzed using Micro-Raman Xplora Plus (HORIBA Scientific® France). This machine is equipped with a closing chamber that isolates the filter from its surrounding, hence minimizing any airborne contamination.

Due to the high amount of time it takes when analyzing filters under micro-Raman, five filters of anchovies and five filters of spiny oysters from each site containing the highest number of potential MPs were identified. Subsamples of sediments and water were analyzed in a way that 25% of particles were analyzed. For MPs polymers identification, two lasers were used with a wavelength of 532 nm and 785 nm with a range of 200-3400 cm⁻¹. Two objectives (Olympus, Rungis, France) were used: x10 and x100. Filters were either analyzed manually or using ParticleFinder module for LabSpec 6. Each particle spectrum is compared to a database polymer identification software (KnowItAll, BioRad®) and a personal library made with specific polymers obtained from Goodfellow (France).

2.6. Data and Statistical analysis

Anchovy Fulton's K condition index was calculated as an indicator of the fish general well-being: $K = (\text{Body weight} / \text{total length}^3) * 100$. The feeding intensity for the fish was calculated using the Gastro Somatic Index (GaSI) (Desai 1970): $\text{GaSI} = \text{stomach weight} / (\text{body weight} - \text{stomach weight})$. The mean ($\pm$ SD) abundances of microplastics were calculated for anchovies (MPs/ind) and spiny oysters (MPs/g). After a Shapiro-Wilk test, data distribution of ingested MPs were non normal and a non-parametric tests (Kruskal-Wallis and Mann-Whitney U tests) were used to determine differences in microplastic numbers among species and sites. GaSI values followed a normal distribution and a one way analysis of variance (ANOVA) test with a post-hoc Tukey test was used. Spearman's rank correlation was used to test any correlation between the number of ingested MPs and the fish condition index or GaSI. Non-metric Multidimensional Scaling (n-MDS) was performed on a Bray-Curtis similarity matrix created from transformed data (square root). This test was used as an ordination method alongside ANOSIM and SIMPER tests to compare the contribution of each observed polymers in sea water, sediments and biota to the similarity and dissimilarity between sites. Significance level was set at $p < 0.05$. All statistical tests were done

using SPSS software (IBM SPSS STATISTICS 20) and PRIMER v7.0 (Primer-E Ltd., Plymouth Marine Laboratory, UK).

3. Results and Discussion

3.1. Microplastics concentrations in sea surface waters and sediments

Microplastics found along the Lebanese coastal waters consisted mainly of fragments (77.5%), fibers (15.25%) and microbeads (7.25%) (Figure 2- A). For all sites, blue and red colors were the most abundant representing 70% of all MPs. Concentration in the surface waters (4.3 MPs/m³) were high compared to other Mediterranean regions and particularly the western part where MPs concentrations vary between 0.17 and 0.62 MPs/m³ (Table 1). In the Eastern Mediterranean, the MPs concentration fluctuates importantly across regions, ranging from 0.7 items/m³ in Northeast Levantine coast of Turkey (Güven et al., 2017) to 7.68 ± 2.38 MP/m³ in the Israeli surface water (van der Hal et al., 2017).

The heterogeneity of the MPs distribution and of its concentration levels could be related to several factors. The mesh size of the net used could play an important role on the number of microplastics caught during sampling. The high concentration of MPs found in the present study maybe due to the low mesh size manta trawl net used (52 µm) compared to other studies (generally between 200 and 500 µm). Also, the different methodological analyses and the lack of a standard protocol could also be the cause of that difference. In fact, the water samples in this study were filtered on GF/A filters and all suspected items were observed; whereas other studies used organic material digestion (Gündoğdu and Çevik, 2017; Güven et al., 2017) or just visual observation (Fossi et al., 2016; Lefebvre et al., 2019) which can lead to an underestimation and a bias in microplastics counting. In addition, the distance of the sampling location from the coast may also influence the concentration of MPs found. Our samples were collected in the coastal waters (about 500 m from the shore). MPs concentration might have differed if the collected samples came from the open sea (Van der Hal et al., 2017; Pedrotti et al., 2016). On the other hand, the influence of hydrodynamic features such as currents, up and down-welling, gyres and fronts could also be responsible for this heterogeneity in MPs

concentrations. The Eastern Mediterranean has been reported for its high contamination of floating plastic debris with a dominance of small fragments < 2.5 cm in comparison to the Western part (Cózar et al., 2015). High fluxes of plastic debris (10-30 kg/km.day) were observed along the Levantine Basin coast from Egypt, Lebanon, Syria and Turkey (Constantino et al., 2019). Differences between the two parts of the Mediterranean Sea can be explained by the different circulation patterns: water currents in the Western part promote circulation; whereas currents in the Eastern part are somewhat attractors leading to the accumulation of floating debris (Mansui et al., 2015). In addition, the complexity of circulation with the existence of mesoscale and sub-basin scale eddies in the Levantine basin (most specifically Shikmona anti-cyclonic eddy) may also explain the high number of MPs found on the Lebanese coast (Menna et al., 2012).

High concentrations of MPs in sediments were also found along the Lebanese coast (2433 ± 2000 MPs/Kg d.w.). These concentrations were higher than those found in others Mediterranean regions (Supplementary Table 2). This important difference can be either caused by the different density separation solution used (ZnCl_2 used in our study is of higher density than NaCl used in other studies and more microplastics can be identified) or by the mismanaged waste along the Lebanese coast where high plastic items were visually observed during sampling. Both micro and macroplastics (MaPs) were present in all collected sediments samples. Lebanon is a potential landfall of plastics along with other Eastern Mediterranean regions (Duncan et al., 2018) explaining the high abundance of macroplastics found in our samples (consisting of half of the observed anthropogenic materials found). All kind of shapes were found: square, rectangular, spherical and cylindrical (Figure 3). White was the dominant color in Sidon and Tripoli (46.3% and 36.8%, respectively); blue color was the most abundant in Beirut (23.7%). Pellets were highly abundant in two of our sampling sites (Tripoli and Sidon) which explain the high abundance of white color in these two locations. Several studies investigating MPs abundance in the Mediterranean sediments also reported the high presence of plastic pellets among their samples (Karkanorachaki et al., 2018; Fanini and Bozzeda, 2018; Kaberi et al., 2013; Turner and Holmes, 2011). The first signaled pellets on the Lebanese coast were due to direct waste discharge of plastic factories on the beaches or in the rivers (Shiber, 1979). To

this day, and with the lack of an appropriate treatment, these industries are still releasing their wastes into the aquatic system and are susceptible to unintentional losses leading to this high accumulation of plastic pellets. All of which indicate their persistence whether in the Eastern or Western Mediterranean and they (pellets and fragments) consist potential harm to the benthic organisms inhabiting sandy beaches (Carson et al., 2011; Nelms et al., 2016).

Along the Lebanese coast, inter-site differences in MPs concentrations were observed. Surface water samples showed that Sidon was the most contaminated site with 6.7 MPs/m³ (Figure 2-A). This site is characterized by a high presence of microbeads with 1.16 microbeads/m³. Beirut had the lowest concentration of MPs with 2.35 MPs/m³ consisting mainly (90%) of fragments. The high number of MPs and pellets observed in the south of Lebanon can be explained by site's localization near the biggest landfill. Although this landfill is closed since 2013, results of our study provide preliminary evidence that landfill isn't the final sink of plastics, but a potential source of microplastics (He et al., 2019). Fragmentation and release of microplastics is a long-term process. Microplastics inside leachates from active and closed landfills have been identified (He et al., 2019). After the stop of post-closure care, microplastics in leachate would be discharged directly into coastal waters. This can pose serious environmental problems since microplastic is not listed as pollutant in the any country's regulations of landfill. This area is also characterized by the presence of the mouth of a major river, Al-Awali River. During sampling, we observed fragments and other wastes being brought into the coastal waters via the river runoff. This river extends for 48 km of length and collects all the urban water from nearby villages due to the non-existence of sewage treatment system. These microplastics are prone to be washed from surface waters onto beaches and, therefore, integrate within the sediments (Barnes et al., 2009; Poeta et al., 2014). The general circulation along the coast of Lebanon is dominant northwards and the water movements are strongly related to surface currents and seasonal metrological factors (Goedicke, 1972). This northwards movement might explain why MPs and MaPs were the most abundant in sediments samples of Tripoli (4.68 MPs/g and 4.08 MaPs/g, respectively) (Figure 2- B). The high levels of MPs found in the surface water in Sidon and Beirut might have been carried by surface circulation ending up washed on Tripoli beaches.

3.2. Microplastics in anchovies and spiny oysters

Microplastics can be ingested by a large variety of marine species and more specifically by filter-feeders (Van Cauwenberghe and Janssen, 2014; Catarino et al., 2018; Collard et al., 2015) raising the alarm for commercial species that are highly consumed by humans. In our samples, 83.3% of European anchovies analyzed had ingested MPs. Such high level of ingestion is in agreement with what was described for anchovy from the Adriatic Sea (90%) (Renzi et al., 2019a). However, in the Western Mediterranean Sea, lower occurrences of MPs were described in anchovy digestive tracts (11 to 40 %) (Collard et al., 2015; Compa et al., 2018; Lefebvre et al., 2019) (Table 2). In the present study, anchovies ingested an average of 2.5 ± 0.3 MPs/individual which was higher than that found in anchovies collected on the Northwestern Mediterranean coasts such as Spanish coasts (0.18 ± 0.20 items/individual) (Compa et al., 2018) and Gulf of Lions (0.11 to 0.85 MPs/individual) (Collard et al., 2015; Lefebvre et al., 2019) but lower than those collected in the Adriatic Sea (6.62 ± 3.67 items/individual) (Renzi et al., 2019a). Anchovies caught in the Western Mediterranean Sea had lower ingested MPs than those caught in the Eastern and Central Mediterranean confirming the higher pollution of these regions (Table 2). For instance, Japanese anchovies, *Engraulis japonicus* from the heavy polluted area of Tokyo bay displayed the same MPs ingestion (2.30 ± 2.50 MP/ind Tanaka and Takada, 2016) as in our study.

Three shapes of plastics were found: fragments, films and fibers (Figure 2 – C; Figure 3). There was a slight dominance of fragments (1.3 ± 0.36 fragments/fish) over of fibers (1.1 ± 0.17 fragment/fish), whereas films had the lowest occurrence (0.1 ± 0.1 films/fish). The items had different colors: black, blue, red, white, transparent, and green. For the three sites, blue fragments were the most abundant representing more than 50%; whereas for the fibers, black was the dominant color. This predominance of blue plastic items has been previously reported in fish and bivalves (Romeo et al., 2015; Güven et al., 2017; Compa et al., 2018; Digka et al., 2018).

Along the Lebanese coast, anchovies from Beirut had ingested the highest number of MPs items (2.9 ± 1.9 MPs/individual) and those from Tripoli the lowest number (2.3 ± 1.6 items/individual) (Supplementary Table 3). However, there were no significant differences between the numbers

of ingested items at the three analyzed sites ($P > 0.05$). No relationship was highlighted between the concentration of MPs in anchovy digestive tracts and in the surface water as it has been described in the western Mediterranean (Compa et al., 2018; Lefebvre et al., 2019). As a pelagic species, anchovy are highly moving fish so they potentially did not ingest MPs at the site they were collected, making correlations harder to underline. Gastro-Somatic Index (GaSI) and Fulton condition index were higher in Beirut yet not significantly different (Supplementary Table 3). There was no correlation between GaSI and the number of ingested items ($r = 0.067$; $P > 0.05$). Fulton condition index and the number of ingested items showed a weak but not significant correlation ($r = 0.316$; $P = 0.089 > 0.05$). In the western Mediterranean Sea, recent studies also revealed that anchovies' MPs ingestion was not related to the body condition (Compa et al. 2018; Lefebvre et al., 2019).

Spiny oyster is a Lessepsian species of high commercial value in Lebanon. The collected individuals had an average of 7.2 ± 1.4 items/individual and 0.45 ± 0.3 items/g ww. To date, this is the first study done on MPs ingestion in *S. spinosus* in the Mediterranean and worldwide. Therefore, our results can only be compared to other oysters or bivalves species. Oysters from the present study had the same average of items found in *Crassostrea gigas* collected from the Atlantic Ocean with an average of 0.47 ± 0.16 particles/g ww (Van Cauwenberghe and Janssen, 2014), yet lower than that found in wild mussels (*Mytilus galloprovincialis*) collected from the central Tyrrhenian Sea (Mediterranean) with an average of 7.2 items/g ww (Renzi et al., 2018a) and from the Adriatic Sea with an average of 1.06–1.3 items/g ww (Gomiero et al., 2019). Our samples were less contaminated than those collected from the Mediterranean which is probably due to the difference in the analyzed species and not the geographical MPs concentration in the environment. Oysters tend to have more flesh than mussels explaining the difference in MPs ingestion per gram of ww.

Oysters from all sites had the presence of fragments, films and fibers with 86.3% of analyzed individuals having suspected MPs in their soft tissue (Figure 2- C). Due to a difference in individual total body soft tissue weight, the number of items was also reported to the number of items per gram. Oysters from Beirut had the highest items concentration with 8.3 ± 4.4

items/individual (or 0.78 ± 0.62 items/g) out of which fibers constituted more than 50%. Sidon had the highest number of fragments/individual reaching 4.3 ± 4.4 fragments/individual. Tripoli had the lowest number of items/g with 0.22 ± 0.13 items/g. Beirut was significantly different when comparing to the other sites ($P < 0.05$). The observed items had six different colors similar to those found in the anchovies; with blue color being dominant in the oysters' samples.

3.3. Size class distribution and polymers characterization

In the water samples, the size class $< 200 \mu\text{m}$ dominated in Tripoli (48.78%) and Sidon (51.8%) whereas for Beirut, the size class $> 1000 \mu\text{m}$ dominated (31.7%) (Figure 3 – A). Suspected items $< 1000 \mu\text{m}$ represented a percentage of 88%, 69.3% and 87% at Tripoli, Beirut and Sidon, respectively. All sediment samples had a high abundance of the size class $> 1000 \mu\text{m}$ representing more than 70% of the samples (Supplementary Figure 1). For the biota, the smallest size class $< 200 \mu\text{m}$ was the most abundant in all sites (Figure3 - B). For anchovies, the ingested size class 200-400 μm was the second most dominant in Tripoli and Beirut with a percentage of 34%. The size class 600-800 μm was only present in Sidon's anchovies, whereas the size class $> 1000 \mu\text{m}$ was absent in Tripoli. These results agree with the feeding ecology of the European anchovy which is a filter-feeder that feeds on planktonic organisms generally < 2 mm (Bacha and Amara, 2009). No relationship between the size of the anchovies and the size of the ingested items was observed. Although Beirut's anchovies were significantly ($P < 0.05$) bigger than those from the other two sites, they ingested items of a similar size than those ingested in Tripoli and Sidon. This indicates that the size of ingested items may be related to the available particles during feeding rather than the morphometric characteristics of organisms (i.e. mouth opening size). Plastics of all size classes were present inside the oyster's body soft tissue. The smallest size class (0-200 μm) was the most abundant for all sites. The two size classes: 200-400 μm and $> 1000 \mu\text{m}$ were the second most abundant between the ingested MPs. Even though there was a significant difference in spiny oyster's net weight ($P < 0.05$) between sites, all MPs items size classes were present in all collected individuals. This confirms that the size of ingested items is not related to the individual weight, but to the available particles existing in the surrounding water.

Among the suspected MPs analyzed, 94% were identified as microplastics. Nine types of polymers were identified in our samples (Figure 4). Six types were common to all three matrices (water, sediments and biota): polypropylene (PP), polyethylene (PE), polystyrene (PS), polyamide (PA), polyethylene terephthalate (PET) and polyurethane (PUR). In addition to these polymers, polyvinyl chloride (PVC) was present in water and sediments; polylactic acid (PLA) was only present in the water whereas acrylonitrile butadiene styrene (ABS) was only present in biota. Each of the three matrices had a specific polymer that dominated over the others: PE was the most abundant in water samples (70.83%, 50% and 47.37% at Tripoli, Beirut and Sidon, respectively). Polyethylene is the most common plastic on a global scale and consequently the most dominant plastic debris in the Mediterranean Sea and worldwide, deriving mainly from plastic bags and bottles (Suaria et al., 2016). It is well documented that plastics as PE float in the water surface, as its density is lower than this of the water. Polymers with a density higher than that of water (PVC, for example) were observed in the surface water. The existence of polymers denser than seawater have been already reported (Suaria et al., 2016; Enders et al., 2015) with no clear mechanism leading to their existence. During sampling, there was strong waves and the water was agitated which might explain their presence due to the high velocity and turbulent mixing which increases the chances of denser polymers to persist on the sea surface (Moritomi et al., 1982; Suaria et al., 2016). Polypropylene was the most abundant in sediments (48.11%, 55.88% and 43.33% at Tripoli, Beirut and Sidon, respectively), and polystyrene dominated in the biota specimens (having a percentage equal to or above 25% for anchovies and spiny oysters). Polypropylene was observed in all anchovies and spiny oysters of all sites except in the anchovies of Beirut where PE was observed instead. Polymers distribution didn't have a pattern with shapes or colors. For example, pellets found were mostly white (several were colored) yet their polymer type was of PE, PA, and PP. The same for fragments, different colors were related to different polymers with no similar pattern.

Comparison of the polymers composition identified in water, sediments and oysters indicated a dissimilarity between groups (ANOSIM, $R=0.796$, significance level = 0.2%). Cluster dendrogram led to the existence of two distinct groups (a and b) regrouping on the one hand the water and sediment samples and on the other hand the biota (anchovies and oysters) (Supplementary

Figure 2). Non-metric multidimensional scaling (n-MDS) showed the dimensional distance separating these groups (a and b) based on their polymer composition. Water and sediments formed 'group a' and were characterized by the high presence of PP and PE. Whereas 'group b' composed by biota samples, and were characterized by the high abundance of PS. Similarity of Percentages (SIMPER) indicated that PE, PP, PA and PUR contributed to 72.12% of dissimilarity between these two groups.

3.4. Microplastics potential entry into the Mediterranean Basin via coastal landfills, and their fate and distribution in the coastal environment

This study showed a high microplastics pollution in areas located in the vicinity of coastal landfills. Even though some of them are no longer active, but their waste continues to find its way into the sea due to mismanagement and lack of proper treatments. Some of the polymers found in our samples such as polyethylene, polypropylene and polystyrene were observed in transparent, white and different colors. They are widely used in packaging, consumer products, pipes and fitting, and are the common polymers found in landfills and their leachates (Kazour et al., 2019; He et al., 2019; Praagh et al., 2018). The high abundance of these types of polymers in sea surface water and sediment samples can be traced back to mismanaged landfills along with accidental garbage disposal. Even though Lebanon is considered as a potential sink of plastics in the Eastern Mediterranean (Duncan et al., 2018), it should also be considered as an important microplastics source in the Mediterranean basin. In most studies, whether in the surface water, sediments or biota, fibers were generally the dominant microplastic shape found (Lefebvre et al., 2019; Renzi et al., 2019b; Compa et al., 2018). In the present study, fragments constituted more than 77%, 90% and 50% of items found in surface water, sediments and biota samples, respectively. The same pattern was found in a study done near an abandoned coastal landfill, where fragments constituted the majority of found items (Kazour et al., 2019). Such dominance of fragments in the analyzed samples highlights on the potential role of Lebanese coastal landfills in fragments input into the Mediterranean Basin. Other factors can also lead to this severe microplastics contamination and, most precisely, on fragments abundance on the

Lebanese beaches. Munari et al. (2017) indicated a high number of fragments (approximately 90% of the samples) was observed near beaches affected by riverine run-off. All of the sampled Lebanese beaches are affected by riverine runoff as well as a significant anthropogenic pressure leading to a major microplastics pollution along the beaches (100 times higher than other reported studies, see Supplementary Table 3). These microplastics are prone to re-enter the surface water with waves hitting the shoreline (directed by wind speed and direction) increasing their chance to end up in nearby regions via the northward general water circulation pattern. The above stated factors increase microplastics availability to marine organisms leading to potential human health risks. In Lebanon, an average of 19.7g of fish and seafood per person is consumed (Nasreddine et al., 2006) making it around 7 kg per caput per annum. Converting the average of ingested MPs/g in oysters to the number of seafood ingested by a Lebanese person, we would have approximately 31, 500 MPs ingested per person (4500 MPs/kg of oysters). This is similar to that obtained by Catarino et al. (2018) (13,731 and 68,415 particles/y/person) yet lower than that obtained by Van Cauwenberghe & Janssen (2014) (11, 000 MPs/y/person). European anchovies are eaten wholly by the Lebanese population as *Bizree* (juvenile pelagic fish caught by purse seiners) as well as spiny oysters that are served in various Lebanese restaurants raising the concern on the human consumption of these contaminated seafood.

4. Conclusion

These results should be considered as a preliminary analysis on MPs levels recorded off the Lebanese coast in three different matrices. This is the first contribution to characterize the area and to understand the degree of its MPs pollution. More studies are necessary to evaluate the ecological impacts of these MPs on the marine biodiversity in the Levantine Basin. The levels of MPs found were alarmingly high in comparison to other Mediterranean regions. The high number of ingested items by both anchovies and oysters could suggest a potential health concern since these two species are wholly consumed (without being degutted). This study highlighted the role of landfills in MPs entry into the coastal marine environment. Coastal landfills and mismanaged wastes are important issues existing in 80% of the Mediterranean

Basin explaining the elevated number of MPs found in this semi-enclosed basin. The whole Mediterranean should be regarded as an important case study for microplastics pollution where more initiative should be taken to insure an appropriate waste management and, hence, decreasing plastics pollution.

Funding

Maria Kazour is financially supported by a PhD fellowship from the National Council for Scientific Research (Lebanon) and Université du Littoral Côte d'Opale (France). This work has been partially financially supported by the European Union (ERDF), the French Government, the Région Hauts-de-France and IFREMER, in the framework of the project CPER MARCO 2015-2020.

Declaration of interests: None

Acknowledgments

The authors would like to thank G. Duflos, A. Dehaut and L. Hermabessiere from the Agence Nationale de Sécurité Sanitaire de L'alimentation, de L'environnement et du Travail (ANSES) – Boulogne-sur-mer for their help and advices during the sampling preparation. The authors would also like to thank Elie Tarek from the National Center of Marine Sciences - Batroun for helping us acquiring our field samples.

References

Abidli, S., Antunes, J.C., Ferreira, J.L., Lahbib, Y., Sobral, P., Trigui El Menif, N., 2018. Microplastics in sediments from the littoral zone of the north Tunisian coast (Mediterranean Sea). *Estuar. Coast. Shelf Sci.* 205, 1–9. <https://doi.org/10.1016/j.ecss.2018.03.006>

477 Alimi, O. S., Farner Budarz, J., Hernandez, L. M., Tufenkji, N., 2018. Microplastics and
 478 nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant
 479 transport. *Environ. Sci. Technol.* 52(4), 1704-1724. <https://doi.org/10.1021/acs.est.7b05559>

480 Andrady, A.L., 2011. Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596–1605.
 481 <https://doi.org/10.1016/j.marpolbul.2011.05.030>

482 Bacha, M., Amara, R., 2009. Spatial, temporal and ontogenetic variation in diet of anchovy
 483 (*Engraulis encrasicolus*) on the Algerian coast (SW Mediterranean). *Estuar. Coast. Shelf Sci.* 85,
 484 257–264. <https://doi.org/10.1016/j.ecss.2009.08.009>

485 Bainsi, M., Fossi, M.C., Galli, M., Caliani, I., Campani, T., Finoia, M.G., Panti, C., 2018. Abundance
 486 and characterization of microplastics in the coastal waters of Tuscany (Italy): The application of
 487 the MSFD monitoring protocol in the Mediterranean Sea. *Mar. Pollut. Bull.* 133, 543–552.
 488 <https://doi.org/10.1016/j.marpolbul.2018.06.016>

489 Barnes, D.K., Galgani, F., Thompson, R.C., Barlaz, M., 2009. Accumulation and fragmentation of
 490 plastic debris in global environments. *Philos. Trans. R. Soc. London. Ser. B*, 364, 1985-1998.
 491 <https://doi.org/10.1098/rstb.2008.0205>

492 Bellas, J., Martínez-Armental, J., Martínez-Cámara, A., Besada, V., Martínez-Gómez, C., 2016.
 493 Ingestion of microplastics by demersal fish from the Spanish Atlantic and Mediterranean coasts.
 494 *Mar. Pollut. Bull.* 109, 55–60. <https://doi.org/10.1016/j.marpolbul.2016.06.026>

495 Blašković, A., Fastelli, P., Čížmek, H., Guerranti, C., Renzi, M., 2017. Plastic litter in sediments
 496 from the Croatian marine protected area of the natural park of Telaščica bay (Adriatic Sea).
 497 *Mar. Pollut. Bull.* 114, 583–586. <https://doi.org/10.1016/j.marpolbul.2016.09.018>

498 Cannas, S., Fastelli, P., Guerranti, C., Renzi, M., 2017. Plastic litter in sediments from the coasts
 499 of south Tuscany (Tyrrhenian Sea). *Mar. Pollut. Bull.* 119, 372–375.
 500 <https://doi.org/10.1016/j.marpolbul.2017.04.008>

501 Carson, H.S., Colbert, S.L., Kaylor, M.J., McDermid, K.J., 2011. Small plastic debris changes water
 502 movement and heat transfer through beach sediments. *Mar. Pollut. Bull.* 62, 1708–1713.
 503 <https://doi.org/10.1016/j.marpolbul.2011.05.032>

504 Catarino, A.I., Macchia, V., Sanderson, W.G., Thompson, R.C., Henry, T.B., 2018. Low levels of
 505 microplastics (MP) in wild mussels indicate that MP ingestion by humans is minimal compared
 506 to exposure via household fibres fallout during a meal. *Environ. Pollut.* 237, 675–684.
 507 <https://doi.org/10.1016/j.envpol.2018.02.069>

508 Cincinelli, A., Martellini, T., Guerranti, C., Scopetani, C., Chelazzi, D., Giarrizzo, T., 2019. A
 509 potpourri of microplastics in the sea surface and water column of the Mediterranean Sea. *TrAC -*
 510 *Trends Anal. Chem.* 110, 321–326. <https://doi.org/10.1016/j.trac.2018.10.026>

511 Collard, F., Gilbert, B., Eppe, G., Parmentier, E., Das, K., 2015. Detection of Anthropogenic
 512 Particles in Fish Stomachs: An Isolation Method Adapted to Identification by Raman
 513 Spectroscopy. *Arch. Environ. Contam. Toxicol.* 69, 331–339. [https://doi.org/10.1007/s00244-](https://doi.org/10.1007/s00244-015-0221-0)
 514 [015-0221-0](https://doi.org/10.1007/s00244-015-0221-0)

515 Collignon, A., Hecq, J.H., Galgani, F., Collard, F., Goffart, A., 2014. Annual variation in neustonic
 516 micro- and meso-plastic particles and zooplankton in the Bay of Calvi (Mediterranean-Corsica).
 517 *Mar. Pollut. Bull.* 79, 293–298. <https://doi.org/10.1016/j.marpolbul.2013.11.023>

518 Compa, M., Ventero, A., Iglesias, M., Deudero, S., 2018. Ingestion of microplastics and natural
 519 fibres in *Sardina pilchardus* (Walbaum, 1792) and *Engraulis encrasicolus* (Linnaeus, 1758) along
 520 the Spanish Mediterranean coast. *Mar. Pollut. Bull.* 128, 89–96.
 521 <https://doi.org/10.1016/j.marpolbul.2018.01.009>

522 Constant, M., Kerherve, P., Sola, J., Sanchez-Vidal, A., Canals, M., Heussner, S., 2018. Floating
 523 microplastics in the Northwestern Mediterranean Sea: temporal and spatial heterogeneities. In
 524 *Proceedings of the International Conference on Microplastic Pollution in the Mediterranean Sea*
 525 (p. 9-15). Springer, Cham. https://doi.org/10.1007/978-3-319-71279-6_2

526 Constant, M., Kerhervé, P., Mino-Vercellio-Verollet, M., Dumontier, M., Sánchez Vidal, A.,
 527 Canals, M., Heussner, S., 2019. Beached microplastics in the Northwestern Mediterranean Sea.
 528 *Mar. Pollut. Bull.* 142, 263–273. <https://doi.org/10.1016/j.marpolbul.2019.03.032>

529 Constantino, E., Martins, I., Salazar Sierra, J.M., Bessa, F., 2019. Abundance and composition of
530 floating marine macro litter on the eastern sector of the Mediterranean Sea. *Mar. Pollut. Bull.*
531 138, 260–265. <https://doi.org/10.1016/j.marpolbul.2018.11.008>

532 Cózar, A., Sanz-Martín, M., Martí, E., González-Gordillo, J.I., Ubeda, B., Ágálvez, J., Irigoien, X.,
533 Duarte, C.M., 2015. Plastic accumulation in the mediterranean sea. *PLoS One* 10, 1–12.
534 <https://doi.org/10.1371/journal.pone.0121762>

535 de Lucia, G.A., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L., Campani, T., Giannetti,
536 M., Coppola, D., Cicero, A.M., Panti, C., Baini, M., Guerranti, C., Marsili, L., Massaro, G., Fossi,
537 M.C., Matiddi, M., 2014. Amount and distribution of neustonic micro-plastic off the western
538 Sardinian coast (Central-Western Mediterranean Sea). *Mar. Environ. Res.* 100, 10–16.
539 <https://doi.org/10.1016/j.marenvres.2014.03.017>

540 Deudero, S., Alomar, C., 2015. Mediterranean marine biodiversity under threat: Reviewing
541 influence of marine litter on species. *Mar. Pollut. Bull.* 98, 58–68.
542 <https://doi.org/10.1016/j.marpolbul.2015.07.012>

543 Digka, N., Tsangaris, C., Torre, M., Anastasopoulou, A., Zeri, C., 2018. Microplastics in mussels
544 and fish from the Northern Ionian Sea. *Mar. Pollut. Bull.* 135, 30–40.
545 <https://doi.org/10.1016/j.marpolbul.2018.06.063>

546 Duncan, E.M., Arrowsmith, J., Bain, C., Broderick, A.C., Lee, J., Metcalfe, K., Pikesley, S.K., Snape,
547 R.T.E., van Sebille, E., Godley, B.J., 2018. The true depth of the Mediterranean plastic problem:
548 Extreme microplastic pollution on marine turtle nesting beaches in Cyprus. *Mar. Pollut. Bull.*
549 136, 334–340. <https://doi.org/10.1016/j.marpolbul.2018.09.019>

550 El-Hoz, M., 2019. A Qualitative–Quantitative Methodological Approach for Sustainable
551 Reclamation of Open Dumps: The Case of the Controlled Dump of Tripoli. In *Waste Valorisation*
552 *and Recycling* (p. 105-118). Springer, Singapore. https://doi.org/10.1007/978-981-13-2784-1_10

553 Ellen MacArthur Foundation, 2016. The New Plastics Economy- Rethinking the Future of Plastics,
554 p. 1-120. [https://www.ellenmacarthurfoundation.org/publications/the-new-plastics-economy-](https://www.ellenmacarthurfoundation.org/publications/the-new-plastics-economy-rethinking-the-future-of-plastics)
555 [rethinking-the-future-of-plastics](https://www.ellenmacarthurfoundation.org/publications/the-new-plastics-economy-rethinking-the-future-of-plastics)

556 Fanini, L., Bozzeda, F., 2018. Dynamics of plastic resin pellets deposition on a microtidal sandy
557 beach: Informative variables and potential integration into sandy beach studies. *Ecol. Indic.* 89,
558 309–316. <https://doi.org/10.1016/j.ecolind.2018.02.027>

559 Fastelli, P., Blašković, A., Bernardi, G., Romeo, T., Čížmek, H., Andaloro, F., Russo, G.F.,
560 Guerranti, C., Renzi, M., 2016. Plastic litter in sediments from a marine area likely to become
561 protected (Aeolian Archipelago's islands, Tyrrhenian sea). *Mar. Pollut. Bull.* 113, 526–529.
562 <https://doi.org/10.1016/j.marpolbul.2016.08.054>

563 Filgueiras, A. V., Gago, J., Campillo, J. A., León, V. M., 2019. Microplastic distribution in surface
564 sediments along the Spanish Mediterranean continental shelf. *Environ. Sci. Pollut. Res.* 1-10.
565 <https://doi.org/10.1007/s11356-019-05341-5>

566 Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R., 2012. Are
567 baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin
568 whale (*Balaenoptera physalus*). *Mar. Pollut. Bull.* 64, 2374–2379.
569 <https://doi.org/10.1016/j.marpolbul.2012.08.013>

570 Fossi, M.C., Marsili, L., Baini, M., Giannetti, M., Coppola, D., Guerranti, C., Caliani, I., Minutoli, R.,
571 Lauriano, G., Finoia, M.G., Rubegni, F., Panigada, S., Bérubé, M., Urbán Ramírez, J., Panti, C.,
572 2016. Fin whales and microplastics: The Mediterranean Sea and the Sea of Cortez scenarios.
573 *Environ. Pollut.* 209, 68–78. <https://doi.org/10.1016/j.envpol.2015.11.022>

574 Giani, D., Baini, M., Galli, M., Casini, S., Fossi, M.C., 2019. Microplastics occurrence in edible fish
575 species (*Mullus barbatus* and *Merluccius merluccius*) collected in three different geographical
576 sub-areas of the Mediterranean Sea. *Mar. Pollut. Bull.* 140, 129–137.
577 <https://doi.org/10.1016/j.marpolbul.2019.01.005>

578 GOEDICKE, T. R., 1972. Submarine canyons on the central continental shelf of Lebanon. The
579 Mediterranean Sea: Natural Sedimentation Laboratory, DJ Stanley Ed., Downen, Hutchinson &
580 Ross, Stroudsburg, PA, 800p.

581 Gomiero, A., Strafella, P., Øysæd, K.B., Fabi, G., 2019. First occurrence and composition
582 assessment of microplastics in native mussels collected from coastal and offshore areas of the

583 northern and central Adriatic Sea. *Environ. Sci. Pollut. Res.* [https://doi.org/10.1007/s11356-019-](https://doi.org/10.1007/s11356-019-05693-y)
584 05693-y

585 Gündoğdu, S., 2017. High level of micro-plastic pollution in the Iskenderun Bay NE Levantine
586 coast of Turkey. *Ege J. Fish. Aquat. Sci.* 34, 401–408.
587 <https://doi.org/10.12714/egejfas.2017.34.4.06>

588 Gündoğdu, S., Çevik, C., 2017. Micro- and mesoplastics in Northeast Levantine coast of Turkey:
589 The preliminary results from surface samples. *Mar. Pollut. Bull.* 118, 341–347.
590 <https://doi.org/10.1016/j.marpolbul.2017.03.002>

591 Güven, O., Gökdağ, K., Jovanović, B., Kıdeys, A.E., 2017. Microplastic litter composition of the
592 Turkish territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal
593 tract of fish. *Environ. Pollut.* 223, 286–294. <https://doi.org/10.1016/j.envpol.2017.01.025>

594 He, P., Chen, L., Shao, L., Zhang, H., Lü, F., 2019. Municipal solid waste (MSW) landfill: A source
595 of microplastics? -Evidence of microplastics in landfill leachate. *Water Res.* 159, 38–45.
596 <https://doi.org/10.1016/j.watres.2019.04.060>

597 Imhof, H.K., Ivleva, N.P., Schmid, J., Niessner, R., Laforsch, C., 2013. Contamination of beach
598 sediments of a subalpine lake with microplastic particles. *Curr. Biol.* 23, R867–R868.
599 <https://doi.org/10.1016/j.cub.2013.09.001>

600 Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., Law,
601 K.L., 2015. Plastic waste inputs from land into the ocean. *Science*, 347, 768-771,
602 <https://doi.org/10.1126/science.1260352>

603 Kaberi, H., Tsangaris, C., Zeri, C., Mousdis, G., Papadopoulos, A., Streftaris, N., 2013.
604 Microplastics along the shoreline of a Greek island (Kea isl., Aegean Sea): types and densities in
605 relation to beach orientation, characteristics and proximity to sources. *Proc. 4th Int. Conf.*
606 *Environ. Manag. Eng. Plan. Econ. SECOTOX Conf.* 197–202.

607 Karkanorachaki, K., Kiparissis, S., Kalogerakis, G.C., Yiantzi, E., Psillakis, E., Kalogerakis, N., 2018.
608 Plastic pellets, meso- and microplastics on the coastline of Northern Crete: Distribution and

609 organic pollution. Mar. Pollut. Bull. 133, 578–589.
610 <https://doi.org/10.1016/j.marpolbul.2018.06.011>

611 Kazour, M., Terki, S., Rabhi, K., Jemaa, S., Khalaf, G., Amara, R., 2019. Sources of microplastics
612 pollution in the marine environment: Importance of wastewater treatment plant and coastal
613 landfill. Mar. Pollut. Bull. 146, 608–618. <https://doi.org/10.1016/j.marpolbul.2019.06.066>

614 Khalil, C., Al Hageh, C., Korfali, S., Khnayzer, R.S., 2018. Municipal leachates health risks:
615 Chemical and cytotoxicity assessment from regulated and unregulated municipal dumpsites in
616 Lebanon. Chemosphere 208, 1–13. <https://doi.org/10.1016/j.chemosphere.2018.05.151>

617 Laglbauer, B.J.L., Franco-Santos, R.M., Andreu-Cazenave, M., Brunelli, L., Papadatou, M.,
618 Palatinus, A., Grego, M., Deprez, T., 2014. Macrodebris and microplastics from beaches in
619 Slovenia. Mar. Pollut. Bull. 89, 356–366. <https://doi.org/10.1016/j.marpolbul.2014.09.036>

620 Lebreton, L.C.M., Greer, S.D., Borrero, J.C., 2012. Numerical modelling of floating debris in the
621 world’s oceans. Mar. Pollut. Bull. 64, 653–661. <https://doi.org/10.1016/j.marpolbul.2011.10.027>

622 Lefebvre, C., Saraux, C., Heitz, O., Nowaczyk, A., Bonnet, D., 2019. Microplastics FTIR
623 characterisation and distribution in the water column and digestive tracts of small pelagic fish in
624 the Gulf of Lions. Mar. Pollut. Bull. 142, 510–519.
625 <https://doi.org/10.1016/j.marpolbul.2019.03.025>

626 Lots, F.A.E., Behrens, P., Vijver, M.G., Horton, A.A., Bosker, T., 2017. A large-scale investigation
627 of microplastic contamination: Abundance and characteristics of microplastics in European
628 beach sediment. Mar. Pollut. Bull. 123, 219–226.
629 <https://doi.org/10.1016/j.marpolbul.2017.08.057>

630 Lusher, A.L., O’Donnell, C., Officer, R., O’conner, I., 2016. Microplastic interactions with North
631 Atlantic mesopelagic fish. ICES J. Mar. Sci. 73, 1214–1225.
632 <https://doi.org/10.1093/icesjms/fsv241>

633 Mansui, J., Molcard, A., Ourmières, Y., 2015. Modelling the transport and accumulation of
634 floating marine debris in the Mediterranean basin. Mar. Pollut. Bull. 91, 249–257.
635 <https://doi.org/10.1016/j.marpolbul.2014.11.037>

636 Menna, M., Poulain, P.M., Zodiatis, G., Gertman, I., 2012. On the surface circulation of the
637 Levantine sub-basin derived from Lagrangian drifters and satellite altimetry data. *Deep. Res.*
638 *Part I Oceanogr. Res. Pap.* 65, 46–58. <https://doi.org/10.1016/j.dsr.2012.02.008>

639 Moritomi, H., Iwase, T., Chiba, T., 1982. A comprehensive interpretation of solid layer inversion
640 in liquid fluidised beds. *Chem. Eng. Sci.* 37, 1751–1757. [https://doi.org/10.1016/0009-](https://doi.org/10.1016/0009-2509(82)80047-X)
641 [2509\(82\)80047-X](https://doi.org/10.1016/0009-2509(82)80047-X)

642 Munari, C., Scoponi, M., Mistri, M., 2017. Plastic debris in the Mediterranean Sea: Types,
643 occurrence and distribution along Adriatic shorelines. *Waste Manag.* 67, 385–391.
644 <https://doi.org/10.1016/j.wasman.2017.05.020>

645 Nasreddine, L., Hwalla, N., Sibai, A., Hamzé, M., Parent-Massin, D., 2006. Food consumption
646 patterns in an adult urban population in Beirut, Lebanon. *Public Health Nutr.* 9, 194–203.
647 <https://doi.org/10.1079/PHN2005855>

648 Nelms, S.E., Duncan, E.M., Broderick, A.C., Galloway, T.S., Godfrey, M.H., Hamann, M., Lindeque,
649 P.K., Godley, B.J., 2016. Plastic and marine turtles: a review and call for research. *ICES J. Mar.*
650 *Sci.* 73, 165–181. <https://doi.org/10.1093/icesjms/fsv165>

651 O'Connor, I.A., Golsteijn, L., Hendriks, A.J., 2016. Review of the partitioning of chemicals into
652 different plastics: Consequences for the risk assessment of marine plastic debris. *Mar. Pollut.*
653 *Bull.* 113, 17–24. <https://doi.org/10.1016/j.marpolbul.2016.07.021>

654 Panti, C., Giannetti, M., Baini, M., Rubegni, F., Minutoli, R., Fossi, M. C., 2015. Occurrence,
655 relative abundance and spatial distribution of microplastics and zooplankton NW of Sardinia in
656 the Pelagos Sanctuary Protected Area, Mediterranean Sea. *Environ. Chem.* 12(5), 618-626.
657 <https://doi.org/10.1071/EN14234>

658 Pedrotti, M.L., Petit, S., Elineau, A., Bruzaud, S., Crebassa, J.C., Dumontet, B., Martí, E., Gorsky,
659 G., Cózar, A., 2016. Changes in the floating plastic pollution of the mediterranean sea in relation
660 to the distance to land. *PLoS One* 11, 1–14. <https://doi.org/10.1371/journal.pone.0161581>

661 Piperagkas, O., Papageorgiou, N., Karakassis, I., 2019. Qualitative and quantitative assessment
 662 of microplastics in three sandy Mediterranean beaches, including different methodological
 663 approaches. *Estuar. Coast. Shelf Sci.* 219, 169–175. <https://doi.org/10.1016/j.ecss.2019.02.016>

664 Phuong, N.N., Zalouk-Vergnoux, A., Kamari, A., Mouneyrac, C., Amiard, F., Poirier, L., Lagarde, F.,
 665 2018. Quantification and characterization of microplastics in blue mussels (*Mytilus edulis*):
 666 protocol setup and preliminary data on the contamination of the French Atlantic coast. *Environ.*
 667 *Sci. Pollut. Res.* 25, 6135–6144. <https://doi.org/10.1007/s11356-017-8862-3>

668 PlasticsEurope, 2018. *Plastics – the Facts 2017, An Analysis of European Plastics Production,*
 669 *Demand and Waste Data*, p. 16, (Brussels – Belgium)

670 Poeta, G., Battisti, C., Acosta, A.T.R., 2014. Marine litter in Mediterranean sandy littorals: Spatial
 671 distribution patterns along central Italy coastal dunes. *Mar. Pollut. Bull.* 89, 168–173.
 672 <https://doi.org/10.1016/j.marpolbul.2014.10.011>

673 Praagh, M. van, Hartman, C., Brandmyr, E., 2018. Microplastics in Landfill Leachates in the
 674 Nordic Countries. <https://doi.org/10.6027/TN2018-557>

675 a) Renzi, M., Guerranti, C., Blašković, A., 2018. Microplastic contents from maricultured and
 676 natural mussels. *Mar. Pollut. Bull.* 131, 248–251.
 677 <https://doi.org/10.1016/j.marpolbul.2018.04.035>

678 b) Renzi, M., Blašković, A., Bernardi, G., Russo, G.F., 2018. Plastic litter transfer from sediments
 679 towards marine trophic webs: A case study on holothurians. *Mar. Pollut. Bull.* 135, 376–385.
 680 <https://doi.org/10.1016/j.marpolbul.2018.07.038>

681 c) Renzi, M., Blašković, A., Fastelli, P., Marcelli, M., Guerranti, C., Cannas, S., Barone, L., Massara,
 682 F., 2018. Is the microplastic selective according to the habitat? Records in amphioxus sands,
 683 Mäerl bed habitats and *Cymodocea nodosa* habitats. *Mar. Pollut. Bull.* 130, 179–183.
 684 <https://doi.org/10.1016/j.marpolbul.2018.03.019>

685 a) Renzi, M., Specchiulli, A., Blašković, A., Manzo, C., Mancinelli, G., Cilenti, L., 2019. Marine
 686 litter in stomach content of small pelagic fishes from the Adriatic Sea: sardines (*Sardina*

687 *pilchardus*) and anchovies (*Engraulis encrasicolus*). Environ. Sci. Pollut. Res. 26, 2771–2781.
688 <https://doi.org/10.1007/s11356-018-3762-8>

689 b) Renzi, M., Čižmek, H., Blašković, A., 2019. Marine litter in sediments related to ecological
690 features in impacted sites and marine protected areas (Croatia). Mar. Pollut. Bull. 138, 25–29.
691 <https://doi.org/10.1016/j.marpolbul.2018.11.030>

692 Rios-Fuster, B., Alomar, C., Compa, M., Guijarro, B., Deudero, S., 2019. Anthropogenic particles
693 ingestion in fish species from two areas of the western Mediterranean Sea. Mar. Pollut. Bull.
694 144, 325–333. <https://doi.org/10.1016/j.marpolbul.2019.04.064>

695 Romeo, T., Pietro, B., Pedà, C., Consoli, P., Andaloro, F., Fossi, M.C., 2015. First evidence of
696 presence of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. Mar. Pollut.
697 Bull. 95, 358–361. <https://doi.org/10.1016/j.marpolbul.2015.04.048>

698 Ruiz-Orejón, L.F., Sardá, R., Ramis-Pujol, J., 2016. Floating plastic debris in the Central and
699 Western Mediterranean Sea. Mar. Environ. Res. 120, 136–144.
700 <https://doi.org/10.1016/j.marenvres.2016.08.001>

701 Schmidt, N., Thibault, D., Galgani, F., Paluselli, A., Sempéré, R., 2018. Occurrence of
702 microplastics in surface waters of the Gulf of Lion (NW Mediterranean Sea). Prog. Oceanogr.
703 163, 214–220. <https://doi.org/10.1016/j.pocean.2017.11.010>

704 Shiber, J.G., 1979. Plastic pellets on the coast of Lebanon. Mar. Pollut. Bull. 10, 28–30.
705 [https://doi.org/10.1016/0025-326X\(79\)90321-7](https://doi.org/10.1016/0025-326X(79)90321-7)

706 Suaria, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte, G., Moore, C.J., Regoli, F.,
707 Aliani, S., 2016. The Mediterranean Plastic Soup: Synthetic polymers in Mediterranean surface
708 waters. Sci. Rep. 6, 1–10. <https://doi.org/10.1038/srep37551>

709 Thompson, R.C., Olson, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D.,
710 Russell, A.E., 2004. Lost at Sea: Where Is All the Plastic? Science 304, 838.
711 <https://doi.org/10.1126/science.1094559>

712 Turner, A., Holmes, L., 2011. Occurrence, distribution and characteristics of beached plastic
 713 production pellets on the island of Malta (central Mediterranean). *Mar. Pollut. Bull.* 62, 377–
 714 381. <https://doi.org/10.1016/j.marpolbul.2010.09.027>
 715 UNEP (2014) UNEP year book 2014. Emerging issues update. Chapter 8: plastic debris in the
 716 ocean. ISBN: 978-92-807-3381-5
 717 Van Cauwenberghe, L., Janssen, C.R., 2014. Microplastics in bivalves cultured for human
 718 consumption. *Environ. Pollut.* 193, 65–70. <https://doi.org/10.1016/j.envpol.2014.06.010>
 719 van der Hal, N., Ariel, A., Angel, D.L., 2017. Exceptionally high abundances of microplastics in the
 720 oligotrophic Israeli Mediterranean coastal waters. *Mar. Pollut. Bull.* 116, 151–155.
 721 <https://doi.org/10.1016/j.marpolbul.2016.12.052>

Figure 1: Localization of the sampling sites (Tripoli, Beirut and Sidon) along the Lebanese coast.

Figure 2: A) Concentration of suspected microplastics (MPs/m³) according to their shapes found in the surface water; B) Suspected microplastics (MPs) and macroplastics (MaPs) concentration (items/g) found in one gram of sublittoral sediments; C) Average number ($\pm$ SD) of ingested suspected microplastics by shape categories in the digestive tract of (E) European anchovies (MP/ind) and in the soft tissues of (O) spiny oysters (MP/gram) at the three sampling sites (* indicate statistically significant difference).

Figure 3: Microplastics of different shapes and colors collected along the Lebanese coast in three different matrices: e.g. A) blue microbead and B) green fiber found in the collected surface water, C) Macro and microplastics collected on the 2 mm sieve and D) white and black pellets collected in Tripoli sediments sample, E) blue fiber and F) blue fragment observed on biota filter.

A)

B)

C)

Figure 4: Proportion of size class composition of suspected microplastics in the A) Water surface, B) anchovies and C) spiny oysters from the different sampling sites

Figure 5: Proportion of different polymers identified by micro-Raman spectroscopy in the three analyzed compartments: A) water; B) Sediments, C) anchovies and (D) spiny oysters.

Table 1: Literature data on microplastics abundance in Mediterranean surface waters and instrumental analysis methods used for detection of MPs. To compare with our results, only studies expressing concentrations in m³ are presented.

Location	Sampling nets	Mesh size (μm)	Identification method	MPs abundances	Reference
NW Mediterranean (the Rhône River and the Têt River)	Manta trawl net	333	Stereomicroscope/FTIR	0.18- 0.19 items/m ³	Constant et al., (2018)
NW Mediterranean Sea (Ligurian Sea and Sardinian Sea)	WP2 standard net	200	Stereomicroscope	0.62 ± 2 items/m ³	Fossi et al., (2012)
NW Mediterranean Sea (Gulf of Lion)	WP2 plankton net	200	Stereomicroscope/FTIR	0.23 ± 0.20 items/m ³	Lefebvre et al., (2019)
NW Mediterranean Sea (Gulf of Asinara and the Sardinian Sea)	Neuston net	200	Stereomicroscope	0.31 ± 1.17 items/m ³	Fossi et al., (2016)
NW Mediterranean Sea (North	WP2 standard ring net	200	Stereomicroscope	0.17 ± 0.32	Panti et al., (2015)

Western of Sardinia; Gulf of Asinara)				items/m ³	
NW Mediterranean Sea (Tuscany coastal water)	Manta trawl net	330	Stereomicroscope/FTIR	0.26/m ³	Baini et al., (2018)
Central-Western Mediterranean Sea (Sardinian Sea)	Manta trawl net	500	Binocular stereomicroscope	0.15 items/m ³	De Lucia et al., (2014)
Eastern Mediterranean (Turkish Mediterranean Coast)	Manta trawl net	333	Microscopic observation/FTIR on 24 items	0.7 items/m ³	Güven et al., (2017)
Eastern Mediterranean (Iskenderun Bay)	Manta trawl net	333	Microscopic observation	7.26 items/m ³	Gündoğdu, (2017)
Eastern Mediterranean Sea (Iskenderun Bay and Mersin Bay)	Manta Trawl net	333	Microscopic observation	2.73 items /m ³	Gündoğdu and Çevik, (2017)

South Eastern Mediterranean Sea (Israeli surface waters)	Manta Trawl net	333	Stereomicroscope	7.68 ± 2.38 items/m ³	van der Hal et al., (2017)
Lebanese coast	Manta Trawl net	52	Stereomicroscope/Raman	4.3 ± 2.2 items/m ³	Present study

Table 2: Average ($\pm$ SD) of ingested items found in the digestive track of European anchovies (*Engraulis encrasicolus*) collected in different areas of the Mediterranean Sea.

Sites	Digestion method	Concentration (items/ind)	Occurrence	Polymers Identification method	Main polymer	Reference
Spanish Western Mediterranean coasts	None. Visual observation	0.07 ($\pm$ 0.35) to 0.33 ($\pm$ 0.26)	15%	FTIR	PET	Compa et al., (2018)
Western Mediterranean Sea (Iberian Peninsula coast and the Balearic Islands)	None Visual observation	0.07 $\pm$ 0.26	2.56%	None	-	Rios-Fuster et al., (2019)
North Western of the Mediterranean Sea (Gulf of Lions)	None. Visual observation	0.11 $\pm$ 0.31	11%	ATR-FTIR	PET	Lefebvre et al., (2019)
Northwestern Mediterranean Sea	NaClO	0.85	40%	Raman spectroscopy	PE, PP, and PET	Collard et al., (2015)
Northern Mediterranean (Adriatic Sea)	H ₂ O ₂	1.25	91%	μ -FT-IR	PVC	Renzi et al., (2019a)
Eastern Mediterranean Sea (Lebanese coast)	KOH 10%	2.5 $\pm$ 0.3	83.4%	Raman Spectroscopy	PS	Present study