



Taxonomic diversity of lianas and vines in forest fragments of southern Togo

Kouami Kokou, Pierre Couteron, Arnaud Martin, Guy Caballe

► To cite this version:

Kouami Kokou, Pierre Couteron, Arnaud Martin, Guy Caballe. Taxonomic diversity of lianas and vines in forest fragments of southern Togo. *Revue d'Écologie*, 2002, 57 (1), pp.3-18. hal-03529961

HAL Id: hal-03529961

<https://hal.science/hal-03529961>

Submitted on 17 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TAXONOMIC DIVERSITY OF LIANAS AND VINES IN FOREST FRAGMENTS OF SOUTHERN TOGO

Kouami KOKOU¹, Pierre COUTERON², Arnaud MARTIN³ & Guy CABALLÉ³⁺

RÉSUMÉ

Ce travail analyse la contribution des plantes grimpantes, ligneuses et herbacées, à la biodiversité des îlots forestiers du sud du Togo. Sur la base d'un inventaire floristique général (649 espèces) couvrant 17,5 ha dans 53 îlots, 207 espèces de lianes, herbacées grimpantes et arbustes grimpants ont été recensées, soit 32 % de la flore (représentant 135 genres et 45 familles). La plupart sont de petite taille, traînant sur le sol ou s'accrochant à des arbres et arbustes ne dépassant pas 8 m de hauteur.

Une analyse factorielle des correspondances a permis de caractériser chacun des trois types d'îlots existants (forêt littorale, forêt semi-caducifoliée et galerie forestière) par plusieurs groupements exclusifs de plantes grimpantes. La dominance des herbacées grimpantes et des arbustes grimpants (132 espèces) sur les lianes *sensu stricto* (75 espèces de ligneuses grimpantes) est révélatrice de forêts plutôt basses, à canopée irrégulière. Environ 60 % des plantes grimpantes du sud Togo sont communes aux forêts tropicales de la côte ouest africaine.

SUMMARY

This work analyses the contribution of climbing plants to the biodiversity of forest fragments in southern Togo, West Africa. Based on a general floristic inventory totalling 17.5 ha of 53 forest fragments, there were found to be a total of 649 species; lianas, vines or climbing shrubs represented 135 genera in 45 families, *i.e.* 207 species or 32 % of the plant diversity. The majority of these climbing plants were small, grew along the ground or were attached to shrubs or trees no more than 8 m in height.

When only climbing plants were considered, it was possible to distinguish three fragment forest types (littoral, semi-deciduous, gallery) using correspondence analysis. The dominance of non-lianas (132 species of vines and climbing shrubs) over lianas *sensu stricto* (75 species) is indicative of relatively low forest having an irregular canopy. Approximately 60 % of climbing plants in southern Togo are common to the tropical forests of the western coast of Africa.

¹ Laboratoire de Botanique et d'Écologie végétale, Faculté des Sciences, Université du Bénin, BP 1515, Lomé, Togo.

² ENGREF, UMR 5120 du CNRS, 648 rue J.F. Breton, 34033 Montpellier cedex 1, France.

³ École Pratique des Hautes Études, UMR 5120 du CNRS, Parc Scientifique Agropolis TA 40/PS 2, boulevard de la Lironde, 34398 Montpellier cedex 5, France.

⁺ To whom correspondence should be addressed. Fax: +33 4 67 61 56 68; E-mail: caballe@cirad.fr

INTRODUCTION

Although lianas (woody climbers) and vines (herbaceous climbers) are characteristic elements of tropical forests, their role in forest dynamics remains poorly understood (Putz, 1984; Gentry & Dodson, 1987; Putz & Chai, 1987; Gentry, 1991; Hegarty & Caballé, 1991; Balfour & Bond, 1993). Paucity of information concerning lianas partly results from the difficulty to count them correctly because of their long, often tangled aerial extensions and multiple rooting (Caballé, 1977, 1980, 1994; Peñalosa, 1984; Garcia Alvez, 1989; Castellanos *et al.*, 1992; Putz, 1995; Troy *et al.*, 1997). Furthermore, numbers and density of lianas vary considerably from continent to continent, forest to forest and even from site to site within the same forest, making it very difficult to interpret results of studies, even if climbing plants are surveyed (Emmons & Gentry, 1983; Caballé, 1984; Putz, 1990; Faber-Langendoen & Gentry, 1991; Gentry, 1993; Siebert, 1993; Phillips & Gentry, 1994; Makana *et al.* 1998).

Liana and vine settlement in a given location requires a sufficient number of suitably-sized supports and full light exposition. In forests, the first of these conditions is often easily met as trees offer an immense array of forms both in terms of size and morphology (Putz, 1985, 1995; Clark & Clark, 1990; Heaney & Proctor, 1990; Coudurier, 1992; Yap *et al.*, 1995); and, the second condition is fulfilled mostly along forest margins or ecotones (Florence, 1981; Caballé, 1986; Lee, 1988; Castellanos, 1991; Gartner, 1991; Lee & Richards, 1991; During *et al.*, 1994). In continuous forest blocks, such situations may be provided by a concentration of treefall gaps and/or by special topographic features (*e.g.* riverbanks or uplands). In contrast, fragmentation of forest blocks allows the diversification of edges and augments both number and area of sites that are favourable for the development of vine and liana communities (Williams-Linera, 1990; Laurance, 1991; Crome, 1993; Jarvis, 1995).

The present study aims at assessing the importance of liana and vine communities for forest fragments located in the "Dahomey gap" that marks an interruption in the continuous forest block of West Africa (Jenik, 1994; Maley, 1996). To our knowledge, this is the first study dealing with a forest environment that offers a high concentration of favourable sites for climber communities. The contribution of lianas and vines to plant diversity is here analysed. Diversity of lianas and vines communities is examined in the light of paleogeographical characteristics of forest fragments in the Dahomey gap.

MATERIALS AND METHODS

VEGETATION OF SOUTHERN TOGO

Togo is situated in the Dahomey gap, which represents the interruption of the West African tropical forest at the level of the Gulf of Guinea (IUCN, 1992). No large tracts of forest are observable in Togo except the mountain forests of the South-West (Akpagana, 1992). However, there are numerous forest fragments, particularly in the south (Kokou, 1998; Kokou *et al.*, 1999b and c). Most of these forest fragments are protected by local people either as sacred sites or as firebreaks. In "sacred groves", local people conduct ceremonies to enter into

contact with gods or deceased ancestors (Kokou *et al.*, 1999a). The “firebreak” forests are close to villages and screen them from bush fires coming from neighbouring savannas.

Climate in southern Togo is sub-equatorial with a main rainy season from March to July (maximum precipitation in June), and a minor rainy season from September to November (maximum precipitation in October). In an average year, there is a geographic gradient of precipitation (L’Hote & Mahé, 1996), with rainfall increasing from the littoral (800 mm/yr.) towards the interior (1 000 - 1 250 mm/yr.). Relative humidity is constantly high and often greater than 80 %. Annual temperatures average 27° C with an annual mean variation of about 4° C.

FOREST FRAGMENTS CHARACTERIZATION

Three main types of forest fragments have been previously identified in southern Togo (Kokou *et al.*, 1999c). The first one (coastal forest) is observed on sands and coastal deposits and is homologous to the “outlier forest” described in south-east Ghana by Hall & Swaine (1981). The second type is referred to as dry semi-deciduous (Hall & Swaine, 1981), and displayed some affinities with the *Celtis* and *Triplochiton* forest type described in Ivory Coast by Guillaumet & Adjanooun (1971). In Togo, such forests are observed on soils developed on metamorphical rocks and also on Tertiary continental deposits. The last type consists in forest galleries which grow on hydromorphic soils located along the river network.

Coastal forests had a sparse canopy reaching 15 and 20 m in height, with an understorey displaying a high density of plants less than 2 m. Dry semi-deciduous forests were taller (above 20 m), with a closer canopy and a sparser understorey. Forest galleries displayed a physiognomy that was highly variable in terms of canopy height and vertical stratification, though the understorey was always very dense.

LIANAS AND VINES CHARACTERIZATION

The description of the principal categories of climbing plants is over a hundred years old (*e.g.* Darwin, 1867; Treub, 1883a and b; Schenck, 1892). Nowadays, classifications used are quite similar, differing only in detail. One important aspect that most modern classifications share is the integration of “dimensions”, functional or adaptive of the different kinds of climbers (*e.g.* Caballé, 1986; Hegarty, 1989 or Putz & Holbrook, 1991).

We classified climbing plants into three categories: woody climbing plants (lianas), herbaceous climbing plants (vines), and climbing shrubs. The latter are usually upright but can be non-self-supporting and climbing (= lianescents); they were treated as lianas (*e.g.* Caballé, 1993, 1998).

GENERAL CENSUS: COLLECTION AND ANALYSIS OF DATA

The interpretation of aerial photographs from missions IGN 1976-79 followed by field verification allowed the mapping of all the 296 forest fragments in southern Togo (under 7° north latitude). From this map, 53 forest fragments (*i.e.*




Figure 1. — Forest fragments of southern Togo.

20 % of the total) were randomly chosen for field investigations (Fig. 1). Their size ranged between 1 ha and 3 658 ha with 23 fragments less than 10 ha, 17 between 10–50 ha, and 13 greater than 50 ha. Field plots (50 m by 10 m) were established along transect lines that were laid out in order to intersect both core and edge areas in each sampled forest fragment. Three hundred and forty-four plots were established with a total area amounting to 17.2 ha.

The presence of all plant species, regardless of size and life form, was recorded in each plot. A list of all angiosperms and pteridophytes found in the plots was compiled (Kokou, 1998; Kokou *et al.*, 1999b). Species in the list were classified according to the four phytogeographical categories (*i.e.* Guineo-Congolese, transition, Sudano-Zambezian, introduced) provided by Aké Assi (1984).

Distribution of lianas and vines in the 53 sampled fragments was investigated through Correspondence Analysis (CA; Benzecri, 1973; Hill 1973a) applied to a species fragment contingency table. Each cell of this table contained the number of plots from a given fragment within which a given species was recorded.

A pairwise analysis of species co-occurrence within plots was used to gain a detailed understanding of climber associations. For each pair of climbing species, the relative frequency of co-occurrence was tested following an approach proposed by Gauthier *et al.* (1977) and based on Fisher's exact test (Siegel, 1956).

Diversity of lianas and vines has been quantified by means of usual indices (Hill, 1973b; Daget, 1980), such as:

- * Species richness (N_0), which represents the total number of species;
- * Shannon diversity index (I_{SH}), given by:

$$I_{SH} = \sum_{i=1}^n pi \log (pi)$$

where n = number of encountered species and pi = probability that a species occurrence within a plot corresponds to species i . In practice, $pi = qi/Q$ where qi is the number of plots wherein species i was observed, and $Q = \sum_{i=1}^n qi$.

RESULTS

CLIMBER DIVERSITY AND FLORISTIC RICHNESS

The complete floristic inventory yielded 649 species of which 207 or 32 % were climbers. Climbing plants of all types numerically outstrips other life forms: 14 % trees, 25 % self-supporting shrubs, 26 % upright herbs and 3 % other. Of the climbing plants, 111 were vines, 75 lianas and 21 climbing shrubs. These last can become climbers and are listed in Table I.

TABLE I

List of climbing shrub species.

<i>Byrsocarpus</i> (= <i>Rourea</i>) <i>coccineus</i> Schum. & Thonn. (Connaraceae)
<i>Cnestis ferruginea</i> DC. (Connaraceae)
<i>Connarus thonningii</i> (DC.) Schellenb. (Connaraceae)
<i>Dalbergia altissima</i> Bak. f. (Leg. Papilionoideae)
<i>Macrosphyra longistyla</i> (SW.) DC. (Rubiaceae)
<i>Maytenus ovatus</i> var. <i>ovatus</i> (Wall. ex Wight & Arn.) Loes. (Celastraceae)
<i>Mimosa pigra</i> L. (Leg. Mimosoideae)
<i>Nauclea latifolia</i> Sm. (Rubiaceae)
<i>Pancovia sessiliflora</i> Hutch. & Dalz. (Sapindaceae)
<i>Plumbago zeylanica</i> L. (Plumbaginaceae)
<i>Premna quadrifolia</i> Schum. & Thonn. (Verbenaceae)
<i>Quisqualis indica</i> L. (Combretaceae)
<i>Ritchiea duchesnei</i> (De Wild.) Keay (Capparaceae)
<i>Ruspolia hypocrateriformis</i> (Vahl.) Milne (Acanthaceae)
<i>Sorindeia warneckei</i> Engl. (Anacardiaceae)
<i>Strychnos usambarensis</i> Gilg (Loganiaceae)
<i>Uvaria chamae</i> P. Beauv. (Annonaceae)
<i>Uvaria doeringii</i> Diels (Annonaceae)
<i>Uvaria ovata</i> subsp. <i>ovata</i> (Dunal) A. DC. (Annonaceae)
<i>Uvaria sofa</i> Sc. Elliot (Annonaceae)
<i>Vangueriopsis spinosa</i> (Schum. & Thonn.) Hepper (Rubiaceae)

With regards to regional phytogeography, 60 % of the liana and vine species are related to forests of tropical Africa (*i.e.* Guineo-Congolese), 32 % are characteristic of the forest-savanna transition, 3 % are typical of savannas that encircle the forest fragments (*i.e.* Sudano-Zambesian), and 5 % represent introduced species.

The 207 climbing plant species are members of 135 genera and 45 families. Leguminosae (Papilionoideae), Asclepiadaceae, Rubiaceae, Cucurbitaceae, Apocynaceae and Convolvulaceae account for the greatest number of genera and species (Table II). For all forest fragments, the most frequent lianas were *Cissus petiolata* (recorded in 197 plots), *Byrsocarpus coccineus* (195), *Secamone afzelii* (180), *Grewia carpinifolia* (170) and *Triclia subcordata* (162). These are all indigenous species.

TABLE II
The ten most common plant families.

Family	Number of	
	genera	species
Leg. Papilionoideae	18	27
Asclepiadaceae	12	18
Rubiaceae	12	13
Cucurbitaceae	10	13
Apocynaceae	6	10
Convolvulaceae	5	17
Leg. Mimosoideae	5	6
Menispermaceae	5	5
Vitaceae	4	10
Annonaceae	3	6
Others	55	82
Total	135	207

Liana and vine richness did not correlate with the number of plots inventoried (Table III) nor with the surface area of the forest fragments (Spearman's rank correlation, $r_s = 0.34$, $n = 53$, $p < 0.0001$), nor even with the distance to small cities and villages ($r_s = 0.26$, $n = 53$, $p < 0.0001$). Gallery forests were the least inventoried (86 plots) and yet they supported the greatest climbing plant richness and highest diversity as measured by the Shannon Index ($I_{SH} = 1.89$). Notwithstanding, the share of each forest fragment type was about one third of the total number of climbing plants.

SIZE AND LAYERING

Among the lianas and vines inventoried, very few (9 %) were large and reached the canopy of trees greater than 8 m in height. Most (65 %) grew attached

TABLE III

Plant species richness, and liana and vine diversity according to forest type.

Forest type	Number of			Shannon diversity index (I_{SH})	Number of		Most common families (number of genera, species)
	plots	climbers	species		genera	families	
Littoral	146	219	69	1.62	58	28	Asclepiadaceae (9, 10) Leg. Papilionoideae (7, 9) Cucurbitaceae (5, 6) Vitaceae (4, 5) Rubiaceae (3, 4) Capparaceae (2, 4)
Semi-deciduous	112	441	142	1.84	97	36	Leg. Papilionoideae (13, 21) Asclepiadaceae (10, 14) Rubiaceae (8, 9) Apocynaceae (5, 7) Vitaceae (4, 9) Convolvulaceae (4, 8) Annonaceae (3, 6)
Gallery	86	499	154	1.89	108	41	Leg. Papilionoideae (13, 20) Rubiaceae (12, 13) Asclepiadaceae (10, 15) Apocynaceae (6, 10) Convolvulaceae (5, 10) Menispermaceae (5, 5) Cucurbitaceae (4, 6) Annonaceae (3, 6)

to or supported by trees and shrubs between 2-8 m high. The remaining (26 %) grew along the ground. This heavy occupation of the understorey by lianas and vines is reflected in the fact that 96 % of forest fragments have relatively closed lower stratas that impeded human penetration. The ground-cover by liana and vine was 70 - 90 % (of total area inventoried), as estimated by orthogonal projection.

FOREST TYPES AND ASSOCIATIONS OF CLIMBING PLANTS

Figure 2 shows the distribution of the 53 forest fragments in the first factorial plane of the correspondence analysis. The first axis (8.13 % of total inertia) discriminates the gallery forests. The second axis (5.96 %) differentiates between littoral and semi-deciduous forests. The three forest types are hence characterized on the sole basis of the climbing plants (207 species) they harbour. Moreover, very similar results were obtained when the correspondence analysis relied on the entire flora (649 species) of the forest fragments (Kokou *et al.*, 1999b, c; Kokou & Caballé, 2000). Furthermore, the relative frequency of the most frequent climber species depends upon forest fragment type (Table IV).

Tests on species co-occurrences enabled a more precise distinction between climber associations within each type of forest fragments (appendix). In coastal




Figure 2. — Results of correspondence analysis of 53 forest fragments (in 1 - 2 factorial plane). Number refers to forest fragment.

TABLE IV

Most frequent plant species according to forest type.

Species (Family)	Littoral	Semi-deciduous	Gallery
<i>Dichapetalum pallidum</i> (Oliv.) Engl. (Dichapetalaceae)	+++	-	-
<i>Grewia carpinifolia</i> Juss. (Tiliaceae)	+++	-	-
<i>Ritchiea reflexa</i> (Thonn.) Gilg & Benedict (Capparaceae)	+++	-	-
<i>Secamone afzelii</i> (Schultes) K. Schum. (Asclepiadaceae)	+++	++	-
<i>Uvaria chamae</i> P. Beauv. (Annonaceae)	+++	-	-
<i>Sorindeia warneckei</i> Engl. (Anacardiaceae)	++	-	+
<i>Byrsocarpus coccineus</i> Schum. & Thonn. (Connaraceae)	-	+++	-
<i>Cissus petiolata</i> Hook. f. (Vitaceae)	-	+++	-
<i>Motandra guineensis</i> A. DC. (Apocynaceae)	-	+++	-
<i>Griffonia simplicifolia</i> (Vahl. ex DC.) Baill. (Leg. Caesalpinoideae)	-	++	++
<i>Paullinia pinnata</i> L. (Sapindaceae)	-	-	+++
<i>Tricilia subcordata</i> Oliv. (Menispermaceae)	-	-	+++
<i>Combretum afzelii</i> Engl. & Diels (Combretaceae)	-	-	++
<i>Cremaspora triflora</i> (Thonn.) K. Schum. (Rubiaceae)	-	-	+
<i>Flagellaria guineensis</i> Schumach. (Flagellariaceae)	-	-	+
<i>Gymnema sylvestre</i> (Retz.) Schultes (Asclepiadaceae)	-	-	+

Species present in inventories: +++ > 50 %, ++ > 40 %, + > 30 %, - < 30 %.

forests, four associations were identified on the basis of 15 species. In semi-deciduous forests, 9 associations were defined on the basis of 40 species. One association containing *Momordica charantia* was identified as peripheral to the fragments. This species often covers trees along edges of fragments or around clearings, and then withers and entirely disappears during the dry season (Kokou & Caballé, 2000). In other semi-deciduous forests, the liana *Cardiospermum grandiflorum* displays the same behaviour. In gallery forests, there were 9 associations defined from 45 climber species (appendix).

DISCUSSION

In southern Togo, climbing plant richness (207 species) is high compared to the overall forest flora (649 species). The only known examples of areas similarly rich in climbing plants are secondary or perturbed forests (Kahn, 1982; Putz & Chai, 1987; Pinard & Putz, 1994; Dewalt *et al.*, 2000). Forest fragments of southern Togo are effectively protected by local people and the multiplication of forest edges is probably the factor that determines the development of climbing plant communities.

Climbing plants of southern Togo represent about one third of the flora, of which lianas *sensu stricto* (75 species) make up only 11.6 % while vines (111 species) contribute 17.1 % and climbing shrubs (21 species) 3.2 %. The comparatively high percentage of vines and climbing shrubs as opposed to lianas is indicative of largely open forest environments lacking a true high, closed canopy. Most of the vines and lianas were climbing (< 8 m) trees or shrubs. Another indication of the openness of the upper strata of fragment forests comes from the prevalence of vines and, to a lesser extent, lianas from the families Asclepiadaceae, Cucurbitaceae, Convolvulaceae and Vitaceae (Table II). Conversely, tall African forests with closed canopies are characterized by few vines and more lianas Apocynaceae, Hippocrateaceae, Dichapetalaceae, Annonaceae, Combretaceae, and Loganiaceae (see Caballé, 1986; Gentry, 1991). The presence of a relatively high number of climbing shrub species is also a characteristic of forests with a low and irregular canopy. In the gallery forests, climbing plant diversity is at a maximum and may be explained by the amplification of edge effects. On the other hand, littoral forest fragments are the least rich (Table III), and their lianescence flora conforms to the general rule that littoral and coastal areas are floristically poor.

The estimation of abundance and richness of climbing plants in an area can vary enormously depending upon whether inventorying included lianas of small diameter (*e.g.* DBH < 2.5 cm), climbing herbaceous plants (vines), and certain forms of hemi-epiphytes and epiphytes. It is evident from published literature that vines and lianas are frequently not recognized as distinct life forms. Furthermore, life forms or growth habits are rarely clearly defined or differentiated (*e.g.* hemi-epiphytes are often confounded with lianas) and usually only plants with DBH \geq 5 or 10 cm are inventoried (Hegarty & Caballé, 1991; Orians *et al.*, 1996). These facts make it difficult to interpret different results and to compare different studies. Future progress in the understanding of climbing plant ecology depends upon resolving these conceptual and technical discrepancies. A universal agreement upon definitions of climbing plant strategies and harmonized inventory protocols is imperatively required.

Jacobs (1976) estimated that lianas compose 8 % of the tropical flora world-wide. Gentry & Dodson (1987) found that of the flora of seven humid tropical forest sites, 19 % of the species were lianas. Denslow (1996) found that lianas and vines made up 10.3 % of deciduous moist forest (Barro Colorado, Panama) and 22.5 % of premontane wet forest (La Selva, Costa Rica). At two dry forest sites in Central America, Lott (1985) reported the composition of lianescence species to be 24 %. In two evergreen forests in the north of the Democratic Republic of Congo (formerly Zaire), Makana *et al.* (1998) estimated liana richness

to be about 30 %. Caballé (1986) found that in the north-east of Gabon in evergreen forest naturally perturbed by numerous treefall gaps, lianas can represent up to 37 % of species having a DBH \geq 5 cm (10 ha inventoried). In the same forest in north-eastern Gabon, Hladik (1974, 1978) found that liana leaves formed 36 % of annual litter production. Conversely, in the Chocó in Colombia, one of the richest forests in the world, Faber-Langendoen & Gentry (1991) found that lianas (11 species/ha) represented less than one per cent of tree species richness (250 species/ha).

Perhaps the most surprising result of this study is that a discrimination between forest types can be achieved on the sole basis of the floristic composition of climbing vegetation. Each forest type can be defined by several exclusive associations of climbers: 4 for littoral, 9 for semi-deciduous and 9 for gallery forest fragments (appendix). Such a result suggests that the dynamism of climbing plant communities is not solely determined by the number of plant interfaces and edges created by perturbation and/or fragmentation of forest areas (Williams-Linera, 1990; Laurance, 1991; Crome, 1993; Phillips & Gentry, 1994; Jarvis, 1995). Albeit that the majority of climbing plants depend upon other plants to develop, they are just as important a life form as any other making up the plant community. Climbing plants are an intrinsic and important component of the biodiversity of tropical areas. This is even more true in the African continent, which possesses the largest number of lianas (Putz & Mooney, 1991).

APPENDIX

Groups of lianas and vines according to three forest types.

Littoral

Species	Sign	Freq	Species	Sign	Freq
<i>Dichapetalum pallidum</i>		89	<i>Opilia amentacea</i>		81
<i>Sorindeia warneckeii</i>	+++	127	<i>Uvaria chamae</i>	+++	138
<i>Uvaria chamae</i>	+++	138	<i>Ancylobothrys scandens</i>	+++	101
<i>Ancylobothrys scandens</i>	+++	101	<i>Ritchiea reflexa</i>	+++	134
<i>Capparis erythrocarpus</i>	+++	87	<i>Dichapetalum pallidum</i>	+++	89
<i>Ritchiea reflexa</i>	+++	134	<i>Sorindeia warneckeii</i>	+	127
<i>Ruspolia hypocrateriformis</i>	+	35	<i>Capparis erythrocarpus</i>	+	87
<i>Strychnos barteri</i>		80	<i>Grewia carpinifolia</i>		170
<i>Ruspolia hypocrateriformis</i>	+++	35	<i>Secamone afzelii</i>	+++	180
<i>Landolphia togolana</i>	+++	94	<i>Ritchiea reflexa</i>	+++	134
<i>Capparis erythrocarpus</i>	+++	87	<i>Uvaria chamae</i>	+	138
<i>Dichapetalum pallidum</i>	+++	89	<i>Uvaria ovata ovata</i>	+	23
<i>Ancylobothrys scandens</i>	++	101	<i>Adenia lobata</i>	+	119

Semi-deciduous

Species	Sign	Freq	Species	Sign	Freq	Species	Sign	Freq
<i>Uvaria sofa</i>		45	<i>Premna quadrifolia</i>		61	<i>Dioscorea preussii</i>		28
<i>Barleria opaca</i>	+++	40	<i>Barleria opaca</i>	+++	40	<i>Motandra guineensis</i>	+++	85
<i>Artobotrys velutinus</i>	+++	48	<i>Motandra guineensis</i>	+++	85	<i>Cryptolepis sanguilenta</i>	+++	19
<i>Motandra guineensis</i>	+++	85	<i>Strophanthus hispidus</i>	+++	37	<i>Mondia withei</i>	+++	35
<i>Strophanthus hispidus</i>	+++	37	<i>Ritchiea capparoides</i>	+++	61	<i>Parquetina nigrescens</i>	+++	61
<i>Griffonia simplicifolia</i>	+++	103	<i>Tragia benthamii</i>	+++	19	<i>Combretum racemosum</i>	+++	49
<i>Momordica charantia</i>	+++	61	<i>Tragia senegalensis</i>	+++	36	<i>Strychnos sourensis</i>	+++	49
<i>Tragia benthamii</i>	+++	19	<i>Triaspis odorata</i>	+++	26	<i>Adenia cissampeloides</i>	+++	59
<i>Hippocratea indica</i>	+++	102	<i>Rhynchosia buettneri</i>	+++	30	<i>Cremaspora triflora</i>	+++	54
<i>Strychnos sourensis</i>	+++	49	<i>Adenia cissampeloides</i>	+++	59	<i>Premna quadrifolia</i>	+++	61
<i>Jasminum pauciflorum</i>	+++	63	<i>Mussaenda elegans</i>	+++	31	<i>Barleria opaca</i>	++	40
<i>Rhynchosia densiflora</i>	+++	51	<i>Artobotrys velutinus</i>	++	48	<i>Artobotrys velutinus</i>	+	48
<i>Mussaenda elegans</i>	+++	31	<i>Parquetina nigrescens</i>	++	61	<i>Cayratia gracilis</i>	+	13
<i>Premna quadrifolia</i>	+++	61	<i>Griffonia simplicifolia</i>	++	103	<i>Asparagus warneckeii</i>		83
<i>Asparagus warneckeii</i>	+++	83	<i>Rhynchosia densiflora</i>	++	51	<i>Barleria opaca</i>	+++	40
<i>Strophanthus sarmentosus</i>	++	23	<i>Mucuna pruriens</i>	+	20	<i>Artobotrys velutinus</i>	+++	48
<i>Triaspis odorata</i>	++	26	<i>Cremaspora triflora</i>	+	54	<i>Motandra guineensis</i>	+++	85
<i>Adenia cissampeloides</i>	++	59	<i>Ritchiea duchesnei</i>		17	<i>Strophanthus hispidus</i>	+++	37
<i>Parquetina nigrescens</i>	+	61	<i>Barleria opaca</i>	+++	40	<i>Ritchiea capparoides</i>	+++	61
<i>Ritchiea capparoides</i>	+	61	<i>Motandra guineensis</i>	+++	85	<i>Jasminum pauciflorum</i>	+++	63
<i>Dioscorea togoensis</i>		60	<i>Ritchiea capparoides</i>	+++	61	<i>Rhynchosia densiflora</i>	+++	51
<i>Uvaria doeringii</i>	+++	12	<i>Strychnos sourensis</i>	+++	49	<i>Premna quadrifolia</i>	+++	61
<i>Cryptolepis sanguilenta</i>	+++	19	<i>Mussaenda elegans</i>	+++	31	<i>Capparis thonningii</i>	++	50
<i>Rhynchosia densiflora</i>	+++	51	<i>Hippocratea indica</i>	++	102	<i>Momordica charantia</i>	+	61
<i>Ampelocissus bombycina</i>	+++	18	<i>Tiliacora funifera</i>	++	106	<i>Triaspis odorata</i>	+	26
<i>Mezoneuron benthamianum</i>	++	34	<i>Asparagus warneckeii</i>	+	83	<i>Tiliacora funifera</i>	+	106
<i>Momordica charantia</i>	++	61	<i>Uvaria sofa</i>	+	45	<i>Adenia cissampeloides</i>	+	59
<i>Abrus canescens</i>	++	30	<i>Adenia cissampeloides</i>		59	<i>Cremaspora triflora</i>	+	54
<i>Cremaspora triflora</i>		54	<i>Motandra guineensis</i>	+++	85	<i>Rhynchosia buettneri</i>		30
<i>Artobotrys velutinus</i>	+++	48	<i>Momordica charantia</i>	+++	61	<i>Barleria opaca</i>	+++	40
<i>Adenia cissampeloides</i>	+++	59	<i>Artobotrys velutinus</i>	+++	48	<i>Motandra guineensis</i>	+++	85
<i>Motandra guineensis</i>	++	85	<i>Barleria opaca</i>	+++	40	<i>Parquetina nigrescens</i>	+++	61
<i>Abrus canescens</i>	++	30	<i>Strophanthus hispidus</i>	+++	37	<i>Griffonia simplicifolia</i>	+++	103
<i>Gongronema latifolium</i>	+	19	<i>Abrus canescens</i>	+++	30	<i>Tragia senegalensis</i>	+++	36
<i>Parquetina nigrescens</i>	+	61	<i>Gongronema latifolium</i>	+++	19	<i>Hippocratea indica</i>	++	102
<i>Mezoneuron benthamianum</i>	+	34	<i>Ritchiea capparoides</i>	++	61	<i>Mondia withei</i>	+	35
<i>Ritchiea capparoides</i>	+	61	<i>Hippocratea africana</i>	++	59			
<i>Tricilia subcordata</i>	+	162	<i>Rhynchosia densiflora</i>	++	51			
			<i>Tragia benthamii</i>	+	19			

Gallery

Species	Sign	Freq	Species	Sign	Freq	Species	Sign	Freq
<i>Paullinia pinnata</i>		66	<i>Flagellaria guineensis</i>		29	<i>Culcasia scandens</i>		27
<i>Gymnema sylvestre</i>	+++	37	<i>Gymnema sylvestre</i>	+++	37	<i>Gymnema sylvestre</i>	+++	37
<i>Tacazzea apiculata</i>	+++	20	<i>Tacazzea apiculata</i>	+++	20	<i>Tacazzea apiculata</i>	+++	20
<i>Tylophora sylvatica</i>	+++	27	<i>Combretum afzelii</i>	+++	30	<i>Tylophora sylvatica</i>	+++	27
<i>Combretum afzelii</i>	+++	30	<i>Ipomoea cairica</i>	+++	8	<i>Combretum zenkeri</i>	+++	4
<i>Combretum smeathmannii</i>	+++	15	<i>Merremia hederacea</i>	+++	8	<i>Connarus thonningii</i>	+++	26
<i>Connarus thonningii</i>	+++	26	<i>Stachyanthus occidentalis</i>	+++	31	<i>Lepistemon owariensis</i>	+++	29
<i>Lepistemon owariensis</i>	+++	29	<i>Triclisia subcordata</i>	+++	162	<i>Merremia hederacea</i>	+++	8
<i>Zehneria capillacea</i>	+++	20	<i>Paullinia pinnata</i>	+++	66	<i>Stachyanthus occidentalis</i>	+++	31
<i>Stachyanthus occidentalis</i>	+++	31	<i>Culcasia scandens</i>	+++	27	<i>Cissampelos mucronata</i>	+++	19
<i>Cissampelos mucronata</i>	+++	19	<i>Connarus thonningii</i>	++	26	<i>Triclisia subcordata</i>	+++	162
<i>Triclisia subcordata</i>	+++	162	<i>Piper guineense</i>	++	8	<i>Abrus pulchellus</i>	+++	30
<i>Entada mannii</i>	+++	16	<i>Combretum smeathmannii</i>	+	15	<i>Centrosema pubescens</i>	+++	14
<i>Abrus pulchellus</i>	+++	30	<i>Zehneria capillacea</i>	+	20	<i>Dioclea reflexa</i>	+++	9
<i>Centrosema pubescens</i>	+++	14	<i>Hippocratea apocynoides</i>	+	3	<i>Paullinia pinnata</i>	+++	66
<i>Diodia scandens</i>	+++	18	<i>Cissampelos mucronata</i>	+	19	<i>Cissus petiolata</i>	+++	197
<i>Ipomoea cairica</i>	++	8	<i>Dioclea reflexa</i>	+	9	<i>Combretum afzelii</i>	++	30
<i>Merremia hederacea</i>	++	8	<i>Cissus petiolata</i>	+	197	<i>Stictocardia beraviensis</i>	+	3
<i>Mucuna sloanei</i>	++	11	<i>Diodia scandens</i>		18	<i>Zehneria capillacea</i>	+	20
<i>Dioclea reflexa</i>	+	9	<i>Tacazzea apiculata</i>	+++	20	<i>Hippocratea apocynoides</i>	+	3
<i>Cissus petiolata</i>		197	<i>Tylophora sylvatica</i>	+++	27	<i>Entada mannii</i>	+	16
<i>Griffonia simplicifolia</i>	+++	103	<i>Combretum afzelii</i>	+++	30	<i>Psophocarpus palustris</i>	+	3
<i>Zehneria capillacea</i>	+++	20	<i>Lepistemon owariensis</i>	+++	29	<i>Abrus pulchellus</i>		30
<i>Cremsapora triflora</i>	+++	54	<i>Cissampelos mucronata</i>	+++	19	<i>Gymnema sylvestre</i>	+++	37
<i>Paullinia pinnata</i>	+++	66	<i>Entada mannii</i>	+++	16	<i>Tylophora sylvatica</i>	+++	27
<i>Parquetina nigrescens</i>	++	61	<i>Centrosema pubescens</i>	+++	14	<i>Tacazzea apiculata</i>	++	20
<i>Combretum afzelii</i>	++	30	<i>Connarus thonningii</i>	++	26	<i>Connarus thonningii</i>	++	26
<i>Tragia senegalensis</i>	++	36	<i>Gymnema sylvestre</i>	+	37	<i>Merremia hederacea</i>	++	8
<i>Gymnema sylvestre</i>	+	37	<i>Combretum smeathmannii</i>	+	15	<i>Zehneria capillacea</i>	++	20
<i>Combretum racemosum</i>	+	49	<i>Passiflora foetida</i>	+	9	<i>Combretum smeathmannii</i>	+	15
<i>Connarus thonningii</i>	+	26	<i>Cissampelos mucronata</i>		19	<i>Lepistemon owariensis</i>	+	29
<i>Rhynchosia buettneri</i>	+	30	<i>Gymnema sylvestre</i>	+++	37	<i>Stictocardia beraviensis</i>	+	3
<i>Premna quadrifolia</i>	+	61	<i>Tacazzea apiculata</i>	+++	20	<i>Cissampelos mucronata</i>	+	19
<i>Rhigiocarya racemifera</i>		7	<i>Tylophora sylvatica</i>	+++	27	<i>Hypselodelphys poggeana</i>		23
<i>Tetraceras alnifolia</i>	+++	8	<i>Combretum afzelii</i>	+++	30	<i>Griffonia simplicifolia</i>	+++	103
<i>Stachyanthus occidentalis</i>	++	31	<i>Combretum smeathmannii</i>	+++	15	<i>Cissus petiolata</i>	+++	197
<i>Sherbournia calycina</i>	++	5	<i>Lepistemon owariensis</i>	+++	29	<i>Culcasia scandens</i>	+++	27
<i>Cissus aralioides</i>	++	15	<i>Merremia hederacea</i>	+++	8	<i>Flagellaria guineensis</i>	+++	29
<i>Smilax kraussiana</i>	++	5	<i>Zehneria capillacea</i>	+++	20	<i>Cardiospermum grandiflorum</i>	++	15
<i>Thunbergia cynanchyfolia</i>	+	8				<i>Connarus thonningii</i>	+	26
<i>Griffonia simplicifolia</i>	+	103				<i>Hippocratea indica</i>	+	102
						<i>Stachyanthus occidentalis</i>	+	31
						<i>Paullinia pinnata</i>	+	66

Names in bold characters are those of species found to form connections in the statistical analysis. They provide the identification of each group. Freq = Frequency. Significantly positive levels (see in Sign column): +++ $p < 0.00001$, ++ $p < 0.0001$, + $p < 0.001$.

ACKNOWLEDGEMENTS

Financial support from the International Foundation for Science (Sweden) for this study is gratefully acknowledged. This paper has been finalized thanks to a 4-month assignment of Kouami Kokou at the École Nationale du Génie Rural des Eaux et des Forêts (ENGREF). The authors are obliged to Dawn Frame for critical review of the manuscript and help with the English text.

REFERENCES

- AKÉ ASSI, L. (1984). — *Flore de la Côte d'Ivoire: étude descriptive et biogéographie, avec quelques notes ethnobotaniques*. Thèse de doctorat, Université d'Abidjan.
- AKPAGANA, K. (1992). — Les forêts denses humides des Monts du Togo et Agou (République du Togo). *Bulletin du Muséum National Histoire Naturelle, sect. B, Adansonia*, 1: 109-172.
- BALFOUR, D.A. & BOND, W.J. (1993). — Factors limiting climber distribution and abundance in Southern African forest. *Journal of Ecology*, 81: 93-99.
- BENZECRI, J.-P. (ed.) (1973). — *L'analyse des données. T.2: l'analyse des correspondances*. Dunod, Paris, 632 p.
- CABALLÉ, G. (1977). — Multiplication végétative en forêt dense du Gabon de la liane *Entada sclerata* (Mimosoideae). *Adansonia*, 17: 215-220.
- CABALLÉ, G. (1980). — Caractéristiques de croissance et multiplication végétative en forêt dense du Gabon de la "liane à eau" *Tetracera alnifolia* Willd. (Dilleniaceae). *Adansonia*, 19: 467-475.
- CABALLÉ, G. (1984). — Essai sur la dynamique des peuplements de lianes ligneuses d'une forêt du nord-est du Gabon. *Revue d'Écologie (Terre Vie)*, 39: 3-35.
- CABALLÉ, G. (1986). — *Sur la biologie des lianes ligneuses en forêt gabonaise*. Thèse de doctorat d'état en sciences, Université des Sciences et Techniques du Languedoc, Montpellier.
- CABALLÉ, G. (1993). — Liana structure, function and selection: a comparative study of xylem cylinders of tropical rainforest species in Africa and America. *Botanical Journal of the Linnean Society*, 113: 41-60.
- CABALLÉ, G. (1994). — Ramet proliferation by longitudinal splitting in the Gabonese rain forest liana *Dalhousia africana* S. Moore (Papilionaceae). *Biotropica*, 26: 266-275.
- CABALLÉ, G. (1998). — Le port autoportant des lianes tropicales: une synthèse des stratégies de croissance. *Canadian Journal of Botany*, 76: 1703-1716.
- CASTELLANOS, A.E. (1991). — Photosynthesis and gas exchange of vines. Pp. 181-204, in: Putz, F.E. & Mooney, H.A. (eds.), *The biology of vines*. Cambridge University Press.
- CASTELLANOS, A.E., DURÁN, R., GUZMÁN, S., BRIONES, O. & FERIA, M. (1992). — Three-dimensional space utilization of lianas: a methodology. *Biotropica*, 24: 396-401.
- CLARK, D.B. & CLARK, D.A. (1990). — Distribution and effects on tree growth of lianas and woody hemiepiphytes in a Costa Rican wet forest. *Journal of Tropical Ecology*, 6: 321-331.
- COUDURIER, T. (1992). — *Sur la place des lianes dans la forêt guyanaise. Une approche qui utilise l'architecture végétale*. Thèse de doctorat d'université, Université de Montpellier II.
- CROME, F.H.J. (1993). — Tropical forest fragmentation: some conceptual and methodological issues. Pp. 67-76, in: Moritz, C. & Kikkawa, H. (eds.), *Conservation biology in Australia and Oceania*, pp. 61-76. Surrey Beatty & Sons, Chipping Norton, Australia.
- DAGET, P. (1980). — Le nombre de diversité de Hill, un concept unificateur dans la théorie de la diversité écologique. *Acta Oecologica/Oecologia Generalis*, 1: 51-70.
- DARWIN, C. (1867). — On the movements and habits of climbing plants. *Journal of the Linnean Society London, Bot.* 9: 1-118.
- DENSLAW, J.S. (1996). — Functional group diversity and responses to disturbance. Pp. 127-151, in: Orians, G.H., Dirzo, R. & Cushman, J.H. (eds.). *Biodiversity and ecosystem processes in tropical forests*. Springer-Verlag, Ecological Studies 122, Berlin.
- DEWALT, S.J., SCHNITZER, S.A. & DENSLAW, J.S. (2000). — Density and diversity of lianas along a chronosequence in a central Panamanian lowland forest. *Journal of Tropical Ecology*, 16: 1-19.
- DURING, H.J., KWANT, R.A. & WERGER, J.A. (1994). — Effects of light on above-ground biomass investment patterns in the vine *Lonicera periclymenum* and the shrub *Lonicera xylosteum*. *Phytocoenologia*, 24: 597-607.
- EMMONS, L.H. & GENTRY, A.H. (1983). — Tropical forest structure and the distribution of gliding and prehensile-tailed Vertebrates. *American Naturalist*, 121: 513-524.

- FABER-LANGENDOEN, D. & GENTRY, A.H. (1991). — The structure and diversity of rain forests at Bajo Calima, Chocó region, Western Colombia. *Biotropica*, 23: 2-11.
- FLORENCE, J. (1981). — *Chablis et sylvigénèse dans une forêt dense humide sempervirente du Gabon*. Thèse 3^e cycle écologie végétale, Université Louis Pasteur, Strasbourg.
- GARCIA ALVES, P. (1989). — *Croissance et distribution spatiale d'une liane ligneuse, Condylarcarpon guianense Desf. (Apocynaceae)*. Mémoire DEA, Université de Montpellier II.
- GARTNER, B.L. (1991). — Structural stability and architecture of vines vs. shrubs of poison oak, *Toxicodendron diversilobum*. *Ecology*, 72: 2005-2015.
- GAUTHIER, B., GODRON, M., HIERNAUX, P. & LE PART, J. (1977). — Un type complémentaire de profil écologique "indicé". *Canadian Journal of Botany*, 55: 2859-2865.
- GENTRY, A.H. (1991). — The distribution and evolution of climbing plants. Pp. 3-47, in: Putz, F.E. & Mooney, H.A. (eds.). *The biology of vines*. Cambridge University Press.
- GENTRY, A.H. (1993). — Diversity and floristic composition of lowland tropical forest in Africa and South America. Pp. 500-547, in: Goldblatt, P. (ed.). *Biological relationships between Africa and South America*. Yale University Press.
- GENTRY, A.H. & DODSON, C. (1987). — Contribution of nontrees to species richness of a tropical rain forest. *Biotropica*, 19: 149-156.
- GUILLAUMET, J.L. & ADIANOHOUN, E. (1971). — La végétation. Pp. 156-263, in: ORSTOM (ed.). *Le milieu naturel de la Côte d'Ivoire*. Mémoire ORSTOM n° 50, Paris.
- HALL, J.B. & SWAINE, M.D. (1981). — *Distribution and ecology of vascular plants in a rain forest vegetation in Ghana*. Geobotany 1, 383 p.
- HEANEY, A. & PROCTOR, J. (1990). — Preliminary studies on forest structure and floristics on Volcán Barva, Costa Rica. *Journal of Tropical Ecology*, 6: 307-320.
- HEGARTY, E.E. (1989). — The climbers-lianes and vines. Pp. 339-353, in: Lieth, H. & Werger, M.J.A. (eds.). *Tropical rain forest ecosystems*. Elsevier Science Publishers B. V., Amsterdam.
- HEGARTY, E.E. & CABALLÉ, G. (1991). — Distribution and abundance of vines in forest communities. Pp. 238-250, in: Putz, F.E. & Mooney, H.A. (eds.). *The biology of vines*. Cambridge University Press.
- HILL, M. (1973a). — Reciprocal averaging: an eigenvector method of ordination. *Journal of Ecology*, 61: 237-249.
- HILL, M. (1973b). — Diversity and evenness: a unifying notation and its consequences. *Ecology*, 45: 427-432.
- HLADIK, A. (1974). — Importance des lianes dans la production foliaire de la forêt équatoriale du Nord-Est du Gabon. *Comptes Rendus de l'Académie des Sciences*, 278: 2527-2530.
- HLADIK, A. (1978). — Phenology of leaf production in rain forest of Gabon: distribution of food for folivores. Pp. 51-71, in: Montgomery, G.G. (ed.). *The ecology of arboreal folivores*. Smithsonian Institution Press.
- IUCN (Sayer, J.A., Harcourt, C.S. & Collins, N.M., eds.) (1992). — *The conservation atlas of tropical forest Africa*. Macmillan Publishers Ltd.
- JACOBS, M. (1976). — The study of lianas. *Flora Malesiana Bulletin*, 29: 2610-2618.
- JARVIS, P.G. (1995). — Scaling processes and problems. *Plant, Cell and Environment*, 18: 1079-1089.
- JENIK, J. (1994). — The Dahomey gap: an important issue in African phytogeography. *Mémoire Société Biogéographie* (3^e série), IV: 125-133.
- KAHN, F. (1982). — *La reconstitution de la forêt tropicale humide, sud-ouest de la Côte d'Ivoire*. ORSTOM, mémoire 97, Paris.
- KOKOU, K. (1998). — *Les mosaïques forestières au sud du Togo: biodiversité, dynamique et activités humaines*. Thèse de doctorat d'université, Université de Montpellier II.
- KOKOU, K., AFIADEMANYO, K. & AKPAGANA, K. (1999a). — Les forêts sacrées littorales du Togo: rôles culturel et de conservation de la biodiversité. *Journal de la Recherche Scientifique, Université du Bénin*, 3: 91-104.
- KOKOU, K. & CABALLÉ, G. (2000). — Les îlots forestiers de la plaine côtière togolaise. *Bois et Forêts des Tropiques*, 263: 39-51.
- KOKOU, K., CABALLÉ, G. & AKPAGANA, K. (1999b). — Analyse floristique des îlots forestiers du sud du Togo. *Acta Botanica Gallica*, 146: 139-144.
- KOKOU, K., CABALLÉ, G., AKPAGANA, K. & BATAWILA, K. (1999c). — Les îlots forestiers au sud du Togo: dynamique et relations avec les végétations périphériques. *Revue d'Écologie (Terre Vie)*, 54: 301-314.
- LAURANCE, W.F. (1991). — Edge effects in tropical forest fragments: application of a model for the design of nature reserves. *Biological Conservation*, 57: 205-219.
- LEE, D.W. (1988). — Simulating forest shade to study the developmental ecology of tropical plants: juvenile growth in three vines. *Journal of Tropical Ecology*, 4: 281-292.

- LEE, D.W. & RICHARDS, J.H. (1991). — Heteroblastic development in vines. Pp. 205-243, in: Putz, F.E. & Mooney, H.A. (eds.). *The biology of vines*. Cambridge University Press.
- L'HOTE, Y. & MAHÉ, G. (1996). — *Afrique de l'Ouest et Centrale: précipitations moyennes annuelles (période 1951-1981)*. ORSTOM, 1 carte + notice.
- LOTT, E.J. (1985). — Listados florísticos de Mexico III. La Estación de Biología Chamela, Jalisco. *Biotropica*, 19: 228-235.
- MAKANA, J.R., HART, T.B. & HART, J.A. (1998). — Forest structure and diversity of lianas and understorey treelets in monodominant and mixed stands in the Ituri forest, Democratic Republic of the Congo. Pp. 429-446, in: Dallmeier, F. & Comiskey, J.A. (eds.). *Forest biodiversity research, monitoring and modelling. Conceptual background and Old world case studies*. Camforth Parthenon Publishing Group.
- MALEY, J. (1996). — The African rain forest - main characteristics of changes in vegetation and climate from the Upper Cretaceous to the Quaternary. *Proceedings of the Royal Society Edinburgh*, 104B: 31-73.
- ORIANS, G.H., DIRZO, R. & CUSHMAN, J.H. (eds.) (1996). — *Biodiversity and ecosystem processes in tropical forests*. Springer-Verlag, Ecological Studies 122, Berlin.
- PEÑALOSA, J. (1984). — Basal branching and vegetative spread in two tropical rain forest lianas. *Biotropica*, 16: 1-9.
- PHILLIPS, O.L. & GENTRY, A.H. (1994). — Increasing turnover through time in tropical forests. *Science*, 263: 954-958.
- PINARD, M.A. & PUTZ, F.E. (1994). — Vine infestation of large remnant trees in logged forest in Sabah, Malaysia: biomechanical facilitation in vine succession. *Journal of Tropical Forest Science*, 6: 302-309.
- PUTZ, F.E. (1984). — The natural history of lianas on Barro Colorado island, Panama. *Ecology*, 65: 1713-1724.
- PUTZ, F.E. (1985). — Woody vines and forest management in Malaysia. *Commonwealth Forestry Review*, 64: 359-365.
- PUTZ, F.E. (1990). — Liana stem diameter growth and mortality rates on Barro Colorado island, Panama. *Biotropica*, 22: 103-105.
- PUTZ, F.E. (1995). — Vines in treetops: consequences of mechanical dependence. Pp. 311-323, in: Lowman, M.D. & Nadkarni, N.M. (eds.). *Forest canopies*. Academic Press, Orlando.
- PUTZ, F.E. & CHAI, P. (1987). — Ecological studies of lianas in Lambir national Park, Malaysia. *Journal of Ecology*, 75: 523-531.
- PUTZ, F.E. & HOLBROOK, N.M. (1991). — Biomechanical studies of vines. Pp. 73-97, in: Putz, F.E. & Mooney, H.A. (eds.). *The biology of vines*. Cambridge University Press.
- PUTZ, F.E. & MOONEY, H.A. (eds.) (1991). — *The biology of vines*. Cambridge University Press.
- SCHENCK, H. (1892). — Beiträge zur Biologie und Anatomie der Lianen, im Besonderen der in Brasilien einheimischen Arten. Pp. 1-253, in: Schimper, A.F.W. (ed.). *Botanische Mittheilungen aus den Tropen* 4. Verlag von G. Fischer, Jena.
- SIEBERT, S.F. (1993). — The abundance and site preferences of rattan *Calamus exilis* and *Calamus zollingeri* in two Indonesian national parks. *Forest Ecology and management*, 59: 105-113.
- SIEGEL, S. (1956). — *Non-parametric tests*. Mc Graw Hill series in psychology, New-York.
- TREUB, M. (1883a). — Sur une nouvelle catégorie de plantes grimpantes. *Annales du Jardin Botanique de Buitenzorg*, 3: 44-75.
- TREUB, M. (1883b). — Observations sur les plantes grimpantes du jardin botanique de Buitenzorg. *Annales du Jardin Botanique de Buitenzorg*, 3: 160-182.
- TROY, A.R., ASHTON, P.M. & LARSON, B.C. (1997). — A protocol for measuring abundance and size of a neotropical liana, *Desmoncus polyacanthos* (Palmae), in relation to forest structure. *Economic Botany*, 51: 339-346.
- WILLIAMS-LINERA, G. (1990). — Vegetation structure and environmental conditions of forest edges in Panama. *Journal of Ecology*, 78: 356-373.
- YAP, S.W., CHAK, C.V., MAJUAKIM, L., ANUAR, M. & PUTZ, F.E. (1995). — Climbing Bamboo (*Dinochloa* spp.) in Deramakot forest reserve, Sabah: biomechanical characteristics, modes of ascent and abundance in a logged-over forest. *Journal of Tropical Forest Science*, 8: 196-202.