



**HAL**  
open science

## Voice-mediated interactions in a megaherbivore

Julie Thévenet, Nicolas Grimault, Paulo Fonseca, Nicolas Mathevon

► **To cite this version:**

Julie Thévenet, Nicolas Grimault, Paulo Fonseca, Nicolas Mathevon. Voice-mediated interactions in a megaherbivore. *Current Biology - CB*, 2022, 32 (2), pp.R70-R71. 10.1016/j.cub.2021.12.017 . hal-03543609

**HAL Id: hal-03543609**

**<https://hal.science/hal-03543609>**

Submitted on 22 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Voice-mediated interactions in a megaherbivore

Julie Thévenet<sup>1,2</sup>, Nicolas Grimault<sup>2</sup>, Paulo Fonseca<sup>3\*</sup> and Nicolas Mathevon<sup>1,4\*</sup>

<sup>1</sup>Equipe de Neuro-Ethologie Sensorielle ENES / CRNL, University of Saint-Etienne, CNRS, Inserm, Saint-Etienne, France.

<sup>2</sup>Equipe Cognition Auditive et Psychoacoustique CAP / CRNL, University Lyon 1, CNRS, Inserm, Lyon, France.

<sup>3</sup> Departamento de Biologia Animal and cE3c—Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal.

<sup>4</sup>Institut universitaire de France.

\*E-mail: [mathevon@univ-st-etienne.fr](mailto:mathevon@univ-st-etienne.fr) or [pjfonseca@fc.ul.pt](mailto:pjfonseca@fc.ul.pt)

By entering the Anthropocene, the planet Earth is becoming increasingly difficult for large animal species to inhabit. Yet, these species are of major importance for the functioning of the biosphere and their progressive disappearance is accompanied by profound negative alterations of ecosystems [1] (Supplemental information). To implement effective conservation measures, it is essential to have a detailed knowledge of the biology of these species, notably regarding their social structures and the interactions between individuals. Here, we show that the hippopotamus *Hippopotamus amphibius*, an iconic African megaherbivore for which little is known about social communication, use vocal recognition to manage relationships between territorial groups. We conducted playback experiments on groups of hippos and observed their response to vocalizations from an individual of the same group (familiar), a group from the same lake (neighbor), and from a distant group (stranger). We found that stranger vocalizations induced a stronger behavioral reaction than the other two stimuli. In addition to showing that hippos are able to identify categories of conspecifics based on vocal signatures, our study demonstrates that hippo groups are territorial entities that behave less aggressively toward their neighbors than toward strangers. These new behavioral data suggest that habituation playbacks prior to conservation translocation operations may help reduce the risk of conflict between individuals that have never seen each other.

31 The megaherbivores -terrestrial mammals that feed on plants and weigh over 1000 kg  
32 (elephants, hippos and rhinos)- are a major source of concern. In African savannah  
33 ecosystems, rhinos are at a survival stage -each species being present in small protected  
34 areas- and, although some populations are increasing at the cost of expensive monitoring,  
35 they remain critically endangered [2]. Elephants are much more numerous and, while they  
36 are also under pressure from human activities, our extensive knowledge of their biology  
37 allows conservationists to devise appropriate management measures, though sometimes  
38 difficult to achieve [3]. While the third type of African megaherbivore, the hippopotamus, is  
39 not yet listed as endangered, its populations have declined dramatically in recent decades.  
40 Habitat loss and unregulated hunting are of increasing concern [4]. This amphibious animal  
41 shares its life between land and water, and has a unique role in the ecosystem mainly  
42 because of its impact on the flux of energy and matter between the two environments [5].  
43 Despite this ecological importance, the biology of the hippopotamus is still mysterious in  
44 many respects, and population management methods remain largely empirical [6].

45 Studying the behavioral biology of hippos in the wild is notably complicated. It is difficult-if  
46 not impossible- to identify and mark individuals, and sometimes highly challenging to locate  
47 them. Hippos are well known for their amphibious habits: while they feed on land mainly at  
48 night and are rather solitary, they gather in groups in the water during the day to avoid  
49 overheating, rest and mate. Hippo groups are socially structured around a dominant male, a  
50 variable number of females and their young, and some peripheral males [7]. However, it is  
51 unclear whether individuals in a pod form a stable group defending a territory, or it is more  
52 like a fission-fusion type organization with individuals moving rather freely from one pod to  
53 another. In any case, the hippo social system appears to rely on communication signals -  
54 hippos are very vocal- whose role and meaning remain almost unexplored [8].

55 In the present study, we focused on the most common hippo vocalization, the wheeze honk,  
56 a loud call heard over long distances that is assumed to be important for social cohesion and  
57 communication between groups, but whose actual function remains unknown (see  
58 Supplemental information). Based on the assumption that hippo groups are territorial  
59 entities, defended by the dominant male but also potentially by other individuals, including  
60 females, we tested the hypothesis that the wheeze honk could signal the identity of the  
61 sender and thus enable behavioral decisions by the receiver individuals.

62 We worked in the *Maputo Special Reserve* (Mozambique, Austral Africa), an area  
63 characterized by the presence of several lakes inhabited by hippos (Figure 1A). For each  
64 group of hippos (minimum number of individuals = 3; maximum = 22), we first recorded  
65 spontaneous vocalizations and then conducted playback experiments (see Supplemental  
66 information for methodological details).

67 We conducted three types of playback tests on groups of hippos: one with a call from the  
68 group, another with a call from a different group present on the same lake, and another with  
69 a call from a distant stranger group (Figure 1B). Of the seven groups tested, five received all  
70 three stimuli. Two groups received only the familiar and stranger stimuli (one group had no  
71 neighbors in its lake and one has not been retested due to experimental constraints; see  
72 Supplementary Table 1 for details on test design). The order of the tests was balanced  
73 among the groups. The signals were played from the shore, around 70-90 meters away from  
74 the group, mimicking the approach of a vocalizing individual (mean duration of a playback  
75 session = 36 minutes, min-max = 15-75 min).

76 The results show that hippos respond to played back call (by calling back, approaching  
77 and/or marking by defecation), but that their response depends on the category of the  
78 stimulus (Figure 1C and 1D; Supplemental information). The overall intensity of the  
79 behavioral response is lowest in response to a call from an individual of the same group and  
80 highest in response to a call from an individual belonging to a stranger group (linear mixed  
81 model, Wald  $X^2 = 17.55$ ,  $p < 0.001$ , see Supplemental information for detailed statistics). The  
82 nature of the response also changes between stimuli. Whereas individuals responded to calls  
83 from any group, marking behavior (dung spraying) is modulated by the category of the calls  
84 (cumulative mixed model: Wald  $X^2 = 11.47$ ,  $p=0.003$ ). Stranger group calls induce more  
85 marking than calls from an individual of the same group (multiple comparisons tests:  $Z=2.41$ ,  
86  $p=0.042$ ), while there is no significant difference between reactions to the calls from the  
87 same group or from a neighboring group (multiple comparisons tests:  $Z=0.40$ ,  $p=0.915$ ).

88 Individuals in territorial animal species often react less aggressively to a known individual  
89 from a neighboring territory than to a stranger ("dear-enemy effect", see Supplemental  
90 References). Sometimes, however, the opposite is observed and it is the neighbor that is  
91 more strongly repelled ("nasty neighbor", see Supplemental References). Our experiments

92 suggest that in hippos, the arrival of a stranger individual is perceived as more threatening  
93 than that of a neighbor.

94 Complementary experiments to assess sound level and sound propagation revealed that the  
95 wheeze honk can propagate more than 1 km away. Although the acoustic features carrying  
96 the vocal signature may be altered during long distance transmission, it is thus likely that  
97 these animals can learn and recognize the voices of neighboring individuals living on the  
98 same lake (Supplemental information).

99 In short, we showed that hippos use vocal recognition to manage their inter-group  
100 relationships, a strategy already observed in other large mammals where competition is  
101 intense and which allows to limit physical fights with often irreparable consequences [9].  
102 While relocating endangered animals to maintain population above critical levels is  
103 increasingly common [10], our results suggest that precautions should be taken during such  
104 relocations with hippos. Before transferring a group of hippos to a new location, a potential  
105 precaution could be to broadcast their voices from a loudspeaker at a distance from the  
106 groups already present so that they get used to them and their level of aggressiveness  
107 gradually decreases. Reciprocity - getting the animals to be relocated accustomed to the  
108 voices of their new neighbors - could also be considered.

109

## 110 **Supplemental information**

111 Supplemental information including experimental procedures, one table, one figure and  
112 references can be found with this article online at XXX.

113

## 114 **Acknowledgements**

115 We thank Maputo Special Reserve and Peace Park Foundation for permission to conduct this  
116 work. We are grateful to Pedro Fonseca and Ernesto Tualufo Simões for logistical assistance,  
117 to Maria Maust-Mohl and Diana Reiss for giving us two recordings of hippo vocalizations, to  
118 Manuel Vieira for his help in computer processing of acoustic data and to David Reby for his  
119 helpful comments on a previous version of the manuscript. Funding was provided by the  
120 Labex CeLyA (France), the University of Saint-Etienne (France), the Centre National de la

121 Recherche Scientifique (CNRS, France) and the Institut universitaire de France. PF was partly  
122 supported by the strategic project UID/BIA/00329/2019 of cE3c.

123

## 124 **References**

125 [1] Enquist BJ et al, 2020. The megabiota are disproportionately important for biosphere functioning.  
126 Nature Communications, 11, 699.

127 [2] Gross M, 2018. Last call to save the rhinos. Current Biology, 28, R1-R16.

128 [3] Wall J et al, 2021. Human footprint and protected areas shape elephant range across Africa.  
129 Current Biology, 31, 2437-2445.

130 [4] Lewison R, Pluháček J. 2017. *Hippopotamus amphibius*. *The IUCN Red List of Threatened Species*  
131 2017: e.T10103A18567364. [https://dx.doi.org/10.2305/IUCN.UK.2017-](https://dx.doi.org/10.2305/IUCN.UK.2017-2.RLTS.T10103A18567364.en)  
132 [2.RLTS.T10103A18567364.en](https://dx.doi.org/10.2305/IUCN.UK.2017-2.RLTS.T10103A18567364.en). Downloaded on 04 July 2021.

133 [5] Schoelynck J et al, 2019. Hippos (*Hippopotamus amphibius*): The animal silicon pump. Science  
134 Advances, 5, eaav0395.

135 [6] Utete B, 2020. A review of some aspects of the ecology, population trends, threats and  
136 conservation strategies for the common hippopotamus, *Hippopotamus amphibius* L, in Zimbabwe.  
137 African Zoology, 55, 187-200.


138 [7] Klingel H. 2013. *Hippopotamus amphibius* Common Hippopotamus; In: Mammals of Africa:  
139 Volume VI: Pigs, Hippopotamuses, Chevrotain, Giraffes, Deer and Bovids. Kingdon J, Hoffmann M.  
140 (eds). Bloomsbury Publishing, London. p. 68-77.

141 [8] Maust-Mohl M et al, 2015. Acoustic and behavioral repertoires of the hippopotamus  
142 (*Hippopotamus amphibius*). Journal of the Acoustical Society of America, 138, 545-554.

143 [9] Mathevon N et al, 2017. Northern elephant seals memorize the rhythm and timbre of their rivals'  
144 voices. Current Biology, 27, 2352-2356.

145 [10] Berger-Tal O et al, 2020. Conservation translocations: a review of common difficulties and  
146 promising directions. Animal Conservation, 23, 121-131.

147  
148  
149



150  
151  
152  
153  
154

**Figure 1. Playback experiments on hippos.**

155 (A) Map of the *Maputo Special Reserve* (Mozambique, Austral Africa) with location of the  
156 tested hippo groups. (B) Examples of hippo vocalizations used for the playback experiments  
157 (three different individuals). (C) Top: typical hippo group. Bottom: approach toward the  
158 loudspeaker and marking (dung spraying followed by threatening display). (D) Behavioral  
159 reaction of hippos to conspecific vocalizations (familiar: call from an individual of the tested  
160 group; neighbor: call from an individual of a group from the same lake; stranger: call from an  
161 individual of a distant group). Left: Response strength represents an integrative measure of  
162 the hippo reaction to calls, calculated using a Principal Component Analysis from all  
163 behavioral variables. Right: Diagram reporting the behavioral scores for each of the three  
164 considered behaviors (approach toward the loudspeaker, number of vocalizations, and  
165 marking behavior by defecation; see Supplemental Information for details).

166

167

168

169 **SUPPLEMENTAL INFORMATION**

170 **The “wheeze honk”.**

171 Hippos have a repertoire of calls (wheeze honk, grunts, bellows, squeals; Eltringham 1999,  
172 Maust-Mohl et al 2015). The wheeze honk is emitted by both adult females and males. It is  
173 an aerial, powerful signal that can be heard at long distances. It can also propagate  
174 underwater (Barklow 2004). However, this underwater propagation remains to be  
175 thoroughly explored as it is unlikely that sound waves propagate at long range in shallow  
176 water. Although the wheeze honk is generally considered as a *contact call* that allows  
177 individuals to signal their presence to one another, its social function is still unclear.

178

179 **Playback experiments.**

180 We recorded hippo vocalizations with a shotgun microphone (Sennheiser MKH8070)  
181 connected to a digital audio recorder (Tascam DR-40; sampling rate = 48 kHz, 16 bit; average  
182 recording distance = 80 m). To serve as stimuli during the playback tests, we selected a single  
183 call per group (two calls for one of the groups), taking care that this call had an excellent  
184 signal-to-noise ratio, and was not blurred by other vocalizations. Each selected call was then  
185 used in playback tests as a familiar call for the group where it was recorded, as a neighbor  
186 call for a group of hippos in the same lake, and as a stranger call for a distant group of hippos  
187 (group of another lake). In addition, two calls recorded from captive animals were used as  
188 stranger calls. Each sound stimulus was low-pass filtered (4.5 kHz) and normalized. As shown  
189 by our measures of sound intensity of actual hippos vocalizing (see below), the wheeze honk  
190 is emitted at 94-110 dB SPL (rel. 20  $\mu$ Pa, N=85 calls), at 1 meter from the animal. We thus  
191 calibrated our playback loudspeaker to emit the sound stimuli at an average intensity of 100  
192 dB SPL at 1 meter, and with a playback distance of 70-90 meters.

193 We observed each group of hippos during a couple of hours before and after the playback  
194 experiments. Five different observers assessed the number of hippos and compared their  
195 assessment. As we conducted several playback experiments over several days on each pod,  
196 we re-assessed the number of hippos each time. The pods were stable in terms of the  
197 number of individuals during the period we performed the study (pod 1: 7 individuals; pod 2:

198 10 individuals; pod 3: 3 individuals; pod 4: 11 individuals; pod 5: 8-11 individuals; pod 6: 22  
199 individuals; pod 7: 7 individuals). The pods did not move much from day to day during the  
200 study, remaining roughly in the same place in their lakes.

201 Before each playback test, we waited until the hippos were calm and quiet, with most  
202 individuals with their head emerging out of water. We played the sound stimuli from a JBL  
203 Boombox2 speaker (HARMAN International) connected via Bluetooth to a cell phone  
204 (Samsung Galaxy XCover 4). If no hippo responded to the playback of the first rendition of  
205 the stimulus, the same call was played again 30 seconds after (this happened in only two  
206 trials: “stranger” trial n°2 of experiment 1 and “familiar” trial n°5 of experiment 6; see  
207 Supplemental Table 1). A minimum delay of 15 minutes was observed between each  
208 playback test. All tests were filmed and recorded.

209 Behavioral responses were scored from the videos according to the following variables, all  
210 measured within 10 minutes after the onset of the stimulus playback (Supplemental Table  
211 1):

212 - Approach toward the speaker (no movement = 0; movement of at least one individual less  
213 than one body length = 1; movement of at least one individual more than one body length =  
214 2).

215 - Marking by dung spraying (no marking = 0; one marking behavior = 1; multiple markings by  
216 one or more hippos = 2).

217 - Vocal production (no vocalization = 0; one vocalizing individual = 1; multiple vocalizing  
218 individuals = 2).

219 For each of these three behaviors, latencies (duration between stimulus onset and observed  
220 behavior onset) were also measured. If we were to repeat the playback, the latency was  
221 measured from the onset of the first stimulus.

222

## 223 **Statistical analysis**

224 We used principal component analysis (PCA, *FactoMineR package*, RStudio v3.6.2) with all  
225 the six behavioral variables measured to create a composite behavioral score. The first  
226 component explained 46.0% of the variables’ variance. The approach score, the marking

227 score and the latency to vocalize were strongly positively correlated with PC1, while the  
228 vocal production, the latency to approach and the latency to mark were strongly negatively  
229 correlated with PC1. Higher positive values of PC1 represented a stronger territorial  
230 response, with individuals approaching and marking more, and more rapidly, and producing  
231 less vocalizations. We called this PC1 the “response strength”. This response strength was  
232 analyzed by using a linear mixed model (*lmer* function of the *lme4* package), with the type of  
233 stimulus (i.e. familiar, neighbor, or stranger) as fixed factor, and the identity of the tested  
234 pod and the sound stimulus as random factors. Post hoc multiple comparisons (Tukey  
235 contrasts, *multcomp* R package) were performed to compare response strength between  
236 types of stimulus. We found that the response strength was significantly higher for stranger  
237 stimuli compared to familiar stimuli (Tukey comparisons:  $\beta_{\text{familiar}} = -2.22$ ,  $SE_{\text{familiar}} = 0.53$ ,  
238  $Z_{\text{familiar}} = -4.19$ ,  $P_{\text{familiar}} < 0.001$ ) and compared to neighbor stimuli (Tukey comparisons:  
239  $\beta_{\text{neighbor}} = -1.55$ ,  $SE_{\text{neighbor}} = 0.64$ ,  $Z_{\text{neighbor}} = -2.43$ ,  $P_{\text{neighbor}} = 0.040$ ). Response strength was not  
240 significantly different between familiar and neighbor calls (Tukey comparisons:  $\beta_{\text{neighbor}} =$   
241  $0.67$ ,  $SE_{\text{neighbor}} = 0.55$ ,  $Z_{\text{neighbor}} = 1.20$ ,  $P_{\text{neighbor}} = 0.450$ ).

242 In addition, three independent statistical tests were conducted on the three principal  
243 behavioral scores (approach, marking, vocal production). For each behavioral score, a  
244 cumulative link mixed model was fitted (*clmm* function of *Ordinal* R package), with the type  
245 of stimulus as fixed factor and the identity of the tested pod and the sound stimulus as  
246 random factors. Post hoc comparisons (Tukey contrasts, *emmeans* R package) were also  
247 conducted to compare the behavioral score for each type of stimulus. The results were the  
248 following:

249 - Approach: no significant effect of the category of stimulus ( $\chi^2$  Wald = 4.09,  $p =$   
250  $0.130$ ); no differences between groups (neighbor and stranger compared to familiar  
251 group :  $\beta_{\text{neighbor}} = 0.61$ ,  $SE_{\text{neighbor}} = 1.33$ ,  $Z_{\text{neighbor}} = 0.46$ ,  $P_{\text{neighbor}} = 0.891$  and  $\beta_{\text{stranger}} =$   
252  $2.64$ ,  $SE_{\text{stranger}} = 1.66$ ,  $Z_{\text{stranger}} = 1.59$ ,  $P_{\text{stranger}} = 0.249$ ; neighbor compared to stranger :  
253  $\beta_{\text{stranger}} = 2.03$ ,  $SE_{\text{stranger}} = 1.51$ ,  $Z_{\text{stranger}} = 1.35$ ,  $P_{\text{stranger}} = 0.371$ ; mean approach score =  
254  $0.75 \pm 0.86$  and  $0.67 \pm 0.52$  for respectively familiar and neighbor stimuli,  $1.08 \pm 0.76$  for  
255 stranger stimuli).

256 - Marking: significant effect of the category of stimulus ( $X^2$  Wald = 11.47,  $p = 0.003$ );  
257 significant difference between familiar and stranger groups ( $\beta_{\text{stranger}} = 3.98$ ,  $SE_{\text{stranger}} =$   
258  $1.65$ ,  $Z_{\text{stranger}} = 2.41$ ,  $P_{\text{stranger}} = 0.042$ ). There is a tendency to have more marking in  
259 stranger group compared to the neighbor group ( $\beta_{\text{stranger}} = 3.37$ ,  $SE_{\text{stranger}} = 1.76$ ,  
260  $Z_{\text{stranger}} = 1.92$ ,  $P_{\text{stranger}} = 0.134$ ). No significant difference between familiar and  
261 neighbor ( $\beta_{\text{neighbor}} = 0.61$ ,  $SE_{\text{neighbor}} = 1.51$ ,  $Z_{\text{neighbor}} = 0.40$ ,  $P_{\text{neighbor}} = 0.915$ ). Mean  
262 marking score =  $0.17 \pm 0.041$  and  $0.77 \pm 0.83$  for respectively neighbor and stranger  
263 stimuli, compared to  $0.13 \pm 0.34$  for familiar stimuli.

264 - Vocal response: no significant effect of the category of stimulus ( $X^2$  Wald = 1.65,  $p =$   
265  $0.438$ ); no differences between groups (neighbor and stranger compared to familiar  
266 group :  $\beta_{\text{neighbor}} = -1.48$ ,  $SE_{\text{neighbor}} = 4.08$ ,  $Z_{\text{neighbor}} = -0.36$ ,  $P_{\text{neighbor}} = 0.930$  and  $\beta_{\text{stranger}} =$   
267  $-4.63$ ,  $SE_{\text{stranger}} = 6.79$ ,  $Z_{\text{stranger}} = -0.68$ ,  $P_{\text{stranger}} = 0.774$ ; neighbor compared to stranger  
268 :  $\beta_{\text{stranger}} = -3.16$ ,  $SE_{\text{stranger}} = 7.05$ ,  $Z_{\text{stranger}} = -0.45$ ,  $P_{\text{stranger}} = 0.896$ ; mean vocal  
269 production score =  $1.75 \pm 0.58$  for familiar stimuli,  $1.67 \pm 0.81$  and  $1.62 \pm 0.77$  for  
270 respectively neighbor and stranger stimuli).

271

## 272 **Measures of sound intensity and estimation of call range.**

273 We used recordings made at distances of 70-150 m from the animals to estimate the sound  
274 pressure level (SPL) of hippo vocalizations ( $N = 85$ ) at 1 m (calibrated audio chain composed  
275 by a Sennheiser shotgun microphone connected to a TASCAM DR-40 recorder). Sound  
276 pressure level estimates averaged  $102 \pm 3.6$  dB re. 20  $\mu\text{Pa}$  (min = 94, max = 110 dB). The  
277 measured ambient noise amplitude was 43 dB SPL, corresponding to 50% SPL percentiles,  
278 i.e. 50 % of the time the noise amplitude was below 43 dB SPL considering all the recordings  
279 from the several lakes (measured from the calibrated recordings considering 1 s segments  
280 and the 0-3 kHz bandwidth; noise SPL ranged 39 to 51 dB with exceptionally quiet moments  
281 as low as 34 dB SPL; notice that recordings were made with a shotgun Sennheiser  
282 microphone, which might lead to underestimate the level of the background noise due to its  
283 anisotropy. However, we assume that our measurements are representative because 1) the  
284 average level of background noise remains approximate due to variations during the day and  
285 with weather conditions, 2) it is not clear which microphone, shotgun or omnidirectional, is

286 closer to the directivity of the hippo's auditory system). As a result, the distance at which the  
287 sound level of the hippo wheeze honk no longer exceeds that of the background noise  
288 according to spherical spreading should be at least 1 km (see black curve in Supplemental  
289 Figure 1A).

290 This estimation was firstly corroborated by recordings of spontaneously emitted hippo  
291 vocalizations along a lake shore (blue curve in Supplemental Figure 1A). The recorders  
292 (Audiomoth v1.2.0 loggers) were placed along the shore near the water of Chingute lake, at  
293 different distances from a hippo pod. The distance from the first recorder to the hippo pod  
294 was estimated at 80 m. The following recorders were placed at 50, 100, 200, 400, 800, 1700  
295 meters from the first recorder. As it was not possible to place the recorders in a straight line  
296 relative to the hippo pod, these distances must be considered as approximative. Hippo calls  
297 appear to attenuate according to the spherical model and were no more detected from  
298 recordings at a distance of 1700 m from the vocalizing animals (which explains why there is  
299 no point at this distance on the figure).

300 This estimation was also corroborated by another analysis integrating signal and noise levels  
301 within standard mammalian auditory bands ( $1/3^{\text{rd}}$  octave bands; Dooling and Blumenrath,  
302 2014; Erbe et al., 2016; Putland et al., 2017). To do so, we averaged the frequency spectrum  
303 of all hippo calls including spontaneous vocalizations and calls recorded during playback  
304 experiments (normalizing for recording distance) and further calculated spectra at different  
305 distances using the spherical propagation model and adding a frequency-dependent excess  
306 attenuation component. To measure the excess attenuation, we did a propagation  
307 experiment using white noise (sequences of 20 seconds of white noise; bandwidth 0-20 kHz;  
308 propagation at 1, 5, 10, 15, 20, 25, 30 and 61 m over an open and flat savannah; sounds  
309 played back from a JBL BoomBox2 loudspeaker and recorded with a Sennheiser MKH8070  
310 microphone and Tascam DR-40 recorder). Besides, we computed the average background  
311 noise spectrum ( $1/3^{\text{rd}}$  octave bands) using 20 files with sound recordings obtained just  
312 before or after the hippo calls considered for this analysis. We chose to consider the  
313 percentiles 5, 20 and 50 since they indicate that the background noise spectrum is  
314 respectively 5%, 20% and 50% of the time below the measured value. The data are displayed  
315 in Supplemental Figure 1B. Considering the frequency bandwidth where the hippo call  
316 exhibits the more energy (40-500 Hz), this call is likely to be above the background noise till

317 1 to 2 km. Since this estimation is based on an average call, louder calls (the loudest being  
318 almost 10 dB above average) may be perceived at longer distances.

319

### 320 **Discussion: The impact of megafauna on ecosystems**

321 This topic was recently highlighted in a paper that demonstrates that megafauna are  
322 disproportionately important for the functioning of the biosphere (Enquist BJ et al, 2020.  
323 Nature Communications, 11, 699). Large herbivorous mammals thus have a major role in the  
324 flux of nutrients and energy in the biosphere through their feces, urine and flesh. The  
325 redistribution of nutrients and the fertilization of ecosystems are highly dependent on them,  
326 and there is evidence that a biosphere without the largest animals is less productive,  
327 contains less biomass, is less fertile. On the other hand, the introduction of a large mammal  
328 as an alien species in an ecosystem where it was not initially present can profoundly alter  
329 the ecosystem. This is what is currently happening with the emerging population of hippos in  
330 Colombia. This population originated from a few animals in captivity, and has been growing  
331 steadily over the last three decades. Their presence is causing eutrophication of the lakes  
332 where they live and may pose a threat to the original ecosystems as well as to human  
333 populations (for the full story, see these two recent papers: Shurin JB et al, 2020. Ecosystem  
334 effects of the world's largest invasive animal. Ecology, 101, e02991; Castelblanco-Martinez  
335 DN et al, 2021. A hippo room: Predicting the persistence and dispersion of an invasive mega-  
336 vertebrate in Colombia, South America).

337

Experience	Trial	Location	Pod	Pod size	Playback	Stimulus	Approach (lat)	Marking (lat)	Vocalization (lat)
1	1	Chingute	1	7	familiar	pod 1	1 (7)	0 (inf)	1 (5.9)
1	2	Chingute	1	7	stranger	captivity	0 (inf)	2 (48)	0 (83.9)
1	3	Chingute	1	7	familiar	pod 1	1 (4)	0 (inf)	2 (7.5)
2	1	Nela	2	10	stranger	captivity	1 (3)	0 (inf)	2 (8.3)
2	2	Nela	2	10	familiar	pod 2	2 (1)	0 (inf)	2 (6.2)
2	3	Nela	2	10	stranger	captivity	1 (2)	0 (inf)	2 (8.2)
2	4	Nela	2	10	familiar	pod 2	0 (inf)	0 (inf)	2 (10.8)
2	5	Nela	2	10	stranger	captivity	0 (inf)	1 (54)	2 (11.2)
3	1	Max	3	3	stranger	captivity	1 (1)	0 (inf)	2 (15.7)
3	2	Max	3	3	familiar	pod 3	1 (1)	0 (inf)	2 (6.7)
3	3	Max	3	3	stranger	captivity	1 (1)	0 (inf)	2 (8.2)
3	4	Max	3	3	familiar	pod 3	2 (1)	0 (inf)	2 (7.5)
4	1	Chingute	4	11	familiar	pod 4	0 (inf)	0 (inf)	2 (0)
4	2	Chingute	4	11	stranger	pod 3	2 (1)	1 (211)	1 (45)
4	3	Chingute	4	11	familiar	pod 4	0 (inf)	0 (inf)	2 (7.1)
4	4	Chingute	4	11	stranger	pod 2	2 (1)	0 (inf)	0 (inf)
4	5	Chingute	4	11	neighbor	pod 1	1 (1)	0 (inf)	0 (inf)
5	1	Chingute	5	11	familiar	pod 5	0 (inf)	0 (inf)	2 (6.7)
5	2	Chingute	5	11	stranger	pod 2	1 (1)	2 (84)	2 (8.2)
5	3	Chingute	5	11	familiar	pod 5	0 (inf)	0 (inf)	2 (7)
5	4	Chingute	5	11	neighbor	pod 4	0 (inf)	0 (inf)	2 (8.6)
6	1	Nela	6	22	neighbor	pod 2	1 (2)	0 (inf)	2 (8.3)
6	2	Nela	6	22	stranger	pod 5	1 (1)	2 (42)	2 (5.2)
6	3	Nela	6	22	familiar	pod 6	0 (inf)	0 (inf)	2 (5.3)
6	4	Nela	6	22	neighbor	pod 2	1 (1)	0 (inf)	2 (6.3)
6	5	Nela	6	22	familiar	pod 6	0 (inf)	0 (inf)	2 (54.7)
7	1	Chingute	1	7	familiar	pod 1	2 (7)	1 (288)	0 (inf)
7	2	Chingute	1	7	familiar	pod 1	2 (2)	1 (382)	1 (7.5)
7	3	Chingute	1	7	stranger	pod 3	2 (8)	1 (168)	2 (10)
8	1	Chingute	5	8	stranger	pod 3	2 (2)	1 (40)	2 (13.7)
9	1	Chingute	7	7	neighbor	pod 5	0 (inf)	0 (inf)	2 (4.5)
9	2	Chingute	7	7	familiar	pod 7	0 (inf)	0 (inf)	2 (4.2)
9	3	Chingute	7	7	stranger	captivity	0 (inf)	0 (inf)	2 (7.5)
10	1	Chingute	1	7	neighbor	pod 5	1 (1)	1 (24)	2 (10.3)
10	2	Chingute	1	7	familiar	pod 1	1 (1)	0 (inf)	2 (66.2)

339

340

341

**Supplemental Table 1. Playback experiments.**

342

The score (0, 1, 2) and the latency (between brackets, in seconds; Inf = "infinite" latency) are indicated for each of the three considered behaviors (Approach, Marking, Vocalization).


343

(Location = name of the lake; pod = hippo group).

344

345

346



347

348

349

350 **Supplemental Figure 1. Propagation range of the hippopotamus wheeze honk. A.**

351 Propagation of hippo vocalizations along the Lake Chingute shore close to the water. The  
352 shaded area in grey indicates the background noise (below 43 dB SPL during 50% of the  
353 time; min = 34 dB SPL, max = 51 dB). Black curve: amplitude of a hippo call as a function of  
354 distance calculated using the spherical propagation model (call amplitude at 1 meter = 102  
355 dB). Blue curve: amplitude of hippo calls recorded at different distances from the vocalizing  
356 individuals. In these recordings, hippo vocalizations were detected up to 880 m but not at  
357 1700 m. **B.** Propagation range of wheeze-honk calls comparing one-third octave spectra of  
358 averaged hippo calls and background sound level. Solid lines represent the frequency  
359 spectra of an averaged hippo call for different propagation distances (values are represented  
360 at the centre frequency of each one-third octave band; the spectrum at 64 meters was  
361 obtained based on recordings from different individuals on several lakes; spectra at other  
362 distances were calculated using the spherical propagation model and adding the excess  
363 attenuation measured in white noise propagation experiments). The dashed lines represent  
364 the background noise spectrum (one-third octave spectrum; averaged over 20 files with  
365 sound recordings obtained in different locations; the three curves represent the 5%, 20%  
366 and 50% percentiles of the distribution of background noise measurements, respectively -  
367 which means that the background noise was below the 0.05 curve during 5% of the total  
368 time of our recordings, below the 0.2 curve during 20% of the total time, and below the 0.5  
369 curve during 50% of the total time). These calculations confirm that the transmission range  
370 of the hippo call can reach more than 1 km.

371

372

373

374 **Supplemental References**

- 375 Barklow, W. E. 1994. Big talkers. *Wildlife Conservation*, 97, 20–29.
- 376 Barklow WE, 1997. Some underwater sounds of the Hippopotamus (*Hippopotamus amphibius*).  
377 *Marine and Freshwater Behaviour and Physiology*, 29, 237-249.
- 378 Barklow WE, 2004. Amphibious communication with sound in hippos, *Hippopotamus amphibius*.  
379 *Animal Behaviour*, 68, 1125-1132.
- 380 Dooling, R.J., Blumenrath, S.H., 2014. Avian sound perception in noise. In: Brumm, H.(Ed.), *Animal*  
381 *Communication in Noise*. Springer Verlag, Heidelberg, pp. 229–250.
- 382 Eksteen J et al, 2016. A conservation assessment of Hippopotamus amphibius. In: Child MF, Roxburg  
383 L, Do Linh San E, Raimondo D, Davies-Mostert HT (Eds), *The Red List of Mammals of South Africa,*  
384 *Swaziland and Lesotho*. South Africa: South African National Biodiversity Institute and Endangered  
385 Wildlife Trust.
- 386 Eltringham SK, 1999. *The Hippos*. T & AD Poyser Ltd, Academic Press, London, UK.
- 387 Erbe, C., Reichmuth, C., Cunningham, K., Lucke, K., Dooling, R., 2016. Communication masking in  
388 marine mammals: a review and research strategy. *Mar. Pollut. Bull.* 103 (1–2), 15–38.
- 389 Karstad EL, Hudson RJ, 1986. Social organization and communication of riverine hippopotami in  
390 southwestern Kenya. *Mammalia*, 50, 153-164.
- 391 Müller CA, Manser M, 2007. ‘Nasty neighbours’ rather than ‘dear enemies’ in a social carnivore.  
392 *Proceedings of the Royal Society B*, 274, 959-965.
- 393 Owen-Smith RN, 1988. Megaherbivores. The influence of very large body size on ecology. Cambridge  
394 University Press.
- 395 Putland, R.L., Merchant, N.D., Farcas, A., Radford, C.A., 2017. Vessel noise cuts down communication  
396 space for vocalizing fish and marine mammals. *Glob. Chang. Biol.* 24 (4), 1708–1721.
- 397 Siracusa ER et al, 2021. Familiar neighbors, but not relatives, enhance fitness in a territorial mammal.  
398 *Current Biology*, 31, 438-445.e3.
- 399 Temeles EJ, 1994. The role of neighbours in territorial systems: when are they ‘dear enemies’?  
400 *Animal Behaviour*, 47, 339-350.
- 401 Tibbetts EA, Dale J, 2007. Individual recognition: it is good to be different. *Trends in Ecology and*  
402 *Evolution*, 22, 529-537.
- 403 Tumulty JP, 2018. Dear Enemy Effect. In J. Vonk, T. K. Shackelford (eds.), *Encyclopedia of Animal*  
404 *Cognition and Behavior*, Springer.

