Col-OSSOS: Probing Ice Line/Colour Transitions within the Kuiper Belt's Progenitor Populations

Laura Buchanan, Megan Schwamb, Wesley Fraser, Michele Bannister, Michel Marsset, Rosemary Pike, David Nesvorny, Johnj Kavelaars, Susan Benecchi, Matthew Lehner, et al.

► To cite this version:

Laura Buchanan, Megan Schwamb, Wesley Fraser, Michele Bannister, Michel Marsset, et al.. Col-OSSOS: Probing Ice Line/Colour Transitions within the Kuiper Belt's Progenitor Populations. Europlanet Science Congress 2020, Sep 2020, virtuel, Germany. pp.EPSC2020-185, 10.5194/epsc2020-185 . hal-03572420

HAL Id: hal-03572420

<https://hal.science/hal-03572420>

Submitted on 14 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destine au dpt et la diffusion de documents scientifiques de niveau recherche, publis ou non, manant des tablissements d'enseignement et de recherche franais ou trangers, des laboratoires publics ou privs.

Distributed under a Creative Commons Attribution| 4.0 International License

Col-OSSOS: Probing Ice Line/Colour Transitions within the Kuiper Belt's Progenitor Populations

Laura Buchanan¹, Megan Schwamb¹, Wesley Fraser², Michele Bannister³, Michel Marsset⁴, Rosemary Pike⁵, David Nesvorn⁶, John J Kavelaars^{2,7}, Susan Benecchi⁸, Matthew Lehner⁹, Shiang-Yu Wang⁹, Audrey Thirouin¹⁰, Nuno Peixinho¹¹, Kathryn Volk¹², Mike Alexandersen⁹, Ying-Tung Chen⁹, Audrey Delsanti¹³, Brett Gladman¹⁴, Stephen Gwyn², and Jean-Marc Petit¹⁵

¹Queen's University Belfast, Astrophysics Research Centre, United Kingdom of Great Britain and Northern Ireland

(lbuchanan14@qub.ac.uk)

²NRC-Herzberg Astronomy and Astrophysics, National Research Council of Canada, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

³School of Physical and Chemical Sciences—Te Kura Mat, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

⁴Department of Earth, Atmospheric and Planetary Sciences, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

⁵Harvard & Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁶Department of Space Studies, Southwest Research Institute, 1050 Walnut St., Suite 300, Boulder, CO, 80302, United States

⁷Department of Physics and Astronomy, University of Victoria, Elliott Building, 3800 Finnerty Road, Victoria, BC V8P 5C2, Canada

⁸Planetary Science Institute, 1700 East Fort Lowell, Suite 106, Tucson, AZ 85719, USA

⁹Institute of Astronomy and Astrophysics, Academia Sinica; 11F of AS/NTU Astronomy-Mathematics Building, No.1, Sec. 4, Roosevelt Rd Taipei 10617, Taiwan, R.O.C

¹⁰Lowell Observatory, 1400 W. Mars Hill Rd., Flagstaff, AZ 86001, USA

¹¹CITEUC — Centre for Earth and Space Science Research of the University of Coimbra, Geophysical and Astronomical Observatory of the University of Coimbra, 3030-004 Coimbra, Portugal

¹²Lunar and Planetary Laboratory, University of Arizona, 1629 E. University Blvd., Tucson, AZ 85721, USA

¹³Aix Marseille Universit, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, F-13388, Marseille, France

¹⁴Department of Physics and Astronomy, University of British Columbia, Vancouver, BC, Canada

¹⁵Institut UTINAM UMR6213, CNRS, Univ. Bourgogne Franche-Comt, OSU Theta F-25000 Besanon, France

The Colours of the Outer Solar System Origins Survey (Col-OSSOS, Schwamb et al., 2019) has examined the surface compositions of Kuiper Belt Objects (KBOs) by way of broadband g-, r- and J-band photometry, using the Gemini North Hawaii Telescope. This survey showed a bimodal distribution in the colours of the objects surveyed, consistent with previous colour surveys (Tegler et al., 2016). These broadband surface colours can be considered a proxy for surface composition of these KBOs, so this survey allows the frequency of different surface compositions within the outer Solar System to be explored. The bimodality of the observed colours suggests the presence of some sort of surface transition within the Kuiper belt, perhaps due to a volatile ice-line transition in the pristine planetesimal disk that existed before Neptune's migration. The Outer Solar System Origins Survey (OSSOS, Bannister et al., 2018), from which Col-OSSOS selected objects brighter than 23.6 r-band magnitude, has well characterised and quantified biases, so allowing for comparisons between the observations and numerical models of the Kuiper belt.

By applying different colour transitions to the primordial planetesimal disk, in this work we explore

the possible positions for ice line/colour transitions within the planetesimal disk that existed before Neptune’s migration. Within Schwamb et al. (2019), a simplified toy model was used to investigate the possible position of this transition. Nesvorny et al. (2020) has investigated the primordial colour fraction, in particular how it can create the inclination distribution that we see in the colours of KBOs today. In this work we use a full dynamical model of the Kuiper belt to more precisely pinpoint the possible location of this transition. We make use of the model by Nesvorny & Vokrouhlicky (2016) of Neptune’s migration from 23 au to 30 au, and the consequent perturbation of the Kuiper belt into its current form. This model allows precise tracking of the objects from their pre-Neptune migration to post-Neptune migration positions, allowing various colour transition positions in the initial disk, an example of which is shown in Figure 1, to be compared with the Col-OSSOS observations of the modern day disk.

Figure 1: An example red/neutral transition at 27 au. The left plots show the objects in the primordial disk, while the right plots show the objects post-Neptune migration from the model of Nesvorny & Vokrouhlicky (2016).

The OSSOS survey simulator (Lawler et al., 2018) can then be used to calculate which of the simulated objects could have been observed by OSSOS, and so selected by Col-OSSOS for surface colour observations. The colour transition within the initial disk, shown in Figure 1, is moved radially outwards through the disk and the corresponding outputs are compared with the Col-OSSOS colour observations to see which initial disk colour transition positions are consistent with the modern day Kuiper belt. We will present results combining an accurate dynamical model of the Kuiper Belt’s evolution by Nesvorny & Vokrouhlicky (2016) with Col-OSSOS photometry. We will explore multiple radial colour distributions in the primordial planetesimal disk and implications for the the positions of ice line/colour transitions within the Kuiper Belt’s progenitor populations.

References

- Bannister, M. T., Gladman, B. J., Kavelaars, J. J., et al. 2018, *ApJS*, 236, 18
- Lawler, S. M., Kavelaars, J. J., Alexandersen, M., et al. 2018, *Front. Astron. Space Sci.*, 5, 14
- Nesvorny, D., Vokrouhlicky, D., Alexandersen, M., et al. 2020, *AJ*, in press
- Nesvorny, D., & Vokrouhlicky, D. 2016, *ApJ*, 825
- Schwamb, M. E., Bannister, M. T., Marsset, M., et al. 2019, *ApJS*, 243, 12
- Tegler, S. C., Romanishin, W., Consolmagno, G. J., & J., S. 2016, *AJ*, 152, 210