



Choice of mixture Poisson models based on Extreme value theory

Samuel Valiquette, Frédéric Mortier, Jean Peyhardi, Gwladys Toulemonde

► To cite this version:

Samuel Valiquette, Frédéric Mortier, Jean Peyhardi, Gwladys Toulemonde. Choice of mixture Poisson models based on Extreme value theory. 2022. hal-03648664

HAL Id: hal-03648664

<https://hal.science/hal-03648664>

Preprint submitted on 22 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CHOICE OF MIXTURE POISSON MODELS BASED ON EXTREME VALUE THEORY

Samuel Valiquette^{a,b,c,d} & Frédéric Mortier^a & Jean Peyhardi^b & Gwladys Toulemonde^{b,c}

^a *CIRAD, UPR Forêts et Sociétés, F-34398 Montpellier, France. Forêts et Sociétés,
Univ Montpellier, CIRAD, Montpellier, France.*

^b *IMAG, CNRS, Université de Montpellier, 34090, Montpellier, France.*

^c *LEMON, Inria, 34095, Montpellier, France*

^d *Université de Sherbrooke, Département de mathématiques, Sherbrooke, Canada, J1K
2R1*

Abstract. Count data are omnipresent in many applied fields, often with overdispersion due to an excess of zeroes or extreme values. With mixtures of Poisson distributions representing an elegant and appealing modelling strategy, we focus here on the challenging problem of identifying a suitable mixing distribution and study how extreme value theory can be used. We propose an original strategy to select the most appropriate candidate among three categories: Fréchet, Gumbel and pseudo-Gumbel. Such an approach is presented with the aid of a decision tree and evaluated with numerical simulations.

Keywords. Poisson mixture, Discrete extreme value theory, Peak-over-threshold.

1 Introduction

Count data are classically observed in many applied fields such as in actuarial science when evaluating risk and the pricing of insurance contracts [e.g., Bartoszewicz, 2005], in genetics to model the number of genes involved in phenotype variability [e.g., Anders and Huber, 2010] or in ecology to model species abundance [e.g., Wenger and Freeman, 2008]. While Poisson models and regression are well established choices for these type of data, they are not suitable for overdispersed data, which typically occur with an excess of zeroes or extreme large values. To overcome such limitations the use of Poisson mixture models has been proposed. This assumes the Poisson's intensity is no longer an unknown fixed value, but a positive random variable. Mixture approach induces overdispersed distributions with more zeroes and high values compared to the classical Poisson model [Shaked, 1980]. A variety of mixture distributions has been already proposed [Karlis and Xekalaki, 2005]. Classical examples include the gamma distribution [Greenwood and Yule, 1920], the lognormal [Bulmer, 1974] or the Bernoulli [Lambert, 1992]. From a general point of view, any distribution with a non negative support, finite or not, can be a potential candidate as a mixing distributions. Some tests exist to verify whether data are overdispersed [Yang et al., 2009] or if they come from a mixed Poisson distribution [Carriere, 1993]. These tests justify the use of Poisson mixtures, but do not make claims on what type of mixing distribution should be selected. To our knowledge, there are

no studies that propose a solution to this problem. This paper aims to propose a new strategy to select an appropriate and efficient mixing distribution family.

Usually one may choose to fit some predetermined Poisson mixtures and keep the best model based on some criteria. However the following example shows the difficulties in the mixing distribution choice. A sample of size $n = 500$ has been simulated from a Poisson-Beta type II with parameters equal to $a = 1$ and $b = 2.2$. This choice ensures finite expectation and variance for the mixed Poisson distribution and are equal to 0.83 and 8.47 respectively. Such a distribution has been used to model accident proneness [Holla and Bhattacharya, 1965]. First and as expected, the use of a simple Poisson model fails to properly fit data (see Figure 1). In particular, it does not capture well the high frequency of zeroes and large values once its parameter is estimated ($\hat{\lambda} = 1.02$).




Figure 1: Fitted Poisson, Poisson-Gamma, Poisson-Inverse-Gamma and Poisson-Lognormal models to data simulated from a Poisson-Beta type II distribution with $a = 1$ and $b = 2.2$ compared to its empirical distribution (Dark Gray)

Alternative latent distributions on λ have been used for this simulation, either the gamma or lognormal as classically proposed in ecology [Tikhonov et al., 2020] or the inverse-gamma applied in actuarial science for liability insurance claims [Tzougas, 2020]. Inference is performed in a Bayesian framework (details in Section 2.2). While the Poisson-gamma, *i.e.* negative binomial, or the Poisson-lognormal are popular choices, the Poisson-inverse-gamma is privileged for this example. Indeed, the *posterior* model probabilities of the negative binomial and Poisson-lognormal are respectively 0% and 38.8% compared to 61.2% for the Poisson-inverse-gamma. The latter mixture distinguishes itself even if the three models behave similarly with regards to Figure 1.

The reason that the inverse-gamma is privileged in this example is due to its tail behavior which is similar to the beta type II distribution. We establish in this paper that an adequate choice for the distribution on λ can be made by analysing such a property. This paper is organized as follow. Section 2 presents a classification of various Poisson mixtures based on their tail behavior using extreme value theory. Using these new classifications, we construct in Section 3 a strategy to choose a family of distributions for λ . This strategy is presented in the form of a decision tree where each step leads to an adequate category of Poisson mixtures. Simulations for each section are also presented to attest the relevance and usefulness of such an approach.

2 Poisson mixture tail behavior

In this section, we present the fundamental results in extreme value theory and the restrictions when it comes to discrete distributions. Following this, we present three categories of Poisson mixtures that characterise different tail behaviors and conclude with simulations to assess the interest of selecting such a category.

2.1 Theoretical foundations and Poisson mixtures categories

The tail behavior of a distribution can be studied using extreme value theory. Such a statistical approach analyses how the maximum of a distribution F stabilizes. The theory says that F belongs to a max domain of attraction if it exists two normalizing sequences $a_n > 0$ and b_n such that $F^n(a_n x - b_n)$ converges to a non-degenerate distribution when $n \rightarrow \infty$ [Resnick, 1987]. There are three possible domains of attraction named Weibull, Gumbel and Fréchet. These domains describe the asymptotic tail behavior of F and correspond respectively to finite, exponential and heavy tailed distributions. In the sequel, they will be denoted by \mathcal{D}_- , \mathcal{D}_0 and \mathcal{D}_+ , and we will write this property by $F \in \mathcal{D}$ where \mathcal{D} is one of the three domains. In the following, we assume that the mixing distribution λ is supported on \mathbb{R}_+ , then only \mathcal{D}_0 and \mathcal{D}_+ are considered. Usual continuous distributions belong to a domain, this is not the case for discrete random variables. Indeed, a necessary

condition for a discrete distribution F to be in a domain of attraction is the property of long tailed [Anderson, 1970] defined by:

$$\lim_{n \rightarrow \infty} \frac{1 - F(n+1)}{1 - F(n)} = 1. \quad (1)$$

In particular well known discrete distributions do not satisfy this property such as among others Poisson, negative binomial or yet geometric distributions. Even though the latter two are Poisson mixtures with a gamma distributed mixing parameter, which belongs to \mathcal{D}_0 , the domain of attraction is not carried over by the mixture distribution. However Anderson [1970] and Shimura [2012] showed that if a discrete distribution verifies

$$\lim_{n \rightarrow \infty} \frac{1 - F(n+1)}{1 - F(n)} = L \in (0, 1) \quad (2)$$

then F is, in a sense, 'close' to the Gumbel domain. More precisely, Shimura [2012] showed that property (2) implies that F is the discretization of a unique continuous distribution belonging to \mathcal{D}_0 . On the other hand, Anderson [1970] showed that there is a sequence b_n such that $F^n(x + b_n)$ has infimum and supremum limits bounded by two different Gumbel distributions, implying that F is not far from this domain.

Because Poisson mixture distributions are discrete distributions, they are constrained to the long tailed property in order to have a domain. Otherwise, they may be close to the Gumbel domain. But Poisson mixtures are uniquely identifiable by the distribution on λ [Feller, 1943], this means that its tail behavior is dependent of the latter. Therefore, we need to understand conditions on λ that allow Poisson mixture distribution to inherit a domain or to be close to the Gumbel one. Perline [1998] established some conditions for such preservation. From this point forward, we note by F and f the cumulative distribution function (cdf) and the probability density function (pdf) for λ and F_M the cdf of the resulting Poisson mixture. Firstly, if $F \in \mathcal{D}_+$ and is such that $\lim_{x \rightarrow \infty} \frac{xf(x)}{1-F(x)} = \alpha$ for some $\alpha > 0$ (1st Von Mises condition), then $F_M \in \mathcal{D}_+$. Secondly, if $F \in \mathcal{D}_0$, $\lim_{x \rightarrow \infty} \frac{d}{dx} \left[\frac{1-F(x)}{f(x)} \right] = 0$ (3rd Von Mises condition) and $\frac{f(x)}{1-F(x)} = o(x^{-\delta})$ as $x \rightarrow \infty$ for some $\delta \geq \frac{1}{2}$, then $F_M \in \mathcal{D}_0$.

These results clarify some conditions for which domain of attraction of the mixing distribution is propagated to the associated Poisson mixture distribution. Very naturally, we denote these situations by two categories of Poisson mixtures: **Fréchet** and **Gumbel**. A broad set of distributions satisfies the 1st Von Mises condition, for example the Fréchet, folded-Cauchy, beta type II, the inverse-gamma or the gamma/beta type II mixture [Irwin, 1968]. Unfortunately, examples are scarce for the Gumbel domain. Indeed the extra condition on the hazard function is quite restrictive. Some examples are the lognormal, the Benktander type I and II [Kleiber and Kotz, 2003] or the Weibull distribution, with

further restrictions on the parameters for the latter two cases. The Gumbel category does not encompass cases like the Poisson-gamma; while the mixing distribution belongs to \mathcal{D}_0 , it does not satisfy the additional condition on the hazard function. In order to categorise such Poisson mixtures, we study distributions on λ that behave like the gamma.

Definition 1 (Willmot [1990]). A density f is a **gamma type** if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{C(x)x^\alpha e^{-\beta x}} = 1 \quad (3)$$

where $C(x)$ is a locally bounded function on \mathbb{R}_+ and slowly varying, i.e. $\lim_{t \rightarrow \infty} \frac{C(tx)}{C(t)} = 1$ for every $x \in \mathbb{R}_+$ (see Bingham et al. [1987]), $\alpha \in \mathbb{R}$ and $\beta > 0$.

Using gamma type distributions in the Poisson mixture context allows us to extend the categorisation to cases where $F \in \mathcal{D}_0$, but $F_M \notin \mathcal{D}_0$. First, we prove that such a mixing distribution belongs to \mathcal{D}_0 (see Proposition 1). Finally Theorem 1 establishes that $F_M \notin \mathcal{D}_0$ and quantifies the closeness to the Gumbel domain.

Proposition 1. If F is a gamma type distribution, then $F \in \mathcal{D}_0$.

Proof. Let \bar{F} be the survival function of a gamma type distribution. A sufficient condition for $F \in \mathcal{D}_0$ is to show that F has an exponential tail, i.e. $\lim_{x \rightarrow \infty} \bar{F}(x+k)/\bar{F}(x) = e^{-\beta k}$ where $k \in \mathbb{R}$ [Shimura, 2012]. Using L'Hôpital's rule and equation (3), the limit becomes

$$\lim_{x \rightarrow \infty} \frac{\bar{F}(x+k)}{\bar{F}(x)} = e^{-\beta k} \lim_{x \rightarrow \infty} \frac{C(x+k)}{C(x)}.$$

It remains to show that the latter limit is equal to 1. Because $C(\cdot)$ is slowly varying, we can use the Karamata representation

$$C(x) = c(x) \exp \left(\int_1^x t^{-1} \eta(t) dt \right),$$

where $c(\cdot)$ and $\eta(\cdot)$ are both functions from \mathbb{R}_+ to \mathbb{R}_+ , $\lim_{x \rightarrow \infty} c(x) = c > 0$ and $\lim_{x \rightarrow \infty} \eta(x) = 0$ [Bingham et al., 1987]. Then the limit equals

$$\lim_{x \rightarrow \infty} \frac{C(x+k)}{C(x)} = \lim_{x \rightarrow \infty} \exp \left(\int_x^{x+k} t^{-1} \eta(t) dt \right).$$

Because $\eta(x) \rightarrow 0$, for any $\varepsilon > 0$ then $0 < \eta(x) < \varepsilon$ for x large enough. Then

$$0 < \int_x^{x+k} t^{-1} \eta(t) dt < \varepsilon \int_x^{x+k} t^{-1} dt = \varepsilon \log \left(\frac{x+k}{x} \right) < \varepsilon \log(1+k)$$

which implies the limit is equal to 1 and establishes the sufficient condition. \square

Theorem 1. *Let F_M be a Poisson mixture with λ distributed according to a gamma type distribution F . Then for any integer $k \geq 1$, $\lim_{n \rightarrow \infty} \frac{1-F_M(n+k)}{1-F_M(n)} = (1+\beta)^{-k} \in (0, 1)$. In particular, F_M is not long tailed ($k = 1$).*

Proof. Let P_M and \bar{F}_M be the probability and survival functions of a Poisson mixture using a gamma type mixing distribution, then Willmot [1990] showed that

$$\lim_{n \rightarrow \infty} \frac{P_M(n)}{C(n)n^\alpha(1+\beta)^{-(n+\alpha+1)}} = 1.$$

Using this result for integer k , we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\bar{F}_M(n+k+1) - \bar{F}_M(n+k)}{\bar{F}_M(n+1) - \bar{F}_M(n)} &= \lim_{n \rightarrow \infty} \frac{P_M(n+k+1)}{P_M(n+1)} \\ &= \left(\frac{1}{1+\beta} \right)^k \lim_{n \rightarrow \infty} \frac{C(n+k+1)}{C(n+1)}, \end{aligned}$$

where the last limit converges to 1 using a similar proof as in Proposition 1. Because \bar{F}_M is monotonically decreasing, the proof can be concluded by applying the Stolz-Cesàro theorem. \square

This result allows to characterize a third category: the **pseudo-Gumbel**. It includes a broad class of mixing distributions among others gamma, gamma/Gompertz, exponential, exponential logarithmic, inverse-Gaussian and its generalization. Perline's result and Theorem 1 lead to consider now three categories for Poisson mixtures allowing the clarification of the mixing distribution choices. For examples, see Table 1 and the supplementary materials for details. Additionally, Theorem 1 also quantifies how 'close' those Poisson mixtures are with the quantity $(1+\beta)^{-1}$ involved in the limit. Indeed, if $\beta \rightarrow 0$, $\frac{1-F_M(n+1)}{1-F_M(n)} \rightarrow 1$, i.e. it approaches a long tailed distribution. Such property can blur the distinction between Gumbel and pseudo-Gumbel for some Poisson mixtures.

2.2 Impact of mixing distribution choice on goodness of fit

Are the categories previously defined useful to distinguish how the Poisson mixtures behaved? How can these categories be efficiently used when it comes to model selection? To answer these questions, we simulated 100 samples of different Poisson mixtures with size $n = 250$ using the gamma, Fréchet and lognormal distributions on λ , each one being a representative of the three categories. For each sample, the Poisson mixture is fitted with the same three distributions and the best model is kept using a Bayesian framework. This can be done with the `rstan` package for the language R [Stan Development Team,

Mixing (λ)	Poisson mixture (P_M)	Category
Fréchet(a, σ)	Poisson-Fréchet	Fréchet
Folded-Cauchy(μ, σ)	Poisson-folded-Cauchy	Fréchet
Inverse-gamma(a, b)	Poisson-inverse-gamma	Fréchet
Beta-II(a, b)	Poisson-beta-II	Fréchet
Gamma/Beta-II-mixture(r, a, b)	Generalized Waring	Fréchet
Lognormal(μ, σ)	Poisson-lognormal	Gumbel
Weibull(a, b)	Poisson-Weibull	Gumbel (if $a < 0.5$)
Benktander-I(a, b)	Poisson-Benktander-I	Gumbel
Benktander-II(a, b)	Poisson-Benktander-II	Gumbel (if $b < 0.5$)
Exponential(a)	Geometric	Pseudo-Gumbel
Gamma(a, b)	Negative binomial	Pseudo-Gumbel
Inverse-Gaussian(μ, σ)	Sichel	Pseudo-Gumbel
Generalized inverse-Gaussian(a, b, p)	PGIG	Pseudo-Gumbel

Table 1: Examples of Poisson mixture and associated categories

2020] to estimate the parameters by MCMC. The best model is then kept using the highest *posterior* model probability. Those probabilities are approximated using the bridge sampling computational technique [Meng and Wong, 1996] and the dedicated R package **Bridgesampling** [Gronau et al., 2020]. All results are based on the following priors: a gamma(1, 1) distribution for positive parameters and a Normal(0, 1) for real parameters. Moreover, we simulated for each sample 4 MCMC with 10000 iterations each in order to ensure reasonable convergence for the parameter estimations and for the *posterior* model probabilities. Results are presented in Table 2 and, in general, the most selected model stood out. The only exception being the gamma(2,2) where the lognormal and the gamma Poisson mixtures are evenly selected throughout the simulations.

Poisson mixture	Fréchet	Lognormal	Gamma
Fréchet(1,1)	95	5	0
Fréchet(2,1)	77	17	6
Lognormal(0,1)	16	63	21
Lognormal(1,1)	6	86	8
Gamma(2,1)	0	23	77
Gamma(2,2)	1	47	52

Table 2: Selected model frequencies for each Poisson mixture simulation with the highest frequency in bold.

This example shows the importance in the comparison of various mixing distributions. However, this approach based on systematic comparisons may suffer from computational limitations and is difficult to use in practice where a selection objective involves too much possibilities for the mixing distribution. Indeed, such an approach requires an appropriate choice of priors, a high number of MCMC iterations, and a study of convergence for each mixing distribution. Here, we assume the same priors and the same number of iterations for each mixing distribution in order to systematically simulate all our categories. In reality, each case must be studied with care and such an approach depends too much on the latent variables of the mixture model. As an alternative, we propose a simple strategy that uses directly the data and allows the user to focus on a specific family of mixing distributions. In doing so, the estimation of the latent variable can be done as a last step. The proposed alternative (see Section 3) relies on the use of a sequential approach by selecting first the most appropriate category (Gumbel, Fréchet or pseudo-Gumbel) and next by comparing only a few representative distributions belonging to this selected domain. For instance, such representative distributions are lognormal/Benktander-I for the Gumbel category, inverse-gamma/folded-Cauchy for the Fréchet category or a gamma/inverse-Gaussian for the pseudo-Gumbel case.

3 Strategy

This section proposes a strategy to choose a mixing distribution on λ using the categories defined in Section 2. As previously mentioned, an excess of zeroes and extreme values create overdispersion in count data which induces a particular tail behavior. The main idea is to choose mixing distributions among the three categories reflecting which ones best fit the empirical tail behavior. Peaks-over-threshold (POT) method [Coles, 2001] is well adapted for this purpose. This technique analyses the distribution of the excesses defined by $Y - u | Y > u$. Pickands [1975], Balkema and de Haan [1974] showed that Y belongs to a domain of attraction if and only if the distribution of the excesses can be uniformly approached by a generalized Pareto distribution (GPD) as u tends to the right endpoint of the distribution of Y . The corresponding cdf is given by

$$H_{\gamma,\sigma}(y) = \begin{cases} 1 - (1 + \gamma \frac{y}{\sigma})^{-1/\gamma} & \text{if } \gamma \neq 0 \\ 1 - \exp(-\frac{y}{\sigma}) & \text{otherwise} \end{cases} \quad (4)$$

with support \mathbb{R}_+ if $\gamma \geq 0$ or $[0; -\frac{\sigma}{\gamma}]$ if $\gamma < 0$ and where $\gamma \in \mathbb{R}$ and $\sigma > 0$ are respectively shape and scale parameters. The sign of the γ parameter is intrinsically related to the domain of attraction. Indeed, the distribution of Y belongs to \mathcal{D}_- , \mathcal{D}_0 or \mathcal{D}_+ if $\gamma < 0$, $\gamma = 0$ or $\gamma > 0$ respectively. Therefore, fitting a GPD to the excesses of the count data can inform whether or not Poisson mixture distribution belongs to a known domain of attraction and, if so, which one.

3.1 Decision tree

Suppose overdispersed count data for which we need to fit a Poisson mixture. Our strategy to select appropriate mixing distributions is based on a decision tree (Figure 2) leading to the three categories defined in Section 2. The first step consists in the selection of a threshold u large enough for the data and to fit a GPD to the excesses. The choice of u can either be based on empirical quantile or on studying the mean residual life plot [Scarrott and MacDonald, 2012] and the GPD parameters can be efficiently estimated using maximum likelihood [Coles, 2001].




Figure 2: Decision tree for Poisson mixtures

Two situations arise: first, if excesses are correctly fitted using the GPD model (left side in Figure 2), then we propose to use a distribution belonging to Gumbel or Fréchet such that the resulting Poisson mixture distribution remains in these domains. Indeed, if the excesses can be approximated by a GPD, then the Poisson mixture must be in a domain of attraction (see Pickands [1975], Balkema and de Haan [1974]). Therefore, the categories defined by Perline [1998] are an adequate set of mixing distributions to choose from. The choice of the specific category is directly based on the estimation of the shape parameter obtained at the first step. Testing whether $\gamma = 0$ or not can be done with the deviance statistic [Coles, 2001]. If it is the case, we should use a Poisson mixture in the Gumbel category. Else if γ is positive, the Fréchet category should be prioritize.

Otherwise, if γ is negative, this strategy cannot assess which mixing distribution should be used. However, no such case has been relevant in our study.

The second situation (right side in Figure 2) corresponds to the case where GPD model is not well adapted. In this case, any mixing distributions such that the Poisson mixture belongs to Gumbel or Fréchet domains of attraction should be avoided. A distribution from the pseudo-Gumbel category should potentially be favoured. As demonstrated by Shimura [2012], such discrete random variables originate from an unique continuous distribution in \mathcal{D}_0 that has been discretized. That is why we transform the excesses to continuous values thanks to a *jittering* technique consisting in the addition of a continuous random noise [Nagler, 2018]. For an application, Coeurjolly and Trépanier [2020] used this technique to study the Poisson’s median and to construct an estimator for λ . In our case, the excesses $Y - u|Y > u$ are discrete and greater or equal to 1. However the GPD with $\gamma \geq 0$ is defined on $(0, \infty)$. To transform the excesses into the same support, we jittered by subtracting a $\text{Uniform}(0, 1)$. With these “jittered” data points, we fit a GPD again by fixing $\gamma = 0$ and testing again if it is adequate. If, in this case, the fit is adequate, we consider that the data are pseudo-Gumbel. Otherwise, one should proceed to another approach in order to choose a mixing distribution. However, we rarely encountered this situation in our simulations.

3.2 Evaluation of a sequential approach for mixing distribution selection

In order to study the performance of the proposed strategy, various Poisson mixtures samples have been simulated. The decision tree is then systematically applied using the `evd` package [Stephenson, 2002] for maximum likelihood estimation of GPD parameters, the modified Anderson Darling test for the goodness-of-fit and deviance statistic for testing the nullity of the shape parameter.

Let X_1, \dots, X_m denote m i.i.d random variables ordered as $X_{(1)} \leq \dots \leq X_{(m)}$. The modified Anderson-Darling test statistic for a distribution H is defined by

$$T(X_1, \dots, X_m) = \frac{n}{2} - 2 \sum_{i=1}^m H(X_{(i)}) - \sum_{i=1}^m \left[2 - \frac{2i-1}{n} \right] \log(1 - H(X_{(i)})).$$

As presented in Chu et al. [2019], this statistic has an asymptotic distribution defined by a weighted sum of χ_1^2 . For this simulation study, the m random variables are the excesses and H is the GPD. We point that such a test works for any distribution H . However, some tests exist specifically when H is a GPD. See Falk et al. [2008] or Villaseñor-Alva and González-Estrada [2009] for examples.

Finally, to test $H_0 : \gamma = 0$ versus $H_1 : \gamma \neq 0$, we fit the two models, that is the complete one and the restricted one, evaluate the corresponding log likelihoods namely \mathcal{L}_1 and \mathcal{L}_0 and we conclude with the deviance statistic $D = 2(\mathcal{L}_1 - \mathcal{L}_0)$ which follows a χ_1^2 under suitable conditions [Coles, 2001].

The following simulation scheme is then applied in the language R:

1. For a fixed sample size n and a Poisson mixture F_M with fixed parameters, simulate the mixed Poisson observations $\mathbf{Y} := (Y_1, \dots, Y_n)$.
2. For a threshold u based on the sample \mathbf{Y} (example: 95th quantile), get the excesses $\mathbf{X} := \mathbf{Y} - u | \mathbf{Y} > u$.
3. Calculate the MLE of γ and σ of the GPD for \mathbf{X} using `evd::fpot` function using the Nelder-Mead optimization method.
4. Test the GPD for \mathbf{X} with the modified Anderson Darling test ($\alpha = 0.05$). The p-values are calculated with a bootstrap approach using 250 iterations (see Chu et al. [2019]).
5. Evaluate which category the sample is classified to using the decision tree with the following outcomes:
 - (a) If the test at step 4 for \mathbf{X} is not rejected, use the deviance statistic to find which domain of attraction \mathbf{X} belongs to. If $\gamma < 0$ is significant, the sample fails to have a category.
 - (b) Else repeat steps 3 and 4 for the jittered excesses $\mathbf{X}^c := \mathbf{X} - \text{Unif}(0, 1)$ and by fixing $\gamma = 0$. If the GPD is not rejected, the sample belongs to pseudo-Gumbel category. Otherwise the sample fails to have a category.
6. Repeat 1000 times the steps 1 to 5.

Distributions from the three categories are tested using these steps with n equal to 1000 or 2000 and the threshold fixed at the 95th and 97.5th empirical quantiles for $n = 1000$ and we add the 98.5th empirical quantile for $n = 2000$. For the Fréchet category, Fréchet and folded-Cauchy mixing distributions are simulated. For the Gumbel category, lognormal and Weibull mixing distributions have been simulated. Finally, gamma and inverse-Gaussian mixing distributions are simulated for the pseudo-Gumbel. Results are presented in Table 3 and also in the Supplementary Materials section (other sets of parameters, different sample sizes and different thresholds).

Mixing distribution	Average excesses	GPD Rejection	Category
Fréchet(1,1)	48.727	0.069	0.917 (Fréchet)
Folded-Cauchy(0,1)	48.243	0.078	0.896 (Fréchet)
Lognormal(1,1)	46.750	0.126	0.720 (Gumbel)
Weibull(0.5, 1)	46.246	0.133	0.754 (Gumbel)
Gamma(2,1)	36.200	0.704	0.635 (Pseudo-Gumbel)
Inverse-Gaussian(1,2)	38.977	0.856	0.709 (Pseudo-Gumbel)

Table 3: Average number of excesses, sample proportion of GPD rejection and of the most frequent category for the simulations with $n = 1000$ and $u = 95$ th empirical quantile

These simulations aim to assess whether the decision tree adequately identifies the Poisson mixture categories. To do so, we calculate the proportion of samples where the GPD is rejected in the first branch and the proportion of the most frequent category where the Poisson mixture is identified. For cases like the Gumbel and Fréchet categories, we should see a low GPD rejection frequency. Conversely, the pseudo-Gumbel category should have a high GPD rejection. For all cases, we should have a high proportion of samples adequately identified to their category.

As presented in Table 3, most of the Poisson mixtures simulated can be adequately identified to the appropriate category. For the mixing distributions Fréchet(1,1), folded-Cauchy(0,1), lognormal(1,1) and Weibull(0.5,1), the GPD is mostly adequate for the excesses due to its low rejection rate. For pseudo-Gumbel mixtures Gamma(2,1) and inverse-Gaussian(1,2), the GPD is mostly rejected and, once the excesses are jittered, the Gumbel domain is found. Moreover, the simulation results considering a Weibull(a,b) as a mixing distribution are relevant with the theory. Indeed, as mentioned in Table 1, the Poisson-Weibull is in the Gumbel category if its parameter a is smaller than 0.5. In comparison to the limit case Weibull(0.5, 1) in Table 3, we also tested the Weibull(1,1) and found that this mixture does not belong to Gumbel category (Table 2 Supp. Material).

Some interesting factors have been identified concerning the categorization. First, as noted by Hitz et al. [2017], the discrete excesses need a certain amount of variability in order to have a smooth adjustment to the GPD. If the variance is not high enough, it can be difficult in practice to adequately identify the mixture category. For example, based on the simulations, the Fréchet(2,1) Poisson mixture belongs to the Fréchet domain only after the excesses are jittered (Table 1 Supp. Material). This distribution has its expectation defined which results in a more stable sample compared to the Fréchet(1,1). Here, both distributions do not have a defined variance because their parameter $a \leq 2$. Still, because the Fréchet(2,1) is the limit case, it leads to a less volatile sample of excesses. This situation is identified in the lognormal case as well. Indeed, the Poisson mixture using the lognormal(0,1) has a smaller variance compare to mixture with a lognormal(1,1). In our simulations, the former case has some difficulty to be adequately identified to the Gumbel category compared to the latter (Table 2 Supp. Material).

Another important aspect is the choice of the threshold u . Indeed, the threshold affects the variance and the domain of attraction inferred by the data. For example, the lognormal(0,1) may have difficulties to be identified due to its variance, but once the u is large enough we do find the Gumbel category. For instance when $n = 1000$ and $u = 95$ th quantile, 257 over 1000 samples are classified to the Gumbel category compared to 476 over 1000 samples in the pseudo-Gumbel category. However, for the same n and $u = 97.5$ th, 790 over 1000 samples are classified to the Gumbel category compared to 64 over 1000 samples in the pseudo-Gumbel category. Therefore a larger threshold u is necessary in this case. However, it can be too large for some distributions like the gamma. In particular, when $n = 1000$ and $u = 95$ th quantile there should be in average 50 excesses, but Table 3 indicates that the gamma(2,1) has 36.2 excesses. This can be explained by the lack of different values in the right tail of the Poisson mixture which leads to an underrepresented sample of excesses. Such discrepancy greatly affect the categorisation. For example, with $n = 2000$ and $u = 98.5$ th quantile, the gamma(2,1) has in average 21.927 excesses and are mostly classified in the Gumbel domain (Table 3 Supp. Material). Clearly this inappropriate classification is due to the lack of excesses.

Finally, some pseudo-Gumbel Poisson mixtures can be very close to the Gumbel domain. For instance, the inverse-Gaussian(1, 2) is adequately classified, but the inverse-Gaussian(2, 1) is mostly classified in the Gumbel category (Table 3 Supp. Material). This can be explained by the 'closeness' property described in Theorem 1. Indeed, the density of a inverse-Gaussian(μ, σ) can be represented by

$$f(x) = C(x)x^{-3/2} \exp\left(-\frac{\sigma}{2\mu^2}x\right)$$

where $C(x) = C \exp\left(-\frac{\sigma}{2x}\right)$ with C the normalizing constant. By equation (3), $\beta = \sigma/2\mu^2$ and substituting the values $\mu = 2$ and $\sigma = 1$ gives $\beta = 1/8$. Therefore the limit defined in Theorem 1 indicates that the resulting Poisson mixture has

$$\lim_{n \rightarrow \infty} \frac{1 - F_M(n+1)}{1 - F_M(n)} = \frac{8}{9}.$$

Because this limit is pretty close to that of a long tailed distribution, i.e. the limit is near to 1, the distinction between pseudo-Gumbel and Gumbel gets blurred. To visualize how this 'closeness' affects the fit, let the parameter $\mu = 2$ be fixed and vary σ from 0.1 to 8 for the inverse-Gaussian. For each value of σ , simulate 500 samples of size $n = 2000$ from the Poisson mixture, fix the threshold u to the 97.5th empirical quantile and calculate the proportion of samples where the GPD is rejected with p-value $\alpha = 0.05$. In theory, the limit from Theorem 1 should approach 0 when σ gets larger, which implies getting further from the Gumbel domain and results into more rejection of the GPD. This result is reflected in Figure 3 when σ approaches 8.




Figure 3: Proportion of inverse-Gaussian($2, \sigma$) Poisson mixture samples (size $n = 2000$) where the GPD has been rejected ($\alpha = 0.05$) for the excesses ($u = 97.5$ th quantile) according to σ .

4 Conclusion and perspectives

Overdispersed count data are commonly observed in many applied fields and Poisson mixtures are appealing to model such data [Karlis and Xekalaki, 2005]. However, the choice of the appropriate mixing distribution is a difficult task relying mainly on empirical approaches related to modelers subjectivity or on intensive computational techniques combined with goodness-of-fit test or information criteria. In this paper, we proposed a new strategy based on the analysis of the tail behavior of the data. We extend the usual Gumbel and Fréchet domains of attraction introducing the pseudo-Gumbel category for Poisson count data. In particular, we show how tail behavior can provide a great amount of information to evaluate the mixing distributions. Based on a sequential strategy and decision tree, we proposed a useful and efficient approach to select the most appropriate category allowing to focus on a more restrictive set of potential candidates. The choice of the most appropriate distribution within a given category is not dealt with in this paper. Some strategies can be proposed helping the choice of such potential candidates. More specifically, it could be based on the simplicity either for the inferential step or for the inclusion of covariates or yet for biological interpretations. Moreover, recently, tremendous

researches have been developed to jointly model count data. For instance, the joint species distribution models are proposed extending classical species distribution models in ecology [Tikhonov et al., 2020] and are often based on the use of the multivariate lognormal distribution [Aitchison and Ho, 1989, Chiquet et al., 2021]. Based on our approach, various and flexible models could be developed combining different mixing distribution belonging to different categories (Gumbel, Fréchet or pseudo-Gumbel) and the use of copulas to model dependencies structures between continuous mixing distributions [Nelsen, 2006].

Acknowledgments

This research was supported by the GAMBAS project funded by the French National Research Agency (ANR-18-CE02-0025) and the french national programme LEFE/INSU. We also thank Éric Marchand for the helpful comments and fruitful discussions.

References

- J. Aitchison and C. H. Ho. The multivariate Poisson-log normal distribution. Biometrika, 76(4):643–653, 1989.
- S. Anders and W. Huber. Differential expression analysis for sequence count data. Genome Biology, 11, 2010.
- C. W. Anderson. Extreme value theory for a class of discrete distributions with applications to some stochastic processes. Journal of Applied Probability, 7(1):99–113, 1970.
- A. A. Balkema and L. de Haan. Residual life time at great age. The Annals of Probability, 2(5):792 – 804, 1974.
- B. Bartoszewicz. Modelling the claim count with Poisson regression and negative binomial regression. In Innovations in Classification, Data Science, and Information Systems, pages 103–110. Springer Berlin Heidelberg, 2005.
- N.H. Bingham, Goldie C.M., and J.L. Teugels. Regular Variation. Cambridge University Press, 1987.
- M. G. Bulmer. On fitting the Poisson lognormal distribution to species-abundance data. Biometrics, 30:101, 1974.
- J. Carriere. Nonparametric tests for mixed Poisson distributions. Insurance: Mathematics and Economics, 12(1):3–8, 1993.
- J. Chiquet, M. Mariadassou, and S. Robin. The Poisson-lognormal model as a versatile framework for the joint analysis of species abundances. Frontiers in Ecology and Evolution, 9:188, 2021.
- J. Chu, O. Dickin, and S. Nadarajah. A review of goodness of fit tests for Pareto distributions. Journal of Computational and Applied Mathematics, 361:13–41, 2019.
- J. F. Coeurjolly and J. Rousseau Trépanier. The median of a jittered Poisson distribution. Metrika, 83:837–851, 2020.
- S. Coles. An Introduction to Statistical Modeling of Extreme Values. Springer, 2001.
- M. Falk, A. Guillou, and G. Toulemonde. A LAN based neyman smooth test for pareto distributions. Journal of Statistical Planning and Inference, 138(10):2867–2886, 2008.
- W. Feller. On a General Class of "Contagious" Distributions. The Annals of Mathematical Statistics, 14(4):389 – 400, 1943.
- M. Greenwood and G.U. Yule. An inquiry into the nature of frequency distributions representative of multiple happenings with particular reference to the occurrence of multiple attacks of disease or of repeated accidents. Journal of the Royal Statistical Society, 83:255–279, 1920.
- Q.F. Gronau, H. Singmann, and E. Wagenmakers. Bridgesampling: An R package for estimating normalizing constants. Journal of Statistical Software, 92(10):1–29, 2020.
- A. Hitz, R. Davis, and G. Samorodnitsky. Discrete extremes, 2017.

- M.S. Holla and S.K. Bhattacharya. On a discrete compound distribution. Annals of the Institute of Statistical Mathematics, 15:377–384, 1965.
- J. O. Irwin. The generalized waring distribution applied to accident theory. Journal of the Royal Statistical Society. Series A (General), 131(2):205–225, 1968.
- D. Karlis and E. Xekalaki. Mixed Poisson distributions. International Statistical Review, 73(1):35–58, 2005.
- C. Kleiber and S. Kotz. Statistical Size Distributions in Economics and Actuarial Sciences. Wiley, 2003.
- D. Lambert. Zero-inflated Poisson regression, with an application to defects in manufacturing. Technometrics, 34(1):1–14, 1992.
- X. L. Meng and W. H. Wong. Simulating ratios of normalizing constants via a simple identity: A theoretical exploration. Statistica Sinica, 6(4):831–860, 1996.
- T. Nagler. A generic approach to nonparametric function estimation with mixed data. Statistics and Probability Letters, 137:326–330, 2018.
- R.B. Nelsen. An Introduction to Copulas. Springer, 2006.
- R. Perline. Mixed Poisson distributions tail equivalent to their mixing distributions. Statistics and Probability Letters, 38(3):229–233, 1998.
- J. Pickands. Statistical inference using extreme order statistics. The Annals of Statistics, 3(1):119–131, 1975.
- S. I. Resnick. Extreme Values, Regular Variation and Point Processes. Springer, 1987.
- C. Scarrott and A. MacDonald. A review of extreme value threshold estimation and uncertainty quantification. Revstat Statistical Journal, 10:33–60, 03 2012.
- M. Shaked. On mixtures from exponential families. Journal of the Royal Statistical Society. Series B (Methodological), 42(2):192–198, 1980.
- T. Shimura. Discretization of distributions in the maximum domain of attraction. Extremes, 15(3):299–317, 2012.
- Stan Development Team. RStan: the R interface to Stan, 2020. URL <http://mc-stan.org/>. R package version 2.21.2.
- A. G. Stephenson. evd: Extreme value distributions. R News, 2(2), 2002. URL <https://CRAN.R-project.org/doc/Rnews/>.
- G. Tikhonov, Ø. H. Opedal, N. Abrego, A. Lehtikainen, M. M. J. de Jonge, J. Oksanen, and O. Ovaskainen. Joint species distribution modelling with the R-package Hmsc. Methods in Ecology and Evolution, 11(3):442–447, 2020.
- G. Tzougas. EM estimation for the Poisson-Inverse-Gamma regression model with varying dispersion: an application to insurance ratemaking. Risks, 8(3), 2020.
- J. A. Villaseñor-Alva and E. González-Estrada. A bootstrap goodness of fit test for the generalized Pareto distribution. Computational Statistics and Data Analysis, 53(11):3835–3841, 2009.
- S. J. Wenger and M. C. Freeman. Estimating species occurrence, abundance, and detection probability using zero-inflated distributions. Ecology, 89(10):2953–2959, 2008.
- G. E. Willmot. Asymptotic tail behaviour of Poisson mixtures with applications. Advances in Applied Probability, 22(1): 147–159, 1990.
- Z. Yang, J. W. Hardin, and C. L. Addy. A score test for overdispersion in Poisson regression based on the generalized Poisson-2 model. Journal of Statistical Planning and Inference, 139(4):1514–1521, 2009.