



HAL
open science

Social participation and mild cognitive impairment in low- and middle-income countries

Lee Smith, Jae-Il Shin, Guillermo Felipe López Sánchez, Hans Oh, Karel Kostev, Louis Jacob, Christopher Tejun Law, Christina Carmichael, Mark A. Tully, Ai Koyanagi

► **To cite this version:**

Lee Smith, Jae-Il Shin, Guillermo Felipe López Sánchez, Hans Oh, Karel Kostev, et al.. Social participation and mild cognitive impairment in low- and middle-income countries. *Preventive Medicine*, 2022, 164, pp.107230. 10.1016/j.ypmed.2022.107230 . hal-03888974

HAL Id: hal-03888974

<https://hal.science/hal-03888974>

Submitted on 20 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Social participation and mild cognitive impairment in low- and middle-income countries

Lee Smith^a, Jae Il Shin^b, Guillermo F. López Sánchez^{c,*}, Hans Oh^d, Karel Kostev^e, Louis Jacob^{f,g}, Christopher Tejun Law^h, Christina Carmichael^a, Mark A. Tullyⁱ, Ai Koyanagi^{f,j}

^a Centre for Health, Performance, and Wellbeing, Anglia Ruskin University, Cambridge, UK

^b Department of Pediatrics, Yonsei University College of Medicine, Seoul, Republic of Korea

^c Division of Preventive Medicine and Public Health, Department of Public Health Sciences, School of Medicine, University of Murcia, Murcia, Spain

^d Suzanne Dworak Peck School of Social Work, University of Southern California, Los Angeles, CA 90007, USA

^e University Clinic of Marburg, Germany

^f Research and Development Unit, Parc Sanitari Sant Joan de Déu, CIBERSAM, ISCIII, Dr. Antoni Pujadas, 42, Sant Boi de Llobregat, 08830, Barcelona, Spain

^g Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines, 78000 Versailles, France

^h Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

ⁱ School of Medicine, Ulster University, Londonderry, Northern Ireland, UK

^j ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

ARTICLE INFO

Keywords:

Cognition
Social engagement
Interpersonal relationships
Epidemiology
Adults
Older adults

ABSTRACT

Social participation may theoretically decrease risk for mild cognitive impairment (MCI). However, to date, no study has specifically investigated the association between social participation and MCI in LMICs, while the mediating role of loneliness is unknown. Thus, we investigated this association in a sample of adults aged ≥ 50 years from six low- and middle-income countries (LMICs; China, Ghana, India, Mexico, Russia, South Africa) using nationally representative datasets. We analyzed cross-sectional, community-based data from the Study on Global Ageing and Adult Health. A social participation score (range 0–10 with higher scores corresponding to greater levels of social participation) was created based on nine questions about involvement in community activities in the last 12 months. The National Institute on Ageing-Alzheimer's Association criteria were used to define MCI. Multivariable logistic regression and mediation analysis was performed. The analytical sample consisted of 32,715 individuals aged ≥ 50 years with preserved functional abilities [mean (SD) age 62.1 (15.6) years; 51.7% females]. In the overall sample, after adjustment for potential confounders, a one-unit increase in the social participation score was associated with a 13% decrease in odds for MCI (OR = 0.87; 95%CI = 0.82–0.93). Loneliness only explained 3.0% of the association. Greater levels of social participation were associated with a reduced odds for MCI, and this was not largely explained by loneliness. It may be prudent to implement interventions in LMICs to increase levels of social participation to aid in the prevention of MCI and ultimately dementia.

1. Introduction

Dementia is a syndrome associated with an ongoing decline of brain functioning (National Health Service, 2020). Globally, dementia is one of the major causes of disability and impaired autonomy among older people, while it leads to significant burden to economic and social systems (Wimo et al., 2017). Approximately, 55 million people live with dementia worldwide, with over 60% living in low- and middle-income countries (LMICs). Moreover, there are nearly 10 million new cases of

dementia every year (World Health Organization, 2021). As the number of older people in the global population is rising, this number is predicted to increase to approximately 78 million in 2030, and 139 million in 2050 (World Health Organization, 2021). As there are no existing treatment methods to cure dementia, identification of risk factors known to predict the onset of dementia are of utmost importance. Specifically, mild cognitive impairment (MCI) is a preclinical transitional state of dementia (Petersen et al., 1999), for which targeted interventions may be possible.

* Corresponding author.

E-mail address: gfls@um.es (G.F. López Sánchez).

<https://doi.org/10.1016/j.ypmed.2022.107230>

Received 28 March 2022; Received in revised form 18 July 2022; Accepted 28 August 2022

Available online 31 August 2022

0091-7435/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

MCI is defined as an early stage of memory loss or other cognitive ability loss (such as language or visual/spatial perception) in individuals who maintain the ability to independently perform most activities of daily living (Alzheimer's Association, 2021), and is associated with an increased conversion rate to dementia (Solfrizzi et al., 2004). Importantly, the prevalence of MCI has been observed to be high in LMICs, and particularly in older adults (McGrattan et al., 2021). Although several potential risk factors for MCI (e.g., low physical activity, stroke, depression) have been identified (Lara et al., 2016), one potentially important but understudied risk factor for MCI is that of social participation.

Social participation is defined as a person's involvement in activities that provide interaction with others in the society or the community, and represents interpersonal interactions outside the home (Aroogh and Shahboulaghi, 2020). Social participation may lead to lower risk for MCI via increased mental stimulation and strategic thinking, as well as better synaptic density and neural growth (Zunzunegui et al., 2003; Tomioka et al., 2018). Furthermore, social participation may lead to reduction in loneliness (Zhang et al., 2018), and loneliness in turn, may lead to MCI by triggering neural responses that may directly influence the development of neurodegenerative conditions. Indeed, loneliness has been identified as a risk factor for MCI (Lara et al., 2019b), also in LMICs (Smith et al., 2021).

However, to date, no research has specifically examined the association between social participation and MCI in LMICs, while the potential mediating role of loneliness in this relationship is unknown. Therefore, the aim of the present study was to investigate the association between social participation and MCI, as well as the potential mediating role of loneliness, in a sample of 32,715 individuals aged ≥ 50 years from six LMICs. We hypothesized that lower levels of social participation will be associated with higher odds for MCI, with the association being partly mediated by loneliness.

2. Methods

The Global Ageing and Adult Health (SAGE) database was used for the present study. The survey methodology has been previously published elsewhere in detail (Kowal et al., 2012). The survey was undertaken between 2007 and 2010 in six countries (China, Ghana, India, Mexico, Russia, South Africa). Importantly, these countries represent varying levels of socio-economic and demographic transition. According to the World Bank classification when the survey was carried out, Ghana was the sole low-income country, whereas China and India were classified as lower middle-income countries. All other countries were upper middle-income countries. SAGE utilized a multistage clustered sampling design to ensure that the sample was nationally representative. The sample consisted of adults aged ≥ 18 years, and people aged ≥ 50 years were oversampled. A standardized questionnaire was utilized by trained interviewers, and face-to-face interviews were conducted (mean interview duration 2.5 h). To achieve comparability between countries, standardized survey translation procedures were implemented. The survey response rates in each country were as follows: China 93%; Ghana 81%; India 68%; Mexico 53%; Russia 83%; and South Africa 75%. To take into account of the population structure as reported by the United Nations Statistical Division and non-response, sampling weights were created. Ethical approval was obtained from the WHO Ethical Review Committee as well as local ethics research review boards: China (Shanghai Municipal Centre for Disease Control and Prevention, Shanghai); Ghana (Ghana Medical School, Accra); India (International Institute of Population Sciences, Mumbai); Mexico (National Institute of Public Health, Cuernavaca); Russia (School of Preventive and Social Medicine, Russian Academy of Medical Sciences, Moscow); South Africa (Human Sciences Research Council, Pretoria). All participants provided written informed consent prior to participation in the survey.

2.1. Mild cognitive impairment

The National Institute on Ageing-Alzheimer's Association recommendations were utilized to determine MCI (Albert et al., 2011). Algorithms to identify MCI employed in previous SAGE publications were used in the present study (Koyanagi et al., 2018; Koyanagi et al., 2019). Those fulfilling all of the following criteria were categorized as having MCI:

- (a) **Concern regarding change in cognition:** This referred to answering 'bad' or 'very bad' to the question "How would you best describe your memory at present?" and/or answering 'worse' to the question "Compared to 12 months ago, would you say your memory is now better, the same or worse than it was then?"
- (b) **Impairment in at least one cognitive domains based on objective evidence:** Objective cognitive function tests included the following: the Consortium to Establish a Registry for Alzheimer's Disease word list immediate and delayed verbal recall (Morris et al., 1989), which measures both learning and episodic memory; the animal naming task (Morris et al., 1989), which assesses verbal fluency; Weschler Adult Intelligence Scale forward and backwards digit span (Tulsky and Ledbetter, 2000), which examines working and attention memory. A result of under -1 SD cut-off after adjustment for level of education, age, and country, for any of these tests corresponded to fulfilling this criterion.
- (c) **Preserved independence in functional abilities:** This was examined by utilizing questions on self-reported difficulties in basic activities of daily living (ADL) over the last 30 days (Katz et al., 1963). The actual questions were: "How much difficulty did you have in getting dressed?" and "How much difficulty did you have with eating (including cutting up your food)?" Response categories were 'none', 'mild', 'moderate', 'severe', and 'extreme (cannot do)'. Those who reported either 'none', 'mild', or 'moderate' to both of these questions were categorized as having preservation of independence in functional activities. All other individuals were removed from the analysis (935 people aged ≥ 50 years).
- (d) **Absence of dementia:** Individuals who were unable to undertake the survey due cognitive impairment were not included in the study.

2.2. Social participation

Following a protocol from previously published SAGE literature, a social participation score was derived utilizing nine questions relating to participant's involvement in community activities over the previous 12 months (e.g., attended religious services, society, club, union etc.). Response options included 'never (coded = 1)', 'once or twice per year (coded=2)', 'once or twice per month (coded=3)', 'once or twice per week (coded=4)', and 'daily (coded = 5)' (Ma et al., 2021). Responses were added and transformed to a scale ranging from 0 to 10 (Cronbach's $\alpha = 0.79$) with 10 corresponding to the highest level of social participation and 0 the lowest (See Table S1 of the supplementary material for all questions). The variable on social participation was also divided into quartiles and used as such in some analyses.

2.3. Control variables

Previous literature was consulted to select control variables (Lara et al., 2019a; Smith et al., 2021). The following variables were included as controls: age, sex, wealth quintiles based on income, education (years), physical activity, alcohol use in the last 30 days, smoking status (never, past, current), depression, diabetes, hypertension, and stroke. Levels of physical activity were derived from the Global Physical

Activity Questionnaire and were classified as low, moderate, and high using conventional cut-offs, which are provided in Table S2 of the Appendix (Bull et al., 2009). Questions derived from the World Mental Health Survey version of the Composite International Diagnostic Interview (Kessler and Üstün, 2004) were used to identify DSM-IV depression (American Psychiatric Association, 2013) (full details are provided in Table S3 of the Appendix). The presence of diabetes and stroke were based on self-reported diagnosis. Hypertension was defined as having either a systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg; or self-reported diagnosis.

2.4. Statistical analysis

Statistical analyses were performed in Stata 14.2 (Stata Corp LP, College station, Texas). MCI is an age-related condition, and thus, the analysis was only carried out on those aged ≥ 50 years. The difference in sample characteristics by level of social participation was tested by Chi-squared tests and Student's *t*-tests for categorical and continuous variables, respectively. Multivariable logistic regression analysis was used to assess the association between social participation (exposure) and MCI (outcome). In order to gain an understanding of the extent to which loneliness may explain the relation between social participation and MCI, we conducted mediation analysis. Loneliness was assessed with the question "Did you feel lonely for much of the day yesterday?" with answer options 'yes' or 'no'. The mediation analysis was conducted using the *kmb* (Karlson Holm Breen) command in Stata (Breen et al., 2013). This method is applicable in logistic regression models and decomposes the total effect of a factor into direct and indirect effects (i.e., the mediational effect). This method allows for the calculation of the mediated percentage, which is the percentage of the main association explained by the mediator. These analyses were also stratified by age groups (i.e., 50–64 and ≥ 65 years) and sex, and these stratified analyses were decided to be undertaken a priori as previous studies have shown that the correlates or risk factors of MCI differs by age or sex (Lara et al., 2016; Lindgren et al., 2019).

Country-wise analysis was also conducted, and to assess the degree of between-country heterogeneity in the association between social participation and MCI, we calculated the Higgin's I^2 based on country-wise estimates. This corresponds to the degree of heterogeneity that is not due to sampling error, with values of 25%, 50%, and 75% frequently considered as heterogeneity levels of low, moderate, and high, respectively (Higgins et al., 2003). Overall estimates were calculated based on country-wise estimates by fixed effects meta-analysis.

All regression analyses including the mediation analysis were adjusted for age, sex, wealth, education, physical activity, alcohol consumption, smoking, depression, diabetes, hypertension, stroke, and country, with the exception of the sex-stratified and country-wise analyses, which were not adjust for sex and country, respectively. To adjust for country, we included dummy variables for each country in the model, as has been done in previous SAGE publications (Koyanagi et al., 2018; Koyanagi et al., 2014). The sample weighting and the complex study design were accounted for in all analyses. Results from the regression analyses are shown as odds ratios (ORs) with 95% confidence intervals (CIs). The level of statistical significance was set at $P < 0.05$.

3. Results

The final analytical sample consisted of 32,715 individuals [China $n = 12,815$ (6812 females and 6003 males); Ghana $n = 4201$ (1999 females and 2202 males); India $n = 6191$ (3033 females and 3158 males); Mexico $n = 2070$ (1234 females and 830 males); Russia $n = 3766$ (2429 females and 1337 males); South Africa $n = 3672$ (2099 females and 1573 males)] aged ≥ 50 years with preservation in functional abilities. The mean (SD) age was 62.1 (15.6) years and 51.7% were females. The overall prevalence (95%CI) of MCI was 15.3% (14.4%–16.3%). The sample characteristics are provided in Table 1. Low levels of social

Table 1
Sample characteristics (overall and by levels of social participation).

Characteristic		Total	Social participation ^a		P-value ^b
			High	Low	
Mild cognitive impairment	No	84.7	88.6	81.4	<0.001
	Yes	15.3	11.4	18.6	
Age (years)	Mean	62.1	60.5	63.5	<0.001
	(SD)	(15.6)	(14.6)	(15.8)	
Sex	Male	48.3	55.6	42.3	<0.001
	Female	51.7	44.4	57.7	
Wealth quintiles	Poorest	16.9	14.0	19.3	<0.001
	Poorer	18.9	18.0	19.6	
	Middle	19.4	19.1	19.5	
	Richer	21.5	23.3	20.0	
Education (years)	Mean	6.1	6.4	5.8	0.002
	(SD)	(8.9)	(9.4)	(8.4)	
Physical activity	High	49.9	57.3	43.9	<0.001
	Moderate	23.0	20.8	24.8	
Alcohol consumption	Low	27.1	21.9	31.3	0.084
	No	80.9	79.9	81.7	
Smoking	Yes	19.1	20.1	18.3	<0.001
	Never	58.7	54.3	62.2	
Depression	Current	34.9	39.3	31.4	0.342
	Past	6.4	6.3	6.5	
Diabetes	No	94.5	94.1	94.9	0.959
	Yes	5.5	5.9	5.1	
Hypertension	No	93.3	93.3	93.3	<0.001
	Yes	6.7	6.7	6.7	
Stroke	No	45.2	47.9	42.9	<0.001
	Yes	54.8	52.1	57.1	
Loneliness	No	97.3	97.9	96.7	<0.001
	Yes	2.7	2.1	3.3	
Loneliness	No	89.3	90.3	88.6	0.015
	Yes	10.7	9.7	11.4	

Abbreviation: SD Standard deviation.

Data are % unless otherwise stated.

^a Low social participation corresponded to the two lowest quartiles while high social participation referred to the two highest quartiles of the social participation score which ranged from 0 to 10.

^b P-value was based on chi-squared tests and Student's *t*-tests for categorical and continuous variables, respectively.

participation were associated with higher prevalence of MCI, female sex, low levels of wealth and physical activity, never smoking, hypertension, stroke, and loneliness, while they were also associated with older age and less years of education. The mean social participation scores (range 0–10) by sample characteristics are provided in Table S4 of the Appendix. The prevalence of MCI decreased with increasing levels of social participation in the overall sample, and also in the samples stratified by age groups and sex (Fig. 1). For example, in the overall sample, among those in the lowest quartile, the prevalence of MCI was 20.9% but this decreased to 10% among those in the highest quartile. After adjustment for potential confounders, a one-unit increase in the social participation score was associated with a 13% decrease in odds for MCI in the overall sample (OR = 0.87; 95%CI = 0.82–0.93) (Table 2). Estimates were similar for those aged 50–64 years and ≥ 65 years as well as males and females. The univariable association between the social participation score or potential confounders and MCI are shown in Table S5 of the Appendix. Country-wise analysis showed that social participation is negatively associated with MCI (i.e., OR < 1) in all countries although statistical significance was not reached in all countries (Fig. 2). The overall estimate was OR = 0.90 (95%CI = 0.86–0.94) with a low level of between-country heterogeneity ($I^2 = 22.9\%$). Finally, mediation analysis showed that loneliness only explained a very small proportion of the association between social participation and MCI (Table 3). Specifically, loneliness explained 3.0% of the association in the overall sample, 4.2% in those aged ≥ 65 years, and 5.9% among females.




Fig. 1. Prevalence of mild cognitive impairment by different levels of social participation. Social participation score ranges by quartile: Quartile I (≤ 1.1); Quartile II (>1.1 to 2); Quartile III (>2 to 2.9); Quartile IV (>2.9).

4. Discussion

4.1. Main findings

This present large multi-country study among middle-aged and older aged adults from multiple LMICs found that after controlling for potential confounders, a one-unit increase in the social participation score (range 0–10) was associated with a 13% decrease in the odds for MCI in the overall sample (OR = 0.87; 95%CI = 0.82–0.93), and a low level of between-country heterogeneity was observed. Interestingly, loneliness explained just 3.0% of the association in the overall sample. However, it is important to note that the present study was cross-sectional in nature, and thus, the direction of the association could not be determined.

4.2. Interpretation of the findings

To the best of the authors' knowledge, this one of the first studies to demonstrate an association between higher levels of social participation and a reduced odds for MCI. There are several plausible pathways that likely explain this association. While loneliness can theoretically be a mediator, we found that it only explained a very small proportion of this association. Therefore, other mechanisms are likely to be more important. First, social engagement may result in preservation of or improvement in cognitive function via increased mental stimulation and strategic thinking, as well as better synaptic density and neural growth (Zunzunegui et al., 2003; Tomioka et al., 2018). Second, social participation may increase levels of social support (Cohen, 2004), and social support in turn, may increase disease resistance owing to improved physiological function (e.g., immune and neuroendocrine systems) (Skoog et al., 1996), consequently reducing the risk of developing cardiovascular diseases, which may increase risk for MCI (Roberts et al., 2013). Finally, perceived social isolation (a lack of social connections),

which may be the consequence of lack of social participation, can induce augmented stress reactivity, which is associated with prolonged activation of the hypothalamic-pituitary-adrenal axis and the sympatho-adrenal system. This represents a disrupted brain response and likely induces sleep deprivation, dysregulation of the immune system, increased levels of oxidative stress and over-expression of pro-inflammatory genes, and these all have the potential to increase risk of MCI (Lara et al., 2019b).

4.3. Implications of the study findings

Findings from the present study suggest that interventions to increase social participation in LMICs may be a useful strategy to aid in the prevention of MCI and ultimately dementia. To achieve this, multi-dimensional initiatives should be considered. For example, those focusing on social (e.g., cultural recreation, volunteering opportunities), psychosocial (e.g., well-being, quality of life), and material (e.g., access to public transportation) aspects could be broadly introduced, all of which could be significant contributors to increasing social participation (Ma et al., 2021). Findings from this study also suggest that interventions should look beyond subjective experiences of loneliness and aim to promote broader social participation in communities.

Finally, interventions that aim to increase group physical activity participation may be particularly beneficial to promote social participation, and to prevent MCI. Such interventions have been shown to increase interpersonal interactions and connections (Franke et al., 2021), while physical activity itself may lead to reduction in risk for MCI or dementia (Gallaway et al., 2017).

4.4. Strengths and limitations

The use of large representative datasets of middle-aged and older

Table 2
Association between social participation (or covariates) and mild cognitive impairment (outcome) estimated by multivariable logistic regression.

		Overall		Age				Sex			
				50–64 years		≥65 years		Male		Female	
Characteristic		OR	95%CI	OR	95%CI	OR	95%CI	OR	95%CI	OR	95%CI
Social participation	Per unit increase	0.87***	[0.82,0.93]	0.91**	[0.86,0.97]	0.84***	[0.76,0.92]	0.85***	[0.79,0.92]	0.90**	[0.84,0.96]
Age	Per one-year increase	1.02***	[1.01,1.03]	1.03**	[1.01,1.05]	1.05***	[1.04,1.06]	1.03***	[1.02,1.04]	1.02***	[1.01,1.03]
Sex	Male	1.00		1.00		1.00					
	Female	1.22*	[1.03,1.44]	1.42***	[1.17,1.73]	1.05	[0.81,1.36]				
	Poorest	1.00		1.00		1.00		1.00		1.00	
Wealth quintiles	Poorer	0.95	[0.80,1.13]	0.93	[0.74,1.16]	0.91	[0.70,1.19]	0.90	[0.71,1.14]	0.99	[0.81,1.20]
	Middle	1.04	[0.86,1.25]	0.89	[0.71,1.13]	1.16	[0.87,1.55]	1.16	[0.87,1.54]	0.93	[0.76,1.14]
	Richer	0.69***	[0.57,0.83]	0.68**	[0.53,0.87]	0.60***	[0.47,0.77]	0.61***	[0.46,0.81]	0.75**	[0.61,0.93]
Education	Richest	0.43***	[0.35,0.53]	0.35***	[0.27,0.46]	0.52***	[0.39,0.70]	0.38***	[0.28,0.51]	0.47***	[0.37,0.61]
	Per one-year increase	0.98**	[0.96,0.99]	0.97**	[0.95,0.99]	0.99	[0.96,1.01]	0.99	[0.96,1.01]	0.97**	[0.95,0.99]
Physical activity	High	1.00		1.00		1.00		1.00		1.00	
	Moderate	0.72***	[0.63,0.82]	0.74***	[0.63,0.87]	0.77**	[0.63,0.93]	0.70***	[0.57,0.86]	0.73***	[0.63,0.85]
Alcohol consumption	Low	1.08	[0.94,1.24]	0.84	[0.69,1.02]	1.37**	[1.11,1.70]	1.09	[0.87,1.35]	1.07	[0.90,1.27]
	No	1.00		1.00		1.00		1.00		1.00	
Smoking	Yes	1.04	[0.88,1.22]	0.99	[0.82,1.20]	1.19	[0.89,1.59]	1.00	[0.82,1.23]	1.14	[0.84,1.57]
	Never	1.00		1.00		1.00		1.00		1.00	
Depression	Current	1.22*	[1.03,1.43]	1.25*	[1.04,1.51]	1.25	[0.93,1.67]	1.14	[0.91,1.43]	1.35**	[1.09,1.67]
	Past	1.22	[1.00,1.50]	1.21	[0.90,1.63]	1.22	[0.91,1.65]	1.09	[0.84,1.42]	1.56*	[1.07,2.29]
Diabetes	No	1.00		1.00		1.00		1.00		1.00	
	Yes	1.16	[0.87,1.56]	1.19	[0.82,1.72]	1.11	[0.73,1.69]	1.48	[0.86,2.53]	0.99	[0.73,1.34]
Hypertension	No	1.00		1.00		1.00		1.00		1.00	
	Yes	1.13	[0.90,1.42]	1.35	[0.96,1.90]	1.02	[0.75,1.37]	0.88	[0.55,1.42]	1.33*	[1.04,1.69]
Stroke	No	1.00		1.00		1.00		1.00		1.00	
	Yes	1.11*	[1.01,1.23]	1.07	[0.95,1.20]	1.19*	[1.00,1.42]	1.08	[0.92,1.28]	1.14	[0.99,1.31]
Stroke	No	1.00		1.00		1.00		1.00		1.00	
	Yes	1.89***	[1.46,2.44]	2.40***	[1.60,3.61]	1.55*	[1.11,2.16]	1.65*	[1.10,2.47]	2.13***	[1.53,2.96]

Abbreviation: OR Odds ratio; CI Confidence interval.

The social participation score ranged from 0 to 10 with higher scores representing higher levels of social participation.

Models are adjusted for all variables in the respective column and country.

* $p < 0.05$.

** $p < 0.01$.

*** $p < 0.001$.




Fig. 2. Country-wise association between social participation and mild cognitive impairment (outcome) estimated by multi-variable logistic regression.

Abbreviation: OR Odds ratio; CI Confidence interval.

The social participation score ranged from 0 to 10 with higher scores representing higher levels of social participation.

Models are adjusted for age, sex, wealth, education, physical activity, alcohol consumption, smoking, depression, diabetes, hypertension, and stroke.

Overall estimate was obtained by meta-analysis with fixed effects.

adults from six LMICs are clear strengths of this study. However, the results should be interpreted in light of the study’s limitations. First, the majority of variables were self-reported, potentially introducing recall

and social desirability bias into the findings. Second, the study was cross-sectional in nature and thus, it is not possible to determine the direction of the association. For example, it is possible that MCI reduces

Table 3

Mediation analysis with loneliness as the mediator in the association between social participation and mild cognitive impairment.

Sample	Effect	OR [95%CI]	P-value	%Mediated ^a
Overall	Total	0.89 [0.84,0.94]	<0.001	3.0
	Direct	0.89 [0.84,0.95]	<0.001	
	Indirect	1.00 [0.99,1.00]	0.015	
Age 50–64 years	Total	0.92 [0.86,0.98]	0.011	NA
	Direct	0.92 [0.86,0.98]	0.013	
	Indirect	1.00 [0.99,1.00]	0.164	
Age ≥ 65 years	Total	0.86 [0.78,0.95]	0.003	4.2
	Direct	0.87 [0.79,0.96]	0.004	
	Indirect	0.99 [0.99,1.00]	0.028	
Male	Total	0.87 [0.80,0.94]	0.001	NA
	Direct	0.87 [0.80,0.94]	0.001	
	Indirect	1.00 [0.99,1.00]	0.181	
Female	Total	0.91 [0.85,0.97]	0.007	5.9
	Direct	0.92 [0.86,0.98]	0.011	
	Indirect	0.99 [0.99,1.00]	0.022	

Abbreviation: OR Odds ratio; CI Confidence interval.

The social participation score ranged from 0 to 10 with higher scores representing higher levels of social participation.

Models are adjusted for age, sex, wealth, education, physical activity, alcohol consumption, smoking, depression, diabetes, hypertension, stroke, and country, with the exception of the sex-stratified analysis which was not adjusted for sex.

^a Mediated percentage was only calculated in the presence of a significant indirect effect ($P < 0.05$).

social participation (Amano et al., 2020), such that cognitive problems make it difficult to organize, find, and attend social events. Third, the sample did not include institutionalized individuals, and therefore, the results are not generalizable to this population. Finally, the ‘MCI’ group was not identified using clinical assessment but through proxy measures. Thus, it is possible for people with mild forms of dementia to have been included in our study sample. Finally, the variable on loneliness only referred to feelings of loneliness in the day prior to the survey. Thus, it is possible for the results to differ if a different timeframe was used.

5. Conclusions

Higher levels of social participation were associated with reduced odds for MCI in a sample of middle-aged and older adults in LMICs, but this association was only minimally explained by loneliness. Future studies of a longitudinal nature are now required to determine the direction of the observed association, and to provide insight into potential causality. If confirmed by longitudinal studies, interventions in LMICs to increase levels of social participation may aid in the prevention of MCI and ultimately dementia.

Funding

Dr. Guillermo F. López Sánchez is funded by the European Union – Next Generation EU.

Author contributions

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

CRedit authorship contribution statement

Lee Smith: Writing – original draft. **Jae Il Shin:** Writing – review & editing. **Guillermo F. López Sánchez:** Writing – review & editing. **Hans Oh:** Writing – review & editing. **Karel Kostev:** Writing – review & editing. **Louis Jacob:** Writing – review & editing. **Christopher Tejun Law:** Writing – review & editing. **Christina Carmichael:** Writing – review & editing. **Mark Tully:** Writing – review & editing. **Ai Koyanagi:** Writing – original draft.

Declaration of Competing Interest

None.

Acknowledgment

This paper uses data from WHO’s Study on Global Ageing and Adult Health (SAGE). SAGE is supported by the U.S. National Institute on Ageing through Interagency Agreements OGH A 04034785, YA1323–08-CN-0020, Y1-AG-1005–01 and through research grants R01-AG034479 and R21-AG034263.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jupmed.2022.107230>.

References

- Albert, M.S., DeKosky, S.T., Dickson, D., Dubois, B., Feldman, H.H., Fox, N.C., et al., 2011. The diagnosis of mild cognitive impairment due to Alzheimer’s disease: recommendations from the National Institute on Aging–Alzheimer’s Association workgroups on diagnostic guidelines for Alzheimer’s disease. *Alzheimers Dement.* 7, 270–279.
- Amano, T., Morrow-Howell, N., Park, S., 2020. Patterns of social engagement among older adults with mild cognitive impairment. *J. Gerontol.* 75, 1361–1371.
- American Psychiatric Association, 2013. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub.
- Aroogh, M.D., Shahboulaghi, F.M., 2020. Social participation of older adults: a concept analysis. *Int. J. Commun. Based Nurs. Midwifery* 8, 55.
- Association, Alzheimer’s, 2021. *Mild Cognitive Impairment (MCI) 2021*. <https://www.alz.org/alzheimers-dementia/what-is-dementia/related-conditions/mild-cognitive-impairment?lang=en-US> (Accessed 09/06/2022).
- Breen, R., Karlson, K.B., Holm, A., 2013. Total, direct, and indirect effects in logit and probit models. *Sociol. Methods Res.* 42, 164–191.
- Bull, F.C., Maslin, T.S., Armstrong, T., 2009. Global physical activity questionnaire (GPAQ): nine country reliability and validity study. *J. Phys. Act. Health* 6, 790–804.
- Cohen, S., 2004. Social relationships and health. *Am. Psychol.* 59, 676.
- Franke, T., Sims-Gould, J., Nettlefold, L., Ottoni, C., McKay, H.A., 2021. “It makes me feel not so alone”: features of the choose to move physical activity intervention that reduce loneliness in older adults. *BMC Public Health* 21, 1–15.
- Gallaway, P.J., Miyake, H., Buchowski, M.S., Shimada, M., Yoshitake, Y., Kim, A.S., et al., 2017. Physical activity: a viable way to reduce the risks of mild cognitive impairment, Alzheimer’s disease, and vascular dementia in older adults. *Brain Sci.* 7, 22.
- Higgins, J.P., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analyses. *BMJ* 327, 557–560.
- Katz, S., Ford, A.B., Moskowitz, R.W., Jackson, B.A., Jaffe, M.W., 1963. Studies of illness in the aged: the index of ADL: a standardized measure of biological and psychosocial function. *JAMA* 185, 914–919.
- Kessler, R.C., Üstün, T.B., 2004. The world mental health (WMH) survey initiative version of the world health organization (WHO) composite international diagnostic interview (CIDI). *Int. J. Methods Psychiatr. Res.* 13, 93–121.
- Kowal, P., Chatterji, S., Naidoo, N., Biritwum, R., Fan, W., Lopez Ridaura, R., et al., 2012. Data resource profile: the World Health Organization study on global AGEing and adult health (SAGE). *Int. J. Epidemiol.* 41, 1639–1649.
- Koyanagi, A., Garin, N., Olaya, B., Ayuso-Mateos, J.L., Chatterji, S., Leonardi, M., et al., 2014. Chronic conditions and sleep problems among adults aged 50 years or over in nine countries: a multi-country study. *PLoS One* 9, e114742.
- Koyanagi, A., Lara, E., Stubbs, B., Carvalho, A.F., Oh, H., Stickley, A., et al., 2018. Chronic physical conditions, multimorbidity, and mild cognitive impairment in low- and middle-income countries. *J. Am. Geriatr. Soc.* 66, 721–727.
- Koyanagi, A., Oh, H., Vancampfort, D., Carvalho, A.F., Veronese, N., Stubbs, B., et al., 2019. Perceived stress and mild cognitive impairment among 32,715 community-dwelling older adults across six low- and middle-income countries. *Gerontology* 65, 155–163.
- Lara, E., Koyanagi, A., Olaya, B., Lobo, A., Miret, M., Tyrovolas, S., et al., 2016. Mild cognitive impairment in a Spanish representative sample: prevalence and associated factors. *Int. J. Geriatr. Psychiatry* 31, 858–867.
- Lara, E., Caballero, F.F., Rico-Urbe, L.A., Olaya, B., Haro, J.M., Ayuso-Mateos, J.L., et al., 2019a. Are loneliness and social isolation associated with cognitive decline? *Int. J. Geriatr. Psychiatry* 34, 1613–1622.
- Lara, E., Martín-María, N., De la Torre-Luque, A., Koyanagi, A., Vancampfort, D., Izquierdo, A., et al., 2019b. Does loneliness contribute to mild cognitive impairment and dementia? A systematic review and meta-analysis of longitudinal studies. *Ageing Res. Rev.* 52, 7–16.
- Lindgren, N., Rinne, J.O., Palviainen, T., Kaprio, J., Vuoksimaa, E., 2019. Prevalence and correlates of dementia and mild cognitive impairment classified with different versions of the modified telephone interview for cognitive status (TICS-m). *Int. J. Geriatr. Psychiatry* 34, 1883–1891.

- Ma, R., Romano, E., Vancampfort, D., Firth, J., Stubbs, B., Koyanagi, A., 2021. Physical multimorbidity and social participation in adult aged 65 years and older from six low-and middle-income countries. *J. Gerontol.* 76, 1452–1462.
- McGrattan, A.M., Zhu, Y., Richardson, C.D., Mohan, D., Soh, Y.C., Sajjad, A., et al., 2021. Prevalence and risk of mild cognitive impairment in low and middle-income countries: a systematic review. *J. Alzheimers Dis.* 1–20.
- Morris, J.C., Heyman, A., Mohs, R.C., Hughes, J.P., van Belle, G., Fillenbaum, G., et al., 1989. The consortium to establish a registry for Alzheimer's disease (CERAD): I. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology* 39 (9), 1159. <https://doi.org/10.1212/WNL.39.9.1159>.
- National Health Service (, 2020. About Dementia 2021. <https://www.nhs.uk/conditions/dementia/about/> (Accessed 09/06/2022).
- Petersen, R.C., Smith, G.E., Waring, S.C., Ivnik, R.J., Tangalos, E.G., Kokmen, E., 1999. Mild cognitive impairment: clinical characterization and outcome. *Arch.Neurol.* 56, 303–308.
- Roberts, R.O., Geda, Y.E., Knopman, D.S., Cha, R.H., Pankratz, V.S., Boeve, B.F., et al., 2013. Cardiac disease associated with increased risk of nonamnesic cognitive impairment: stronger effect on women. *JAMA Neurol.* 70, 374–382.
- Skoog, I., Nilsson, L., Persson, G., Lernfelt, B., Landahl, S., Palmertz, B., et al., 1996. 15-year longitudinal study of blood pressure and dementia. *Lancet* 347, 1141–1145.
- Smith, L., Bloska, J., Jacob, L., Barnett, Y., Butler, L., Trott, M., et al., 2021. Is loneliness associated with mild cognitive impairment in low-and middle-income countries? *Int. J.Geriatr.Psychiatry.* 36 (9), 1345–1353. <https://doi.org/10.1002/gps.5524>.
- Solfrizzi, V., Panza, F., Colacicco, A.M., D'introno, A., Capurso, C., Torres, F., et al., 2004. Vascular risk factors, incidence of MCI, and rates of progression to dementia. *Neurology* 63, 1882–1891.
- Tomioka, K., Kurumatani, N., Hosoi, H., 2018. Social participation and cognitive decline among community-dwelling older adults: a community-based longitudinal study. *J. Gerontol.* 73, 799–806.
- Tulsky, D.S., Ledbetter, M.F., 2000. Updating to the WAIS-III and WMS-III: considerations for research and clinical practice. *Psychol.Assess.* 12, 253.
- Wimo, A., Guerchet, M., Ali, G., Wu, Y., Prina, A.M., Winblad, B., et al., 2017. The worldwide costs of dementia 2015 and comparisons with 2010. *Alzheimers Dement.* 13, 1–7.
- World Health Organization, 2021. Dementia 2020. <https://www.who.int/news-room/fact-sheets/detail/dementia> (Accessed 09/06/2022).
- Zhang, W., Liu, L., Tang, F., Dong, X., 2018. Social engagement and sense of loneliness and hopelessness: findings from the PINE study. *Gerontol. Geriatr. Med.* 4 (2333721418778189).
- Zunzunegui, M., Alvarado, B.E., Del Ser, T., Otero, A., 2003. Social networks, social integration, and social engagement determine cognitive decline in community-dwelling Spanish older adults. *J. Gerontol. Ser. B Psychol. Sci. Soc. Sci.* 58, S93–S100.