

Shape Coexistence In Odd And Odd-Odd Nuclei In The $A \sim 100$ Region

J. A. Pinston¹, J. Genevey¹, G. Simpson², and W. Urban³

¹*Laboratoire de Physique Subatomique et de Cosmologie,
F-38042 Grenoble Cedex, France*

²*Institut Laue-Langevin, F-38042 Grenoble Cedex, France*

³*Institute of Experimental Physics, Warsaw University
Ul. Hoza 69, PL-00-681 Warszawa, Poland*

Abstract. In the even-even nuclei around $A=100$ a transition from spherical to deformed shapes occurs from $N=58$ to $N=60$. The isotones with $N=59$ are of special interest, because they are just at the border between the two regions. Very recently, we have studied odd-neutrons and odd-odd nuclei with $N=59$, by means of prompt γ -ray spectroscopy of the spontaneous fission of ^{248}Cm , using the EUROGAM 2 multi-detector, and by measurements of μs isomers produced by fission of $^{239,241}\text{Pu}$ with thermal neutrons at ILL (Grenoble). In the latter case, the detection is based on time correlation measurements between fission fragments detected by the LOHENGRIN mass spectrometer and γ -rays or conversion electrons from the isomer decay. It was found that three shapes coexist in the odd ^{97}Sr and ^{99}Zr and two shapes coexist in the odd-odd ^{96}Rb . A simple explanation of the shape-coexistence mechanism is proposed. It is based upon the Nilsson diagram and stresses the fundamental importance of the unique parity states.

Keywords: Exotic nuclei, Shape coexistence.

PACS: 23.20.Lv, 25.85.Ca, 25.85.Ec, 21.60.Ev

INTRODUCTION

The region of neutron-rich nuclei near $A=100$ is distinctive for the sudden change in the ground-state (g.s.) properties of nuclei. In particular, for the even ^{98}Sr and ^{99}Zr isotopes a sudden onset of strong deformation at $N=60$ is observed, whereas the lighter isotopes up to $N=58$ are rather spherical. Consequently, the isotones with $N=59$ are of special interest because they are just at the border between the two regions. A good knowledge of the spectroscopic properties of these nuclei should allow a better understanding of the origin of the deformation and the nature of the shape-coexistence phenomena in this mass region. Unfortunately, the $N=59$ isotones of interest are far from the stability line and are rather difficult to study. Consequently, for several decades, only β -decay experiments were performed and only low-spin states were measured. The main progress in this mass region was recently obtained by Urban *et al.* [1] who was able to observe for the first time well developed rotational bands in ^{97}Sr and ^{99}Zr . In this work, the $N=59$ isotones were produced in the spontaneous fission of ^{248}Cm and the prompt γ -rays were measured using the EUROGAM 2 Ge array. The level scheme of ^{99}Zr is shown in figure 1. These data confirmed that the g.s. and first

the prompt γ rays were detected with the EUROGAM 2 array, while in the second one, microsecond isomers in ^{95}Kr , ^{97}Sr and ^{96}Rb , were produced by fission of $^{239,241}\text{Pu}$ and studied with the LOHENGRIN spectrometer at ILL. In the last experiments, the detection is based on time correlation between fission fragments selected by the spectrometer and the γ rays and conversion electrons from the isomers. More details on this experimental setup can be found in Ref. [2, 3].




FIGURE 2. The level schemes of the strongly deformed $[404\ 9/2^+]$ bands in ^{97}Sr , ^{99}Zr and ^{101}Zr . The levels below, mark the position of the other intrinsic configurations of moderate deformations (from [5]).

THE [404 9/2⁺] BAND IN ⁹⁹Zr, ⁹⁷Sr AND ¹⁰¹Zr

A third rotational band was latter observed in ⁹⁹Zr from the data of EUROGAM 2 experiments [4]. The band head at 1038.8 keV excitation is a *K* isomer with a half life $T_{1/2}=54$ ns, which decays to several lower energy states. The spin and parity assignment of the band g.s., $I^\pi=9/2^+$, was derived from the intensities of the various partial branching ratios and angular-correlation measurements. For the first time, the [404 9/2⁺] band was observed in this mass region. Soon after, this band was also observed in ⁹⁷Sr [5,6,7] and ¹⁰¹Zr[5]. In these three nuclei, the band head is a *K* isomer. However, for ⁹⁷Sr, where the half life was re-measured very recently, the new value found (526(13) ns) was substantially longer than the one reported by Hwang et al. (265(27) ns) [6].

In figure 2, the level schemes of the three [404 9/2⁺] bands are shown and the intrinsic excitations of the 3/2⁺, 3/2⁻ and 5/2⁻ levels, of moderate deformation, are also indicated [5]. The intraband-transition energies show very strong similarities in these three nuclei and the energy spacings in these bands have a strongly coupled character. The quadrupole moments Q_0 deduced from the experimental γ branching ratio $\Delta I=1$ to $\Delta I=2$ for intraband transitions lead to a mean deformation of $\beta_2=0.41(3)$, for these three nuclei.




FIGURE 3. Systematic of deformations for various configurations in Sr and Zr isotopes. For $N=59$, isotones three shape coexist simultaneously. Above $N=59$, the deformation saturates at a value $\beta_2=0.41$. Note that this value is comparable to the deformation found for the [404 9/2⁺] band in ⁹⁷Sr, ⁹⁹Zr and ¹⁰¹Zr (from [5]).

SHAPE COEXISTENCE IN $N=59$ ISOTONES

In figure 3, the β_2 values for the three known $[404\ 9/2^+]$ bands in ^{99}Zr , ^{97}Sr and ^{101}Zr are compared with the values found for the other bands in strontium and zirconium isotopes in $A=100$ mass region [5]. This picture shows that three shapes coexist for $N=59$ isotones. In these nuclei, the ground state and first two excited states are the neutrons $s_{1/2}$, $d_{3/2}$ and $g_{7/2}$ shell-model levels. At about 600 keV, two rotational bands, based upon the $\nu[411\ 3/2^+]$ and $\nu[541\ 3/2^-]$ configurations are present and have a mean deformation $\beta_2=0.32(2)$. In contrast, the $[404\ 9/2^+]$ bands at about 1 MeV are strongly deformed and their deformations are comparable with the maximum value found in this mass region $\beta_2=0.41(2)$. This large value is observed for several even and odd Sr and Zr nuclei above $N=59$ and is remarkably constant.

The two-neutron unique-parity states, $\nu g_{9/2}$ of the $N=4$ shell and $\nu h_{11/2}$ of the $N=5$ shell, play a dominant role in the mechanism of shape coexistence observed in this region. Figure 4 shows a portion of the Nilsson diagram, around $N=58$ region [5]. It is interesting to note that the occupation of the $[404\ 9/2^+]$ orbital, with its strong upwards slope, will favor a spherical equilibrium shape, whereas the state originating from the $g_{9/2}$ level slopes downward and thus favors deformation. For the moderate deformation bands at $N=59$, the odd unpaired neutron has the configurations $\nu[541\ 3/2^-]$ or $\nu[411\ 3/2^+]$. Consequently, in both bands the core has the same structure with one pair in the deformation-driving $\nu[550\ 1/2^-]$ orbital and the other pair in the up-sloping $\nu[404\ 9/2^+]$ orbital, working against deformation. This configuration produces the moderate deformation measured in ^{99}Zr and ^{97}Sr nuclei.




FIGURE 4. a) Portion of the Nilsson neutrons levels, b) Schematic representation of the deformed configurations in Sr and Zr isotopes (from[5]).

Promotion of one neutron of the $\nu[404\ 9/2^+]^2$ pair into a $\nu[541\ 3/2^-]$ orbital changes dramatically the situation. In this case, the core has four neutrons occupying the down

sloping $\nu[550\ 1/2^-]$ and $\nu[541\ 3/2^-]$ orbitals while the $\nu[404\ 9/2^+]$ is completely depleted. Now, the neutron spectator orbiting around the core is in the $\nu[404\ 9/2^+]$ orbital. In this mass region, this configuration produces the deformation limit, already at $N=59$. A comparable mechanism was proposed a long time ago by Kleinheinz *et al.*, [8,9] to explain shape coexistence in the lanthanide region. In this case, the unique parity neutron states $\nu h_{11/2}$ and $\nu i_{13/2}$ play a role analogous to the one of $\nu g_{9/2}$ and $\nu h_{11/2}$ in the $A=100$ region. However, it is interesting to note that the effect is simpler in the latter region because just a few levels are active and the deformation change is also more dramatic.

SHAPE COEXISTENCE IN ODD-ODD NUCLEI

To increase nuclear-structure information in the $N=59$ isotones, the odd-odd nuclei were also investigated. For this reason, ^{98}Y was recently revisited by Brant *et al.* [10] and we have also reinvestigated ^{96}Rb . The latter nucleus was previously measured with the LOHENGRIN spectrometer [11]. However, the efficiency of the γ detection was too weak to build a reliable level scheme. More recently, this efficiency was strongly improved and the γ and conversion electrons de-exciting the isomer were studied. The new level scheme is shown in figure 5. This very neutron-rich nucleus has a structure comparable to the previously known ^{98}Y . Both nuclei show rather spherical levels at low energy, while deformed states appear at about 500 keV excitation. The $(\pi[431\ 3/2^+]\ \nu[541\ 3/2^-])3^-$ and $(\pi[422\ 5/2^+]\ \nu[541\ 3/2^-])4^-$ intrinsic configurations were found for ^{96}Rb and ^{98}Y nuclei, respectively. All these neutron and proton orbitals originate from the $\pi g_{9/2}$ and $\nu h_{11/2}$ spherical unique parity states. A lower limit, $\beta_2 > 0.28$, was deduced for these two nuclei, from the experimental γ branching ratio $\Delta I=1$ to $\Delta I=2$ for intraband transitions. Moreover, the comparable behavior observed for the decay of the isomer in both nuclei, suggests that they have the same $(\pi g_{9/2}\ \nu h_{11/2})10^-$ configuration. Consequently, in the same nucleus, these unique-parity states are present in spherical and deformed configurations. The presence of a spherical yrast trap in competition with spherical levels is the consequence of a strongly attractive n-p interaction, because n and p are in coplanar orbitals. Very recently, we observed an $I^\pi=17^-$ isomer in ^{98}Zr [12], of spherical origin and about 6.5 MeV excitation. This result shows that the competition between spherical and deformed states, along the yrast line, is still active at high excitation energy.




FIGURE 5. Level scheme of the $2.0 \mu\text{s}$ isomer in ^{96}Rb . The g.s. with spin and parity $I^\pi=2^-$, the low-lying levels and the 1135 keV isomer have rather spherical configurations, while a well developed rotational band is seen at 461.6 keV excitation.

CONCLUSIONS

In this work odd and odd-odd $N=59$ isotones were investigated, using a combination of two experimental techniques. It is now well established that three shapes coexist in odd ^{97}Sr and ^{99}Zr , while two different shapes were seen in odd-odd ^{98}Y and ^{96}Rb . These new data demonstrate that the spectroscopy of odd- A and odd-odd nuclei, provides much more information on the structure of the different shapes than even-even nuclei. The theoretical interpretation of these collective excitations is based on the Nilsson diagram in the $N=58$ region and it shows that the spherical unique-parity state plays a very important role in the shape coexistence mechanism. In conclusion, a great wealth of nuclear structure information was recently gained for these odd and odd-odd $N=59$ isotones and we hope that these new results will trigger new calculations for this mass region.

ACKNOWLEDGMENTS

We thank Pr. Asghar for stimulating discussions and a careful reading of the manuscript.

REFERENCES

1. Urban W. et al., Nucl. Phys. A **689**, 605 (2001).
2. Pinston J. A. and Genevey J., J. Phys. G **30**, R57 (2004).
3. Genevey J. et al., Eur. Phys. J. A **9**, 191 (2000).
4. Urban W. et al., Eur. Phys. J. A **16**, 11 (2003).
5. Urban W. et al., Eur. Phys. J. A **22**, 241 (2004).
6. Hwang J.K. et al., Phys. Rev. C **67**, 054304 (2003).
7. Wu C. Y. et al., Phys. Rev. C **70**, nn064312 (2004).
8. Kleinheinz P. et al., Phys. Rev. Lett. **32**, 68 (1974).
9. Kleinheinz P. et al., Nucl. Phys. A **283**, 189 (1977).
10. S. Brant et al., Phys. Rev. C **69**, 034327 (2004).
11. Genevey J. et al., Phys. Rev. C **59**, 82 (1999).
12. Simpson G. et al., talk presented at this conference.