



Cognitive Barriers to Readiness to Change in Alcohol-Dependent Patients.

Anne-Pascale Le Berre, François Vabret, Cauvin Céline, Karine Pinon, Philippe Allain, Anne Lise Pitel, Francis Eustache, Hélène Beaunieux

► To cite this version:

Anne-Pascale Le Berre, François Vabret, Cauvin Céline, Karine Pinon, Philippe Allain, et al.. Cognitive Barriers to Readiness to Change in Alcohol-Dependent Patients.. Alcoholism: Clinical and Experimental Research, 2012, epub ahead of print. 10.1111/j.1530-0277.2012.01760.x . inserm-00665745

HAL Id: inserm-00665745

<https://inserm.hal.science/inserm-00665745>

Submitted on 2 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cognitive barriers to readiness to change in alcohol-dependent patients

Anne-Pascale Le Berre^{a,b,c,d}, François Vabret^{a,b,c,e}, Céline Cauvin^{a,b,c,e}, Karine Pinon^f, Philippe Allain^f, Anne-Lise Pitel^{a,b,c,d}, Francis Eustache^{a,b,c,d} and Hélène Beaunieux^{a,b,c,d}

^a INSERM, U1077, Caen, France

^b Université de Caen Basse-Normandie, UMR-S1077, Caen, France

^c Ecole Pratique des Hautes Etudes, UMR-S1077, Caen, France

^d CHU de Caen, U1077, Caen, France

^e CHU de Caen, Service d'Addictologie, Caen, France

^f Université d'Angers, UPRES EA 2646, Angers, France

Corresponding author: Dr Hélène Beaunieux, Unité de Recherche U1077, Laboratoire de Neuropsychologie, CHU Côte de Nacre, 14033 Caen Cedex, France. Phone: +33 (0)2 31 06 51 97; Fax: +33 (0)2 31 06 51 98; e-mail: helene.beaunieux@unicaen.fr

Funding

This research was funded by Inserm and the Basse-Normandie Regional Council (R07012EE).

Declaration of Interest

None of the authors has any conflicts of interest with this work, either financial or otherwise.

Abstract

Background

Patients' personal investment and readiness to change have proved to be a prerequisite for the successful treatment of alcohol addiction. The aim of this study was to determine the contribution of cognitive functions to the motivation process to abandon maladjusted behaviour in favour of a healthier lifestyle.

Methods

An adapted version of the 'readiness to change' questionnaire was completed by 31 alcohol-dependent patients after detoxification and at alcohol treatment entry. This tool is designed to assess the three main stages of motivation to change regarding alcohol consumption: precontemplation (substance abuse and no intention to stop drinking), contemplation (strong intention to change habits but ambivalent behaviour) and action (cessation of excessive alcohol consumption and behavioural changes for healthier habits) stages. Patients and 37 healthy controls also underwent an extensive neuropsychological battery including episodic memory, metamemory, executive functions and decision-making assessment.

Results

When alcohol-dependent patients were considered as a group, the mean score on the action subscale was significantly higher than the precontemplation and contemplation ones. Nevertheless, when the stage of change reached by each patient was considered individually, we found that some alcohol-dependent patients were still in the earlier precontemplation and contemplation stages. Stepwise regression analysis revealed links between impaired memory and executive functions and low motivation, and between good decision-making skills and high motivation.

Conclusions

Our results suggest that a set of complementary cognitive abilities is needed to achieve awareness and resolve ambivalence towards alcohol addiction, which is essential for activating the desire to change problematic behaviour.

Keywords: Alcohol dependence; Motivation; Episodic memory; Metamemory; Executive functions; Decision-making

1. Introduction

The psychological treatment of excessive alcohol consumption requires the initiation and development of intrinsic motivation to change addictive behaviour. This can be achieved through motivational interviewing, a client-centered and directive therapeutic style that helps patients to explore and resolve their ambivalence to enhance their desire to change their behaviour (Hettema et al., 2005; Miller and Rose, 2009). High motivation is the key to patients' engagement in clinical treatment (DiClemente et al., 1999). Prochaska and DiClemente's 'stages of change' (Prochaska and DiClemente, 1983; DiClemente, 2007) model comprises different stages of motivation, including 'precontemplation' (substance abuse and no intention of stopping drinking), 'contemplation' (strong intention of changing addictive habits but ambivalent attitude) and 'action' (cessation of excessive alcohol consumption and adoption of previously envisaged healthier lifestyle). To progress through the different stages and adopt permanent and sustainable healthier behavioural changes, patients must implement processes of change, which are covert and overt activities and experiences in which individuals engage in order to attempt to modify risky behaviour such as consciousness raising (increasing information about unhealthy drinking behaviour) and dramatic relief (experiencing and expressing feelings about drinking behaviour). It is well known to clinicians that alcohol-dependent patients who seek treatment can be at different levels of motivation, some of them being ready to change behaviour and others being still ambivalent and resistant to modify their unhealthy habits (e.g., Carney and Kivlahan, 1995; DiClemente and Hughes, 1990; Edens and Willoughby, 2000). This heterogeneity in motivational level at treatment entry could be explained by personal and contextual factors including demographic variables, perceived stress, perceived severity of alcohol dependence and reported drinking consequences, psychiatric comorbidity, environmental quality of life and positive treatment outcome expectancies (DiClemente et al., 2009); In addition to these

1 previous findings, the contribution of cognitive functions to motivation to change should be
2 also investigated.

3 Even though efficient cognitive functioning would appear to promote abstinence, the
4 results in the literature are heterogenous. Some investigations have reported that poor
5 cognitive functioning may be predictive of difficulty in maintaining long-term abstinence
6 from alcohol (Abbott and Gregson, 1981; Allsop et al., 2000; Glenn and Parsons, 1991;
7 Gregson and Taylor, 1977; Miller, 1991; Noel et al. 2002; Parsons, 1994; Parsons et al., 1990;
8 Rourke and Grant, 1999), while others have failed to find any such link (Donovan et al., 1984;
9 Jin et al., 1998; Macciocchi et al., 1989; Moriyama et al., 2002; Pitel et al., 2009). However,
10 cognitive impairments could indirectly influence clinical outcomes (abstinence or relapse) by
11 affecting the influence of readiness to change, as several studies have highlighted a strong
12 relationship between motivation to change and drinking outcome (Penberthy et al., 2007;
13 Project MATCH Research Group, 1997, 1998). While several studies have investigated a
14 potential association between cognitive functioning and treatment outcome (abstinence or
15 relapse), few researchers have considered the relationships between cognitive functioning and
16 motivation to change. Intellectual functioning, memory and executive skills have predicted
17 greater motivation to change substance use including alcohol use among people with severe
18 co-occurring mental disorders (Blume et al., 1999) and among college students with drinking
19 problems (Blume et al., 2000). In addition, Severtson et al. (2010) demonstrated an
20 association between executive functioning and motivation to enter treatment among regular
21 users of heroin and/or cocaine. Although episodic memory (Fama et al., 2004; Pitel et al.,
22 2007), executive functions (Sullivan et al., 2000; Moselhy et al., 2001; Pitel et al., 2007),
23 metamemory (Le Berre et al., 2010) and decision-making (Goudriaan et al., 2005; Noel et al.,
24 2007) are all known to be impaired in alcohol-dependent patients, only one study (Blume et
25 al., 2005) examined the relationships between episodic memory, executive functions and the

1 readiness to change drinking behaviour, as assessed by the 'readiness to change'
2 questionnaire (RTC, Rollnick et al., 1992), in alcohol-dependent patients who met either the
3 criteria for alcohol abuse or dependence but were not in treatment. This preliminary study
4 revealed links between motivation stages and verbal memory scores but did not show a
5 relationship with executive abilities, in accordance with another study conducted within the
6 same clinical population (Blume et al., 2009), Further studies are now required to determine
7 the specific memory components (learning, encoding, retrieval or recollection) which might
8 be associated with readiness to change .in alcohol-dependent patients at treatment entry.

9 The aim of the present study was therefore to define the cognitive factors that might, at
10 least partially, predict the motivational evolution to change drinking behaviour. To this end,
11 we measured motivation with the RTC and assessed a range of neuropsychological functions,
12 including episodic memory and executive functions and higher level cognitive functions such
13 as metamemory and decision making.

15 **2. Materials and Methods**

16 **2.1. Participants**

17 Two groups of participants were included in this study: 31 chronic alcohol-dependent
18 patients (mean age: 43.9 ± 7 years; range: 31-56) and 37 controls (mean age: 45.5 ± 6.1 years;
19 range: 31-60) matched for age and education. There were 25 men in the control group and 26
20 men in the alcohol-dependent group. Alcohol-dependent patients were early in abstinence
21 from alcohol (mean days of sobriety before inclusion: 12.64 ± 7.16 ; range: 7-40).

22 Age, education level and Mini Mental State Examination (MMSE, Folstein et al.,
23 1975), depression level and anxiety level scores are summarized in Table 1.

24 Performances on the MMSE were significantly poorer in the chronic alcohol-
25 dependent patients than in the controls. The depression level, measured with the self-

administered Beck Depression Inventory (BDI, Beck et al., 1961), was higher in the alcohol-dependent group than in the control group. Similarly, anxiety, measured with the State-Trait Anxiety Inventory (STAI, Spielberger et al., 1983), was higher in the alcohol-dependent group. Demographic and clinical data for patients and controls are summarized in Table 1.

Insert Table 1

Alcohol-dependent participants were recruited by clinicians on the basis of the DSM-IV criteria for alcohol dependence (American Psychiatric Association, 1994) while they were receiving treatment for alcohol dependence as in-patients at Caen University Hospital. Patients had no history of other types of substance dependence (except tobacco), and they had already completed detoxification when they were included in the study. They were interviewed to constitute a review of their alcohol use history and typical use patterns prior to treatment including the length of alcohol-dependence and their usual daily alcohol intake over the six months prior to treatment (Table 1). None of the participants was taking psychotropic medication, displayed psychiatric problems or had any history of pathology (head injury, coma, epilepsy, Wernicke's encephalopathy, cirrhosis or depression) that might have affected their cognitive functions. Control participants were interviewed to confirm they did not drink more than 21 (men) or 14 (women) weekly standard drinks and no more than four at a time (World Health Organization criteria). Prior to inclusion, all the participants filled in a health questionnaire including questions about alcohol and drug consumption. None of the controls had a history of drug abuse. All the participants gave their informed consent to the study, which was approved by the local ethics committee.

2.2. Materials and procedure

2.2.1. ‘Readiness to change’ (RTC) questionnaire

We used an adapted version of the RTC (Rollnick et al., 1992) self-report questionnaire to assess motivation and readiness to change alcohol abuse behaviour. Based on the stages of change model (Prochaska and DiClemente, 1983), it is made up of 12 questions divided into three subscales, each corresponding to a stage of change (4 questions per subscale). These three subscales correspond to the precontemplation stage (“It’s a waste of time thinking about my drinking”), the contemplation stage (“I enjoy my drinking but sometimes I drink too much”) and the action stage (“Anyone can talk about wanting to do something about their drinking but I am actually doing something about it”). Patients were asked to rate their agreement with statements about their alcohol consumption on a 5-point Likert-type scale ranging from ‘strongly disagree’ (-2) to ‘strongly agree’ (+2). We took the patients’ scores on each subscale (ranging from -8 to 8) as our variables. The subscale with the highest score indicated which motivational stage the patient had reached at the time of the testing. When two or more subscale scores were equal, we considered that the scale farthest along the continuum of change (precontemplation-contemplation-action) represented the patient’s stage of change. The interpretation of the motivation measures must be done as follow. A higher score on action subscale than on precontemplation and contemplation subscales refers to high level of motivation. Inversely, a higher score on precontemplation and contemplation subscales than on action subscale refers to low level of motivation

2.2.2. Neuropsychological assessments (a description of each task is provided in the online Table SM1)

Episodic memory

We assessed episodic memory with the Free and Cued Selective Reminding Test (FCSRT) (Grober and Buschke, 1987; Grober et al., 1988) for verbal learning abilities. The state of consciousness associated with memories, together with encoding and retrieval skills,

was measured with the Spondee test (Pitel et al., 2007), which includes the Remember-Know paradigm (Gardiner et al., 2002).

Metamemory

We assessed metamemory with the “Feeling-of-Knowing” (FOK) task (Pinon et al., 2005; Le Berre et al., 2010). After learning 20 cue-target pairs and prior to a recognition task, participants were asked to say whether they thought they would recognize the target afterwards. An episodic FOK accuracy measure was then calculated (Goodman-Kruskal gamma, Nelson, 1984) to assess the agreement between predicted and actual recognition performance.

Executive functions and decision-making

We assessed executive functions with a verbal fluency task (Cardebat et al., 1990; adapted version by the Groupe de Réflexion pour l’Evaluation des Fonctions EXécutives (GREFEX), Godefroy O, 2001) for spontaneous flexibility abilities and strategic search in memory, the Trail-Making Test (Reitan, 1958; adapted GREFEX version) Part B for reactive flexibility, the Stroop test (Stroop, 1935; adapted GREFEX version) for inhibition, the Hayling test (Burgess and Shallice, 1996; French adapted version by Meulemans et al., 2001) for semantic inhibition, the Modified Card Sorting Test (MCST) (Nelson, 1976; adapted GREFEX version) for cognitive flexibility, working memory, problem-solving and set-shifting abilities, and the Iowa Gambling Task (Bechara et al., 1994) for decision-making. These are well-established tests and detailed descriptions of them are widely available (Lezak et al., 2004).

2.3. Statistical analysis

We analysed the data in four steps. First, we conducted a group analysis to describe the motivational features of the alcohol-dependent group as a whole, applying a paired-samples *t*-test to the three RTC subscale scores. In addition, we analysed individual

motivation profiles to find out which stage of change each patient was, according to the RTC subscale with the highest score. Second, we investigated the effects of alcohol-dependence on episodic memory, metamemory, executive functions and decision-making skills, using an unpaired two-sample *t*-test. Third, in order to identify the potential effect of confounding factors (age, education, drinking history MMSE score and anxiety or depression levels), we calculated Pearson's correlation coefficients in the alcohol-dependent group between these factors and each motivation subscale score. Factors identified as being significantly correlated ($p < .05$) with motivational data were included as covariates in a subsequent stepwise regression analysis. Fourth, a stepwise regression analysis was carried out in the alcohol-dependent group to examine the best cognitive predictors of each RTC subscale score among episodic memory, metamemory, executive and decision-making scores. A probability level of .05 was adopted for all analyses.

3. Results

3.1. Readiness to change

Group analysis: The mean score on the action stage subscale was higher than the mean scores on both the precontemplation, $t(30) = -11.01, p < 0.001$, and contemplation stage subscales, $t(30) = -4.18, p < 0.001$ (Table 2).

Insert Table 2

Individual analyses: 21 alcohol-dependent patients scored highest on the action subscale, nine on the contemplation subscale and one on the precontemplation subscale.

3.2. Effects of alcohol-dependence on episodic memory, metamemory, executive functions and decision-making faculties

Results revealed that alcohol-dependent patients performed more poorly than controls on all the verbal episodic memory tasks (learning, encoding, retrieval and recollection), as

well as on the metamemory measure, the executive tasks (reactive flexibility, inhibition, cognitive flexibility, working memory, problem-solving and set-shifting) and the decision-making measure. Data are summarized in Table 3.

Insert Table 3

3.3. Confounding variables

As age, education, MMSE scores, psychiatric scores and drinking variables did not correlate with the precontemplation and contemplation subscale scores in the alcohol-dependent group, these variables were not included in the stepwise regression analyses for these stages. Only alcohol intake was entered as a covariate in the stepwise regression analysis conducted with the action stage score, as this drinking variable was significantly correlated with it ($r = -0.40$, $p = 0.03$). Detailed results of the correlations are provided in Table 4.

Insert Table 4

3.4. Cognitive predictors of readiness to change

Precontemplation subscale: The only cognitive predictor for the precontemplation subscale score was the level of consciousness associated with recollection (mean percentage of ‘Remember’ answers), which accounted for 20% of the variance. A lower state of consciousness significantly predicted higher precontemplation subscale scores ($r = -0.44$, $p = 0.01$) (Fig. 1A and Table 5).

Contemplation subscale: In the first step, 23% of the variance in contemplation subscale score was account for by the time spent on the Stroop interference task. The second cognitive predictor making a significant contribution to contemplation subscale score was the number of errors for the Modified Card Sorting Test (MCST) (accounting for 15% of the variance). Higher Stroop Color-Word time score and higher MCST number of errors scores

significantly predicted higher contemplation subscale scores ($r = 0.48, p = 0.006$] and [$r = 0.46, p = 0.01$] respectively) (Figure 1B and Table 5).

Action subscale: The only cognitive predictor for the action subscale score was the decision-making measure, which accounted for 19% of the variance. Higher decision-making scores significantly predicted higher action subscale scores ($r = 0.43, p = 0.01$) (Fig. 1C and Table 5).

Insert Fig. 1 and Table 5

4. Discussion

The main objective of our study was to improve our understanding of how cognitive abilities are involved in the motivation to change addictive behaviour. More specifically, we investigated the basic and higher-level cognitive functions that are known to be impaired in alcohol-dependence to explain – at least in part - why some patients have difficulty shifting from a low level of motivation to a high level of motivation. To this end, we focused our attention on the effect of episodic memory, metamemory, executive function and decision-making abilities on motivation in alcohol-dependent patients early in abstinence and at clinical treatment entry.

Concerning readiness to change for the alcohol-dependent group as a whole, the mean score on the action subscale was significantly higher than the precontemplation and contemplation mean scores. This observation is not surprising, given that those patients were at alcohol treatment entry. Nevertheless, when we adopted a more qualitative approach, investigating the stage of change reached by each individual patient, we found that some patients were still in the earlier precontemplation and contemplation stages. This finding is in accordance with clinical observation that not every patient is highly motivated to change at treatment entry (e.g., Carney and Kivlahan, 1995; DiClemente and Hughes, 1990; Edens and

Willoughby, 2000). Moreover, alcohol-dependent patients performed more poorly than controls on the cognitive tasks, in agreement with previous studies showing episodic memory impairment (Beatty et al., 1995; Fama et al., 2004; Goldstein et al., 2004; Pitel et al., 2007), executive dysfunction (Fama et al., 2004; Noel et al., 2001; Pitel et al., 2007; Tedstone et Coyle, 2004), metamemory decline (Le Berre et al., 2010) and decision-making deficit (Goudriaan et al., 2005; Noel et al., 2007) in these patients. Our results also revealed a relationship between verbal episodic memory impairment, more specifically a deficit of conscious recollection (Pitel et al., 2007), and high precontemplation scores, as well as relationships between weak executive performances and high contemplation scores, and between good decision-making skills and high action scores.

These findings suggest that patients with higher episodic memory abilities would have greater intention of changing their high-risk behaviour in the near future. More particularly, our data showed an association with a specific episodic memory score: the ‘R answers’. This measure refers to conscious recollection involving retrieval of a personal event and features about its encoding context (Gardiner et al., 2002). We can assume that higher episodic memory functioning promotes patients’ engagement in the ‘consciousness-raising’ process. This hypothesis is supported by the findings of Blume et al. (2005) suggesting that intact verbal memory performance favours awareness of difficulties arising from alcohol consumption. ‘Consciousness-raising’ refers to an increase in the level of awareness about the origins, consequences and treatment possibilities of inappropriate behaviour such as excessive alcohol consumption. It is achieved when patients are able to access information about alcohol-dependence to supplement their own knowledge about their disease.

Moreover, during the contemplation stage, patients need to complete a ‘decisional balance’ (Janis and Mann, 1977) evaluating the perceived advantages and disadvantages of consuming alcohol or quitting drinking. This ‘decisional balance’ enables them move on to

1 the next stages of change. Alcohol-dependent patients are ambivalent and oscillate between
2 their impulsive ‘limbic’ behaviour and more controlled behaviour subtended by frontal lobe
3 activity (Crews and Boettiger, 2009). Our study revealed that patients with lower motivation
4 to change also had lower executive functioning including deficits of inhibition, mental
5 flexibility, problem-solving and rule abstraction. These findings suggest that efficient
6 executive functioning is necessary to go through “decisional balance”.

7 Finally, our statistical analysis demonstrated a relationship between higher decision-
8 making abilities and higher action stage scores. This observation could imply that patients
9 early in abstinence who have problems making decisions (Noel et al., 2007) experience
10 greater difficulty resolving decisional balance and moving towards a strong desire for change.
11 They must hone their decision-making faculties in order to detect risks related to alcohol in
12 the everyday life situations. They also have to glean information about causes and outcomes
13 from past experiences and project themselves into the future. However, some patients who are
14 addicted to alcohol or other substances and who have decision-making disorders may have
15 difficulty anticipating the future. They prefer choices that bring an immediate benefit, even if
16 they are harmful in the long term (loss of work and family or social deterioration) (Bechara et
17 al., 2001; Bechara et al., 2002; Goudriaan et al., 2005; Noel et al., 2007).

18 To sum up, cognitive performance may predict the level of motivation of alcohol-
19 dependent patients at treatment entry and must be taken into account in clinical decisions.
20

21 *Clinical implications and limitations*

22 High cognitive functioning including efficient episodic memory, executive functions
23 and decision-making abilities seems essential to enhance readiness to change drinking
24 behaviour. The neuropsychological screening of possible cognitive impairments may prove to
25 be extremely useful at alcohol treatment entry and have to be taken into account to make

1 relevant clinical decisions and choose the most appropriate treatment according to the
2 patients' level of motivation. As a consequence of their cognitive deficits, some alcohol-
3 dependent patients may not be able to undergo the standard treatment in an addictology
4 department because of their low motivation. In alcohol-dependence, several studies have
5 shown that cognitive disorders are at least partly reversible with abstinence (Rourke and
6 Grant, 1999; Pitel et al., 2009). Therefore, once patients have physically completed
7 detoxification, it may be a good idea to favour the neuropsychological recovery of those who
8 display cognitive deficits by extending the period they spend without alcohol (e.g., in-patient
9 follow-up rehabilitation) before being admitted to the addictology department. It would also
10 be interesting to promote spontaneous neuropsychological recovery by using rehabilitation
11 techniques used with brain-injured patients.

12 One of the limitations of this study is that it was carried out with a very particular
13 sample of alcohol-dependent patients, as our aim was to explore readiness to change at a
14 specific point in the clinical course of alcohol dependence (i.e., at alcohol treatment entry,
15 early in abstinence). Although we are sure that cognitive performance was not influenced by
16 acute alcohol consumption, our findings cannot, nevertheless, be extended to the general
17 clinical population of alcohol-dependent patients. Therefore, other studies are needed to
18 provide information about the link between motivation and cognitive skills in patients with
19 alcohol abuse or dependence who are not receiving clinical treatment and who continue to
20 drink alcohol. Another limitation is that as the gender distribution of our alcohol-dependent
21 group reflected the gender distribution that prevails in most alcohol treatment departments in
22 France, our results are mainly representative of alcohol-dependent men. Moreover, although
23 the lack of motivation could partly be explained by cognitive impairment, the move towards
24 higher readiness to change relies on many processes of change which, in turn, rely not only on
25 higher-level cognitive functions but also on emotional and social skills including theory of

1 mind, empathy and self. Further studies are therefore needed to assess the possible
2 impairment of social cognition in alcohol-dependence and the latter's links – if any - with the
3 multidimensional phenomenon of motivation.

4

1 **Acknowledgments**

2 The authors would like to thank Julia Rivier and Nezha Bissara for their help in
3 collecting the data.

4

Reference List

- Abbott MW, Gregson RA (1981) Cognitive dysfunction in the prediction of relapse in alcoholics. *J Stud Alcohol* 42: 230-43.
- Allsop S, Saunders B, Phillips M (2000) The process of relapse in severely dependent male problem drinkers. *Addiction* 95: 95-106.
- American Psychiatric Association (1994) *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association, Washington.
- Beatty WW, Katzung VM, Moreland VJ, Nixon SJ (1995) Neuropsychological performance of recently abstinent alcoholics and cocaine abusers. *Drug Alcohol Depend* 37: 247-253.
- Bechara A, Damasio AR, Damasio H, Anderson SW (1994) Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* 50: 7-15.
- Bechara A, Dolan S, Denburg N, Hindes A, Anderson SW, Nathan PE (2001) Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia* 39: 376-389.
- Bechara A, Dolan S, Hindes A (2002) Decision-making and addiction (part II): myopia for the future or hypersensitivity to reward? *Neuropsychologia* 40: 1690-1705.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J (1961) An inventory for measuring depression. *Arch Gen Psychiatry* 4: 561-571.

- 1 Blume AW, Davis JM, Schmaling KB (1999) Neurocognitive dysfunction in dually-
2 diagnosed patients: a potential roadblock to motivating behavior change. J
3 Psychoactive Drugs 31: 111-115.
- 4 Blume AW, Marlatt GA (2009) The role of executive cognitive functions in changing
5 substance use: what we know and what we need to know. Ann Behav Med 37: 117-
6 125.
- 7 Blume AW, Marlatt GA, Schmaling KB (2000) Executive cognitive function and heavy
8 drinking behavior among college students. Psychol Addict Behav 14: 299-302.
- 9 Blume AW, Schmaling KB, Marlatt GA (2005) Memory, executive cognitive function, and
10 readiness to change drinking behavior. Addict Behav 30: 301-314.
- 11 Burgess PW, Shallice T (1996) Response suppression, initiation and strategy use following
12 frontal lobe lesions. Neuropsychologia 34: 263-272.
- 13 Cardebat D, Doyon B, Puel M, Goulet P, Joanette Y (1990) Formal and semantic lexical
14 evocation in normal subjects. Performance and dynamics of production as a function
15 of sex, age and educational level. Acta Neurol Belg 90: 207-217.
- 16 Carney MM, Kivlahan DR (1995) Motivational subtypes among veterans seeking substance
17 abuse treatment: profiles based on stages of change. Psychol Addict Behav 9: 135-
18 142.
- 19 Crews FT, Boettiger CA (2009) Impulsivity, frontal lobes and risk for addiction. Pharmacol
20 Biochem Behav 93: 237-247.
- 21 DiClemente CC (2007) Mechanisms, determinants and processes of change in the
22 modification of drinking behavior. Alcohol Clin Exp Res 31: 13s-20s.

- 1 DiClemente CC, Bellino LE, Neavins TM (1999) Motivation for change and alcoholism
2 treatment. *Alcohol Res Health* 23: 86-92.
- 3 DiClemente CC, Doyle SR, Donovan D (2009) Predicting treatment seekers' readiness to
4 change their drinking behaviour in the COMBINE study. *Alcohol Clin Exp Res* 33:
5 879-892.
- 6 DiClemente CC, Hughes SO (1990) Stages of change profiles in outpatient alcoholism
7 treatment. *J subst Abuse* 2: 217-235.
- 8 Donovan DM, Kivlahan DR, Walker RD (1984) Clinical limitations of neuropsychological
9 testing in predicting treatment outcome among alcoholics. *Alcohol Clin Exp Res* 8:
10 470-5.
- 11 Edens JF, Willoughby FW (2000) Motivational patterns of alcohol dependent patients: A
12 replication. *Psychol Addict Behav* 14: 397-400.
- 13 Fama R, Pfefferbaum A, Sullivan EV (2004) Perceptual learning in detoxified alcoholic men:
14 contributions from explicit memory, executive function, and age. *Alcohol Clin Exp*
15 *Res* 28: 1657-1665.
- 16 Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for
17 grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12: 189-198.
- 18 Gardiner JM, Ramponi C, Richardson-Klavehn A (2002) Recognition memory and decision
19 processes: a meta-analysis of remember, know, and guess responses. *Memory* 10: 83-
20 98.
- 21 Glenn SW, Parsons OA (1991) Prediction of resumption of drinking in posttreatment
22 alcoholics. *Int J Addict* 26: 237-54.

- 1 Godefroy O (2001) Groupe de Réflexion sur l'Evaluation des Fonctions Exécutives
2 (GREFEX). L'évaluation des fonctions exécutives en pratique clinique. Rev
3 Neuropsychol 11: 383-434.
- 4 Goldstein RZ, Leskovjan AC, Hoff AL, Hitzemann R, Bashan F, Khalsa SS, Wang GJ,
5 Fowler JS, Volkow ND (2004) Severity of neuropsychological impairment in
6 cocaine and alcohol addiction: association with metabolism in the prefrontal cortex.
7 Neuropsychologia 42: 1447-1458.
- 8 Goudriaan AE, Oosterlaan J, de Beurs E, van den Brink W (2005) Decision making in
9 pathological gambling: a comparison between pathological gamblers, alcohol
10 dependents, persons with Tourette syndrome, and normal controls. Brain Res Cogn
11 Brain Res 23: 137-151.
- 12 Gregson RA, Taylor GM (1977) Prediction of relapse in men alcoholics. J Stud Alcohol 38:
13 1749-60.
- 14 Grober E, Buschke H (1987) Genuine memory deficits in dementia. Dev Neuropsychol 3: 13-
15 36.
- 16 Grober E, Buschke H, Crystal H, Bang S, Dresner R (1988) Screening for dementia by
17 memory testing. Neurology 38: 900-903.
- 18 Hettema J, Steele J, Miller WR (2005) Motivational interviewing. Annu Rev Clin Psychol 1:
19 91-111.
- 20 Janis IL, Mann L (1977) Emergency decision making: a theoretical analysis of responses to
21 disaster warnings. J Human Stress 3: 35-45.

- 1 Jin H, Rourke SB, Patterson TL, Taylor MJ, Grant I (1998) Predictors of relapse in long-term
2 abstinent alcoholics. *J Stud Alcohol* 59: 640-6.
- 3 Le Berre AP, Pinon K, Vabret F, Pitel AL, Allain P, Eustache F, Beaunieux H (2010) Study
4 of metamemory in patients with chronic alcoholism using a feeling-of-knowing
5 episodic memory task. *Alcohol Clin Exp Res* 34: 1888-1898.
- 6 Lezak MD, Howieson DB, Loring DW, Hannay HJ, Fischer JS (2004) *Neuropsychological*
7 *Assessment*. 4th ed. Oxford University Press, New York.
- 8 Macciocchi SN, Ranseen JD, Schmitt FA (1989) The relationship between
9 neuropsychological impairment in alcoholics and treatment outcome at one year. *Arch*
10 *Clin Neuropsychol* 4: 365-70.
- 11 Meulemans T, Steyaert M, Vincent E (2001, May) *Evaluation des déficits d'inhibition chez*
12 *les patients traumatisés crâniens à l'aide du test de Hayling*. Poster session presented
13 at the Journées de Printemps of the Société de Neuropsychologie de Langue Française,
14 Lyons, France.
- 15 Miller L (1991) Predicting relapse and recovery in alcoholism and addiction:
16 neuropsychology, personality, and cognitive style. *J Subst Abuse Treat* 8: 277-91.
- 17 Miller WR, Rose GS (2009) Toward a theory of motivational interviewing. *Am Psychol* 64:
18 527-537.
- 19 Moriyama Y, Mimura M, Kato M, Yoshino A, Hara T, Kashima H, Kato A, Watanabe A
20 (2002) Executive dysfunction and clinical outcome in chronic alcoholics. *Alcohol Clin*
21 *Exp Res* 26: 1239-44.

1 Moselhy HF, Georgiou G, Kahn A (2001) Frontal lobe changes in alcoholism: a review of the
2 literature. *Alcohol Alcohol* 36: 357-368.

3 Nelson HE (1976) A modified card sorting test sensitive to frontal lobe defects. *Cortex* 12:
4 313-324.

5 Nelson TO (1984) A comparison of current measures of the accuracy of feeling-of-knowing
6 predictions. *Psychol Bull* 95: 109-133.

7 Noel X, Bechara A, Dan B, Hanak C, Verbanck P (2007) Response inhibition deficit is
8 involved in poor decision making under risk in nonamnesic individuals with
9 alcoholism. *Neuropsychology* 21: 778-786.

10 Noel X, Sferrazza R, Van der Linden M, Paternot J, Verhas M, Hanak C, Pelc I, Verbanck P
11 (2002) Contribution of frontal cerebral blood flow measured by (99m)Tc-Bicisate
12 spect and executive function deficits to predicting treatment outcome in alcohol-
13 dependent patients. *Alcohol Alcohol* 37: 347-54.

14 Noel X, Van der Linden M, Schmidt N, Sferrazza R, Hanak C, Le Bon O, De Mol J,
15 Kornreich C, Pelc I, Verbanck P (2001) Supervisory attentional system in
16 nonamnesic alcoholic men. *Arch Gen Psychiatry* 58: 1152-1158.

17 Parsons OA (1994) Neuropsychological measures and event-related potentials in alcoholics:
18 interrelationships, long-term reliabilities, and prediction of resumption of drinking. *J*
19 *Clin Psychol* 50: 37-46.

20 Parsons OA, Schaeffer KW, Glenn SW (1990) Does neuropsychological test performance
21 predict resumption of drinking in posttreatment alcoholics? *Addict Behav* 15: 297-
22 307.

1 Penberthy JK, Ait-Daoud N, Breton M, Kovatchev B, DiClemente CC; Johnson BA (2007)
2 Evaluating readiness and treatment seeking effects in a pharmacotherapy trial for
3 alcohol dependence. *Alcohol Clin Exp Res* 31: 1538-1544.

4 Pinon K, Allain P, Kefi MZ, Dubas F, Le Gall D (2005) Monitoring processes and
5 metamemory experience in patients with dysexecutive syndrome. *Brain Cogn* 57: 185-
6 188.

7 Pitel AL, Beaunieux H, Witkowski T, Vabret F, Guillery-Girard B, Quinette P, Desgranges B,
8 Eustache F (2007) Genuine episodic memory deficits and executive dysfunctions in
9 alcoholic subjects early in abstinence. *Alcohol Clin Exp Res* 31: 1169-1178.

10 Pitel AL, Rivier J, Beaunieux H, Vabret F, Desgranges B, Eustache F (2009) Changes in the
11 episodic memory and executive functions of abstinent and relapsed alcoholics over a
12 6-month period. *Alcohol Clin Exp Res* 33: 490-498.

13 Project MATCH Research Group (1997) Project MATCH secondary a priori hypotheses.
14 *Addiction* 92: 1671-1698.

15 Project MATCH Research Group (1998) Matching alcoholism treatments to client
16 heterogeneity: project MATCH three-year drinking outcomes. *Alcohol Clin Exp Res*
17 22:1300-1311.

18 Prochaska JO, DiClemente CC (1983) Stages and processes of self-change of smoking:
19 toward an integrative model of change. *J Consult Clin Psychol* 51: 390-395.

20 Reitan RM (1958) The validity of the Trail Making Test as an indicator of organic brain
21 damage. *Percept Mot Skills* 8: 271-276.

- 1 Rollnick S, Heather N, Gold R, Hall W (1992) Development of a short 'readiness to change'
2 questionnaire for use in brief, opportunistic interventions among excessive drinkers.
3 Br J Addict 87: 743-754.
- 4 Rourke SB, Grant I (1999) The interactive effects of age and length of abstinence on the
5 recovery of neuropsychological functioning in chronic male alcoholics: a 2-year
6 follow-up study. J Int Neuropsychol Soc 5: 234-246.
- 7 Severtson SG, von Thomsen S, Hedden SL, Latimer W (2010) The association between
8 executive functioning and motivation to enter treatment among regular users of heroin
9 and/or cocaine in Baltimore, MD. Addict Behav 35: 717-720.
- 10 Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA (1983) Manual for the State-
11 Trait Anxiety Inventory (form Y). Consulting Psychologists Press, Palo Alto, CA.
- 12 Stroop J (1935) Studies of interference in serial verbal reactions. J Exp Psychol 18: 643-662.
- 13 Sullivan EV, Rosenbloom MJ, Pfefferbaum A (2000) Pattern of motor and cognitive deficits
14 in detoxified alcoholic men. Alcohol Clin Exp Res 24: 611-621.
- 15 Tedstone D, Coyle K (2004) Cognitive impairments in sober alcoholics: performance on
16 selective and divided attention tasks. Drug Alcohol Depend 75 277-286.
- 17

Figure Legend

Fig. 1. Scatterplot illustrating the relationship between (A) the Precontemplation subscale score and the Recollection score (%); (B) the Contemplation subscale score and the inhibition score (Stroop test Colour-Word time); and (C) the Action subscale score and the decision-making measure in the alcohol-dependent group.

1 Table Legends

2 Table 1. Participants' main demographic and clinical features

3

4 Table 2. Patients' readiness to change

5

6 Table 3. Assessment of episodic memory, metamemory, executive functions and decision-
7 making in controls and alcohol-dependent patients

8

1 Table 1. Participants' main demographic and clinical features

	Controls (N = 37)	Alcohol-dependent patients (N = 31)	t	p value
Age	45.49 (6.07)	43.87 (6.97)	-1.02	0.31
<i>Range</i>	31-60	31-56		
Years of education	12.11 (3.56)	10.77 (2.14)	-1.82	0.07
<i>Range</i>	5-20	6-15		
MMSE	29.46 (0.77)	27.35 (1.68)	-6.81	0.001**
Beck Depression Inventory	1.54 (2.01)	7.68 (3.49)	/	0.001*
State-Trait Anxiety Inventory "State anxiety" (STAI Y-A)	29.22 (9.12)	33.84 (9.89)	/	0.05*
State-Trait Anxiety Inventory "Trait anxiety" (STAI Y-B)	33.22 (7.20)	50.77 (14.41)	/	0.001*
Days of sobriety before inclusion	/	12.64 (7.16)	/	/
<i>Range</i>	/	7-40	/	/
Years of alcohol-dependence	/	8.26 (8.26)	/	/
<i>Range</i>	/	0.5-33	/	/
Quantity per day (in units of alcohol)	/	21.95 (11.70)	/	/
<i>Range</i>	/	3.5-53.57	/	/

2 Data are shown as means (standard deviation).

3 ** Significant difference between alcohol-dependent patients and controls ($p < 0.05$) for unpaired parametric
4 Student's paired *t*-test

5 * Significant difference between alcohol-dependent patients and controls ($p < 0.05$) for nonparametric
6 Kolmogorov-Smirnov test
7

1 Table 2. Patients' readiness to change

Precontemplation subscale score	-2.13 (3.40)
Range	-8-5
Contemplation subscale score	1.03 (5.28)
Range	-8-8
Action subscale score	5.35 (2.83)
Range	-4-8

2 Data are shown as means (standard deviation).

3

1 Table 3. Assessment of episodic memory, metamemory, executive functions and decision-
2 making in controls and alcohol-dependent patients

	Tasks	Controls	Alcohol-dependent patients	<i>t</i>	<i>p</i> value
Verbal Episodic Memory					
Learning	sum of the three free recall scores, FCSRT	21.07 (4.98)	30.03 (5.01)	-4.29	0.001*
Encoding	recognition in the 'spontaneous' condition, Spondee test	0.86 (0.11)	0.77 (0.13)	-3.18	0.005*
Retrieval	free recall in the 'deep' condition, Spondee test	0.60 (0.16)	0.46 (0.18)	-3.08	0.005*
Recollection	R answers, Spondee test	0.77 (0.17)	0.66 (0.25)	-2.04	0.05*
Metamemory					
	FOK gamma correlation	0.25 (0.50)	-0.01 (0.54)	2.09	0.05*
Executive Functions					
Spontaneous flexibility	Verbal fluency	61.57 (12.28)	56.06 (14.95)	-1.67	0.10
Inhibition	Stroop test Colour-Word time	98.27 (22.60)	115.13 (29.51)	2.66	0.01*
	Stroop test Colour-Word errors	0.78 (1.16)	2.68 (3.65)	2.98	0.005*
	Hayling Part B time	64.76 (34.38)	76.39 (40.17)	1.29	0.20
	Hayling Part B penalties	6.67 (16.09)	7.77 (4.14)	0.40	0.71
Reactive flexibility	TMT B time	62.05 (20.50)	77.10 (34.51)	2.23	0.05*
	TMT B errors	0.24 (0.49)	3.77 (18.06)	1.19	0.24
Set-shifting Problem-solving Rule abstraction	MCST time	268.24 (98.22)	357.81 (165.88)	2.76	0.01*
	MCST categories	5.73 (0.61)	4.93 (1.55)	-2.87	0.005*
	MCST errors	5.51 (2.40)	11.06 (5.78)	2.94	0.005*
Decision-Making					
	Iowa Gambling Task	18.06 (26.33)	6.45 (17.81)	2.15	0.05*

3 Data are shown as means (standard deviation).

4 * Significant difference between alcohol-dependent patients and controls ($p < .05$) for unpaired parametric
5 Student's *t*-test

1 Table 4. Correlations between age, years of education, MMSE, psychiatric data (BDI and
2 STAI), drinking variables and ‘readiness to change’ subscales

	precontemplation subscale	contemplation subscale	action subscale
age	0.27	0.07	0.06
Years of education	-0.03	-0.16	0.01
MMSE	-0.06	-0.24	0.13
Beck Depression Inventory	0.27	0.02	-0.17
State-Trait Anxiety Inventory “State anxiety” (STAI Y-A)	0.17	0.03	0.04
State-Trait Anxiety Inventory “Trait anxiety” (STAI Y-B)	0.02	0.13	-0.24
Days of sobriety before inclusion	0.11	-0.15	0.09
Years of alcohol-dependence	0.02	0.03	-0.24
Quantity per day (in units of alcohol)	-0.14	-0.09	-0.40*

* p<0.05

1 Table 5. Cognitive predictors of ‘readiness to change’ subscale in alcohol-dependent patients

Dependent variable	Independent variable	β coefficient	Standard error of β	<i>p</i> value
precontemplation subscale	R answers	-0.44	0.17	0.013
contemplation subscale	Stroop test Colour-Word time	0.42	0.15	0.009
	MCST errors	0.39	0.15	0.015
action subscale	decision-making measure (Iowa Gambling Task)	0.43	0.17	0.015

2