



A reappraisal of instrumental magnetic measurements made in Western Europe before AD 1750: confronting historical geomagnetism and archeomagnetism

Maxime Le Goff, Yves Gallet

► To cite this version:

Maxime Le Goff, Yves Gallet. A reappraisal of instrumental magnetic measurements made in Western Europe before AD 1750: confronting historical geomagnetism and archeomagnetism. *Earth Planets and Space*, 2017, 69, pp.32. 10.1186/s40623-017-0616-4 . insu-01472186

HAL Id: insu-01472186

<https://insu.hal.science/insu-01472186>

Submitted on 20 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FULL PAPER

Open Access



A reappraisal of instrumental magnetic measurements made in Western Europe before AD 1750: confronting historical geomagnetism and archeomagnetism

Maxime Le Goff*  and Yves Gallet

Abstract

We present a new compilation and analysis of historical geomagnetic measurements made in Western Europe before AD 1750. The dataset in its ensemble provides a coherent evolution of magnetic field directions. Several data points excluded from previous analyses actually appear very consistent with most of the present compilation. A new average historical curve is computed for Paris, which is in very good agreement with the archeomagnetic data obtained in France, while significantly differing from the directional curve expected for Paris before AD 1675 based on the gufm1 model (Jackson et al. in *Philos Trans R Soc Lond A* 358:957–990, 2000). This finding suggests that the older segment of the gufm1 model lacks reliability and should be improved. Similarly, the historical part of the regional geomagnetic field model built for Europe by Pavón-Carrasco et al. (*Geochem Geophys Geosyst* 10:Q03013, 2009) should be revised because it erroneously incorporates directions derived from the gufm1 model.

Keywords: Historical magnetism, Archeomagnetism, Secular variation, Geomagnetic directions, Western Europe

Introduction


Since the dawn of archeomagnetic research, data obtained in Europe from archeological artifacts and volcanic deposits were used to build regional-scale geomagnetic directional secular variation curves (e.g., Chevalier 1925; Aitken and Hawley 1966; Tanguy 1969; Thellier 1966, 1981). An issue rapidly arose regarding the continuity between archeo-/paleomagnetic results and the most ancient instrumental or “direct” measurements of declinations (D) and inclinations (I). In Western Europe, magnetic declinations have been directly measured since the middle of the sixteenth century, whereas only a very small number of inclination data became available more than a century later (e.g., Jackson et al. 2000; Jonkers et al. 2003). These series of pioneering D and I measurements from the sixteenth and seventeenth centuries, especially those obtained in London and in Paris, were the subject

of several compilations (e.g., Malin and Bullard 1981; Alexandrescu et al. 1996, 1997). Comprehensive datasets further incorporating measurements reported in ship logbooks by mariners during their navigation around the world have been used to construct global geomagnetic field models encompassing the past four centuries, in the form of Gauss coefficient time series. At present, the gufm1 model developed by Jackson et al. (2000) is used most frequently. It enabled, in particular, studies to decipher the evolution of the main core flow patterns at the core surface (e.g., Hulot et al. 2002; Holme 2007; Finlay et al. 2016). Note that the first examples of global reconstruction of the ancient field date back to the nineteenth century; one can, for instance, mention the work of the Swedish scholar Carlheim-Gillensköld (1896), whose computations went back to the sixteenth century. More recently, a regional field model was also constructed for Europe using a spherical cap harmonic (SCHA) analysis technique incorporating both archeomagnetic and historical geomagnetic data (Pavón-Carrasco et al. 2009).

*Correspondence: legoff@ipgp.fr
Institut de Physique du Globe de Paris, Sorbonne Paris Cité, UMR 7154
CNRS, Université Paris Diderot, Paris, France

In addition to the construction of global or regional geomagnetic field models, instrumental measurements were also used to extend the directional geomagnetic field variation curves derived from archeomagnetic data. Using a compilation of direct data from Paris, London and Rome, Thellier (1981) traced the oval characterizing the directional variations in Western Europe during the historical period (back to 1600; dark-blue curve in Fig. 1), which according to Thellier was “bien connu depuis plus d’un siècle” (well known for over a century). In Thellier’s work, this oval revealed a near-perfect continuity with the French archeomagnetic data then available for the sixteenth and seventeenth centuries. This agreement was strengthened by additional archeomagnetic data obtained by Bucur (1994), who published the last compilation of French archeomagnetic data spanning the past two millennia. The orange curve in Fig. 1 displays the mean archeomagnetic directions between AD ~1000 and 1550, as computed by

Bucur (1994), whose confidence ovals at 95% were omitted for a better readability (the individual archeomagnetic data presently available between AD ~1500 and 1700 are grayed out in Fig. 1; note further that some of them are still unpublished). Its continuity with the instrumental curve argues for a large clockwise loop of the geomagnetic directions since the middle of the fourteenth century. A striking observation, however, is that this good continuity is lost when the gufm1 historical field model is used to compute the expected directional variations in Western Europe during the seventeenth century (white curve in Fig. 1). While the declination results are similar, the gufm1 inclinations appear too high with respect to the mean archeomagnetic variation curve computed by Bucur (1994). Two other historical variation curves are also reported in Fig. 1: a brown curve derived from a global field model previously constructed by Thomson and Barraclough (1982), and a black curve computed by Malin and Bullard (1981) from



D and I values measured in London and its surroundings. These two curves happen to be in better agreement with the French archeomagnetic data than the expected curve from gufm1.

The objective of this study is to better constrain the continuity and/or compatibility between the earliest Western European instrumental directional data and the available French archeomagnetic results. For this purpose, we present a new analysis of the historical geomagnetic data spanning the AD 1500–1750 time interval. This analysis therefore provides a reliability test of the directions expected in Europe from gufm1 and, as a consequence, for that of the older segment of the gufm1 model.

Instrumental magnetic directional measurements in Western Europe before AD 1750

The catalogs of historical instrumental magnetic measurements are subject to sporadic reviews. For the Western European region, practically all references can be found in the recent article of D'Ajello Caracciolo et al. (2011), whereas all records obtained worldwide contained in the global database used for the construction of the gufm1 model (Jackson et al. 2000) are available on the Web (<ftp://ftp.nmh.ac.uk/geomag/Shipborne/>).

Instrumental inclination data are scarce before AD 1750, and perhaps approximate given the rudimentary design of instruments used at that time (see illustration, for instance, in Cabeo 1628 or Kircher 1641). Note also that the ancient writings often lacked precision on dates and/or locations of measurements (see below). However, it seems certain that the observers tried to make measurements as accurate as possible. Here we took into account these results without any unfavorable consideration. Whenever possible, we returned to the original documents written by the scholars of that time.

In the present study, we associated to the pre-1750 inclination data gathered in Western Europe the declinations simultaneously measured (if existing) or the values interpolated from the declination dataset available for the same location. Where inclinations were measured at an isolated site, declinations corresponding to the site provided by the gufm1 model were assigned, assuming a good fit of declination data, which were already relatively numerous worldwide during the seventeenth century. We acknowledge that this approach is not flawless, but it appears reasonable, and there is no other means of estimating declinations at isolated locations during the sixteenth- to seventeenth-century time interval. All pre-1750 Western European data considered in the present study are reported in Table 1. Data were separated into two groups. The first group comprises values used by Emile Thellier (archives of his 1981 article) including

datasets from London, Paris and Rome. The second group includes all other direct measurements found in the gufm1 global database.

Examination of the available records

All data listed in Table 1 were reported on an equal-area projection diagram after their reduction to a common site using virtual geomagnetic poles (Fig. 2); here the chosen site is Paris ($\lambda = 48.9^\circ\text{N}$, $\phi = 2.3^\circ\text{E}$). Furthermore, in order to take into account possible small misalignment, these data are shown with an arbitrary angular error (α_{95}) of 1.0° and 0.7° for records pre- and post-dating AD 1700, respectively. These rough values are consistent with previous errors estimated by Jackson et al. (2000) and Pavón-Carrasco et al. (2009). The error was increased to 1.5° for only two particular data that were derived from aurora curtains (see below).

Group 1 data from London compiled by Malin and Bullard (1981) are the most numerous and arguably the best documented (yellow circles in Fig. 2a with corresponding dates preceded by “L” for London). Most of the older London data were previously given very low or no weight for the construction of an average curve, mainly due to an ignorance of the context in which values were recorded (see Table 3 in Malin and Bullard 1981). However, one can see that apart from L1661 and L1684 their consistency is very satisfactory with a clear directional evolution toward the West and with slightly increasing inclinations. The French data compiled by Alexandrescu et al. (1996) are much less numerous (pink circles in Fig. 2a with dates preceded by “P” for Paris). P1660 and P1668 were disregarded by Alexandrescu et al. (1996) because they did not appear in agreement with the general directional trend defined by data from London (and also following a similar comment made by J. Raulin; see §6 in Alexandrescu et al. 1996), whereas P1671 was retained. However, from Fig. 2a, one could easily argue for excluding P1671 while retaining P1660 and P1668. The unique and well-known measurement made in Rome by Athanase Kircher and consensually dated at 1640 (red circle labeled R1640 in Fig. 2a) is also in particularly good agreement with the directional trend seen from the London data. We note that in Kircher (1641), the inclination value is $65^\circ 50'$ on page 401, but $65^\circ 40'$ on page 410. Nevertheless, regarding the distribution of the data, we do not see any reason to reject this particular datum, in contrast with doubt raised by Lanza et al. (2005) and D'Ajello Caracciolo et al. (2011). The Italian dataset does not contain any other inclination value until the measurements made by Alexander von Humboldt in 1805–1806 (e.g., Cafarella et al. 1992).

Group 2 contains 16 other inclination data from Western Europe found in the global database and reported

Table 1 Synthesis of the historical directional measurements from Western Europe compiled in the present study


Location	Date	Name	Long	Lat	Local Decl.	Local Meas. Incl.	Reduced Decl.	Reduced Incl.	Refs.
Group 1									
London	1540	L1540	0	51.5	7	69.4	6.5	67.7	Malin and Bullard (1981)
London	1560	L1560	0	51.5	10	70.4	9.2	68.8	Malin and Bullard (1981)
London	1576	L1576	0	51.5	11.2	71.8	10.2	70.3	Malin and Bullard (1981)
London	1600	L1600	0	51.5	6	72	5.2	70.4	Malin and Bullard (1981)
London	1610	L1610	0	51.5	6	73	5.1	71.4	Malin and Bullard (1981)
London	1613	L1613	0	51.5	6	72.5	5.2	70.9	Malin and Bullard (1981)
London	1657	L1657	0	51.5	0	73.9	-0.5	72.3	Malin and Bullard (1981)
London	1661	L1661	0	51.5	0	70	-0.1	68.2	Malin and Bullard (1981)
London	1673	L1673	0	51.5	-2	73.7	-2.4	72	Malin and Bullard (1981)
London	1676	L1676	0	51.5	-4	73.5	-4.2	71.8	Malin and Bullard (1981)
London	1677	L1677	0	51.5	-4	73.5	-4.2	72.3	Malin and Bullard (1981)
London	1684	L1684	0	51.5	-5	71.8	-4.9	70	Malin and Bullard (1981)
London	1700	L1700	0	51.5	-8	74.5	-8	72.8	Malin and Bullard (1981)
London	1719	L1719	0	51.5	-10	73.7	-9.8	71.9	Malin and Bullard (1981)
London	1720	L1720	0	51.5	-11	75.3	-10.8	73.6	Malin and Bullard (1981)
London	1723	L1723	0	51.5	-11.2	74.5	-10.9	72.8	Malin and Bullard (1981)
London	1749	L1749	0	51.5	-17	73.5	-16.2	71.7	Malin and Bullard (1981)
London	1754	L1754	0	51.5	-18	73.3	-17.1	71.4	Malin and Bullard (1981)
Paris	1660	P1660	2.3	48.9	0.5	70	0.5	70	Alexandrescu et al. (1996)
Paris	1668	P1668	2.3	48.9	-1	72.25	-0.9	72.3	Alexandrescu et al. (1996)
Paris	1671	P1671	2.3	48.9	-1.5	75	-1.4	75	Alexandrescu et al. (1996)
Paris	1726	P1726	2.3	48.9	-13	73.66	-12.9	73.7	Alexandrescu et al. (1996)
Paris	1752	P1752	2.3	48.9	-17	72.25	-16.9	72.3	Alexandrescu et al. (1996)
Rome	1640	R1640	12.5	41.9	2	65.83	4.2	70.5	Kircher (1641)
Group 2									
Uraniborg (S)	1584	Tych1584	12.7	55.9	10.5	73.42	9.5	68.1	Tycho Brahe (Mairan 1733)
Kaufbeuren (D)	1607	Kepl1607	10.62	47.88	7.4	70	8.9	70.2	Kepler (Mairan 1733)
Ferrare (It)	1629	Ferr1629	11.63	44.83	3	62	2.8	65	Cabeo (1628)
Malta	1639	Malt1639	14.53	35.9	0.3	59.25	1.8	69.3	Kircher (1641)
Palermo (It)	1639	Pal1639	13.38	38.13	0.62	59.25	1.3	67.7	Kircher (1641)
Messina (It)	1639	Mess1639	15.55	38.22	0	59.25	0.7	67.7	Kircher (1641)
Napoli (It)	1639	Nap1639	14.25	40.83	0.5	59.25	0.3	65.8	Kircher (1641)
Lisbonne (Pt)	1639	Lisb1639	-9.12	38.73	3.5	62.75	2.4	70.5	Kircher (1641)
La Flèche (Fr)	1639	LaFl1639	0	47.38	3.9	70	3.6	71.1	Grandamicus (1645)
Rouen (Fr)	1639	Rou1639	1.08	49.43	3.6	72	3.3	71.7	Grandamicus (1645)
South Wales (GB)	1666	GB1666	-5.21	49.93	0	73.03	-2	72.3	
Bristol (GB)	1667	Brist1667	-2.57	51.45	-0.2	73.85	-1.4	72.2	
Copenhagen (Dk)	1672	Co1672	12.57	55.72	-4.1	76.3	-0.4	72.3	
Bruxelles (B)	1680	Bx1680	4.75	51	-3.6	72.72	-2.8	71.4	
Amsterdam (NL)	1686	Am1686	4.73	52.55	-5	74.53	-3.7	72.3	
South Eire	1711	Ir1711	-9.82	50.13	-9.5	74.5	-12.9	72.8	

Carthem-Gyllensköld (1896)

These data are separated into two groups. The first group contains the datasets from London, Paris and Rome, whereas the second group comprises additional data found in the gufm1 global database. All declinations from the second group were estimated using the gufm1 model (see text). Reduced decl. and incl. are the declinations and inclinations after transfer to Paris of the local directions using virtual geomagnetic poles. The data written in italics, and whose names are marked off, were eliminated from the computation in Paris of a new average historical geomagnetic directional variation curve valid for Western Europe

in Fig. 2b. Geographically, they are located in an area bounded by Malta to the south, Sweden (from the observatory Uraniborg created by Tycho Brahe) to the north, Ireland to the west and Bavaria, in Germany, to the East. In Fig. 2b, all declinations associated with the

inclination data were estimated using the gufm1 model. It is worth mentioning that two inclinations dated at 1584 and 1607 obtained by astronomers Tycho Brahe and Johannes Kepler were deduced from the continuation to the Earth's surface of rays of aurora curtains



aligned following the local direction of the geomagnetic field (Mairan 1733; see page 29 in Carlheim-Gyllensköld 1896). It is remarkable to see how these two results lie close to the overall trend defined by most available data (Fig. 2a, b). In contrast, four data from Italy appear significantly distant from this trend. One of these records was reported from Ferrare by Cabeo (1628). It is curiously dated 1629, thus 1 year after Cabeo's publication. The three others are part of a group of four strictly identical inclination values (59.25°) all dated 1639 and reported by Kircher (1641) from Malta, Palermo, Messina and Napoli. None of them were listed in the Italian catalog of Cafarella et al. (1992), but were considered by Carlheim-Gyllensköld (1896). Their distinct


positions in Fig. 2b originate from the effect of reduction to Paris because of the different latitudes of these cities. The original paragraph in which these values have been mentioned is ambiguous: Kircher (1641) wrote on page 401 that he went to Malta to perform a measurement, thus finding an inclination of 59.25° , but he did not say whether he went to the three other Italian cities to achieve other measurements. He only wrote that the three values were almost the same as the one measured in Malta, and from all the observations he knew at that time, he concluded that the inclinations were varying as a function of latitude, thus in contradiction with the data in question. Owing to this clear ambiguity, the present compilation retains only the inclination from Malta.

Discussion and concluding remarks

Figure 3 exhibits together the 41 historical geomagnetic directions discussed above. Among these directions distributed between AD 1540 and 1754, seven values are excluded following the observations made in “[Examination of the available records](#)” section (dashed circles in Fig. 3 and data in italics in Table 1). The 34 remaining values define a very consistent directional evolution. Given the large geographical distribution of the data, such good consistency indicates that any non-dipole field effect across Western Europe most probably was very limited during the historical period. Otherwise, the reduction to Paris of the D and I values via virtual geomagnetic poles would have introduced a significant scatter in the transferred values. This evolution conforms to the curve previously proposed by Thellier (1981). This is further confirmed by the computation of an average curve using the method developed by Le Goff (1990) and Le Goff et al. (2002), which is shown in black-red in Fig. 3 (see results in Additional file 1: Fig. S1 and Table S1). It is worth recalling that this method relies on bivariate statistics and on sliding windows whose durations vary as a function of the temporal distribution of data. The

only difference with Thellier’s (1981) curve concerns a shift in time of the mean directions estimated during the sixteenth and the beginning of the seventeenth century, which is likely due to the fact that Emile Thellier essentially established a hand-drawn curve and considered a smaller number of data. Conversely, our new curve is significantly distant from the directional evolution expected at Paris from the gufm1 model. Differences essentially concern the inclinations, as they are larger than the error bars estimated for the instrumental measurements, while they are weak enough to remain within the error bars estimated for the declination data (Additional file 1: Fig. S1). It is interesting to note that adding the seven data excluded from the computation would not reconcile the two curves. This likely indicates that the gufm1 model does not faithfully reproduce the geomagnetic field evolution in Western Europe between the late sixteenth century and the seventeenth century and only does so from approximately 1675 onward (Fig. 3).

When now considering the European SCHA model constructed by Pavón-Carrasco et al. (2009) that combines both archeomagnetic and historical geomagnetic data, the directional evolution expected at Paris



(light-green curve in Fig. 4) is in excellent agreement with our new curve, but again with a slight time delay for the directions during the sixteenth century (Fig. 4). It still shows a clear divergence with the gufm1 curve (light-blue curve in Fig. 4). On the other hand, it is very intriguing to see that the curve derived at Rome from the regional model and transferred to the latitude of Paris using virtual geomagnetic poles (dark-green curve in Fig. 4) is in better agreement with the gufm1 curve derived at Paris and therefore in disagreement with most of the data discussed in “[Examination of the available records](#)” section. Such a difference between the Paris and Rome directional variation curves would indicate the existence of a significant non-dipole field effect between these two regions, while the gufm1 model does not show any such effect (Fig. 4; e.g., Casas and Incoronato 2008). In fact, this difference arises from a mistake made for the computations of the regional model, previously highlighted by

D’Ajello Caracciolo et al. (2011). Pavón-Carrasco et al. (2009) indeed used a series of “instrumental” data from Viterbo, i.e., from the Italian magnetic observatory (see their Fig. 3 but with no reference), which actually corresponds to the values expected at this location from the gufm1 model. It was calculated by Lanza et al. (2005) to compensate for the very small number of Italian historical data between AD 1600 and 1805. This mistake therefore introduces a harmful interference from gufm1 into the SCHA model for the Italian region. Finally, Pavón-Carrasco et al. (2009) emphasized the existence of a possible jerk around AD 1800 (see their Fig. 10). This date is precisely at the junction between the series of “pseudo data” from Viterbo and the first real Italian geomagnetic measurements used to calculate the regional field model. The reality of a jerk at AD ~1800 therefore appears doubtful.

As concluding remarks, we highlight the following points:




Fig. 4 Geomagnetic directional field variations in Western Europe as constrained from historical data and from two geomagnetic field models (Jackson et al. 2000; Pavón-Carrasco et al. 2009). The individual archeomagnetic results available in France between AD 1500 and 1700 are grayed out, whereas the selected instrumental measurements have a reddish tint. The average instrumental curve computed in the present study is reported in black-red. The light-blue (gufm1 Paris) and light-green (SCHA Paris) curves display the directions expected at Paris from the gufm1 model (Jackson et al. 2000) and from the regional European geomagnetic field model constructed by Pavón-Carrasco et al. (2009), respectively. Similarly, the dark-blue (gufm1 Rome reduc. to Paris) and dark-green (SCHA Rome reduc. to Paris) curves show the directions expected at Rome after their transfer to the latitude of Paris using virtual geomagnetic poles

- The geomagnetic directional measurements made in Western Europe before AD ~1750 show an overall good consistency and define a coherent directional evolution in spite of their relatively large geographical coverage. Only a few discrepant values are identified.
- The new historical geomagnetic curve computed for Western Europe shows good agreement with the most recent segment of the French archeomagnetic directional variation curve (Bucur 1994). We therefore confirm the good consistency previously underlined by Thellier (1981) between the oldest historical magnetic measurements and the French archeomagnetic data.
- The directions expected from the gufm1 model do not agree with the historical geomagnetic field measurements available in Western Europe for the sixteenth and most of the seventeenth century, nor with the French archeomagnetic data. This likely indicates that the older segment of gufm1 lacks reliability and that this model should be revised. A similar inference was previously made from the comparison of archeomagnetic data obtained in La Réunion Island and gufm1-derived directions (Tanguy et al. 2011). Moreover, Genevey et al. (2009) and Hartmann et al. (2011) reached the very same conclusion from archeointensity analyses carried out on ceramics from France and Brazil.
- In their current form, neither the older part of the gufm1 model nor the recent sixteenth–eighteenth-century segment of the regional magnetic field model constructed by Pavón-Carrasco et al. (2009) can be used for archeomagnetic dating purposes or for studies of the geomagnetic field in the past millennium.

Additional file

Additional file 1. Table S1: New averaged historical geomagnetic directions computed in Paris between the sixteenth and the mid eighteenth century. **Figure S1.** New historical Western European geomagnetic directional variation curve, and its comparison with the gufm1 model.

Authors' contributions

MLG designed the study and performed the data analysis. MLG and YG discussed the data and interpretation and wrote the paper together.

Acknowledgements

We thank Jean-Claude Tanguy for stimulating recent and past discussions, Eva Fareau for her help in finding ancient documents, Suzette Roux and Michel Rival for their assistance in Latin translation. We are grateful to France Lagroix and Vincent Courtillot for a careful reading of the manuscript. We also thank F. D'Ajello Caracciolo and an anonymous reviewer for their constructive comments. This is IGP contribution no. 3817.

Competing interests

The authors declare that they have no competing interests.

Received: 3 December 2016 Accepted: 6 February 2017

Published online: 17 February 2017

References

- Aitken MJ, Hawley HN (1966) Magnetic dating—III: further archaeomagnetic measurements in Britain. *Archaeometry* 9:187–197
- Alexandrescu M, Courtillot V, Le Mouél J-L (1996) Geomagnetic field direction in Paris since the mid-sixteenth century. *Phys Earth Planet Inter* 98:321–360
- Alexandrescu M, Courtillot V, Le Mouél J-L (1997) High-resolution secular variation of the geomagnetic field in western Europe over the last 4 centuries: comparison and integration of historical data from Paris and London. *J Geophys Res* 102(B9):20245–20258
- Bucur I (1994) The direction of the terrestrial magnetic field in France during the last 21 centuries. Recent progress. *Phys Earth Planet Inter* 87:95–109
- Cabeo N (1628) *Philosophia magnetica*. Ferrare
- Cafarella L, De Santis A, Meloni A (1992) The historical Italian geomagnetic data catalogue. ING, Rome, p 160
- Carlheim-Gillensköld V (1896) Sur la forme analytique de l'attraction magnétique de la Terre exprimée en fonction du temps. J. G. Wolbach Library. Provided by the NASA Astrophysics Data System
- Casas L, Incoronato A (2008) Application of the relocation-error distribution on geomagnetic databases. Analyse on the "Historical Italian Geomagnetic Data Catalogue". *Ann Geophys* 51(4):553
- Chevalier R (1925) L'aimantation des laves de l'Etna et l'orientation du champ terrestre en Sicile. Ph.D. thesis, University of Paris, Paris, France
- D'Ajello Caracciolo F, Pignatelli A, Speranza F, Meloni A (2011) A re-evaluation of the Italian historical geomagnetic catalogue: implications for paleomagnetic dating at active Italian volcanoes. *Solid Earth* 2:65–74
- Finlay CC, Aubert J, Gillet N (2016) Gyre-driven decay of the Earth's magnetic dipole. *Nat Commun* 7:10422
- Genevey A, Gallet Y, Rosen J, Le Goff M (2009) Evidence for rapid geomagnetic field intensity variations in Western Europe over the past 800 years from new French archeomagnetic data. *Earth Planet Sci Lett* 284:132–143
- Grandamicus PJ (1645) Nova demonstratio immobilitatis terrae petita ex virtute magnetica. Societatis I E S V, La Flèche, France
- Hartmann GA, Genevey A, Gallet Y, Trindade RIF, Le Goff M, Najjar R, Etchevarne C, Afonso Holme MC (2011) New historical archeointensity data from Brazil: evidence for a large regional non-dipole field contribution over the past few centuries. *Earth Planet Sci Lett* 306:66–76
- Holme R (2007) Large-scale flow in the core. In: Olson P (ed) *In: Treatise on Geophysics*, vol 8. Elsevier, Amsterdam, pp 107–130
- Hulot G, Eymin C, Langlais B, Mandea M, Olsen N (2002) Small-scale structure of the geodynamo inferred from Oersted and Magsat satellite data. *Nature* 416:620–623
- Jackson A, Jonkers A, Walker M (2000) Four centuries of geomagnetic secular variation from historical records. *Philos Trans R Soc Lond A* 358:957–990
- Jonkers ART, Jackson A, Murray A (2003) Four centuries of geomagnetic data from historical records. *Rev Geophys* 41:1006
- Kircher A (1641) *Magnes, sive de Arte Magnetica, Opus tripartitum*. Rome, Italy (at least two other editions in 1643 and 1654)
- Lanza R, Meloni A, Tema E (2005) Historical measurements of the Earth's magnetic field compared with remanence directions from lava flows in Italy over the last four centuries. *Phys Earth Planet Inter* 148:97–107
- Le Goff M (1990) Lissage et limites d'incertitude des courbes de migration polaire: pondération des données et extension bivariable de la statistique de Fisher. *Comptes Rendus Acad Sci Sér II* 311:1191–1198
- Le Goff M, Gallet Y, Genevey A, Warmé N (2002) On archeomagnetic secular variation curves and archeomagnetic dating. *Phys Earth Planet Inter* 134:203–211
- Mairan D (1733) *Traité physique et historique de l'aurore boréale*. Imprimerie Royale, Paris, France
- Malin SRC, Bullard EG (1981) The direction of the Earth's magnetic field at London, 1570–1975. *Philos Trans R Soc Lond* 299:357–423
- Pavón-Carrasco FJ, Osete ML, Torta JM, Gaya-Piqué LR (2009) A regional archeomagnetic model for Europe for the last 3000 years, SCHA.DIF.3K: applications to archeomagnetic dating. *Geochem Geophys Geosyst* 10:Q03013

- Tanguy J-C (1969) Sur l'aimantation des laves de l'Etna et l'orientation du champ magnétique terrestre en Sicile, de 1300 à 1800. *Comptes Rendus Acad Sci Paris* 268:197
- Tanguy J-C, Bachèlery P, Le Goff M (2011) Archeomagnetism of Piton de la Fournaise: bearing on volcanic activity at La Réunion Island and geomagnetic secular variation in Southern Indian Ocean. *Earth Planet Sci Lett* 303:361–368
- Thellier E (1966) Le champ magnétique terrestre fossile. *Nucleus* 7:1–35
- Thellier E (1981) Sur la direction du champ magnétique terrestre, en France, durant les deux derniers millénaires. *Phys Earth Planet Inter* 24:89–132
- Thomson R, Barraclough DR (1982) Geomagnetic secular variation based on spherical harmonic and cross validation analysis of historical and archaeomagnetic data. *J Geomagn Geoelectr* 34:245–263

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

Date	Declin (°)	Inclin (°)	Alpha95 (°)	1/2 window (yr)
1555	8.8	68.8	1.8	30
1580	8.6	69.4	1.9	25
1600	6.9	70.2	1.7	20
1620	4.3	70.7	1.1	20
1630	2.8	70.6	1.4	10
1640	2.8	70.6	1.3	10
1655	-0.5	71.7	1.3	15
1665	-1.1	71.9	1.0	10
1675	-2.4	72.1	0.9	10
1685	-3.8	72	1.4	10
1705	-9.4	72.7	1.1	20
1715	-11.2	72.8	1.3	10
1725	-11.2	73	1.2	10
1740	-16.2	72.3	1.2	15
1750	-17	71.9	1.0	10

Table S1. New historical Western European geomagnetic directional variation curve computed in Paris using the method developed by Le Goff (1990) and Le Goff et al. (2002). The computations rely on auto-adjustable sliding windows, with a minimum of 4 data in each interval and with a minimal duration of 20 years. The columns are the following: Date, Declination in Paris, Inclination in Paris, radius of the 95% confidence circle, and half duration of the sliding window over which the direction was computed.




Figure S1

Figure S1. New historical Western European geomagnetic directional variation curve computed in Paris between the sixteenth and the middle of the eighteenth century AD using the method developed by Le Goff (1990) and Le Goff et al. (2002). (a) Evolution in declination. Red diamonds: individual data reduced to Paris with their error bars in shaded red; thick blue curve: mean variations, with their angular error bars and windows widths in thin blue. (b) Inclination evolution. Same design as in (a) but with magenta and green colors, respectively. Red dashed lines show the Declination (b) and Inclinations (a) values derived at Paris from the gufm 1 model.