AN INTERACTION BETWEEN THE REPETITIVELY PULSED XeCl LASER RADIATION AND METALS

V.M. BORISOV, A.M. KOVALEVICH and Yu.Yu. STEPANOV

The Troitsk Kurchatov Research Institute, Moscow Region, Troitsk 142092, USSR

Results are described of experiments and a theoretical analysis of the processes of interaction between the excimer radiation with the intensity of $\sim 1 \text{ GW/cm}^2$ and a metal surface. The dynamics has been investigated of the formation, heating and expansion of the plasma in the vicinity of the surface under the atmospheric pressure. The spectroscopic measurements of the luminescence from the metal vapor made it possible to determine pressure, temperature and concentration of the plasma.

An exposure of metals to the repetitively pulsed XeCl laser radiation with the intensity of 10^6 – 10^8 W/cm^2 was studied experimentally and calculated. The main channels of energy losses have been elucidated which are associated with metal evaporation, heat radiation and convective cooling by an ambient gas. An existence is shown of optimal conditions for heating of metal samples which depend on the laser radiation intensity. The optimal conditions for each of the metals under study are governed by its optical and thermophysical properties.

ENERGY PER MELTING 1g OF
METAL IN DEPENDENCE ON LASER
INTENSITY

* -Al-250 Hz	x -Al-150 Hz
△ -Cu-250 Hz	◇ -Cu-150 Hz
○ -Ti-250 Hz	□ -Ti-150 Hz