Solid state excimer lasers

G. ZERZA, G. SLIWINSKI* and N. SCHWENTNER

Institut für Experimentalphysik, FU Berlin, Arnimallee 14, 14195 Berlin, Germany

** Polish Academy of Sciences, IF-FM, Fiszerza 14, 80952 Gdansk, Poland*

Doped rare gas crystals represent an interesting group of laser materials for which stimulated emission and lasing has been demonstrated in the UV and visible spectral range [1,2]. We report the results of an extensive investigation of Ar and Ne solids binary doped with Xe and F₂ for laser applications [2,3]. The XeF population is photochemically prepared with high quantum efficiency and excited state densities in the range of 10¹⁸/cm³ are generated. The emission lines are slightly red shifted in comparison with the gas phase due to the dielectric interaction with the host. The XeF cross sections for stimulated emission of 10⁻¹⁶ – 10⁻¹⁸ cm² together with high inversion densities result in large gain coefficients, e.g. for the D → X (286 nm), B → X (411 nm) and C → A (540 nm) transitions. Values of 11,6 and 7 cm⁻¹ are measured by means of various methods. Experiments with optimized cavities allow for extraction of low divergence beams (3 mrad) with quantum efficiencies up to 14%. Inhomogeneously broadened spectral profiles and relaxation oscillations are observed in the C→A laser emission of XeF in solid Ar.

References:

- /1/ N. Schwentner, V.A. Apkarian; *Chem.Phys.Lett.* **154**,413,(1989)
- /2/ G. Zerza, G. Sliwinski and N. Schwentner; *Appl.Phys.* **B55**,331,(1992)
- /3/ G. Zerza, G. Sliwinski and N. Schwentner; *Appl.Phys.* **A56**,156,(1993)