

Theoretical Study of Localized Spin Waves in a Ferrimagnet with Impurities

O. Nagai, M. Kang, M. Tomita, Y. Yamada, H. Diep

▶ To cite this version:

O. Nagai, M. Kang, M. Tomita, Y. Yamada, H. Diep. Theoretical Study of Localized Spin Waves in a Ferrimagnet with Impurities. Journal de Physique IV Proceedings, 1997, 07 (C1), pp.C1-639-C1-640. 10.1051/jp4:19971264 . jpa-00254974

HAL Id: jpa-00254974 https://hal.science/jpa-00254974

Submitted on 4 Feb 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Theoretical Study of Localized Spin Waves in a Ferrimagnet with Impurities

O. Nagai, M. Kang, M. Tomita, Y. Yamada* and H.T. Diep**

Department of Physics, Kobe University, Kobe 657, Japan

* Department of Management, Kansai Jogakuin Women's College, Miki 673-05, Japan

** Groupe de Physique Statistique, Université de Cergy-Pontoise, 2 avenue Adolphe Chauvin,

95302 Cergy-Pontoise cedex, France

Abstract. In this paper, we theoretically investigate the impurity state in a ferrimagnet. We consider a bodycentered cubic crystal based on the model of two sublattices, A-sublattice and B-sublattice. The impurity spin (S_i) is assumed to be not located at the interstitial but located at the A-sublattice point or B-sublattice point. Confining our interests to the case of one impurity and one spin-wave, we solve our problem exactly. We have found several interesting features for our system. For example, the localized spin-wave appears in the band gap only when S_i is located at A-sublattice point if $S_A < S_B$ and the exchange coupling between the impurity spin and the neighboring S_B spin is antiferromagnetic.

1. Introduction

30

It is known that we can obtain the information concerning the physical states of the perturbation such as the surface by theoretically analyzing the experimental data on the localized spin waves [1,2]. In this paper, we develop a theory of spin waves in a ferrimagnet with impurities. Although both the experimental and theoretical investigations of ferro- and antiferromagnets including localized impurities have been widely performed [2,3], the theoretical study of ferrimagnets with impurities has not yet been made as far as we know. One of the main differences between ferrimagnetism and antiferromagnetism will be the existence of the gap between two bands. It will be worth to study how the localized spin waves at the impurity appear outside the bands, especially in the band gap.

2. Model and Calculation

Let us consider a body-centered-cubic ferrimagnet based on the model of two sublattices, up-sublattice and down-sublattice. The former and the latter are called A-sublattice and B-sublattice respectively. We confine our interests to the case of one impurity and one magnon. This will be experimentally relevant if the impurity concentration is very low and theoretically our problem can be solved exactly. We assume that the impurity spin is localized at one of the A-sublattice points. Our Hamiltonian is written as

$$H = 2J \sum_{j,n} \vec{S}_j \cdot \vec{S}_n (1 - p c_0^{\dagger} c_0)$$

where $\sum_{j,n}$ stands for the summation over the nearest neighbor site pairs. The lattice site 0 belongs to the A-sublattice. j denotes the A-sublattice sites and n denotes the B-sublattice sites throughout this paper. c^{\dagger} and c are the creation and annihilation operators for the impurity at the lattice site 0. These operators follow the Fermi commutation relations. p is a numerical parameter. J is the exchange coupling between neighboring host spins and J(1-p) = J' is the exchange coupling between the impurity and the neighboring B spin. The impurity spin is antiparallel to the neighboring spins if p < 1 and it is parallel to the neighboring spins if p > 1 in the ground state. When p = 1, our system is composed of isolated spin and a ferrimagnetic system with nonmagnetic hole. The magnitude of the spin operator S_j (S_n) is denoted by S_A (S_B). The magnitude of the impurity spin, S_i , is denoted by $S_i = S_A(1-q)$, where q is a numerical constant with 0 < q < 1.

By using Holstein-Primakoff method, the spin operators are rewritten by the spin deviation operators; $S_j^z = S_A - a_j^{\dagger}a_j, S_j^- = \sqrt{2S_A}a_j^{\dagger}, S_j^+ = \sqrt{2S_A}a_j, S_n^z = -S_B + b_n^{\dagger}b_n, S_n^+ = \sqrt{2S_B}b_n^{\dagger}, S_n^- = \sqrt{2S_B}b_n$. Here a, a^{\dagger}, b and b^{\dagger} are Bose operators. For the impurity spin, $S_0^z = S_i - a_0^{\dagger}a_0, S_0^- = \sqrt{2S_i}a_0^{\dagger}$ and $S_0^+ = \sqrt{2S_i}a_0$ when p < 1 and $S_0^z = -S_i + a_0^{\dagger}a_0, S_0^+ = \sqrt{2S_i}a_0^{\dagger}$ and $S_0^- = \sqrt{2S_i}a_0$ when p > 1. Let us consider a one-magnon and one-impurity state which is defined by

$$\Psi = \left(\sum_{j} A_{j} a_{j}^{\dagger} + \sum_{n} B_{n} b_{n}\right) c_{0}^{\dagger} \mid 0 >$$

Here | 0 > denotes the spin wave ground state for the ferrimagnet in which neither magnons nor impurities exist.

Using Ψ and H in the Schrödinger equation, we obtain the secular equations for A_j and B_n . For the case p < 1, the secular equations are written as

$$(-U + 2JS_B z)A_j - 2JS_r \sum_{\rho} B_{j+\rho} + \delta(j,0)(-2JzS_B pA_j + 2JS_r(p+p'q')\sum_{\rho} B_{j+\rho}) = 0$$

$$(-U - 2JS_A z)B_n + 2JS_r \sum_{\rho} A_{n+\rho} + \delta(n,j+\rho_1)(2JS_A(p+p'q)B_n - 2JS_r(p+p'q')A_{n+\rho_1}) = 0$$

where U represents the energy measured from the ground state energy. z is the number of nearest neighbors and δ is the usual Kronecker's delta function. $S_r = \sqrt{S_A S_B}$ and ρ , ρ_1 denote the vectors directed between nearest neighbors. p' = 1 - p and $q' = 1 - \sqrt{1 - q}$. Introducing the Green functions, we can solve the above secular equations. The energy bands, $u^{\pm} = U^{\pm}/2JzS$, is obtained as

$$u^{\pm} = \frac{1}{2} \left[\sqrt{\beta} - \frac{1}{\sqrt{\beta}} \pm \sqrt{(\sqrt{\beta} - \frac{1}{\sqrt{\beta}})^2 + 4(1 - \gamma_k^2)} \right]$$

where $\beta = S_B/S_A$ and $\gamma_k = (1/z) \sum_{\rho} e^{ik \cdot \rho}$. In the following, we show the impurity-magnon bound states in the band gap for the case $\beta = 2$.



Fig.1 Impurity-magnon bound states in the band gap for the case of $\beta = 2$.

In this work, we have investigated localized spin wave states outside the continuous bands for various parameters. If the experiment for the localized spin waves at the impurity were performed, we may obtain useful information about the physical states of the impurity.

Acknowledgments

The author wish to thank Prof. E. Ilisca for useful discussion.

References

- P.E.Wigen and H.Puszkarski, Solid State Commun. 18 (1976) 363. I.Harada, O.Nagai and T.Nagainiya, Phys. Rev. B16 (1977) 4882.
- [2] For example, V.Cannella and J.A.Mydosh, Phys. Rev. B6 (1972) 4220.
- [3] T.Wolfram and J.Callaway, Phys. Rev. 130 (1963) 2207.

C1-640