EXAFS Study of Zn and ZnEDTA Sorption at the Goethite (α -FeOOH)/Water Interface

M. Schlegel, A. Manceau, L. Charlet

► To cite this version:

M. Schlegel, A. Manceau, L. Charlet. EXAFS Study of Zn and ZnEDTA Sorption at the Goethite (α -FeOOH)/Water Interface. Journal de Physique IV Proceedings, 1997, 7 (C2), pp.C2-823-C2-824. 10.1051/jp4:1997248 . jpa-00255327

HAL Id: jpa-00255327

<https://hal.science/jpa-00255327>

Submitted on 1 Jan 1997

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXAFS Study of Zn and ZnEDTA Sorption at the Goethite (α -FeOOH)/Water Interface

M.L. Schlegel, A. Manceau and L. Charlet

Environmental Geochemistry Group, LGIT-IRIGM, UJF, BP. 53, 38041 Grenoble cedex 9, France

Abstract: EXAFS spectroscopy has been used to compare the sorption mechanism of Zn and ZnEDTA on goethite (α -FeOOH). In the absence of EDTA, Zn is found to form an inner-sphere surface complex. Based on Zn-O and Zn-Fe distances, Zn is inferred to be located in a distorted oxygen octahedron which shares edges and/or corners with surface Fe octahedra. EXAFS spectra of ZnEDTA in solution and sorbed on goethite are similar, indicating that the ZnEDTA complex does not dissociate following its sorption. The absence of an Fe shell within the first 4 Å precludes the existence of Zn-O-Fe bonds and allows us to reject a ligand exchange sorption mechanism involving the formation of Zn-O-Fe linkages.

1. INTRODUCTION

The fate of trace metals in natural aquatic systems is influenced by adsorption processes at the oxide-water interface. The nature of these reactions and the amount of sorbed species can be modified by the presence of natural or anthropogenic organic ligands. For instance, complexation of Zn by the synthetic chelating agent EDTA ($\text{C}_{10}\text{H}_{12}\text{O}_8\text{N}_2$) has been shown to enhance its mobility in aquifers [1]. The sorption of Me-EDTA complexes on different oxide surfaces has been studied by means of batch chemical experiments ([2,3] and ref. therein) : MeEDTA exhibit a ligand-like, ionic strength-dependent sorption behavior. Contrasting molecular mechanisms (i.e. outer-sphere complexation [2] versus inner-sphere ligand exchange [3]), have been proposed to account for similar chemical results. To gain further insight on the influence of EDTA in the sorption mechanism of Zn, Zn-EXAFS experiments were performed on Zn- and ZnEDTA-sorbed goethite.

2. EXPERIMENTAL

All chemicals were ACS reagent grade. Adsorption experiments were conducted at $25.0 \pm 0.1^\circ\text{C}$ in a thermostated glass vessel, at a constant background electrolyte concentration (NaNO_3 0.1 mol.l⁻¹). Humidified argon was bubbled through a 15 g.l⁻¹ stirred suspension of goethite ($49 \pm 5 \text{ m}^2.\text{g}^{-1}$) to maintain a CO_2 -free atmosphere. The proton site concentration in the suspension was 2.10^{-3} mol.l⁻¹. Zn sorption at 10% proton site coverage was realized by adding an acid aliquot of $\text{Zn}(\text{NO}_3)_2$ ($\text{TotZn} = 2.10^{-4}$ M) to the suspension at pH 4, and then titrating the suspension with NaOH 0.1 mol.l⁻¹ up to pH 7. The suspension was equilibrated at pH 7 for 24 h. ZnEDTA sorption on goethite at 30% site coverage was realized by titrating a ZnEDTA rich suspension ($\text{TotZnEDTA} = 1.8.10^{-3}$ M) from pH 9 to pH 4.5, and then equilibrating the suspension at pH 4.5 for 30 min. Both samples were filtered (Sartorius 0.1 μm), washed with pH-adjusted NaNO_3 0.1 mol.l⁻¹, air-dried and loaded as powders in sample holders. Washing may only remove the most weakly sorbed ions.

Zn K-EXAFS spectra were recorded in fluorescence mode at the SRS synchrotron facility in Daresbury UK, on the EXAFS 9.2 station. Interatomic distances were determined on Fourier-filtered spectra by using phases and amplitudes functions either extracted from model compounds (ZnO for the Zn-O atomic pair, γ -FeOOH (Fe_L)) or calculated [4].

3. RESULTS AND DISCUSSION

3.1 Zn sorbed goethite

$\chi(k)$ spectra and RDFs of Zn sorbed goethite (#a), ZnEDTA in solution (#b) and ZnEDTA sorbed goethite (#c) are displayed in Fig. 1 and 2. The RDF for #a presents one main peak at 1.7 Å, which corresponds to the contribution of the Zn-O coordination shell, and a broad double-peak extending from 2.5-3.0 Å. Two Zn-O distances were determined from the fitting of the Fourier-filtered Zn-O contribution: $d_{\text{Zn-O}1} = 2.01$ Å ($N=3.5$) and $d_{\text{Zn-O}2} = 2.19$ Å ($N=3.2$). $\langle d_{\text{Zn-O}} \rangle$ is equal to 2.10 Å, which is typical of six-coordinated Zn (2.12 Å, [5]). $d_{\text{Zn-O}1}$ can hardly be attributed to four-fold Zn, as $\langle d_{\text{Zn-O}} \rangle = 1.97$ Å, but is found in distorted Zn octahedra [6]. Thus, Zn is inferred to be located in a distorted octahedron.

The broad peak at 2.5-3.0 Å on the RDF is absent for solvated Zn, and points to the presence of Zn-metal interactions. Since the suspension contained only Zn and Fe, only these atoms can contribute to these RDF peaks. A comparison of the EXAFS spectrum for #a with the spectrum of amorphous $\text{Zn}(\text{OH})_2$ (Fig. 1) indicates that the local structures of $\text{Zn}(\text{OH})_2$ and Zn-sorbed goethite are different. This observation allows us to reject the precipitation of Zn hydroxide. Therefore one can

assume that these metal shell peaks come from contributions of surface Fe atoms, i.e. that Zn forms an inner-sphere surface complex. Zn-Fe contributions were modeled by two Fe shells at 3.00 Å ($N=0.9$) and 3.20 Å ($N=1.2$). These distances are in good agreement with those determined for Cd-sorbed goethite ($d_{\text{Cd-Fe}} = 3.26$ and 3.48 Å, [7]), taking into account the difference of ionic radii between 6-fold Zn and Cd ($r_{\text{Zn}} = 0.74$ Å and $r_{\text{Cd}} = 0.95$ Å; [5]). As for cadmium, the short Zn-Fe distance is attributed to an edge linkage between Zn and surface Fe octahedra, and the longer distance to a corner linkage.

In conclusion, Zn octahedra are found to share edges and corners with surface Fe octahedra. By analogy with Cd [7], these two types of linkage are believed to correspond to different surface sites.

3.2 ZnEDTA sorbed goethite

The strong organometallic complex formed by Zn and EDTA in solution may adopt several conformations similar to those determined in crystals. For example, in $\text{Mg}[(\text{H}_2\text{O})_4\text{ZnEDTA}]\cdot 2\text{H}_2\text{O}$ [8], Zn is coordinated to 4 acetyl and 2 amide groups of the EDTA molecule. In solution, this hexadentate conformation is in equilibrium with a pentadentate one, the coordination sphere of zinc being completed by a water oxygen. In both conformations, Zn is coordinated to N and O atoms, and has 10 next-nearest C neighbors. The RDF of ZnEDTA in solution (#b) displays 3 major peaks centered at 1.7 Å, 2.5 Å and 3.5 Å. The first two peaks correspond to (N,O) and C contributions, respectively. They were modeled by two (N, O) subshells at 2.07 Å ($N=3.4$) and 2.20 Å ($N=2.1$) ($\langle d_{\text{Zn-(O, N)}} \rangle = 2.12$ Å), and two C subshells at 2.82 Å ($N=3.8$) and 2.95 Å ($N=7.1$) ($\langle d_{\text{Zn-C}} \rangle = 2.90$ Å). These mean distances are close to those found in solid structures ($\langle d_{\text{Zn-(O, N)}} \rangle = 2.11$ Å and $\langle d_{\text{Zn-C}} \rangle = 2.89$ Å; [8]). Due to the absence of heavy metal near Zn, the third RDF peak cannot be assigned to a cationic shell. *Ab initio* calculations performed on ZnEDTA clusters derived from the $\text{Mg}[(\text{H}_2\text{O})_4\text{ZnEDTA}]\cdot 2\text{H}_2\text{O}$ crystalline structure [8], using FEFF6 [9] showed that this third RDF peak predominantly corresponds to multiscattering contributions.

Fig. 2 shows that ZnEDTA in solution (#b) and sorbed on goethite (#c) have very similar EXAFS spectra and RDFs, and that they are markedly different from Zn-sorbed goethite. This result indicates that the structure of ZnEDTA was preserved during sorption. A quantitative spectral analysis of #c yielded interatomic distances which were, to the precision of the method, identical to those obtained for #b. The third peak for sample #c is shifted toward shorter Å values. This may result from slight structural distortions. No Fe-Zn pairing could be identified below 4 Å. Solely external oxygens of the EDTA molecule, that are not bound to the Zn atom, may possibly interact with surface Fe octahedra, but in this case, the Fe atoms would be at much higher distance than 4 Å from Zn and would not be detected by EXAFS spectroscopy.

In conclusion, EXAFS spectroscopy indicates that the conformation of the ZnEDTA complex is preserved during sorption on goethite and precludes a ligand exchange sorption mechanism involving the formation of Zn-O-Fe linkages. This conclusion may be generalized to several other MeEDTA (Me = Ca, Zn, Cu, Co(II), Pb) complexes, inasmuch as these complexes show similar macroscopic sorption behaviors on goethite [3].

Fig. 1: $k^2\chi(k)$ spectra for $\text{Zn}(\text{OH})_2$, Zn sorbed goethite (#a), ZnEDTA in solution (#b) and ZnEDTA sorbed goethite (#c)

Fig. 2: RDF for Zn sorbed goethite (#a), ZnEDTA in solution (#b) and ZnEDTA sorbed goethite (#c)

Acknowledgement

We thank the Daresbury staff for providing the X-Ray beam and for their technical assistance.

References

- [1] Kent D.B., Davis J.A., Anderson L.D. and Rea B.A., "Ligand enhanced transport of strongly adsorbing ions in the groundwater environment", Water-Rock interactions (Karaka Y.K and Maes A.S., Eds, Balkema, Rotterdam, 1992)
- [2] Bowers A.R. and Huang C.P., *J. Coll. Interf. Sci.*, **110** (1986) 575-590.
- [3] Nowack B. and Sigg L., *J. Coll. Interf. Sci.*, **177** (1996) 106-121.
- [4] McKale A.G., Veal B.V., Paulikas A.P., Chan S.K. and Knapp J.S., *J. Am. Chem. Soc.*, **110** (1988) 3763-3768.
- [5] Shannon R.D., *Acta Cryst.*, **A32** (1975) 751-767.
- [6] Wadsley A.D., *Acta Cryst.*, **8** (1955) 165-172
- [7] Spadini L., Manceau A., Schindler P.W., Charlet L., *J. Coll. Interf. Sci.*, **167** (1994) 73-86.
- [8] Solans X., Font-Alba M., Oliva J. and Herrera J., *Acta Cryst.*, **C39** (1983) 1435-1438.
- [9] Rehr J.J., Albers R.C. and Zabinsky S.L., *Phys. Rev. Lett.*, **69** (1992) 3397-3400.